Canadian Science Publishing | Canadian Journal of Forest Research

# Early influences of tamarack (Larix laricina) on black spruce (Picea mariana) and its immediate environment in plantations

| | |
|---|---|
| Journal: | *Canadian Journal of Forest Research* |
| Manuscript ID | cjfr-2023-0220.R2 |
| Manuscript Type: | Article |
| Date Submitted by the Author: | 18-Jan-2024 |
| Complete List of Authors: | Roy Proulx, Samuel; Centre d'Étude de la Forêt, Sciences biologiques; Centre de foresterie des Laurentides, Centre Canadien sur la fibre de bois<br>Leduc, Alain; Centre d'Étude de la Forêt, Sciences biologiques<br>Thiffault, Nelson; Centre de foresterie des Laurentides, Canadian Wood Fiber Centre; Centre d'Étude de la Forêt, Sciences biologiques<br>Laganière, Jérôme; Centre d'Étude de la Forêt; Centre de foresterie des Laurentides, Service Canadien des Forêts |
| Is the manuscript for consideration in a Special Issue or Collection?: | Not applicable (regular submission) |
| Keyword: | mixed plantation, young plantation, neighbouring tree effect, boreal forest, non-crop competition |
| | |

SCHOLARONE™ Manuscripts

© The Author(s) or their Institution(s)

# Early influences of tamarack (*Larix laricina*) on black spruce (*Picea mariana*) and its immediate environment in plantations

*Samuel Roy Proulx[1, 2], Alain Leduc[1], Nelson Thiffault [1, 2] and Jérôme Laganière[1, 3]

[1]Centre d'étude de la Forêt, Université du Québec à Montréal, C.P. 8888, Succ. Centre-Ville, Montréal, QC H3C 3P8, Canada;

[2]Centre Canadien sur la Fibre de bois, Service Canadien des Forêts, Ressources naturelles Canada, 1055 rue du P.E.P.S, C.P. 10380, Succ. Sainte-Foy, Québec, QC G1V 4C7, Canada

[3]Centre de Foresterie des Laurentides, Service Canadien des Forêts, Ressources naturelles Canada, 1055 rue du P.E.P.S, C.P. 10380, Succ. Sainte-Foy, Québec, QC G1V 4C7, Canada

* Corresponding author: roy_proulx.samuel@courrier.uqam.ca

Draft

## Abstract

Mixed plantations are garnering increased attention due to their potential to provide a broader array of benefits compared to monocultures. Although numerous studies have indicated promising complementarity between black spruce (*Picea mariana*) and tamarack (*Larix laricina*), few have delved into individual tree growth interactions to thoroughly assess early growth complementarity. We sampled 119 planted black spruce and their immediate environment to quantify and qualify any differences between two conditions in young plantations: mixed tamarack ("mixtures") and black spruce monocultures ("monocultures") within young plantations. We investigated the effect of neighbouring under four perspectives: tree competition, microenvironment, foliar nutrients, and

© The Author(s) or their Institution(s)

soil nutrients. Our results showed increased values for black spruce foliar nitrogen total concentration, soil pH and canopy closure in mixtures compared to monocultures. Furthermore, black spruce stem volume was increased by 38.1% in mixture compared to monoculture. Black spruce stem volume was negatively affected (86% decrease) by the combined effect of shrubs and non-crop trees under high competition pressure, despite the plantation being mechanically released in 2017. Collectively, our results suggest that black spruce growing in mixtures holds a greater growth potential than black spruce in monocultures.

Keywords: mixed plantation, young plantation, neighbouring tree effect, boreal forest, non-crop competition

## Introduction

Draft

Sustainable forest management encompasses diverse benefits for social, economic, and ecological purposes, achieved in part by using reforestation with multiple tree species and initiating mixed stands (Bolte et al. 2009, Felton et al. 2016). Eventually, mixed stands can exhibit an increased resilience and stabilize yields in the face of extreme climate, provide additional protection against insect defoliation and increase biodiversity (Jactel and Brockerhoff 2007, Bielak et al. 2014, Felton et al. 2016). Notably, trees in mixed stands often demonstrate better resource use (soil and light) than in monoculture (Liang et al. 2016, Pretzsch and Schütze 2016), potentially leading to over-yielding of mixed stands (Zhang et al. 2012, Lu et al. 2016). However, managing mixed stands is more complex than monocultures, requiring more planning for harvesting and adapted machinery (Yilmaz and Akay 2008). Interactions in multi-species plantations can be complex, and when the mixing ratio, site conditions, and/or tree spacing are sub-optimal, under-yielding can occur (Toïgo et al. 2015, Drossler et al. 2018). Therefore, identifying compatible tree species should be a primary

© The Author(s) or their Institution(s)

focus of forest management, followed by studying growth relationships to determine adequate species combinations and under which circumstances.

Studying planted trees during early years provides insights on individual growth trajectories and the future productivity of plantations (Bianchi et al. 2021). Positive growth interactions can emerge in the early stages (10-15 years) of mixed plantations, enhancing the growth of one or both species compared to monoculture (Dijkstra et al. 2009; Nord-Larsen and Meilby 2016). Mixing shade tolerant with shade intolerant species often lead to good growth complementarity and potentially to over-yielding (Cordonnier et al. 2018). However, early positive effects don't guarantee over-yielding over a whole rotation, as this situation can reverse during subsequent stages of stand development (Drossler et al. 2018). Moreover, interactions between crop trees, shrubs, and non-crop trees can have lasting impacts on individual tree growth and alter stand structure, particularly if the affected crop tree is shade intolerant (Bell and Pitt 2005).

The Canadian boreal forest is characterized by a cold climate and relatively low productivity (Saucier et al. 2015), but also serves as the primary source of wood fibre for the forest industry (Beaudoin et al. 2014). Multiple actions are needed to ensure plantation productivity in the boreal forest, such as mechanical release of competing vegetation (Wiensczyk et al. 2011), and mixed plantations could help ensure productivity, through better space and resources acquisitions (Forrester 2014, Pretzsch 2014). The selection of compatible species should be based on different shade tolerance, resource requirements and root system depth, as well as compatible crown architecture (Pretzsch 2014, Coll et al. 2018). Furthermore, growth compatibility between species could be modulated by site conditions such as water and/or nutrient availability (Laganière et al. 2015). Diversifying the species chosen for reforestation in the boreal forest could possibly enhance plantation productivity (Thiffault et al. 2010). Currently, spruces dominate reforestation in Canada,

Draft

© The Author(s) or their Institution(s)

constituting 44.3% of all planted species (CCFM 2020). In the boreal forest, black spruce (*Picea mariana* [Mill.] B.S.P.) is the most used spruce species in plantations. In contrast, tamarack (*Larix laricina* [Du Roi] K. Koch) is a marginal species in reforestation programs, accounting for only 1.5% of planted trees nationwide (CCFM 2020). Yet numerous studies highlight physiological complementarity between black spruce and tamarack (Islam and Macdonald 2004, Boyden et al. 2009). Tamarack is a fast-growing, shade-intolerant and deciduous conifer, while black spruce is a slow-growing, shade-tolerant evergreen conifer (Strong and Roi 1983). The annual litter input from tamarack is rich in nitrogen and phosphorus, which could benefit soil fertility (Moore et al. 2010). This annual litter input can change soil macronutrient content, microbial communities, and pH (Dijkstra et al. 2009, Prescott and Grayston 2013), potentially enhancing neighbouring tree growth and overall stand yield by increasing macronutrient availability (Forrester et al. 2006, Laganière et al. 2017). Only a few studies have reported about the early growth relations between tamarack and black spruce (Boyden et al. 2009). Assessing the microenvironment is valuable to untangle the effects of shrubs, soil composition and neighbouring species on crop tree growth (Baraloto et al. 2005). Furthermore, examining tree leaf nutritional status is useful to better predict crop tree long-term growth trajectory (Alva et al. 2006).

Our objective was to compare the neighbouring effects of tamarack and black spruce on soil characteristics (macronutrient concentration and pH), competing vegetation, and the stem volume of black spruce, eight years after planting in the Canadian boreal forest. Our specific aims were to (1) detect differences in microenvironment associated with the neighbouring of tamarack or black spruce, (2) assess the effects of neighbouring crop tree species and competing vegetation on black spruce dimensions after 8 years, (3) identify potential changes in early canopy closure driven by tamarack and black spruce neighbours, and (4) investigate the influence of neighbouring species

Draft

© The Author(s) or their Institution(s)

effect on soil and black spruce foliar nutrient concentrations. We hypothesized that the presence of tamarack as a neighbouring species would significantly alter soil macronutrient concentrations and pH levels, compared to areas dominated exclusively by black spruce. We anticipated that the stem volume of black spruce would be differently influenced by the proximity of tamarack versus black spruce, owing to variations in their microenvironmental effects. We predicted that early canopy closure will be distinctly affected by tamarack and black spruce, potentially leading to varied competitive dynamics among plant species. Finally, we expected that the neighbouring species, be it tamarack or black spruce, would have a measurable yet distinct impact on both soil and foliar nutrient concentrations in black spruce.

# Materials and methods

## Site description and experimental design

Draft

The experimental site was in the Lake Duparquet Research and Teaching Forest (LDRTF) in the northwestern boreal forest of Québec, Canada (48º29'7" N – 48º29'5" N; 79º25'5" W – 79º26'0" W), within the balsam fir (*Abies balsamea* (L.) Mill.) – white birch (*Betula papyrifera* Marsh) bioclimatic domain (Saucier et al. 2009). This region is known for extensive clay deposits left by the proglacial Lake Barlow-Ojibway (Vincent and Hardy 1977). The climate is cold and continental with average temperatures of 0 °C and mean annual precipitation of 900 mm (Environment Canada 2019). The tree species dominating early succession are trembling aspen (*Populus tremuloides* Michx.), white birch, and jack pine (*Pinus banksiana* Lamb.). Late successional stands are typically dominated by balsam fir, white spruce (*Picea glauca* [Moench] Voss), black spruce, and eastern white cedar (*Thuja occidentalis* L.) (Bergeron 2000). The understory composition is generally characterized by ericaceous shrubs (Thiffault et al. 2015), along with species such as *Gaultheria*

© The Author(s) or their Institution(s)

111 *hispidula*, *Coptis groenlandica*, *Cornus canadensis*, and *Dryopteris spinulosa* depending on soil
112 fertility and overstory dominance (Légaré et al. 2001). Soils in this area are generally classified as
113 Luvisols that originated from silty clay deposits from proglacial Lakes Barlow and Ojibway, with
114 a mor humus layer ranging from 0 to 50 cm in thickness. In natural stands of this region, tamarack
115 and black spruce usually co-occur on peatlands or on thick organic soils with complex drainage
116 (Perala 1971, Viereck et al. 1990).

117 The site was clearcut during the winter of 2011–2012. In spring 2013, the site was mechanically
118 prepared using a double-pass forest harrow. The experimental design included two blocks, each
119 measuring 5000 m² (100 m × 50 m). The blocks were divided into two different treatments: black
120 spruce monospecific and 50/50 mixture covering 2500 m² (50 m × 50 m) each. All treatments were
121 established with a density of 2500 stems/ha, with seedlings planted at an approximate spacing of 2
122 m x 2 m. The mixed plantation treatment involved alternating rows of planted tamarack and black
123 spruce (Fig. 1). This design allowed us to investigate the effects of neighbouring tree species and
124 competing vegetation on black spruce growth.

125 In 2017, the plantation underwent mechanical vegetation control using motor-manual brushsaws
126 to remove competing vegetation, leaving only black spruce and tamarack. At the time of sampling,
127 the competing vegetation had resprouted and was primarily composed of willow (*Salix* spp.), pin
128 cherry (*Prunus pennsylvanica* L.f.), trembling aspen and red raspberry (*Rubus idaeus* L.).

129 ## Tree measurements

130 In June and July 2021 (8 years after planting), we selected 121 planted black spruce as target trees,
131 69 neighboured only by other black spruce saplings and 52 neighboured by tamarack and black
132 spruce sapling (Fig. 1). The target trees were randomly distributed across all experimental units,
133 irrespective of the treatment applied (Fig. 1). Our selection criteria focused on height (equal or

Draft

© The Author(s) or their Institution(s)

134 superior to other crop tree height), absence of scars or injuries, and overall healthy foliage
135 excluding chlorotic trees. This approach aimed to target the trees most likely to become dominant
136 or codominant in the future (Jobidon 2000). To assess the neighbouring tree effects, we established
137 40 m² circular plots (radius = 3.57 m) centred on each target black spruce tree. Within these plots,
138 we recorded and measured all planted trees present (Fig 1.). In the mixed planting environment,
139 this allowed us to investigate the effect of neighbouring trees at a ratio of 2 black spruce trees for
140 6 tamarack trees, as the plot is centred on target black spruce (Fig. 1, Table 1).

Draft

© The Author(s) or their Institution(s)




141

142 Figure 1: Schematic representation of the plantation design in pure and mixed plots, with the list of variables assessed.

© The Author(s) or their Institution(s)

Target trees and neighbouring crop tree competitors were measured for total height, ground-level diameter (GLD), and diameter at breast height (DBH, measured at 130 cm above the ground level) (Table 1). All target trees had a height greater than 1.3 m; we therefore calculated stem volume index during the 8th growing year as:

$$Vol_8 = \frac{13\,\pi}{1200} \times (GLD^2 + GLD \times DBH + DBH^2) + \frac{\pi(DBH)^2(H-130)}{12000} \quad \text{Eq. 1}$$

where $Vol_8$ is the stem volume index (dm³) based on the volume of a cylinder from 0 m to 1.3 m in height and the volume of a cone from 1.3 m up, GLD, DBH and total tree height (H) are all measured in centimetres.

Table 1 : Plot characteristics as a function of competing neighbours (black spruce or tamarack). DBH = diameter at breast height (130 cm), GLD = ground-level diameter, BA = basal area, LAI = leaf area index, ATT = above target tree, BCNT = between closest neighbouring tree. Values are presented as mean ± standard deviation.

| | **Black spruce neighbour** | **Tamarack neighbour** |
|---|---|---|
| **Soil characteristics** | | |
| Organic layer depth (cm) | 8.7 ± 5.8 | 9.2 ± 4.9 |
| C/N mineral | 15.0 ± 2.4 | 14.7 ± 2.3 |
| C/N organic | 27.6 ± 7.4 | 25.2 ± 7.1 |
| pH mineral | 5.00 ± 0.42 | 5.17 ± 0.62 |
| pH organic | 5.52 ± 0.42 | 5.81 ± 0.59 |
| **Plot level characteristics** | | |
| Tamarack BA (m² / ha) | 0.02 ± 0.05 | 0.80±0.48 |
| Black spruce BA (m² / ha) | 0.17 ± 0.19 | 0.003 ± 0.028 |
| LAI ATT | 0.15 ± 0.34 | 0.34 ± 0.32 |
| LAI BCNT | 0.79 ± 0.48 | 1.07 ± 0.45 |
| Number of crop trees | 7.5 ± 1.8 | 6.3 ± 1.8 |
| **Black spruce dimensions and foliar nutrition** | | |
| DBH (cm) | 2.53 ± 0.64 | 2.47 ± 0.67 |

Draft

© The Author(s) or their Institution(s)

| | | |
|---|---|---|
| GLD (cm) | 5.17 ± 0.88 | 4.91 ± 1.06 |
| Height (cm) | 239.1 ± 30.8 | 241.4 ± 36.3 |
| Stem volume index ($dm^3$) | 35.7 ± 21.6 | 33.8 ± 23.6 |
| C/N | 60.7 ± 5.9 | 56.7 ± 5.6 |
| Total N (%) | 0.86 ± 0.10 | 0.91 ± 0.09 |
| Total P (%) | 2.28 ± 1.72 | 2.68 ± 1.81 |
| Total K (%) | 8.06 ± 6.51 | 9.14 ± 6.52 |

## Plot characteristics measurements

To determine the leaf area index (LAI) of the canopy in each plot, we captured images with a fisheye camera (CI-110 Plant canopy Imager, CID Bio-Science, Inc. Camas, WA, USA) at the beginning of July 2021, during cloudy days. The first image was taken perpendicular to the ground at mid-height of the target black spruce trees, achieved by bending down the tree. This measurement is referred to as 'LAI above target tree'. A second image was taken 45 cm above the ground, at the midpoint between the target tree and the neighbouring nearest competitor crop tree. This measurement is referred to as 'LAI between closest neighbouring tree'. The images were analyzed with CID Bio-Science software (Camas, WA, USA) to compute the LAI. Microtopography at the base of each target tree was classified into three categories: flat, mound, and pit. This visual estimation considers the surrounding microtopography to estimate if the water would accumulate or not at the base of each tree during the snow melt period or after a prolonged rain event.

## Soil and leaf sampling and processing

From June 21-25, 2021, we collected soil samples from both organic and mineral layers using a trowel to analyze their C/N ratio and measure their pH. The sampling was conducted 1 m south

© The Author(s) or their Institution(s)

171 from each target tree trunk, at a maximum depth of 20 cm as the root system of black spruce is
172 shallow with roots from mature trees rarely exceeding 30 cm depth (Strong and Roi 1983). If the
173 organic horizon exceeded a thickness of 20 cm, we did not collect mineral soil samples since tree
174 roots are not reaching the mineral soil. The depth of the organic layer was recorded at the sampling
175 location. All samples were frozen within 8 hours. In the lab, soil samples were cleaned of any
176 impurity (stone, root, etc.), air-dried for 2-week, oven-dried at 50 °C for 1 week and sieved on a 2
177 mm mesh screen. Total C and total N (%) were determined by dry combustion on a LECO CNS
178 928 (LECO Corporation, St-Joseph, MI, USA). Soil pH was determined using a 1:2 soil:water ratio
179 for mineral soil samples and a 1:10 ratio for organic samples (Fisher Scientific Accumet 50; Denver
180 Instrument, Bohemia, New York).

181 Between June 20-26, 2021, we collected foliar samples from each target black spruce tree. The
182 samples consisted of 1-year-old twigs (from 2020) located in the upper third of the trees and facing
183 South. After collection, the samples were kept in a cooler and frozen within 8 hours. The foliar
184 samples were dried using the same method as the soil samples. Needles were separated from the
185 twigs and then ground using a Retsch cutting mill SM 200 (Verder scientific, Vleuten, Netherlands)
186 operating at 10,000 rpm. Total N concentration was determined by dry combustion as above; P and
187 K concentrations were measured following the method outlined by Yash (1998) using a Perkin-
188 Elmer Optima 7300DV (PerkinElmer, Inc. Waltham, MA, USA).

## 189 Competition indices

190 We calculated the Hegyi competition index (Hegyi 1974) to estimate the effects of neighbouring
191 crop tree competitors on target trees as follows:

192 $$Hegyi\ competition\ index\ = \sum_{j=1\ j\neq i}^{n} \frac{d_j}{d_i L_{ij}} \quad \text{Eq.2}$$

© The Author(s) or their Institution(s)

where $d_j$ and $d_i$ are respectively the diameters of the neighbouring competitors and of target trees, and $L_{ij}$ is the distance between neighbouring competitors and target trees. We considered any planted tree with a height equal to or greater than the target trees and located within a 3.57 m radius as significant neighbouring competitors (White et al. 2014).

We identified and measured the height of all competing woody shrubs and non-crop trees reaching at least two thirds of the height of the target trees and located within a 1.2-meter radius from the target trees (White et al. 2014). We used the sum of competing vegetation angles (Prévosto 2005) as a competition index to quantify the competition of shrubs and non-crop trees on target trees (Eq. 3). The angles of each shrub and non-crop tree were taken from the base of the target tree to the top of each competing shrubs. This competition index is distance dependent, influenced by the height of competing vegetation, independent from the target tree height, and calculated as:

$$Sum\ of\ competing\ vegetation\ angles = \sum_{j=1\, j\neq i}^{n} \Phi_i \quad \text{Eq. 3}$$

where $\Phi_i$ is the angle from the base of the target tree to the top of each neighbour.

## Statistical analyses

### Supervised forward model selection

We used a supervised forward model selection approach with the linear model function *lm* in the 'stats' package (R Core team Development Team 2022) [Objective 1]. This analysis allowed us to explore the relationships between the stem volume of black spruce at age 8 and various environmental variables. The variables included in our analysis were the number of crop trees per plot, depth of the organic layer, microtopography, Hegyi competition index and the sum of competing vegetation angles around the target trees. We used the linear model:

$$Vol^{8th}_{k} = \beta_0 + \beta_1(x1) + \ldots + \beta k(xk) \quad \text{Eq. 4}$$

© The Author(s) or their Institution(s)

where β is a vector of k fixed effects.

We included a global model and a null model as part of our candidate models to test for the lack of effect of all variables and the null hypothesis, respectively. We checked the homoscedasticity and normality of residuals to ensure that the linear model assumptions were met. We initially used generalized linear model to test the spatial autocorrelation with treatments nested within blocks as random factors, but since these factors did not explain any variance of black spruce stem volume, we excluded them from further analysis and used linear models instead (Bates et al. 2015). With the list of environmental variables (i.e. number of crop trees, depth of the organic layer, microtopography, Hegyi competition index and the sum of competing vegetation angles around the target trees), we conducted the supervised forward model selection based on the Akaike Information Criterion corrected for small samples (*AICc*) (Akaike 1998) and *ΔAICc* (the difference in *AICc* score from the top model) to determine the inclusion or exclusion of models. The *AICc* score for each model was calculated using the *AICcmodavg* function in the 'AICcmodavg' package (Mazerolle 2023). The *AICc* threshold value and the model selection process at each step followed the approach of Portier et al. (2016). In the first step, we built univariate models to test each variable individually. We retained each model with a $\Delta AICc < 6$ of the first model (Symonds and Moussali 2011). In the second step, we added a second variable to each selected model and performed *AICc* model selection. If the model with two variables showed a lower *AICc* score of 2 or less compared to the related univariate model, we kept the model with two variables and discarded the univariate model. We repeated the process of adding variables until the model could no longer be improved by additional variables. At the end of this procedure, each model with a *ΔAICc* distance of 2 or more from the best model was considered less parsimonious and discarded (Anderson and Burnham 2002). Furthermore, each added variable in the model had to be uncorrelated with each

Draft

© The Author(s) or their Institution(s)

other (Pearson coefficient < 0.7) and have a significant effect ($p < 0.05$) on black spruce stem volume to avoid overfitting the model. Finally, we compared the *AICc* of the selected model to a null model to ensure the improvement of the model. The model parameter estimates and unconditional 95% confidence intervals, excluding 0, were considered influential of the 8th-year stem volume of black spruce. The QR decomposition matrix was used to estimate the parameters for each model and the adjusted $R^2$ was calculated with the 'performance' package (Lüdecke et al. 2021).

Neighbouring species impact on black spruce

To evaluate the neighbouring species effect of tamarack and black spruce [Objective 2], we included the neighbouring species as a factor (black spruce or tamarack) into the best model selected from the supervised forward model selection for predicting the $8^{th}$-year stem volume of black spruce. We then compared the best model with and without the neighbouring species effect to examine its impact on black spruce stem volume using $AIC_c$. We verified the normality of residuals and checked for homogeneity of variances for each model using Levene's test (Gastwirth et al. 2009).

All statistical analyses were performed within the R programming environment (R Core team Development Team 2022). We conducted analyses of variance (ANOVA) to examine the effects of neighbouring tree species on both measured LAI, foliar nutrient concentrations (N, P and K), soil nutrients (C/N ratio) and soil pH (organic and mineral) [Objectives 1, 3 and 4]. We assessed the normality of residuals and checked for homogeneity of variances using Levene's test (Gastwirth et al. 2009). In order to meet the assumption of normality for LAI above target trees, we applied a log transformation to the response variable. Analyses were conducted on transformed data, but we

Draft

© The Author(s) or their Institution(s)

present values (and differences) on their original scale for the sake of clarity. We used $\alpha = 0.05$ as a threshold of statistical significance.

# Results

## Effects of environmental variables on black spruce volume

The supervised forward model selection, using only environmental variables, resulted in the selection of a single model comprising microtopography, the sum of competing vegetation angles, and the Hegyi competition index by the Akaike criterion (Table 2). The second-ranked model was composed of competing vegetation angles, and the Hegyi competition index (Table 2). With 58% of *AICc* weight and explaining 22.9% of the variance, the first-ranked model was deemed the best fit for the data as no other model was best fitting for black spruce stem volume (Table 2). None of the other candidate models had a $\Delta AIC_c$ within 2; they were not kept for further analysis.

Draft

Table 2 : Most parsimonious models explaining black spruce 8th-year stem volume, including the number of parameters (K), delta $AIC_c$ compared to the highest-ranked model ($\Delta AIC_c$), model weight ($AIC_cWt$), and model predictive power (Adjusted $R^2$). The second half of the table shows parameters influencing black spruce 8th years stem volume and 95% confidence intervals excluding 0. For microtopography, mound was used as the reference level. Model in bold indicate retained model ($\Delta AIC_c < 2$). We included the null model as a reference level for *AICc* score.

| Candidate Models | *K* | $\Delta AIC_c$ | $AIC_cWt$ | Adjusted $R^2$ |
|---|---|---|---|---|
| **8th-year stem volume ~ microtopography + Hegyi competition index + Sum of competing vegetation angles** | **6** | **0** | **0.58** | **0.229** |
| 8th-year stem volume ~ Hegyi competition index + Sum of competing vegetation angles | 4 | 2.24 | 0.19 | 0.197 |

© The Author(s) or their Institution(s)

| | | | | |
|---|---|---|---|---|
| 8th years stem volume ~ 1 | 2 | 24.3 | < 0.001 | - |

| Parameters | Lower 95% CI | Averaged estimate (β) | Upper 95% CI | Adjusted $R^2$ |
|---|---|---|---|---|
| Microtopography | | | | |
| Pit | 0.84 | 14.1 | 27.36 | 0.06 |
| Flat | 2.05 | 11.87 | 21.68 | |
| Sum of competing vegetation angles | -0.06 | -0.04 | -0.02 | 0.138 |
| Hegyi competition index | -9.30 | -5.67 | -2.05 | 0.07 |

The largest effect on black spruce stem volume was observed from the competition of surrounding shrubs and no-crop trees (Sum of competing vegetation angles) (Fig. 2A, Table 2). Competing vegetation detrimentally impacted black spruce at maximum observed competition (angles > 600°) reducing predicted stem volume of black spruce of 86% (-26.9 $dm^3$) relative to conditions where competing vegetation was absent (Fig. 2A). The predicted size decline associate with intermediate competition values (200 to 400 °) ranged between -8 and -16 $dm^3$ respectively (Fig. 2A). The explained variance of the competing vegetation was the highest among tested parameters at 13.8% (Table 2). The main contributors to this competing vegetation were willow, pin cherry and trembling aspen, in order of abundance. When competition from other crop-trees was high (Hegyi > 4), our model predicted stem volume that were 25 $m^3$ lower than the highest predicted mean stem volume under low competition (44 $dm^3$) (Fig. 2B). This corresponded to a 43.2% decrease of stem volume. However, there were a limited number of observations for high values of Hegyi competition index, which resulted in wider confidence intervals at these extreme values (Fig. 2B).

Draft

© The Author(s) or their Institution(s)




Figure 2: Predicted black spruce stem volume at the 8th years ($dm^3$) based on sum of competing vegetation angles (°) (A) and Hegyi competition index (B). Averaged predictions using the retained model are included with 95% CI. The *p-values* shows the level of significance and grey circles correspond to the original observations.

The flat microtopography exhibited a positive effect of black spruce stem volume compared to mounds (Table 2; Appendix A, Fig. A1). In contrast, the presence of a pit at the base of crop trees had a similar effect as mound and flat microtopography, with no significant increase or reduction in the predicted stem volume (Fig. A1).

## Effects of neighbouring species on black spruce volume

With the inclusion of the neighbouring species effect in the best and most parsimonious environmental model, the explained variance of black spruce stem volume increased from 22.9 to 25.4% (adjusted $R^2$). The model, incorporating the neighbouring species effect, retained 76% of the $AIC_cWt$ compared to the model without the neighbouring species effect with $\Delta AIC_c$ greater than 2, which indicate it was less parsimonious than the model with the neighbouring effect (Table 3).

Table 3 : Most parsimonious models explaining black spruce 8th-year stem volume with and without the neighbouring species effect, including the number of parameters (K), delta $AIC_c$ compared to the highest-ranked model ($\Delta AIC_c$), model weight ($AIC_cWt$), and model predictive power (adjusted $R^2$). Model in bold indicate retained model ($\Delta AIC_c < 2$).

© The Author(s) or their Institution(s)

307 The second half of the table shows parameters influencing black spruce 8th years stem volume and 95% confidence
308 intervals excluding 0.

| Candidate Models | *K* | $\Delta AIC_c$ | $AIC_cWt$ | Adjusted $R^2$ |
|---|---|---|---|---|
| **8th year stem volume ~ microtopography + Hegyi competition index + Sum of competing vegetation angles + neighbouring species** | **7** | **0** | **0.76** | **0.254** |
| 8th year stem volume ~ microtopography + Hegyi competition index + Sum of competing vegetation angles | 6 | 2.3 | 0.24 | 0.229 |

| Parameters | Lower 95% CI | Averaged estimate (β) | Upper 95% CI | Adjusted $R^2$ |
|---|---|---|---|---|
| Neighbouring species (Tamarack) | 0.58 | 9.72 | 18.86 | 0.008 |

Draft

309

310 The predicted stem volume of black spruce exhibited a 38.1% increase (31.5 $dm^3$) when
311 neighboured by tamarack compared to only other black spruce neighboured (predicted stem volume
312 was = 19.5 $dm^3$; Fig. 3). However, this positive effect of tamarack was weak as the variance
313 explained by the neighbouring species variable alone was low ($R^2$ = 0.008). The neighbouring
314 species effect alone would not predict a significant portion of stem volume variance. The observed
315 effect, though significant (p = 0.037), was not particularly strong, as the upper confidence interval
316 for black spruce neighbours and mean prediction estimate for tamarack neighbours nearly
317 overlapped (30.4 $dm^3$ and 31.5 $dm^3$, respectively) (Fig. 3).

© The Author(s) or their Institution(s)




318

319 Figure 3 : Predicted black spruce stem volume ($dm^3$) based upon neighbouring species effect (black spruce or
320 tamarack). Averaged predictions using the retained model are included with 95% CI. Grey circles correspond to the
321 original observations, and the *p-value* shows the level of significance between groups.

322 Effects of neighbouring species on immediate growing environment

323 The ANOVA revealed significant differences between tamarack and black spruce neighbouring for

324 both measured leaf area indices, total foliar N concentration, and pH of mineral and organic layers

© The Author(s) or their Institution(s)

(Appendix A; Table A1), with *p-values* ≤ 0.02 (Table A1). However, neighbouring tree species did not have an impact on the other tested variables, i.e., P and K total foliar concentration, C/N ratio of organic and mineral layers (Appendix A; Table A1).

Having tamarack as a neighbour resulted in a 25.6% increase in LAI between closest neighbouring tree (Fig. 4A). The LAI above target trees increased by 55% (Fig. 4B). These increases in canopy closure were consistently observed across both measurements, and the effect of tamarack was highly significant, with *p-values* of 0.002 and < 0.001 (Fig. 4). Furthermore, there was no difference in the mean competing vegetation cover and the number of planted crop trees (p = 0.15; p = 0.08 respectively) between plots testing the effect of neighbouring species, which enhances our confidence in the observed effect of tamarack on canopy closure (Table 1).




Figure 4 : Effect of black spruce neighbouring species for leaf area index at mid distance between the closest crop tree at 45 cm from ground level and above target tree at its mid height. The *p-values* shows the differences between groups.

© The Author(s) or their Institution(s)

338 Tamarack neighbours slightly increased pH of both soil layers, thereby reducing soil acidity (Fig.
339 5). Specifically, tamarack raised pH from 5.53 to 5.81 and from 5.00 to 5.17 for the mineral and
340 organic layers, respectively (Table A1; Fig. 5).




341

342 Figure 5 : Effect of black spruce neighbouring species on the pH of organic and mineral soil. The *p-values* show the
343 differences between groups.

344 Lastly, we observed a very slight difference in the total foliar N concentration of black spruce when
345 neighbouring tamarack versus black spruce (Appendix A; Table A1). The variation in total foliar
346 N concentration in black spruce was 0.91% with tamarack neighbouring, compared to 0.87% with
347 black spruce neighbouring.

## 348 Discussion

349 We found an early, positive effect of neighbouring tamarack on the 8$^{th}$-year stem volume of black
350 spruce. It is unusual to detect such an effect at this stage of stand development since positive effects

© The Author(s) or their Institution(s)

of neighbouring species usually occur at later stages, under closed canopy (Forrester 2014, Feng et al. 2022). The positive effect of tamarack on black spruce stem volume can be attributed to growth complementarity for resource acquisition and minimal interference. This effect could be the result of added effects of tamarack on pH and total N foliar concentrations enhancing soil conditions. Furthermore, the candidate model without the neighbouring species term was still plausible with a $\Delta AIC_c$ just over 2 (Table 3). Despite the modest strength of tamarack effect, the difference in black spruce stem volume was ecologically meaningful (mean prediction of 19.5 $dm^3$ for black spruce neighbours vs 31.5 $dm^3$ for tamarack neighbours). This shows the early high potential of mixed plantations of black spruce and tamarack to achieve over-yielding through the stand rotation. Our results are in agreement with Boyden et al. (2009), who observed higher tree volume after 11 years for both tamarack and black spruce mixed plantations compared to monocultures of both species. They attributed this over-yielding to a better resource acquisition through optimal canopy stratification. In our case, the higher stem volume of black spruce in mixed stands could be explained by low competitive interactions during the early years of plantation due to the mechanical release and black spruce's shade tolerance, even with tamarack creating higher canopy closure. The light and space availability were not constraining resources, as larger black spruce trees were observed in mixture, even though LAI measurements suggest that tamarack intercepted more sun light than black spruce neighbouring trees. Other work suggests overall foliage density is probably lower for tamarack than for black spruce (Lambert et al. 2005). On the other hand, we observed only a few black spruce trees facing high competition intensity from both shrubs and crop trees (Fig. 2). Even if tamarack grows faster and has a larger crown than black spruce (Lambert et al. 2005), it is a deciduous conifer filtering more light in spring to its neighbours than other conifer species. This can explain why we did not detect any negative shading effects on black spruce in the sampled plots even if the canopy was more closed in the mixtures. However, light and space

Draft

© The Author(s) or their Institution(s)

will become less available in the future and the potential for over-yielding should be assessed at this time. Moreover, yield is measured at the stand level, whereas our study focuses on the individual tree level. While our study shows good potential for mixed stand, we can't extrapolate on mixed plantation yield, but only on individual tree growth 8 years after planting. Finally, we sampled the dominant black spruce as they are likely to be a part of the mature stand, as the oppressed trees will probably die (Cyr and Thiffault 2009). The direct upscaling of our results to the stand level must be done with caution.

In mixed stands, the net positive effect of mixture on growth typically occurs later, peaking between 20 and 40 years (Forrester 2014, Feng et al. 2022). The growth relation effect arises during canopy closure, which increases competition for resources, thus accentuating the growth differences between monoculture and mixed plantations (Jucker et al. 2020). When a species mixture results in an overall lower level of competition for resources between different tree species, the stand yield is usually enhanced (Ratcliffe et al. 2015). The early effect observed in this study could be the beginning of a clear positive difference between monoculture and mixed plantations or could be reversed at canopy closure when competition becomes more intense (Drossler et al. 2018). Tamarack, being fast-growing and shade intolerant, exhibits good growth complementarity for space occupancy when mixed with black spruce (slow-growing shade tolerant) (Strong and Roi 1983, Jucker et al. 2015, Lu et al. 2018). Slow-growing shade tolerant mixed with shade intolerant fast-growing trees usually lead to over-yielding (Cordonnier et al. 2018). Furthermore, tamarack growth can also be enhanced by neighbouring black spruce, as shown by Dijkstra et al. (2009). Although our study only focused on black spruce size and did not sample pure tamarack plantations, it is worth noting the possible neighbouring positive effect in the other direction. Mixed stands can enhance stand yield (Pretzsch et al. 2015), but also improve growth stability through

Draft

© The Author(s) or their Institution(s)

stand rotation (Del Rio et al. 2017, Aussenac et al. 2019). While our study focuses on early stand development, future research should investigate how increasing stand density affects interspecific competition and growth dynamics in mixed planted boreal forests.

Higher canopy closure surrounding young crop trees offers some benefits, including protection from extreme temperature variations, increased soil moisture retention; it could also help to outcompete other vegetation (Balandier et al. 2009, Pitt et al. 2009, Swanson et al. 2011, Martinez Pastur et al. 2023). Additionally, canopy cover protects from frost damage during leaf out in spring (Marquis et al. 2021). Our results suggest that this difference in canopy closure is a result of the tamarack effect, as we measured competing vegetation cover and height and did not observe differences between black spruce monoculture and mixture plots (data not shown). However, there is a delicate balance between the positive effects of canopy closure and the negative effects of competition for soils resources and light. As trees become bigger, competition for light intensifies, leading to self-thinning if left unmanaged (Drossler et al. 2018). Furthermore, during the canopy closure process, competition between trees can influence the architecture of both fast and slow-growing, and shade tolerant and intolerant trees (Van de Peer et al. 2017). In our study, we observed that intense competition from crop trees and shrubs resulted in a significant reduction in black spruce stem volume. Despite the mechanical release of the plantation in 2017, we still observed a negative effect of shrubs, indicating the high potential of shrubs and non-crop trees to influence crop tree growth. The main groups forming competing vegetation were tall shrubs and intolerant hardwoods (willow, pin cherry and aspen), which can be effectively managed with mechanical release as they mainly compete for light (Balandier et al. 2006, Thiffault et al. 2014). It is possible that we mainly observed the effect of past competition conditions from competing vegetation on black spruce, which continues to influence its current growth. Early mechanical release from non-

Draft

© The Author(s) or their Institution(s)

crop species in black spruce plantations is essential for productivity since the basal area of crop trees linearly decreases as the basal area of non-crop species increases (Jobidon et al. 2004, Sharma et al. 2010). Thiffault et al. (2014) highlighted the importance of early mechanical release to prevent the overtopping of competing vegetation on crop trees, as overtopped trees may experience up to 96% decline in growth and face reduced survival. Our results corroborate this, but also highlight the negative competitive effect from other crop trees. Furthermore, the competitive effect differed when black spruce was neighboured by tamarack or black spruces. The combine results of all vegetation effect on black spruce allow us to create a gradient of competitive effects on black spruce stem volume. This gradient starts with non-crop tree having the strongest negative effect, followed by other black spruce trees with an intermediate negative effect and tamarack neighbouring with the least negative effect. Thus, a combination of mechanical release and mixed planting would maximize black spruce individual growth.

In the sampled mixed stand, the presence of tamarack slightly reduced soil acidity; higher soil pH can promote macronutrient cycling and increase their availability in soils (Binkley and Fisher 2019). However, pH difference of 0.3 and 0.5 units are common and generally have a minimal ecological impact on tree growth (Binkley and Fisher 2019), which is in the range of the pH difference we observed. Additionally, we did not observe any difference in soil C/N ratio associated with the neighbouring species effect. Therefore, it is surprising that we detected a difference in total foliar N concentration associated with the presence of tamarack. However, the difference is minimal and probably not biologically significant for black spruce. The relations between soil macronutrients and leaf nutrient concentrations can be complex and take longer than 8 years to be detected a biologically significant effect after plantation establishment. For instance, Dijkstra et al. (2009) observed a reduction in N mineralization when black spruce was mixed with tamarack

Draft

© The Author(s) or their Institution(s)

compared to black spruce monoculture, ten years after planting. This reduction was attributed to the interaction between tamarack's N-rich litter and black spruce's lignin-rich litter (Reich et al. 1998), forming a complex that is resistant to microbial degradation (Dijkstra et al. 2009). Despite this litter interaction, the above-ground biomass of mixed plantations equalled the monocultures for both tamarack and black spruce (Dijkstra et al. 2009). Tamarack litter is known for its high N content (Reich et al. 1998), which enhance the N mineralization rate compared to litter in black spruce monoculture (Moroni et al. 2009). Finding the optimal species ratio and spacing between trees can help balance the acceleration of N mineralization from tamarack litter while reducing the input of lignin-rich black spruce litter. In our study, we tested a 50-50% mixture ratio with 2 m spacing, while Dijkstra et al. (2009) also tested a 50% ratio, but with three densities (0.25, 0.5 and 1 m spacing) without observing enhanced N mineralization by tamarack compared to black spruce monoculture. We did not measure N mineralization, but we observed a slight increase in total N foliar from black spruce in mixtures. It could be the spacing which was higher in our study preventing the formation of the lignin-rich and microbial-resistant complex, but this would need to be formally measured and followed on a longer time period. Under an optimal ratio, tamarack litter could act as nutrient pumps beneficial for neighbouring trees, but this has yet to be demonstrated. An improved nutrient cycling in mixed stands compared to monoculture can be a key mechanism to improve growth and potentially stand yield (Forrester et al. 2006).

The mean density of crop trees observed was 1850/ha, despite the initial density being 2500/ha, with a maximum sampled density of 2325/ha and a minimum of 1125/ha (Table 1). These variations could be attributed to mortality or the removal of smaller crop trees during mechanical release. However, as the plantation ages, the radius of influence will expand, changing the number of trees and the species ratio that have a competitive effect on black spruce. The observed species

Draft

© The Author(s) or their Institution(s)

ratio is due to the plot being centred on target black spruce and to the competitive radius of influence only being 3.57 m considering equal or taller neighbouring trees (White et al. 2014). Furthermore, the mixed plantation was established with alternating rows of each species, thus limiting the number of crop tree influencing growth at the time of the study. Exploring various plantation densities and patterns could evidence other early interactions that influence the growth trajectory of the plantation. Boyden et al. (2009) demonstrated that high-density planting (64000/ha; 64 trees $m^{-2}$) was beneficial for tamarack – black spruce mixture during the $10^{th}$ year of growth, surpassing both monospecific plantation of tamarack and black spruce. However, this density will lead to high stand self-thinning and is not a viable option for forest management. Finally, mixed plantations with alternating species rows are easy to carry out, but as more complex plantation patterns can be difficult to consistently achieve. Although our study was conducted at a single location, the mixed-species plantation exhibited numerous encouraging signs for individual tree productivity when compared to black spruce monoculture. Further investigation should therefore aim to include more sites and locations with different competing vegetation to test our hypotheses in a larger context.

# Conclusion

Our study offers valuable insights into the drivers of black spruce growth and soil characteristics during early plantation establishment and highlights the potential of tamarack as neighbouring species to improve black spruce growth. We observed positive impacts of tamarack neighbours on black spruce attributes as well as on soil characteristics, though to a lesser extent. Even if these effects were not all ecologically significant, the emergence of numerous early differences between neighbouring species eight years after planting is both encouraging and surprising (Boyden et al. 2009). Black spruce and tamarack mixture showed great potential to be a viable management

Draft

© The Author(s) or their Institution(s)

option for wood production through the positive effect of tamarack on black spruce stem volume. However, our results showed the importance of at least one mechanical release, as shrubs and non-crop trees have shown considerable negative impacts on black spruce growth, even if the studied plantation was released in 2017.

Assessing if mixed stand of black spruce – tamarack support greater biodiversity compared to their respective monocultures could add substantial weight to the case for this forest management strategy. Other associated advantages of mixed stands, such as heightened resistance and resilience to disturbances, drought, insects, and diseases (Kelty 2006, Jactel and Brockerhoff 2007, Poeydebat et al. 2021), should also be assessed for tamarack-black spruce stands. To ascertain the inference potential and extent of our results, we recommended that our study design be replicated in other bioclimatic domains, including diverse soil conditions, and monitored over the long-term. Such studies would strengthen the knowledge and validate, or possibly challenges, our findings regarding the mixed stand of tamarack and black spruce.

Draft

## Acknowledgments

We thank M. Madore for her help in field work, D. Charron for coordinating field work, M. Létourneau for conducting soil and leaf elemental analyses, B. Harvey for plantation design and O. Villemaire-Côté for comments on an earlier version of this manuscript. We are also indebted to two anonymous reviewers and the Associate Editor for providing constructive comments. Funding for this work was provided by the Natural Sciences and Engineering Research Council of Canada (RDCPJ 525064-18), in collaboration with Scierie Landrienne, GreenFirst Forest Products, the Ministère des Ressources naturelles et des Forêts du Québec (3329-2017-1423321543), and the Canadian Wood Fibre Centre of Natural Resources Canada.

© The Author(s) or their Institution(s)

## Author contributions

AL and NT acquired funding. SRP, NT and AL conceptualized the project. SRP wrote the original draft. SRP collected data. SRP, NT, AL and JL conducted the analyses. NT, AL and JL read, commented, edited and approved the final manuscript.

## Data availability

Data will be made available on request.

## Supporting information

The details on the supporting information are presented below:

## Conflicts of Interest

The authors declare that they have no competing interests.

Draft

## Reference

Akaike, H. 1998. Information Theory and an Extension of the Maximum Likelihood Principle. In: Parzen, E., Tanabe, K., Kitagawa, G. (eds) Selected Papers of Hirotugu Akaike. Springer Series in Statistics. Springer, New York, NY. pp. 199–213.

Alva, A.K., Paramasivam, S., Obreza, T.A., and Schumann, A.W. 2006. Nitrogen best management practice for citrus trees: I. Fruit yield, quality, and leaf nutritional status. Sci. Hortic. **107**(3): 233–244. doi:10.1016/j.scienta.2005.05.017.

© The Author(s) or their Institution(s)

Anderson, D.R., and Burnham, K.P. 2002. Avoiding pitfalls when using information-theoretic methods. J. Wild. Manage. **66**(3): 912–918. doi:10.2307/3803155.

Aussenac, R., Bergeron, Y., Gravel, D., Drobyshev, I., and Baltzer, J. 2019. Interactions among trees: a key element in the stabilising effect of species diversity on forest growth. Funct. Ecol. **33**(2): 360–367. doi:10.1111/1365-2435.13257.

Balandier, P., Collet, C., Miller, J.H., Reynolds, P.E., and Zedaker, S.M. 2006. Designing forest vegetation management strategies based on the mechanisms and dynamics of crop tree competition by neighbouring vegetation. Forestry. **79**(1): 3–27. doi:10.1093/forestry/cpi056.

Balandier, P., Frochot, H., and Sourisseau, A. 2009. Improvement of direct tree seeding with cover crops in afforestation: microclimate and resource availability induced by vegetation composition. For. Ecol. Manage. **257**(8): 1716–1724. doi:10.1016/j.foreco.2009.01.032.

Baraloto, C., Goldberg, D.E., and Bonal, D. 2005. Performance trade-offs among tropical tree seedlings in contrasting microhabitats. Ecology. **86**(9): 2461–2472. doi:10.1890/04-1956.

Bates, D., Kliegl, R., Vasishth, S., and Baayen, H. 2015. Parsimonious mixed models. *arXiv* preprint arXiv:1506.04967.

© The Author(s) or their Institution(s)

552 Beaudoin, A., Bernier, P.Y., Guindon, L., Villemaire, P., Guo, X., Stinson, G., Bergeron, T.,
553 Magnussen, S., and Hall, R.J. 2014. Mapping attributes of Canada's forests at moderate resolution
554 through kNN and MODIS imagery. Can. J. For. Res. **44**: 521–532. doi:10.1139/cjfr-2013-0401.
555
556 Bergeron, Y. 2000. Species and stand dynamics in the mixed woods of Quebec's southern boreal
557 forest. Ecology. **81**(6): 1500–1516. doi:10.1890/0012-9658(2000)081[1500:SASDIT]2.0.CO;2.
558
559 Bianchi, S., Huuskonen, S., Hynynen, J., Oijala, T., Siipilehto J., and Saksa, T. 2021. Development
560 of young mixed Norway spruce and Scots pine stands with juvenile stand management in Finland.
561 Scand. J. For. Res. **36**(5): 374–388. doi:10.1080/02827581.2021.1936155.
562
563 Bielak, K., Dudzińska, M., and Pretzsch, H. 2014. Mixed stands of Scots pine (*Pinus sylvestris* L.)
564 and Norway spruce (*Picea abies* (L.) Karst) can be more productive than monocultures. Evidence
565 from over 100 years of observation of long-term experiments. For. Syst. **23**(3): 573–589.
566 doi:10.5424/fs/2014233-06195.
567
568 Binkley, D., and Fisher, R.F. 2019. Ecology and Management of Forest Soils. John Wiley & Sons
569 Ltd., Glasgow, UK.
570
571 Bolte, A., Ammer, C., Löf, M., Nabuurs, G.J., Schall, P., and Spathelf, P. 2009. Adaptive Forest
572 Management: A Prerequisite for Sustainable Forestry in the Face of Climate Change. In
573 Sustainable Forest Management in a Changing World: a European Perspective. Spathelf, P. (ed)
574 Springer Dordrecht. pp. 115–139.
575

© The Author(s) or their Institution(s)

576 Boyden, S.B., Reich, P.B., Puettmann, K.J., and Baker, T.R. 2009. Effects of density and ontogeny
577 on size and growth ranks of three competing tree species. J. Ecol. **97**(2): 277–288.
578 doi:10.1111/j.1365-2745.2008.01477.x.
579
580 CCFM. 2020. Canadian Council of Forest Ministers, National Forestry Database. February 02
581 2023. Available from http://nfdp.ccfm.org/en/data/regeneration.php [accessed 2023-07-26].
582
583 Coll, L., Ameztegui, A., Collet, C., Löf, M., Mason, B., Pach, M., Verheyen, K., Abrudan, I.,
584 Barbati, A., Barreiro, A., Bielak, K., Bravo-Oviedo, A., Ferrari, B., Govedar, Z., Kulhavy, J.,
585 Lazdina, D., Metslaid, M., Mohren, F., Pereira, M., Peric, S., Rasztovits, E., Short, I., Spathelf, P.
586 Sterba, H., Stojanovic, D., Valsta, L., Zlatanov, T., and Ponette Q. 2018. Knowledge gaps about
587 mixed forests: What do European forest managers want to know and what answers can science
588 provide? For. Ecol. Manage. **407**: 106–115. doi:10.1016/j.foreco.2017.10.055.
589
590 Cordonnier, T., Kunstler, G., Courbaud, B., and Morin, X. 2018. Managing tree species diversity
591 and ecosystem functions through coexistence mechanisms. Ann. For. Sci. **75**: 65.
592 doi:10.1007/s13595-018-0750-6
593
594 Del Río, M., Pretzsch, H., Ruíz-Peinado, R., Ampoorter, E., Annighöfer, P., Barbeito, I., Bielak,
595 K., Brazaitis, G., Coll, L., Drössler, L., Fabrika, M., Forrester, D.I., Heym, M., Hurt, V., Kurylyak,
596 V., Löf, M., Lombardi, F., Madrickiene, E., Matovic, B., Mohren, F., Motta, R., Ouden, J., Pach,
597 M., Ponette, Q., Schütze, G., Skrzyszewski, J., Sramek, V., Sterba, H., Stojanović, D., Svoboda,
598 M., Zlatanov, T.M., Bravo- Oviedo, A., and Hector, A. 2017. Species interactions increase the

Draft

© The Author(s) or their Institution(s)

599 temporal stability of community productivity in *Pinus sylvestris–Fagus sylvatica* mixtures across
600 Europe. J. Ecol. **105**(4): 1032–1043. doi:10.1111/1365-2745.12727.
601
602 Cyr, G., and Thiffault, N. 2009. Long-term black spruce plantation growth and structure after
603 release and juvenile cleaning: A 24-year study. For. Chron. **85**(3): 417–426. doi:10.5558/tfc85417-
604 3.
605
606 Dijkstra, F.A., West, J.B., Hobbie, S.E., and Reich, P.B. 2009. Antagonistic effects of species on
607 C respiration and net N mineralization in soils from mixed coniferous plantations. For. Ecol.
608 Manage. **257**(3): 1112-1118. doi:10.1016/j.foreco.2008.11.014.
609
610 Drossler, L., Agestam, E., Bielak, K., Dudzinska, M., Koricheva, J., Liziniewicz, M., Löf, M.,
611 Mason, B., Pretzsch, H., Valkonen, S., and Wellhausen, K. 2018. Over-and underyielding in time
612 and space in experiments with mixed stands of Scots pine and Norway spruce. Forests. **9**(8): 495.
613 doi:10.3390/f9080495.
614
615 Environment Canada. 2021. Canadian Daily Climate Data. National Climate Data and Information
616 Archive, Ottawa, ON, Canada. Available from
617 https://climate.weather.gc.ca/historical_data/search_historic_data_e.html [accessed November 6,
618 2022].
619
620 Felton, A., Nilsson, U., Sonesson, J., Felton, A.M., Roberge, J.M., Ranius, T., Ahlstrom, M., Bergh,
621 J., Bjorkman, C., Boberg, J., Drossler, L., Fahlvik, N., Gong, P., Holmstrom, E., Keskitalo, E.C.H.,
622 Klapwijk, M.J., Laudon, H., Lundmark, T., Niklasson, M., Nordin, A., Pettersson, M., Stenlid, J.,

© The Author(s) or their Institution(s)

623 Stens, A., and Wallertz, K. 2016. Replacing monocultures with mixed-species stands: Ecosystem
624 service implications of two production forest alternatives in Sweden. Ambio. **45**(Suppl 2): 124–
625 139. doi:10.1007/s13280-015-0749-2.
626
627 Feng, Y., Schmid, B., Loreau, M., Forrester, D.I., Fei, S., Zhu, J., Tang Z., Zhu, J., Hong, P., Ji,
628 C., Shi, Y., Ji Su, H., Xiong, X., Xiao, J., Wang, S., and Fang, J. 2022. Multispecies forest
629 plantations outyield monocultures across a broad range of conditions. Science. **376**(6595): 865–
630 868. doi:10.1126/science.abm6363.
631
632 Forrester, D.I., Bauhus, J., Cowie, A.L., and Vanclay, J.K. 2006. Mixed-species plantations of
633 *Eucalyptus* with nitrogen-fixing trees: a review. For. Ecol. Manage. **233**(2–3): 211–230.
634 doi:10.1016/j.foreco.2006.05.012.
635
636 Forrester, D.I. 2014. The spatial and temporal dynamics of species interactions in mixed-species
637 forests: From pattern to process. For. Ecol. Manage. **312**: 282–292.
638 doi:10.1016/j.foreco.2013.10.003.
639
640 Gastwirth, J.L., Gel, Y.R., and Miao, W. 2009. The impact of Levene's test of equality of variances
641 on statistical theory and practice. Stat. Sci. **24**(3): 343–360. doi: 10.1214/09-STS301.
642
643 Hegyi, F. 1974. A Simulation Model for Managing Jack Pine Stands. Growth Models for Tree
644 and Stand Simulation. Royal College of Forestry. pp. 74-90.
645

Draft

© The Author(s) or their Institution(s)

Islam, M.A., and Macdonald. S.E. 2004. Ecophysiological adaptations of black spruce (*Picea mariana*) and tamarack (*Larix laricina*) seedlings to flooding. Trees. **18**: 35–42. doi:10.1007/s00468-003-0276-9.

Jactel, H., and Brockerhoff, E.G. 2007. Tree diversity reduces herbivory by forest insects. Ecol. Lett. **10**(9): 835–848. doi:10.1111/j.1461-0248.2007.01073.x.

Jobidon, R. 2000. Density-dependent effects of northern hardwood competition on selected environmental resources and young white spruce (*Picea glauca*) plantation growth, mineral nutrition, and stand structural development – a 5-year study. For. Ecol. Manage. **130**(1–3): 77–97. doi:10.1016/S0378-1127(99)00176-0.

Jobidon, R., Cyr, G., and Thiffault, N. 2004. Plant species diversity and composition along an experimental gradient of northern hardwood abundance in *Picea mariana* plantations. For. Ecol. Manage. **198**(1–3): 209–221. doi:10.1016/j.foreco.2004.04.020.

Jucker, T., Koricheva, J., Finér, L., Bouriaud, O., Iacopetti, G., and Coomes, D.A. 2020. Good things take time–diversity effects on tree growth shift from negative to positive during stand development in boreal forests. J. Ecol. **108**(6): 2198–2211. doi:10.1111/1365-2745.13464.

Kelty, M.J. 2006. The role of species mixtures in plantation forestry. For. Ecol. Manage. **233**: 195–204. doi:10.1016/j.foreco.2006.05.011.

Draft

© The Author(s) or their Institution(s)

669 Lambert, M.-C., Ung, C.H., and Raulier, F. 2005. Canadian national tree aboveground biomass
670 equations. Can. J. For. Res. **35**(8): 1996–2018. doi:10.1139/x05-112
671
672 Laganière, J., Cavard, X., Brassard, B. W., Paré, D., Bergeron, Y., and Chen, H.Y. 2015. The
673 influence of boreal tree species mixtures on ecosystem carbon storage and fluxes. For. Ecol.
674 Manage. **354**: 119–129. doi:10.1016/j.foreco.2015.06.029.
675
676 Laganière, J., Boča, A., Van Miegroet, H., and Paré, D. 2017. A tree species effect on soil that is
677 consistent across the species' range: the case of aspen and soil carbon in North
678 America. Forests. **8**(4): 113. doi:10.3390/f8040113.
679
680 Légaré, S., Bergeron, Y., Leduc, A., and Paré, D. 2001. Comparison of the understory vegetation
681 in boreal forest types of southwest Quebec. Can. J. Bot. **79**(9): 1019–1027. doi: 10.1139/cjb-79-9-
682 1019.
683
684 Liang, J., Crowther, T.W., Picard, N., Wiser, S., Zhou, M., Alberti, G., and Reich, P.B. 2016.
685 Positive biodiversity-productivity relationship predominant in global forests. Science. **354**(6309).
686 doi:10.1126/science.aaf8957.
687
688 Lu, H., Mohren, G.M., den Ouden, J., Goudiaby, V., and Sterck, F.J. 2016. Overyielding of
689 temperate mixed forests occurs in evergreen–deciduous but not in deciduous–deciduous species
690 mixtures over time in the Netherlands. For. Ecol. Manage. **376**: 321–332.
691 doi:10.1016/j.foreco.2016.06.032.
692

Draft

© The Author(s) or their Institution(s)

693 Lu, D., Wang, G.G., Zhang, J., Fang, Y., Zhu, C., and Zhu, J. 2018. Converting larch plantations
694 to mixed stands: Effects of canopy treatment on the survival and growth of planted seedlings with
695 contrasting shade tolerance. For. Ecol. Manage. **409**: 19–28. doi:10.1016/j.foreco.2017.10.058.
696
697 Lüdecke, D., Ben-Shachar, M.S., Patil, I., Waggoner, P., and Makowski, D. 2021. performance:
698 An R package for assessment, comparison and testing of statistical models. J. Open Source
699 Softw. **6**(60). doi:10.21105/joss.03139.
700
701 Marquis, B., Bergeron, Y., Simard, M., and Tremblay, F. 2021. Disentangling the effect of
702 topography and microtopography on near-ground growing-season frosts at the boreal-temperate
703 forest ecotone (Québec, Canada). New For. **52**: 1079–1098. doi:10.1007/s11056-021-09840-7.
704
705 Martínez Pastur, G., Rodríguez-Souilla, J., Lencinas, M.V., Cellini, J.M., Chaves, J.E., Aravena-
706 Acuña, M.C., Roig, F.A., and Peri, P.L. 2023. Microclimatic conditions restrict the radial growth
707 of *Nothofagus antarctica* regeneration based on the type of forest environment in Tierra del Fuego.
708 Sustainability. **15**: 8687. doi:10.3390/su15118687.
709
710 Mazerolle, M.J. 2023. AICcmodavg: Model Selection and Multimodel Inference Based on
711 (Q)AIC(c). R package 2.3-2. Available from https://cran.r-project.org/package=AICcmodavg
712 [accessed on 8 March 2023].
713
714 Moore, T.R., Trofymow, J.A., Prescott, C.E., Titus, B.D., and CIDET Working Group. 2011.
715 Nature and nurture in the dynamics of C, N and P during litter decomposition in Canadian
716 forests. Plant Soil. **339**: 163–175. doi:10.1007/s11104-010-0563-3.

Draft

© The Author(s) or their Institution(s)

Moroni, M.T., Thiffault, N., Titus, B.D., Mante, C., and Makeschin, F. 2009. Controlling *Kalmia* and reestablishing conifer dominance enhances soil fertility indicators in central Newfoundland, Canada. Can. J. For. Res. **39**(7): 1270–1279. doi:10.1139/X09-061.

Nord-Larsen, T., and Meilby, H. 2016. Effects of nurse trees, spacing, and tree species on biomass production in mixed forest plantations. Scand. J. For. Res. **31**(6): 592–601. doi:10.1080/02827581.2015.1131845.

Perala, D.A. 1971. Growth and Yield of Black Spruce on Organic Soils in Minnesota. North Central Forest Experiment Station, Forest Service, US Department of Agriculture.

Pitt, D.G., Wagner, R.G., and Towill, W.D. 2004. Ten years of vegetation succession following ground-applied release treatments in young black spruce plantations. North. J. Appl. For. **21**(3): 123–134. doi:10.1093/njaf/21.3.123.

Pitt, D.G., and Bell, F.W. 2005. Juvenile response to conifer release alternatives on aspen-white spruce boreal mixedwood sites. Part I: Stand structure and composition. For. Chron. **81**(4): 538–547. doi:10.5558/tfc81538-4.

Pitt, D.G., Morneault, A., Parker, W.C., Stinson, A., and Lanteigne, L. 2009. The effects of herbaceous and woody competition on planted white pine in a clearcut site. For. Ecol. Manage. **257**(4): 1281–1291. doi:10.1016/j.foreco.2008.11.039.

Draft

© The Author(s) or their Institution(s)

741 Poeydebat, C., Castagneyrol, B., Halder, I.V., and Jactel, H. 2021. Changes in host basal area
742 explain associational resistance of mixed forests to primary pests. For. Ecol. Manage. **495**: 119374.
743 doi:10.1016/j.foreco.2021.119374.
744
745 Portier, J., Gauthier, S., Leduc, A., Arseneault, D., and Bergeron, Y. 2016. Fire regime along
746 latitudinal gradients of continuous to discontinuous coniferous boreal forests in eastern
747 Canada. Forests. **7**(10): 211. doi:10.3390/f7100211.
748
749 Prescott, C.E., and Grayston, S.J. 2013. Tree species influence on microbial communities in litter
750 and soil: current knowledge and research needs. For. Ecol. Manage. **309**: 19–27. doi:
751 10.1016/j.foreco.2013.02.034.
752
753 Pretzsch, H. 2014. Canopy space filling and tree crown morphology in mixed-species stands
754 compared with monocultures. For. Ecol. Manage. **327**: 251–264. doi:
755 10.1016/j.foreco.2014.04.027.
756
757 Pretzsch, H., and Schütze, G. 2016. Effect of tree species mixing on the size structure, density, and
758 yield of forest stands. Eur. J. For. Res. **135**: 1–22. doi : 10.1007/s10342-015-0913-z.
759
760 Prévosto, B. 2005. Les indices de compétition en foresterie: exemples d'utilisation, intérêts et
761 limites. Rev. For. Fran. **57**(5): 413–430. doi: 10.4267/2042/5062.
762
763 R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for
764 Statistical Computing, Vienna, Austria. Available from https://www.R-project.org/.

Draft

© The Author(s) or their Institution(s)

765

766 Ratcliffe, S., Holzwarth, F., Nadrowski, K., Levick, S., and Wirth, C. 2015. Tree neighbourhood
767 matters–Tree species composition drives diversity–productivity patterns in a near-natural beech
768 forest. For Ecol. Manage. **335**: 225–234. doi: 10.1016/j.foreco.2014.09.032.

769

770 Reich, P.B., Walters, M.B., Tjoelker, M.G., Vanderklein, D., and Buschena, C. 1998.
771 Photosynthesis and respiration rates depend on leaf and root morphology and nitrogen
772 concentration in nine boreal tree species differing in relative growth rate. Funct. Ecol. **12**(3): 395–
773 405. doi:10.1046/j.1365-2435.1998.00209.x.

774

775 Saucier, J.-P., Baldwin, K., Krestov, P., and Jorgenson, T. 2015. Boreal Forests. In Routledge
776 Handbook of Forest Ecology. Peh, K.S.H., Corlett, R.T., and Bergeron, Y. (eds) Routledge
777 Handbooks, Oxford, UK, pp. 23–45.

778

779 Saucier, J.-P., Robitaille, A., and Grondin, P. 2009. Cadre bioclimatique du Québec. In Manuel de
780 foresterie, 2nd edition. Doucet R., and Côté, M. (eds) Ordre des ingénieurs forestiers du Québec,
781 Éditions Multimondes. pp. 186–205.

782

783 Sharma, M., Bell, F.W., White, R.G., Morneault, A., and Towill, W.D. 2010. Seedling size and
784 woody competition most important predictors of growth following free-to-grow assessments in
785 four boreal forest plantations. For. Chron. **86**(2): 213–224. doi:10.5558/tfc86213-2.

786

787 Strong, W., and Roi, G.L. 1983. Root-system morphology of common boreal forest trees in Alberta,
788 Canada. Can. J. For. Res. **13**(6):1164–1173. doi:10.1139/x83-155.

Draft

© The Author(s) or their Institution(s)

789

790 Symonds, M.R.E. and Moussalli, A. 2011. A brief guide to model selection, multimodel inference
791 and model averaging in behavioural ecology using Akaike's information criterion. Behav. Ecol.
792 Sociobiol. **65**: 13–21. doi:10.1007/s00265-010-1037-6.

793

794 Swanson, M.E., Franklin, J.F., Beschta, R.L., Crisafulli, C.M., DellaSala, D.A., Hutto, R.L.,
795 Lindenmayer, D.B., and Swanson, F.J. 2011. The forgotten stage of forest succession:
796 early-successional ecosystems on forest sites. Front. Ecol. Environ. **9**(2): 117–125.
797 doi:10.1890/090157.

798

799 Thiffault, N., Titus, B.D., and Moroni, M.T. 2010. Silviculture and planted species interact to
800 influence reforestation success on a *Kalmia*-dominated site–a 15-year study. For. Chron**. 86**(2):
801 234–242. doi:10.5558/tfc86234-2.

802

803 Thiffault, N., Hébert, F., Charette, L., and Jobidon, R. 2014. Large spruce seedling responses to
804 the interacting effects of vegetation zone, competing vegetation dominance and year of mechanical
805 release. Forestry. **87**(1): 153–164. doi:10.1093/forestry/cpt048.

806

807 Thiffault, N., Grondin, P., Noël, J., and Poirier, V. 2015. Ecological gradients driving the
808 distribution of four Ericaceae in boreal Quebec, Canada. Ecol. Evol. **5**(9): 1837–1853.
809 doi:10.1002/ece3.1476.

810

Draft

© The Author(s) or their Institution(s)

Toïgo, M., Vallet, P., Perot, T., Bontemps, J.-D., Piedallu, C., and Courbaud, B. 2015. Overyielding in mixed forests decreases with site productivity. J. Ecol. **103**(2): 502–512. doi:10.1111/1365-2745.12353.

Van de Peer, T., Verheyen, K., Kint, V., Van Cleemput, E., and Muys, B. 2017. Plasticity of tree architecture through interspecific and intraspecific competition in a young experimental plantation. For. Ecol. Manage. **385**: 1–9. doi: 10.1016/j.foreco.2016.11.015.

Viereck, L.A., Johnston, W.F., Burns, R., and Honkala, B. 1990. *Picea Mariana* (Mill.) BSP, Black Spruce. Silvics of North America, Vol. 1. Forest Service, US Department of Agriculture, Handbook pp. 227–237.

Vincent, J.-S., and Hardy, L. 1977. L'évolution et l'extension des lacs glaciaires Barlow et Ojibway en territoire québécois. Géo. Phys. Quat. **31**(3–4): 357–372. doi:10.7202/1000283ar.

White, R.G., Bell, F.W., Lennon, K., Morneault, A., Sharma, M., Subedi, N., and Towill, W.D. 2014. Free-growing assessments: evaluating assessment criteria using Ontario's Vegetation Management Alternatives Program (VMAP) database. Min. Nat. Resour. For. Tech. Rep. TR-149. Science and Research Branch, Northwest Biodiversity and Monitoring, Ontario Ministry of Natural Resources and Forestry.

Wiensczyk, A., Swift, K., Morneault, A., Thiffault, N., Szuba, K., and Bell, F.W. 2011. An overview of the efficacy of vegetation management alternatives for conifer regeneration in boreal forests. For. Chron. **87**(2): 175–200. doi:10.5558/tfc2011-007.

Draft

© The Author(s) or their Institution(s)

835

836 Yilmaz, M., and Akay, A. 2008. Stand damage of a selection cutting system in a uneven aged
837 mixed forest of Cimendagi in Kahramanmaras-Turkey. Int. J. Nat. Eng. Sci. **2**(1): 77–82.

838

839 Zhang, Y, Chen, H.Y.H., and Reich, P.B. 2012. Forest productivity increases with evenness,
840 species richness and trait variation: a global meta-analysis. J. Ecol. **100**(3):742–749.
841 doi:10.1111/j.1365-2745.2011.01944.

Draft

© The Author(s) or their Institution(s)

# Appendix A

Table A1 ANOVA results (*p-values*) testing the neighbouring tree species effect between black spruce (*Picea mariana*) and tamarack (*Larix laricina*), significant effects are presented in bold.

| Tested variables | Residual degree of freedom | Mean value as function of the neighbour | | *p*-values |
|---|---|---|---|---|
| | | Black spruce | Tamarack | |
| Leaf area index | | | | |
| **Above target** | **118** | **0.15** | **0.34** | **<0.001*** ** |
| **Between target tree and nearest neighbour** | **112** | **0.79** | **1.07** | **0.002**** |
| Black spruce foliar nutrients | | | | |
| **N total (%)** | **111** | **0.87** | **0.91** | **0.02*** |
| P total (%) | 112 | 2.25 | 2.70 | 0.19 |
| K total (%) | 112 | 7.88 | 9.04 | 0.31 |
| Soil nutrients | | | | |
| C/N total mineral | 104 | 15.0 | 14.74 | 0.589 |
| C/N total organic | 95 | 27.60 | 25.22 | 0.122 |
| **pH mineral** | **104** | **5.53** | **5.81** | **0.006**** |
| **pH organic** | **104** | **5.00** | **5.17** | **0.006**** |

Draft

© The Author(s) or their Institution(s)




847

848 Figure A1 : Predicted black spruce stem volume at the 8th years ($dm^3$) based upon average microtopography effect
849 (Mound, Pit and Flat). Averaged predictions using the retained model are included with 95% CI. Red open circles
850 correspond to the original observations. The *p value* and letters show the differences between groups.

Draft

© The Author(s) or their Institution(s)

Table 1 : Plot characteristics as a function of competing neighbours (black spruce or tamarack). DBH = diameter at breast height (130 cm), GLD = ground-level diameter, BA = basal area, LAI = leaf area index, ATT = above target tree, BCNT = between closest neighbouring tree. Values are presented as mean ± standard deviation.

| | **Black spruce neighbour** | **Tamarack neighbour** |
|---|---|---|
| **Soil characteristics** | | |
| Organic layer depth (cm) | 8.7 ± 5.8 | 9.2 ± 4.9 |
| C/N mineral | 15.0 ± 2.4 | 14.7 ± 2.3 |
| C/N organic | 27.6 ± 7.4 | 25.2 ± 7.1 |
| pH mineral | 5.00 ± 0.42 | 5.17 ± 0.62 |
| pH organic | 5.52 ± 0.42 | 5.81 ± 0.59 |
| **Plot level characteristics** | | |
| Tamarack BA ($m^2$ / ha) | 0.02 ± 0.05 | 0.80±0.48 |
| Black spruce BA ($m^2$ / ha) | 0.17 ± 0.19 | 0.003 ± 0.028 |
| LAI ATT | 0.15 ± 0.34 | 0.34 ± 0.32 |
| LAI BCNT | 0.79 ± 0.48 | 1.07 ± 0.45 |
| Number of crop trees | 7.5 ± 1.8 | 6.3 ± 1.8 |
| **Black spruce dimensions and foliar nutrition** | | |
| DBH (cm) | 2.53 ± 0.64 | 2.47 ± 0.67 |
| GLD (cm) | 5.17 ± 0.88 | 4.91 ± 1.06 |
| Height (cm) | 239.1 ± 30.8 | 241.4 ± 36.3 |
| Stem volume index ($dm^3$) | 35.7 ± 21.6 | 33.8 ± 23.6 |
| C/N | 60.7 ± 5.9 | 56.7 ± 5.6 |
| Total N (%) | 0.86 ± 0.10 | 0.91 ± 0.09 |

Draft

© The Author(s) or their Institution(s)

| | | |
|---|---|---|
| Total P (%) | 2.28 ± 1.72 | 2.68 ± 1.81 |
| Total K (%) | 8.06 ± 6.51 | 9.14 ± 6.52 |

Draft

© The Author(s) or their Institution(s)

Table 2 : Most parsimonious models explaining black spruce 8th-year stem volume, including the number of parameters (K), delta $AIC_c$ compared to the highest-ranked model ($\Delta AIC_c$), model weight ($AIC_cWt$), and model predictive power (Adjusted $R^2$). The second half of the table shows parameters influencing black spruce 8th years stem volume and 95% confidence intervals excluding 0. For microtopography, mound was used as the reference level. Model in bold indicate retained model ($\Delta AIC_c < 2$). We included the null model as a reference level for *AICc* score.

| Candidate Models | *K* | $\Delta AIC_c$ | $AIC_cWt$ | Adjusted $R^2$ |
|---|---|---|---|---|
| **8th-year stem volume ~ microtopography + Hegyi competition index + Sum of competing vegetation angles** | **6** | **0** | **0.58** | **0.229** |
| 8th-year stem volume ~ Hegyi competition index + Sum of competing vegetation angles | 4 | 2.24 | 0.19 | 0.197 |
| 8th years stem volume ~ 1 | 2 | 24.3 | < 0.001 | - |

| Parameters | Lower 95% CI | Averaged estimate (β) | Upper 95% CI | Adjusted $R^2$ |
|---|---|---|---|---|
| Microtopography | | | | |
| Pit | 0.84 | 14.1 | 27.36 | 0.06 |
| Flat | 2.05 | 11.87 | 21.68 | |
| Sum of competing vegetation angles | -0.06 | -0.04 | -0.02 | 0.138 |
| Hegyi competition index | -9.30 | -5.67 | -2.05 | 0.07 |

Draft

© The Author(s) or their Institution(s)

Table 3 : Most parsimonious models explaining black spruce 8th-year stem volume with and without the neighbouring species effect, including the number of parameters (K), delta $AIC_c$ compared to the highest-ranked model ($\Delta AIC_c$), model weight ($AIC_cWt$), and model predictive power (adjusted $R^2$). Model in bold indicate retained model ($\Delta AIC_c < 2$). The second half of the table shows parameters influencing black spruce 8th years stem volume and 95% confidence intervals excluding 0.

| Candidate Models | *K* | $\Delta AIC_c$ | $AIC_cWt$ | Adjusted $R^2$ |
|---|---|---|---|---|
| **8th year stem volume ~ microtopography + Hegyi competition index + Sum of competing vegetation angles + neighbouring species** | **7** | **0** | **0.76** | **0.254** |
| 8th year stem volume ~ microtopography + Hegyi competition index + Sum of competing vegetation angles | 6 | 2.3 | 0.24 | 0.229 |
| **Parameters** | **Lower 95% CI** | **Averaged estimate (β)** | **Upper 95% CI** | **Adjusted $R^2$** |
| Neighbouring species (Tamarack) | 0.58 | 9.72 | 18.86 | 0.008 |

Draft

© The Author(s) or their Institution(s)




Figure 1: Schematic representation of the plantation design in pure and mixed plots, with the list of variables assessed.

© The Author(s) or their Institution(s)




Figure 2: Predicted black spruce stem volume at the 8th years ($dm^3$) based on sum of competing vegetation angles (°) (A) and Hegyi competition index (B). Averaged predictions using the retained model are included with 95% CI. The *p-values* shows the level of significance and grey circles correspond to the original observations.

Draft

© The Author(s) or their Institution(s)




Figure 3 : Predicted black spruce stem volume ($dm^3$) based upon neighbouring species effect (black spruce or tamarack). Averaged predictions using the retained model are included with 95% CI. Grey circles correspond to the original observations, and the *p-value* shows the level of significance between groups.

© The Author(s) or their Institution(s)




Figure 4 : Effect of black spruce neighbouring species for leaf area index at mid distance between the closest crop tree at 45 cm from ground level and above target tree at its mid height. The *p-values* shows the differences between groups.

Draft

© The Author(s) or their Institution(s)




Figure 5 : Effect of black spruce neighbouring species on the pH of organic and mineral soil. The *p-values* show the differences between groups.

Draft

© The Author(s) or their Institution(s)

# Appendix A

Table A1 ANOVA results (*p-values*) testing the neighbouring tree species effect between black spruce (*Picea mariana*) and tamarack (*Larix laricina*), significant effects are presented in bold.

| Tested variables | Residual degree of freedom | Mean value as function of the neighbour | | *p*-values |
|---|---|---|---|---|
| | | Black spruce | Tamarack | |
| Leaf area index | | | | |
| **Above target** | **118** | **0.15** | **0.34** | **<0.001*** ** |
| **Between target tree and nearest neighbour** | **112** | **0.79** | **1.07** | **0.002**** |
| Black spruce foliar nutrients | | | | |
| **N total (%)** | **111** | **0.87** | **0.91** | **0.02*** |
| P total (%) | 112 | 2.25 | 2.70 | 0.19 |
| K total (%) | 112 | 7.88 | 9.04 | 0.31 |
| Soil nutrients | | | | |
| C/N total mineral | 104 | 15.0 | 14.74 | 0.589 |
| C/N total organic | 95 | 27.60 | 25.22 | 0.122 |
| **pH mineral** | **104** | **5.53** | **5.81** | **0.006**** |
| **pH organic** | **104** | **5.00** | **5.17** | **0.006**** |

Draft

© The Author(s) or their Institution(s)




Figure A1 : Predicted black spruce stem volume at the 8th years ($dm^3$) based upon average microtopography effect (Mound, Pit and Flat). Averaged predictions using the retained model are included with 95% CI. Red open circles correspond to the original observations. The *p value* and letters show the differences between groups.

Draft

© The Author(s) or their Institution(s)