

Novel discoveries of Xylariomycetidae (Ascomycota) taxa from peat swamp forests and other terrestrial habitats in Thailand

Omid Karimi^{1,2}, Naghmeh Afshari^{2,3}, Raheleh Asghari^{1,2}, Qirui Li^{4,5}, K. W. Thilini Chethana^{1,2}, Kevin D. Hyde^{1,2,6}, Fatimah O. Alotibi⁶

¹ School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

² Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

³ Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

⁴ State Key Laboratory of Functions and Applications of Medicinal Plants, Guizhou Medical University, Guiyang 550004, China

⁵ The High Efficacy Application of Natural Medicinal Resources Engineering Center of Guizhou Province (The Key Laboratory of Optimal Utilization of Natural Medicine Resources), School of Pharmaceutical Sciences, Guizhou Medical University, Guiyang, China

⁶ Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 22452, Riyadh 11495, Saudi Arabia

Corresponding author: K. W. Thilini Chethana (kandawatte.thi@mfu.ac.th)

This article is part of:

Exploring the Hidden Fungal Diversity: Biodiversity, Taxonomy, and Phylogeny of Saprobic Fungi

Edited by Samantha C. Karunarathna, Danushka Sandaruwan Tennakoon, Ajay Kumar Gautam



Academic editor:

Samantha C. Karunarathna

Received: 17 May 2024

Accepted: 8 July 2024

Published: 7 August 2024

Citation: Karimi O, Afshari N, Asghari R, Li Q, Chethana KWT, Hyde KD, Alotibi FO (2024) Novel discoveries of Xylariomycetidae (Ascomycota) taxa from peat swamp forests and other terrestrial habitats in Thailand. MycoKeys 107: 219–247. <https://doi.org/10.3897/mycokeys.107.127749>

Copyright: © Omid Karimi et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

In a comprehensive survey of fungi conducted in the northern (Chiang Rai Province) and southern (Narathiwat Province) regions of Thailand, several xylariales-like specimens were discovered. Through the integration of molecular phylogeny and morphological analyses, one previously undocumented taxon, *Oxydothis narathiwatensis* **sp. nov.**, was identified, along with *Xylaria bawanglingensis* and *Hypoxylon hypomiltum* as new host and geographical records from *Azelia xylocarpa*, and *Dalbergia cana*, respectively. In addition, *Annulohypoxylon thailandicum* was identified as a new host record from *Swietenia macrophylla* in Thailand. The morphological characters, including ascomata, asci, and ascospores, were compared with known *Oxydothis*, *Xylaria*, *Hypoxylon*, and *Annulohypoxylon* species. Multi-locus phylogenetic analyses based on ITS, LSU, and SSU (for Oxydothaceae), ITS, *rpb2*, *tub2*, and *act* (for Xylariaceae), and ITS, LSU, *rpb2*, and *tub2* (for Hypoxylaceae) gene regions were carried out to refine the taxonomic classifications of these specimens further. This research contributes to understanding fungal diversity in these ecologically significant regions, highlighting insights into the relationships among xylariales-like species.

Key words: Fungal diversity, multi-gene phylogeny, novel species, Sordariomycetes, taxonomy

Introduction

Xylariomycetidae, introduced by Eriksson and Winka (1997), is one of the largest subclasses in Ascomycota and belongs to the class Sordariomycetes (Hyde et al. 2020a). This subclass encompasses three orders and more than 35 families (Wijayawardene et al. 2022). Among these, Xylariaceae and Hypoxylaceae stand out as two particularly significant families (Wendt et al. 2018; Voglmayr et al. 2019; Hyde et al. 2020a, 2020b; Sun et al. 2021; Hernández-Restrepo et

al. 2022; Sugita et al. 2022, 2024; Wijayawardene et al. 2022), while Oxydothiaceae is one of the poorly represented families in terms of sequence data, with fewer than 20 species having sequence data (Konta et al. 2016; Hyde et al. 2020b; Senanayake et al. 2023).

Xylariaceae stands out as one of the largest and most diverse families within the Xylariales, a fact highlighted by various studies (Ju and Rogers 1996; Fröhlich and Hyde 2000; Tang et al. 2009; Stadler et al. 2013; Koyani et al. 2016; Daranagama et al. 2018; Konta et al. 2020; Samarakoon et al. 2022; Li et al. 2024a, 2024b). *Xylaria* is the largest genus in Xylariaceae, with *Xylaria hypoxylon* as the type species (Peršoh et al. 2009; Wijayawardene et al. 2022). The majority of xylariaceous species function as endophytes or saprobes, thriving on fallen woods, leaves, fruits, seeds, dung, soil, and termite nests. Notably, a few are recognized as plant pathogens (Rogers 2000; Okane et al. 2008; Stadler et al. 2013; Husbands et al. 2018; Pourmoghaddam et al. 2022).

The family Hypoxylaceae, initially proposed by de Candolle (cf. de Lamarck and de Candolle 1805), was later synonymized under Xylariaceae but resurrected and emended by Wendt et al. (2018). It encompasses approximately 422 species across 19 genera, notably *Hypoxylon* and *Annulohypoxylon* (Hyde et al. 2020a; Wijayawardene et al. 2022), thriving in diverse climates and primarily associated with dead dicotyledonous wood. *Hypoxylon*, the type genus introduced by Bulliard (1791), consists of 235 species, often found in warmer regions, particularly the neotropics (Ju and Rogers 1996; Daranagama et al. 2018). Traditionally, *Hypoxylon* was characterized by nodulisporium-like anamorphs and specific stromatal features (Ju and Rogers 1996), but modern understanding integrates morphology with molecular phylogeny and stromatal pigment profile (Hsieh et al. 2005; Kuhnert et al. 2015). As a result, *Hypoxylon* sect. *Annulata* sensu evolved into the genus *Annulohypoxylon* following phylogenetic analyses (Hsieh et al. 2005), with subsequent revisions and species descriptions (Kuhnert et al. 2017). *Annulohypoxylon*, predominantly characterized by ostioles encircled by an annulated disc, comprises 69 species (<https://www.speciesfungorum.org>), exhibiting diverse ecological roles (Pažoutová et al. 2013; Daranagama et al. 2018; Wendt et al. 2018; Cruz et al. 2020). Both *Hypoxylon* and *Annulohypoxylon* are known for harboring a diverse array of secondary metabolites, some of which exhibit promising agricultural and medicinal properties (Scherlach et al. 2010; Helaly et al. 2018; Becker and Stadler 2021).

Oxydothiaceae was introduced by Konta et al. (2016) to accommodate *Oxydothis* species. *Oxydothis* was introduced by Penzig and Saccardo (1897) with the type species *O. grisea* and two other species, *O. nigricans* and *O. maculosa*, placed in the family Amphisphaeriaceae (sensu Eriksson and Hawksworth 1991). Hyde (1993c) reviewed the genus and proposed that *Oxydothis* should be transferred from Amphisphaeriaceae to the Hyponectriaceae based on ascus, ascospore, and peridium morphologies. He also emphasized the consistency of ascus and ascospore morphology, which is essential for identifying species, and compared it with the closely related genera *Ceriospora*, *Frondispora*, *Lasiobertia* and *Leiosphaerella* (Hyde 1993c). Kang et al. (1999) transferred the genus to Clypeosphaeriaceae, but Jeewon et al. (2003) suggested that it was related to *Leiosphaerella* (Xylariales, genera incertae sedis) based on DNA sequence data. Konta et al. (2016) transferred *Oxydothis* to Oxydothiaceae (Xylariales). Besides confirming *Oxydothis* placement in Oxydothiaceae, the

family was accepted in Amphisphaeriales by Samarakoon et al. (2022) and Wijayawardene et al. (2022). Currently, *Oxydothis* comprises 80 species listed in the Species Fungorum (<https://www.speciesfungorum.org>, accessed in June 2024). *Oxydothis* species are predominantly found on bamboo, palms, and *Pandanus*, primarily as saprobes (Hyde 1993a, 1993b, 1994a, 1994b; Wang and Hyde 1999; Fröhlich and Hyde 2000; Wong and Hyde 2001; Taylor and Hyde 2003; Shenoy et al. 2005; Hidayat et al. 2006; Tibpromma et al. 2018), with occasional occurrences as pathogens (Fröhlich and Hyde 1994) and endophytes (Hyde 1994b; Taylor et al. 1999).

This paper describes one new species, *Oxydothis narathiwatensis*, and presents three new records, *Xylaria bawanglingensis*, *Hypoxylon hypomiltum*, and *Annulohypoxylon thailandicum*. Additionally, molecular sequence data and phylogenetic information are provided for these taxa.

Materials and methods

Sampling and morphological studies

Xylariaceae and Hypoxylaceae specimens were collected in 2022 from dead wood of *Azelia xylocarpa*, *Swietenia macrophylla*, and *Dalbergia cana* at Doi Tung National Park and Mae Fah Luang University, Chiang Rai, Thailand. Oxydothidaceae specimens were collected from submerged rachis of *Eleiodoxa conferta* from a peat swamp forest in Narathiwat Province, Thailand, in 2023. Morphological observations and single spore isolations followed the procedures outlined in Senanayake et al. (2020). The samples were cultured in different media, including potato dextrose agar (PDA), malt extract agar (MEA), and oatmeal agar (OA). Herbarium specimens were deposited in the Herbarium of Mae Fah Luang University (MFLU), Chiang Rai, Thailand. Living cultures are deposited in the Mae Fah Luang University Culture Collection (MFLUCC), Chiang Rai, Thailand. Faces of fungi and Index Fungorum numbers are registered as described in Jayasiri et al. (2015) and Index Fungorum (<https://www.indexfungorum.org>), respectively. Descriptions and collection details are added to the Greater Mekong Subregion database (Chaiwan et al. 2021).

DNA extraction, PCR amplification, and Phylogenetic analyses

Fresh mycelia were used to extract genomic DNA using the Mega Genomic DNA Extraction Kit, following the manufacturer's instructions (Omega Bio-tek Inc, The United States). The PCR amplifications were conducted for six loci, including Internal transcribed spacer (ITS), large subunit rDNA (LSU), nuclear small subunit rDNA (SSU), RNA polymerase II subunit (*rpb2*), beta-tubulin (*tub2*), and actin (*act*) (Table 1). Sequence data of related taxa (Xylariaceae, Hypoxylaceae, and Oxydothidaceae) were downloaded from the GenBank database (Suppl. material 1: tables S1–S3). The phylogenetic analyses based on maximum likelihood (ML) and Bayesian inferences were conducted for both the individual datasets and the concatenated dataset of six loci. The CIPRES Science Gateway platform (Miller et al. 2010) was used for performing the maximum likelihood (ML) phylogenetic analysis, utilizing the RAXMLHPC2 tool on the XSEDE (v. 8.2.10) platform (Stamatakis 2014). The analysis employed the

Table 1. Primers and PCR conditions used for each gene region in the current study.

Gene region	Primers	PCR Condition	References
ITS	ITS5/ITS4	Initial denaturation: 94 °C (3 min); 35 cycles of denaturation: 95 °C (1 min), annealing: 53 °C (55 sec) and extension 72 °C (2 min); Final extension: 72 °C (10 min)	White et al. (1990)
LSU	LR0R/LR5	Initial denaturation: 94 °C (5 min); 35 cycles of denaturation: 94 °C (30 sec), annealing: 55 °C (50 sec) and extension 72 °C (2 min); Final extension: 72 °C (10 min)	Vilgalys and Hester (1990); Rehner and Samuels (1994)
SSU	NS1/NS4	Initial denaturation: 94 °C (3 min); 35 cycles of denaturation: 94 °C (30 sec), annealing: 55 °C (50 sec) and extension 72 °C (2 min); Final extension: 72 °C (10 min)	White et al. (1990)
<i>rpb2</i>	fRPB2-5F/ fRPB2-7cR	Initial denaturation: 95 °C (5 min); 35 cycles of denaturation: 95 °C (1 min), annealing: 52 °C (1 min) and extension 72 °C (2 min); Final extension: 72 °C (10 min)	Liu et al. (1999)
<i>tub2</i>	T1/T22	Initial denaturation: 95 °C (2 min); 35 cycles of denaturation: 95 °C (1 min), annealing: 54 °C (1.5 min) and extension 72 °C (2 min); Final extension: 72 °C (10 min)	O'Donnell et al. (1998); Hsieh et al. (2005)
<i>act</i>	ACT-512F/ ACT-738R	Initial denaturation: 95 °C (5 min); 35 cycles of denaturation: 95 °C (1 min), annealing: 55 °C (30 sec) and extension 72 °C (1 min); Final extension: 72 °C (10 min)	Carbone and Kohn (1999)

GTRGAMMA substitution model and the rapid bootstrap analysis algorithm for 1000 replicates. The Bayesian analysis was performed using MrBayes v. 3.2.6 on XSEDE at the CIPRES Science Gateway, with the analysis set to two parallel runs, four Markov chains, and run for 10,000,000 generations. The pairwise homoplasy index (PHI) test was conducted using the combined sequence dataset comprising ITS, LSU, and SSU genes of closely related species using Split Tree version 4.18.2 (Huson and Bryant 2006) to evaluate the recombination level. The generated phylograms were visualized using FigTree v. 1.4.0 (Rambaut 2015), and annotations were added using Adobe Photoshop CS6 Extended version 10.0 software (Adobe Systems, United States).

Results

Phylogenetic analysis of Xylariaceae

The combined ITS, *rpb2*, *tub2*, and *act* dataset consisted of 213 isolates belonging to 151 Xylariaceae taxa, with *Hypoxylon fragiforme* (MUCL 51264), *Hypoxylon monticulosum* (MUCL 54604), and *Daldinia loculatoidea* (CBS:113279) as outgroup taxa (Suppl. material 1: table S1). The final alignment comprised 3,726 characters (ITS: 606 bp, *rpb2*: 1,191 bp, *tub2*: 1,627 bp, *act*: 302 bp), including gaps. The final ML optimization likelihood value of the best RAxML tree (Fig. 1) was -136336.552252, and the matrix had 2,610 distinct alignment patterns, with 32.19% undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.234039, C = 0.279027, G = 0.241491, T = 0.245443; substitution rates AC = 1.325774, AG = 5.176867, AT = 1.193683, CG = 1.131857, CT = 6.488876, GT = 1.000000; gamma distribution shape parameter α = 0.409124. The RAxML and Bayesian analyses yielded almost similar tree topologies. The topology of our phylogenetic tree is nearly identical to previous publications (Friebes et al.

2022; Pan et al. 2022; Ju et al. 2023; Karimi et al. 2023; Li et al. 2024b) with minor differences, which may be due to different taxon sampling, as well as different studies that have explored the taxonomy of *Xylaria* using various combinations of gene regions, such as ITS, *act*, *tub2*, and *rpb2* combination (Garcia-Aroca et al. 2021), *act* and *tub2* combination (Wangsawat et al. 2021), *tub2*, *act*, and *rpb2* combination (Hsieh et al. 2020, 2022) or ITS, *tub2*, and *rpb2* combination (Pan et al. 2022).

Taxonomy

Xylaria bawanglingensis Y.P. Wu & Q.R. Li, J. Syst. Evol. 16 (2024)

Index Fungorum: IF849696

Facesoffungi Number: FoF16035

Figs 1, 2

Description. *Saprobic* on dead wood of *Afzelia xylocarpa* (Fabaceae). **Sexual morph:** *Stromata* 0.5–2 × 0.1–0.3 cm ($\bar{x}$ = 1.2 × 0.17 cm, n = 15), solitary or in a small group, not branched or rarely branched at the base, cylindrical to obclavate, narrowing towards the apex, and sometimes with a tomentum; surface rough, wrinkled, slightly cracked, gray to blackish and interior yellow. **Perithecia** 165–380 µm diam., ($\bar{x}$ = 285 µm, n = 15), immersed, subglobose, ostioles inconspicuous. **Paraphyses** 5–10 µm width ($\bar{x}$ = 8 µm, n = 20), cylindrical, septate. **Asci** 68–153 × 4–5.5 µm ($\bar{x}$ = 109 × 4.8 µm, n = 30), 8-spored, cylindrical, and spore bearing part 67–87 µm long. Apical apparatus 0.2–2 × 1–2 µm ($\bar{x}$ = 1.3 × 1.5 µm, n = 15), bluing in Melzer's iodine reagent. **Ascospores** 8–12 × 2–5 µm ($\bar{x}$ = 10 µm × 4 µm, n = 70), uniseriate, ellipsoid, subhyaline, light brown to dark brown, smooth, guttulate, germ slit apparent, sigmoid, almost full length of spore. **Asexual morph:** Undetermined.

Culture characters. Colonies grown on PDA, reaching 30 mm in diameter after 15 days at 25 °C, under dark conditions, mycelium superficial to immersed, without pigment diffusion, sporulation and concentric zones, white on the top and back views.

Material examined. THAILAND, Doi Tung National Park, Chiang Rai, on dead wood of *Afzelia xylocarpa*, 27 September 2022, N. Afshari 5C3T2R2 (MFLU 24-0018), living culture MFLUCC 24-0087.

Known distribution. China (Li et al. 2024b), and Thailand (This study).

Known hosts. *Afzelia xylocarpa* (This study).

Notes. Our collection (MFLU 24-0018) morphologically resembles the *Xylaria bawanglingensis* (GMB1023) in having solitary, fertile inflated, roughened surface and gray stromata with interior yellow, subglobose ostiolate perithecia, cylindrical asci with a J+ apical ring and unicellular ascospores with spore-length sigmoid germ slit and almost similar sized stromata, perithecia, asci and ascospores. In the phylogenetic analyses (Fig. 1), our collection (MFLUCC 24-0087) clustered with *Xylaria bawanglingensis* (GMB1023, GMB1162) with 100% ML bootstrap support and 1.00 posterior probability support. In this study, we report our isolate (MFLU 24-0018) as a new host and geographical record of *Xylaria bawanglingensis* on *Afzelia xylocarpa* from Thailand.




Figure 1. RAxML tree based on the analysis of a combined ITS, *rpb2*, *tub2*, and *act* dataset. ML bootstrap supports (MLBS) equal to or higher than 65% and Bayesian posterior probabilities (BYPP) equal to or greater than 0.95 are given near the nodes. Newly generated isolate of the current study is in red, and ex-types are in bold. The tree is rooted to *Dal-dinia loculatoidea*, *Hypoxylon fragiforme* and *Hypoxylon monticulosum*.




Figure 1. Continued.




Figure 2. *Xylaria bawanglingensis* (MFLU 24-0018, new host and geographical record) **a** stromata on host **b, c** stromata **d** longitudinal section through the stroma **e–h** asci **i** paraphyses **j** apical ring bluing in Melzer's reagent **k, l** ascospores **m** upper view and reverse view of the one-week-old colony on PDA. Scale bars: 1 mm (**b, d**); 20 μ m (**e–g**); 70 μ m (**h**); 20 μ m (**i**); 5 μ m (**j**); 10 μ m (**k, l**).

Phylogenetic analysis of Hypoxylaceae

The combined phylogenetic dataset of ITS, LSU, *rpb2*, and *tub2* contains 164 isolates belonging to 116 Hypoxylaceae taxa (Suppl. material 1: table S2). After trimming, the analyzed dataset comprised 3,499 characters, including gaps (ITS = 605 bp, LSU = 740 bp, *rpb2* = 1,082 bp, *tub2* = 1,072 bp). The final ML optimization likelihood value of the best RAxML tree was -76207.411440 (Fig. 3), and the matrix had 1933 distinct alignment patterns, with 33.77% undetermined characters or gaps. In our phylogenetic tree, we have labeled the clades as follows: *Annulohypoxylon*: A, *Daldinia*: D, *Entonaema*: E, *Hypomontagnella*: Hm, *Hypoxylon*: Hyp, *Jackrogersella*: J, *Pyrenopolyporus*: P, *Rhopalostroma*: Rho, *Rostrophoxylon*: Ros, *Ruwenzoria*: Ruw and *Thamnomycetes*: T. The topologies of phylogenetic trees based on the combined dataset generated from ML and BI analyses were almost identical, while the statistical supports showed slight differences. Our phylogenetic tree is nearly identical to previously published studies (Wendt et al. 2018; Becker et al. 2020; Lambert et al. 2021; Ma et al. 2022; Cedeño-Sánchez et al. 2023). *Hypoxylon papillatum* clustered at a basal position (Hyp1) with 100% ML bootstrap support and 1.00 posterior probability support. The clade Hyp2 was segregated from the backbone of the tree with 90% ML bootstrap support and 1.00 posterior probability support. Members of Hyp3 were clustered with 69% ML bootstrap support and 0.99 posterior probability support, while the sister clade Hyp4 clustered with 67% ML bootstrap support. All the clades of *Annulohypoxylon* (A1-A5), *Hypomontagnella*, *Jackrogersella*, *Rostrophoxylon*, *Pyrenopolyporus*, *Entonaema*, *Ruwenzoria*, *Thamnomycetes*, *Rhopalostroma*, and *Daldinia* displayed well-supported segregation in the phylogeny (Fig. 3). Our strain of *A. thailandicum* (MFLUCC 24-0086) clustered closer to *A. thailandicum* (MFLUCC 13-0118) with 97% ML bootstrap support and 1.00 posterior probability support. Our collection (MFLUCC 24-0088) of *H. hypomiltum* clustered with *H. hypomiltum* (MUCL 51845) with 100% ML bootstrap support and 1.00 posterior probability support.

Annulohypoxylon thailandicum Daranag. & K.D. Hyde, Fungal Diversity 72: 53 (2015)

Index Fungorum: IF550799

Facesoffungi Number: FoF00373

Figs 3, 4

Description. *Saprobic* on a dead branch of *Swietenia macrophylla* (Meliaceae). **Sexual morph:** *Ascostromata* 0.8–2 × 0.5–1.5 mm ($\bar{x}$ = 1.5 × 1 mm, n = 15), spherical to hemispherical, superficial, effused-pulvinate, conglomerate, solitary or catenated, with conspicuous perithecial mounds, carbonaceous, black surface, blackish granules beneath surface and between perithecia, releasing greenish-olivaceous pigments in 10% KOH. **Perithecia** 0.35–0.8 × 0.4–1.0 mm ($\bar{x}$ = 0.5 × 0.6 mm, n = 15), immersed, spherical to pyriform, encased in carbonaceous tissue. **Ostioles** coarsely papillate, truncatum-type disc, 0.25–0.4 mm diam. **Paraphyses** not observed. **Asci** 79–100 × 3.5–4.9 µm ($\bar{x}$ = 91 × 4.9 µm, n = 20), unitunicate, cylindrical, uniseriate, rarely overlapping uniseriate, 8-spored, apical apparatus faintly bluing in Melzer's reagent.




Figure 3. RAxML tree based on the analysis of the combined ITS, LSU, *rpb2*, and *tub2* dataset. ML bootstrap supports (MLBS) equal to or higher than 65%, and the Bayesian posterior probabilities (BYPP) equal to or greater than 0.90 are given near the nodes. The ex-types are in bold. The two new sequences are shown in red font. The phylogenetic analyses constituted *Xylaria hypoxylon* (Outgroup) and 11 genera of Hypoxylaceae: *Annulohypoxylon* (A), *Daldinia* (D), *Entonaema* (E), *Hypomontagnella* (Hm), *Hypoxylon* (Hyp), *Jackrogersella* (J), *Pyrenopolyporus* (P), *Rhodalostroma* (Rho), *Rostrophoxylon* (Ros), *Ruwenzoria* (Ruw) and *Thamnomycetes* (T).




Figure 3. Continued.



Figure 4. *Annulohypoxyton thailandicum* (MFLU 24-0019, new host record) **a** host **b** stromatal habit on host **c** ostioles with ostiolar discs **d** stromata in horizontal section showing perithecia **e** pigments in KOH **f–h** asci **i** ascus apical apparatus in Melzer's reagent **j–m** ascospores **o, p** colonies on PDA after two weeks. Scale bars: 1 mm (**b**); 500 μ m (**c, d**); 30 μ m (**f–h**); 10 μ m (**i**); 5 μ m (**j–m**).

Ascospores 7–10 $\times$ 3.5–5.0 μ m ($\bar{x}$ = 8.5 $\times$ 4.5 μ m, n = 30), unicellular, ellipsoid-in-equilateral, narrowly to broadly rounded ends, hyaline at immaturity, olivaceous brown to brown at maturity, straight spore-length germ slit, episporium smooth, perispore dehiscent in 10% KOH. **Asexual morph:** Undetermined.

Culture characters. Colonies on PDA reaching 4 cm after one week, white, effuse, erose, medium, dull, rhizoid edge, no sporulation, top white, reverse white at first, turning light brown after few days (4–5 days).

Material examined. THAILAND, Mae Fah Luang University, Chiang Rai, on a dead branch of *Swietenia macrophylla* (Meliaceae), 20 November 2022, Raheleh Asghari MOF4 (MFLU 24-0019), living culture MFLUCC 24-0086.

Known distribution. China (Zhang et al. 2023), Indonesia (Yurnaliza et al. 2021), and Thailand (Liu et al. 2015, this study).

Known hosts. *Elaeis guineensis* (Yurnaliza et al. 2021) and *Swietenia macrophylla* (This study).

Notes. Our collection (MFLU 24-0019) resembles *A. thailandicum* in having effused-pulvinate black ascostromata, releasing pigment in KOH, papillate perithecia, with truncatum-type disc, cylindrical asci with J+ apical ring, ellipsoid-inequilateral brown ascospores with germ slit and dehiscent perispore in 10% KOH as detailed in Liu et al. (2015). Our specimen differs slightly in having larger perithecia and smaller ascospores (Table 2). Sharing similar morphological characteristics, such as ascospores, ostiolar disc, and the release of pigments in KOH, *A. thailandicum* (MFLU 24-0019) can be compared to *A. archeri* and *A. microcarpum* (Liu et al. 2015). However, it is crucial to note that the latter two species exhibit reddish ascostromata with a smaller truncatum-type disc, distinguishing them from *A. thailandicum*. In the multi-loci phylogenetic tree, *A. thailandicum* isolate MFLUCC 24-0086 clustered with *A. thailandicum* (MFLUCC 13-0118) with 100% ML bootstrap support and 1.00 posterior probability support (Fig. 3). In this study, we report our collection (MFLU 24-0019) as a new host record on *Swietenia macrophylla* from Thailand.

Table 2. Synopsis of morphological characteristics of *Annulohypoxylon thailandicum*, *A. archeri*, and *A. microcarpum*.

Name	stromata size (mm)	Perithecia (mm)	Asci (µm)	Apical ring (µm)	Ascospore size (µm)	Germ slit
<i>A. thailandicum</i> MFLU 13-0441 (Liu et al. 2015)	0.5–0.7 × 0.8–1.7 × 0.8 ($\bar{x}$ = 0.7 × 1.5 × 0.8)	0.35–0.5 × 0.3–0.7 ($\bar{x}$ = 0.4 × 0.5)	89–100.8 × 4.5–6.5 ($\bar{x}$ = 97 × 5.6)	globose, 1.5 × 2	6–11.5 × 4–6.5 ($\bar{x}$ = 12.5 × 5.6)	Straight
<i>A. thailandicum</i> MFLU 24-0019 (This study)	0.8–2 × 0.5–1.5 ($\bar{x}$ = 1.5 × 1)	0.35–0.8 × 0.4–1.0 mm ($\bar{x}$ = 0.5 × 0.6 mm)	79–100 × 3.5–4.9 µm ($\bar{x}$ = 91 × 4.9 µm)	discoid, wedge, 1.4–1.8 × 0.7–1.1	7–10 × 3.5–5.0 µm ($\bar{x}$ = 8.5 × 4.5 µm)	Straight
<i>A. archeri</i> (Ju and Rogers 1996)	–	0.1 mm diam	–	–	9–10.5 µm × 4–5 µm	–
<i>A. microcarpum</i> (Ju and Rogers 1996)	–	0.15–0.2 mm diam	–	–	7–8 × 3–4 µm	–

Hypoxylon hypomiltum Mont., *Annls Sci. Nat., Bot., sér. 2* 13: 356 (1840)

Index Fungorum: IF158066

Facesoffungi Number: FoF06137

Figs 3, 5

Description. *Saprobic* on dead wood of *Dalbergia cana* (Fabaceae). **Sexual morph:** *Stromata* superficial, sessile, effused, pulvinate, conspicuous, hemispherical, dense, forming compact mass, discoid, black, KOH extractable pigments yellow. *Perithecia* 360–400 × 300–350 µm ($\bar{x}$ = 350 × 300 µm, n = 15),




Figure 5. *Hypoxylon hypomiltum* (MFLU 24-0043, new host and geographical record) **a** host **b** stromatal habit on host **c** stromata in vertical section showing perithecia **d, e** asci **f** pigments in KOH **g** ascilar apical apparatus in Melzer's reagent **h-i** ascospore (**i** ascospore with perispore **j** ascospore with germ slit) **k** germinated ascospore **l** colony on PDA after three weeks. Scale bars: 500 μ m (**b, c**); 20 μ m (**d, e**); 5 μ m (**g-j**); 10 μ m (**k**).

hemispherical to spherical. **Asci** cylindrical, 8-spored, uniseriate, spore-bearing part $43\text{--}55 \times 4\text{--}6 \mu\text{m}$ ($\bar{x} = 48 \times 5 \mu\text{m}$, $n = 20$), and stipes $37\text{--}50$ ($\bar{x} = 45$, $n = 20$), with amyloid apical apparatus bluing in Melzer's reagent, discoid, $0.4\text{--}0.6 \times 1.3\text{--}1.5 \mu\text{m}$ ($\bar{x} = 5 \times 1.4 \mu\text{m}$, $n = 10$). **Ascospores** $6\text{--}8 \times 3\text{--}4 \mu\text{m}$ ($\bar{x} = 7 \times 3.5 \mu\text{m}$, $n = 20$), brown, ellipsoidal with rounded ends, uniseriate, aseptate, guttulate, straight germ slit. **Asexual morph**: Undetermined.

Culture characters. Colonies grown on PDA, reaching 55 mm in diameter after 20 days at 25 °C, under dark conditions, circular, flat, smooth, entire edge, medium dense, mycelium superficial to immersed, no sporulation, yellow pigment diffusion, light brown on the top and reverse pale yellowish-brown.

Material examined. THAILAND, Doi Tung National Park, Chiang Rai, on dead wood of *Dalbergia cana* (Fabaceae), 27 September 2022, N. Afshari 4C3T2R3a (MFLU 24-0043), living culture MFLUCC 24-0088.

Known distribution. French Guiana (Montagne 1840), Japan (Abe 1986), Panama (Standley 1933), Pakistan (Ahmad 1978), Sri Lanka (Palapathwala et al. 2019), Thailand (This study), USA (Ju and Rogers 1996), and India (Sarbhoy et al. 1971).

Known hosts. *Fagus crenata* (Abe 1986), *Morus alba* (Ahmad 1978), *Mangifera indica* (Sarbhoy et al. 1971), and *Dalbergia cana* (This study).

Notes. Our collection (MFLU 24-0043) shares similar characteristics with *Hypoxylon hypomiltum* (Montagne 1840), including effuse, pulvinate, conspicuous, hemispherical stromata, KOH extractable pigments, hemispherical to spherical perithecia, cylindrical, uniseriate asci, brown, ellipsoidal, unicellular ascospores with a straight germ slit, and mostly similar sized stromata, perithecia, asci and ascospores. In the phylogenetic analyses (Fig. 3), our collection (MFLUCC 24-0088) clustered with *Hypoxylon hypomiltum* (MUCL52887) with 100% ML bootstrap support and 1.00 posterior probability support. In this study, we report our isolate (MFLU 24-0043) as a new host and geographical record of *Hypoxylon hypomiltum* on *Dalbergia cana* from Thailand.

Phylogenetic analysis of Oxydothidaceae

The combined ITS, LSU, and SSU dataset consisted of 20 isolates belonging to Oxydothidaceae taxa, with *Vialaea mangiferae* (MFLUCC 12-0808) and *Vialaea minutella* (BRIP 56959) as outgroup taxa (Suppl. material 1: table S3). The final alignment comprised 2,657 characters (ITS: 452 bp, LSU: 1,200 bp, SSU: 1,021 bp), including gaps. The final ML optimization likelihood value of the best RaxML tree was -9733.301849 (Fig. 6), and the matrix had 1,637 distinct alignment patterns, with 30% undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.256773, C = 0.222793, G = 0.273403, T = 0.247031; substitution rates AC = 1.198924, AG = 2.053309, AT = 1.196302, CG = 1.103201, CT = 3.773909, GT = 1.000000; gamma distribution shape parameter $\alpha = 0.253842$. The topology of our phylogenetic tree is identical to previous publications (Konta et al. 2016; Hu et al. 2022; Senanayake et al. 2023). Our collection of the new species *Oxydothis narathiwatensis* (MFLUCC 24-0085) is from a distinct clade with *Oxydothis hoehnelii* (HKUCC 3854) and *Oxydothis* sp. (IFO 32218) with 99% ML bootstrap support (Fig. 6).

***Oxydothis narathiwatensis* O. Karimi & K.D. Hyde, sp. nov.**

Index Fungorum: IF902133

Facesoffungi Number: FoF16036

Figs 6–8

Etymology. The epithet “*narathiwatensis*” refers to Narathiwat Province, where the holotype was collected.

Holotype. MFLU 24-0044.

Description. *Saprobic* on the submerged rachis of *Eleiodoxa conferta* (Arecaceae). **Sexual morph:** *Ascomata* 170–320 µm diam., ($\bar{x}$ = 250 µm diam., n = 15), mostly in small groups, immersed, erumpent, with the non-blistering area on the host, subglobose or pyriform. **Peridium** 17–30 µm ($\bar{x}$ = 22 µm, n = 10), thick, dark brown to black, textura angularis. **Paraphyses** 40–80 × 3–6 µm ($\bar{x}$ = 62 × 4 µm, n = 20), cylindrical, fragmented, hyaline, branched or non-branched. **Asci** 171–257 × 7–11 µm ($\bar{x}$ = 225 × 9 µm, n = 20), 8-spored, cylindrical, unitunicate, short pedicellate, smooth-walled, with a J+, wedge-shaped, subapical ring. **Ascospores** 95–121 × 3–5 µm ($\bar{x}$ = 110 × 4 µm, n = 20), 2–3-seriate, hyaline, filiform, straight, curved or flexuous, rounded ends, centrally uniseptate, guttulate with smooth walls. **Appressoria** 10–20 × 9–10 µm ($\bar{x}$ = 13 × 9.5 µm, n = 10), irregular, hyaline to green, thick-walled, verrucose. **Asexual morph:** Undetermined.

Culture characters. Colonies on PDA, reaching 55 mm in diameter after 30 days at 25–27 °C, under dark conditions, medium dense, mycelium superficial to immersed, circular, flat, raised in the center with aerial mycelium, dull surface, entire edge, velvety, without pigment diffusion and sporulation, dark brown on the top and reverse-side black.

Material examined. THAILAND, Narathiwat, peat swamp forest, on the submerged rachis of *Eleiodoxa conferta* (Arecaceae), 3 August 2023, O. Karimi, 19-W (MFLU 24-0044, holotype); Ex-type living culture MFLUCC 24-0085.

Notes. Morphologically, *Oxydothis narathiwatensis* (MFLU 24-0044) shares similar characteristics with *O. gigantea* (BRIP 21921) and *O. maquilingiana* (3975) in having cylindrical asci with J+, wedge-shaped, subapical ring and filiform ascospores (Hyde 1994b). However, *O. narathiwatensis* (MFLU 24-0044) differs from *O. gigantea* (BRIP 21921) in having longer and narrower asci (171–257 × 7–11 µm vs. 240 × 20 µm), and shorter and narrower ascospore (95–121 × 3–5 µm vs. 100–150 × 6.5–7.5 µm). *Oxydothis narathiwatensis* (MFLU 24-0044) differs from *O. maquilingiana* (3975) in having longer and narrower asci (171–257 × 7–11 µm vs. 140–150 × 12–14 µm), longer and narrower ascospore (95–121 × 3–5 µm vs. 85–95 × 5–6 µm) and longer ascal ring (1.5–5 × 1–3 µm vs. 2.6–3.5 × 1.6–2.4 µm). However, due to the lack of sequence data for *O. gigantea* and *O. maquilingiana*, a phylogenetic comparison with *O. narathiwatensis* was not possible. Phylogenetically, *O. narathiwatensis* (MFLUCC 24-0085) formed a robust subclade (100% ML) basal to *O. hoehnelii* (KDH 1837). Morphologically, *O. narathiwatensis* differs from *O. hoehnelii* in having shorter and narrower asci (171–257 × 7–11 µm vs. 250–290 × 12–14 µm), fusiform ascospores against filiform ascospores in *O. narathiwatensis* (MFLU 24-0044) and longer and narrower ascospores (95–121 × 3–5 µm vs. 72–86 × 7–10 µm). The result of the pairwise homoplasy index (PHI) test revealed no significant recombination (Φ_w = 0.4) between




Figure 6. RAxML tree is based on the analysis of a combined dataset of ITS, LSU, and SSU sequence data. Maximum likelihood bootstrap support (MLBS) values equal to or higher than 60%, and the Bayesian posterior probability (BYPP) equal to or greater than 0.95 are given near the nodes. The ex-types are in bold. The new sequence is shown in yellow font. The tree is rooted with *Vialaea mangiferae* and *Vialaea minutella*.

O. narathiwatensis (MFLUCC 24-0085) and its closely related species (Fig. 8). Therefore, we introduced *Oxydothis narathiwatensis* (MFLU 24-0044) as a novel species based on morphological evidence and phylogenetic analyses (Figs 6–8).




Figure 7. *Oxydothis narathiwatensis* (MFLU 24-0044, holotype) **a** host substrate **b** close up of ascomata **c** section of ascoma **d** peridium **e** paraphyses **f** + reaction of apical ring in Melzer's reagent **g, h** asci **i-l** ascospores **m** germinating ascospore **n** appressoria **o, p** colony on PDA after two weeks. Scale bars: 500 μm (**b**); 50 μm (**c, d, g, h**); 20 μm (**e, m**); 5 μm (**f**); 25 μm (**i-l**); 10 μm (**n**).




Figure 8. The split diagram resulting from the pairwise homoplasy index (PHI) test was constructed using the combined ITS, LSU, and SSU sequence data of closely related taxa. The PHI test (Φ_w) < 0.05 indicates significant recombination within the dataset. The newly identified taxon is represented in blue.

Discussion

In this study, xylariales-like specimens were collected from Doi Tung National Park and Mae Fah Luang University in Chiang Rai Province, located in northern Thailand, and from Narathiwat Province in southern Thailand. Our investigation, employing a polyphasic approach that combines both morphological and molecular analyses (Figs 1–8), resulted in the identification of one novel species, *Oxydothis narathiwatensis*, and documented a new host record for *Annulohypoxyton thailandicum* from *Swietenia macrophylla* and two new host and geographical records of *Xylaria bawanglingensis* and *Hypoxyton hypomiltum* from *Afzelia xylocarpa* and *Dalbergia cana*, respectively, in Thailand.

Our *Xylaria* species were collected from the dead wood of *Afzelia xylocarpa*, and phylogenetic analysis reveals that it forms a sister clade with the wood-inhabiting species, *Xylaria bawanglingensis* (GMB1023 and GMB1162) and *Xylaria feejeensis* (HAST 92092013 and JRD 180). However, *Xylaria phyllocharis* (HAST 528), collected on fallen leaves (Dennis 1956) is separated from these two wood-inhabiting species in our phylogenetic tree; this finding agrees and is comparable with previous studies (Hsieh et al. 2010; Pan et al. 2022).

Annulohypoxyton thailandicum (MFLU 13-0441) was described by Daranagama & K.D. Hyde in Liu et al. (2015) from Chiang Mai. In the phylogenetic analyses, the closest species to our isolate (MFLUCC 24-0086) was found to be *A. thailandicum* (MFLUCC 13-0118), with slight differences in the ascostromata size. The relatedness is also statistically well-supported in our phylogenetic analyses (Fig. 3). *Annulohypoxyton thailandicum* (MFLU 13-0441) has also been reported from Guizhou, China (Zhang et al. 2023), and North Sumatra, Medan City, Indonesia (Yurnaliza et al. 2021). The climate of Doi Suthep, Chiang Mai, Chiang Rai (north of Thailand), Guizhou (China), and Medan City, North Sumatra (Indonesia) are mostly similar, showing a preference for Hypoxylaceae for

tropical to subtropical climates. Hence, we believe that conducting more studies in tropical to subtropical climatic areas will enhance our understanding and lead to the discovery of more undescribed species of *Annulohypoxyton*.

The delimiting characteristics of the sexual morph of *Hypoxyton* include never erecting unipartite stromata, with solid and homogenous basal tissue underneath the perithecial layer (Ju and Rogers 1996). Based on the morphological assessment, our collection fits well with *Hypoxyton hypomiltum* (Montagne 1840). In our phylogenetic analysis (Fig. 3), there is a notable clustering of all four isolates of *H. griseobrunneum*, which diverges from the findings in recent publications (Ma et al. 2022; Song et al. 2022; Yang et al. 2022). This discrepancy may stem from the utilization of sequences with incorrect names. For instance, Hsieh et al. (2005) provided β -tubulin and actin sequences for the *Hypoxyton anthochroum* isolate BCRC 34050 and incorporated them into their phylogenetic tree. However, Kuhnert et al. (2015) later updated the status of the BCRC 34050 isolate, assigning it to *H. griseobrunneum* after conducting morphological and phylogenetic analyses that revealed a close relationship with *H. griseobrunneum* isolates. Subsequently, this sequence was employed for the latter species. In contrast, recent publications (Ma et al. 2022; Song et al. 2022; Yang et al. 2022) have utilized the sequence of the BCRC 34050 isolate as *H. anthochroum*. Therefore, due to frequent changes in taxonomy, it is crucial to carefully choose accurate sequences and species names.

Oxydothis species belong to the family Oxydothidaceae and comprise approximately 80 species, predominantly associated with palms (Arecaceae) in various habitats, including peat swamps, marine, and terrestrial environments, where they are primarily found as saprobes (Hyde 1993a, 1993b, 1994b; Wang and Hyde 1999, 2001; Fröhlich and Hyde 2000; Taylor and Hyde 2003; Shenoy et al. 2005; Hidayat et al. 2006). Despite the substantial diversity within this genus, only 15 species have sequence data. Among these, some species lack crucial genetic markers, such as ITS sequences (e.g., *O. calamicola*, and *O. rhapsidicola*) or SSU or LSU sequences (e.g., *O. chinensis*, *O. cyrtostachicola*, and *O. fortunei*), which impacts the accuracy of phylogenetic reconstructions (Hidayat et al. 2006; Konta et al. 2016; Hu et al. 2022; Senanayake et al. 2023). Therefore, obtaining sequence data for previously described species is imperative to enhance our understanding of the phylogenetic relationships among *Oxydothis* species. In our study, we proposed a new species collected from submerged rachides of *Eleiodoxa conferta* in Narathiwat Province. We provided sequence data (ITS, LSU, and SSU) with detailed morphological descriptions. Notably, our investigation revealed the production of appressoria by germinating ascospores in our saprobic species, *O. narathiwatensis*, is similar to *O. garethjonesii*, *O. metroxylonicola*, *O. metroxylonis*, and *O. palmicola* (Konta et al. 2016). Appressoria in saprobic fungi raises intriguing questions about their ecological role and evolutionary adaptation. Some *Oxydothis* species are endophytes, and several produce appressoria, indicating an endophytic lifestyle (Taylor et al. 1999; Konta et al. 2016). This may suggest that these species are likely host-specific endophytes that change their lifestyle and become early saprobes when palm fronds die (Chethana et al. 2021). This supports the hypothesis that many saprobes initially start as endophytes (Chethana et al. 2021; Bhunjun et al. 2024). Additionally, the production of appressoria can help the fungi attach to their substrate in aquatic habitats (Au et al. 1996). Further exploration into the functional role of appressoria in *Oxydothis* species is warranted and could yield valuable insights into their ecology, interactions with host plants, and broad-

er evolutionary strategies. This research has the potential to significantly enhance our understanding of fungal ecology and evolution in diverse ecosystems, shedding light on the intricate relationships between their host plants.

Acknowledgements

Omid Karimi and Raheleh Asghari would like to thank Mae Fah Luang University for the award of the Partial Scholarship for the doctoral degree program and the Mushroom Research Foundation. Naghmeh Afshari would like to thank Chiang Mai University for providing the Presidential Scholarship and Mae Fah Luang University for the research collaboration. Naghmeh Afshari would also like to thank Martin van de Bult, Narong Apichai and the Doi Tung Development Project for sample collection (permission number 7700/17142 with the title “The diversity of saprobic fungi on selected hosts in forest northern Thailand”), Chiang Rai, Thailand. Omid Karimi would like to thank Dr. Saithong Kaewchai for her help with sample collection from a peat swamp forest in Narathiwat, Thailand. Also, thank the National Research Council of Thailand for providing permission to survey, study, and conduct research or academic experiments in the National Parks (No. 0907.4/5133) and Princess Sirindhorn Wildlife Sanctuary (Pa Phru To Daeng), Narathiwat (No. 0401/5816). Qirui Li would like to thank the National Natural Science Foundation of China (32000009 and 31960005). The authors extend their appreciation to the Researchers Supporting Project number (RSP2024R114), King Saud University, Riyadh, Saudi Arabia.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was funded by the Basic Research Fund (Fundamental Fund) of the National Science, Research, and Innovation Fund (Grant No. 652A01006 and 662A01003) entitled ‘Taxonomy, Phylogeny and chemo-profiling of selected families in Xylariales’.

Author contributions

Supervision: KWTC, QL, KDH, FOA, Writing – original draft: OK, NA, RA. Review and editing: KWTC, QL, KDH

Author ORCIDs

Omid Karimi  <https://orcid.org/0000-0001-9652-2222>

Naghmeh Afshari  <https://orcid.org/0009-0000-0740-6327>

Raheleh Asghari  <https://orcid.org/0009-0006-4897-5327>

Qirui Li  <https://orcid.org/0000-0001-8735-2890>

K. W. Thilini Chethana  <https://orcid.org/0000-0002-5816-9269>

Kevin D. Hyde  <https://orcid.org/0000-0002-2191-0762>

Fatimah O. Alotibi  <https://orcid.org/0000-0003-3629-5755>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Abe Y (1986) Notes on some common xylariaceous and diatrypaceous fungi on hardwoods in Japan II. *Hypoxylon hypomiltum* and its small-spored variety. Transactions of the Mycological Society of Japan 27: 51–56.
- Ahmad S (1978) Ascomycetes of Pakistan Part I. Biological Society of Pakistan Monograph 7: 1–236.
- Au DWT, Jones EBG, Moss ST, Hodgkiss IJ (1996) The role of mucilage in the attachment of conidia, germ tubes, and appressoria in the saprobic aquatic Hyphomycetes *Lemonniera aquatica* and *Mycocentrospora filiformis*. Canadian Journal of Botany 74: 1789–1800. <https://doi.org/10.1139/b96-216>
- Becker K, Stadler M (2021) Recent progress in biodiversity research on the Xylariales and their secondary metabolism. The Journal of Antibiotics 74(1): 1–23. <https://doi.org/10.1038/s41429-020-00376-0>
- Becker K, Wessel AC, Luangsa-Ard JJ, Stadler M (2020) Viridistratins A–C, antimicrobial and cytotoxic benzo[*j*]fluoranthenes from stromata of *Annulohypoxylon viridistratum* (Hypoxylaceae, Ascomycota). Biomolecules 10(5): 805. <https://doi.org/10.3390/biom10050805>
- Bhunjun CS, Phukhamsakda C, Hyde KD, McKenzie EH, Saxena RK, Li Q (2024) Do all fungi have ancestors with endophytic lifestyles? Fungal Diversity 125(1): 73–98. <https://doi.org/10.1007/s13225-023-00516-5>
- Bulliard P (1791) Histoire des champignons de la France. Paris, 384 pp.
- Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. Mycologia 91(3): 553–556. <https://doi.org/10.1080/00275514.1999.12061051>
- Cedeño-Sánchez M, Charria-Girón E, Lambert C, Luangsa-ard JJ, Decock C, Franke R, Brönstrup M, Stadler M (2023) Segregation of the genus *Parahypoxylon* (Hypoxylaceae, Xylariales) from *Hypoxylon* by a polyphasic taxonomic approach. MycoKeys 95: 131–162. <https://doi.org/10.3897/mycokeys.95.98125>
- Chaiwan N, Gomdola D, Wang S, Monkai J, Tibpromma S, Doilom M, Wanasinghe DN, Mortimer PE, Lumyong S, Hyde KD (2021) an online database providing updated information of microfungi in the Greater Mekong Subregion. Mycosphere 12(1): 1513–1526. <https://doi.org/10.5943/mycosphere/12/1/19>
- Chethana KWT, Jayawardena RS, Chen YJ, Konta S, Tibpromma S, Phukhamsakda C, Abeywickrama PD, Samarakoon MC, Senwanna C, Mapook A, Tang X, Gomdola D, Marasinghe DS, Padaruth OD, Balasuriya A, Xu J, Lumyong S, Hyde KD (2021) Appressorial interactions with host and their evolution. Fungal Diversity 110(1): 75–107. <https://doi.org/10.1007/s13225-021-00487-5>
- Cruz KD, Lima MC, de Jesus MA, de Souza AQ, Sales-Campos C (2020) *Annulohypoxylon* (Hypoxylaceae, Ascomycota) from Amazonian-forest of Brazil, with a description of one new species. New Zealand Journal of Botany 59(2): 267–284. <https://doi.org/10.1080/0028825X.2020.1846570>
- Daranagama DA, Hyde KD, Sir EB, Thambugala KM, Tian Q, Samarakoon MC, McKenzie EH, Jayasiri SC, Tibpromma S, Bhat JD, Liu X, Stadler M (2018) Towards a natural classification and backbone tree for Graphostromataceae, Hypoxylaceae, Lopadosto-

- mataceae and Xylariaceae. *Fungal Diversity* 88(1): 1–165. <https://doi.org/10.1007/s13225-017-0388-y>
- de Lamarck JB, de Candolle AP (1805) *Flore française*, vol 5, edn 3 (Paris) 2, 280 pp.
- Dennis RWG (1956) Some xylariales of tropical America. *Kew Bulletin* 11(3): 401–444. <https://doi.org/10.2307/4109126>
- Eriksson OE, Hawksworth DL (1991) Outline of the Ascomycetes–1990. *Systema Ascomycetum* 9: 39–271.
- Eriksson OE, Winka K (1997) Supraordinal taxa of Ascomycota. *Myconet* 1: 1–16.
- Friebes G, Gallé A, Hsieh HM, Ju YM (2022) *Xylaria rohrensii*, a new species inhabiting nutshells of *Juglans regia* in Austria. *Nova Hedwigia* 115(1–2): 133–142. https://doi.org/10.1127/nova_hedwigia/2022/0700
- Fröhlich J, Hyde KD (1994) New *Oxydothis* species associated with palm leaf spots in North Queensland, Australia. *Mycological Research* 98(2): 213–218. [https://doi.org/10.1016/S0953-7562\(09\)80188-3](https://doi.org/10.1016/S0953-7562(09)80188-3)
- Fröhlich J, Hyde KD (2000) Palm Microfungi. *Fungal Diversity Research Series* 3, 1–375.
- Garcia-Aroca T, Price PP, Tomaso-Peterson M, Allen TW, Wilkerson TH, Spurlock TN, Faske TR, Bluhm B, Conner K, Sikora E, Guyer R, Kelly H, Squiers BM, Doyle VP (2021) *Xylaria necrophora*, sp. nov., is an emerging root-associated pathogen responsible for taproot decline of soybean in the southern United States. *Mycologia* 113(2): 326–347. <https://doi.org/10.1080/00275514.2020.1846965>
- Helaly SE, Thongbai B, Stadler M (2018) Diversity of biologically active secondary metabolites from endophytic and saprotrophic fungi of the ascomycete order Xylariales. *Natural Product Reports* 35(9): 992–1014. <https://doi.org/10.1039/C8NP00010G>
- Hernández-Restrepo M, Decock CA, Costa MM, Crous PW (2022) Phylogeny and taxonomy of *Circinotrichum*, *Gyrothrix*, *Vermiculariopsiella* and other setose hyphomycetes. *Persoonia* 49(1): 99–135. <https://doi.org/10.3767/persoonia.2022.49.03>
- Hidayat I, Jeewon R, To-Anua C, Hyde KD (2006) The genus *Oxydothis*: New palmicolous taxa and phylogenetic relationships within the Xylariales. *Fungal Diversity* 23: 159–179.
- Hsieh HM, Ju YM, Rogers JD (2005) Molecular phylogeny of *Hypoxylon* and closely related genera. *Mycologia* 97(4): 844–865. <https://doi.org/10.1080/15572536.2006.11832776>
- Hsieh HM, Lin CR, Fang MJ, Rogers JD, Fournier J, Lechat C, Ju YM (2010) Phylogenetic status of *Xylaria* subgenus *Pseudoxylaria* among taxa of the subfamily Xylarioideae (Xylariaceae) and phylogeny of the taxa involved in the subfamily. *Molecular Phylogenetics and Evolution* 54(3): 957–969. <https://doi.org/10.1016/j.ympev.2009.12.015>
- Hsieh HM, Chou JC, Ju YM (2020) *Xylaria insolita* and *X. subescharoidea*: two newly described species collected from a termite nesting site in Hua-lien, Taiwan. *Botanical studies* 61: 1–9. <https://doi.org/10.1186/s40529-020-00287-1>
- Hsieh HM, Ju YM, Lechat C, Fournier J, Huat D (2022) New ecological, morphological, cultural and molecular phylogenetic insights into *Xylaria guepini* (Xylariaceae). *Ascomycete. Org* 14: 177–184.
- Hu HM, Liu LL, Zhang X, Lin Y, Shen XC, Long SH, Kang JC, Wijayawardene NN, Li QR, De Long Q (2022) New species and records of *Neomassaria*, *Oxydothis* and *Roussoella* (Pezizomycotina, Ascomycota) associated with palm and bamboo from China. *MycKeys* 93: 165–191. <https://doi.org/10.3897/mycokeys.93.89888>
- Husbands DR, Urbina H, Lewis SM, Aime MC (2018) *Xylaria karyophthora*: A new seed-inhabiting fungus of Greenheart from Guyana. *Mycologia* 110(2): 434–447. <https://doi.org/10.1080/00275514.2018.1457349>

- Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution* 23(2): 254–267. <https://doi.org/10.1093/molbev/msj030>
- Hyde KD (1993a) Fungi from palms. VI. Reflections on *Oxydothis* and related genera. *Sydowia* 45: 204–225.
- Hyde KD (1993b) Fungi from palms. VII. The genus *Oxydothis* from rachides of palms in north Queensland, including five new species. *Sydowia* 45: 226–240.
- Hyde KD (1993c) Fungi from palms. VI. Reflections on *Oxydothis* and related genera. *Sydowia* 45: 15–20.
- Hyde KD (1994a) Fungi from palms. XI. *Appendispora frondicola* gen. et sp. nov. from *Oncosperma horridum* in Brunei. *Sydowia* 46: 29–34.
- Hyde KD (1994b) Fungi from palms. XIII. The genus *Oxydothis*, a revision. *Sydowia* 46: 265–314.
- Hyde KD, Norphanphoun C, Maharachchikumbura SSN, Bhat DJ, Jones EBG, Bundhun D, Chen YJ, Bao DF, Boonmee S, Calabon MS, Chaiwan N, Chethana KWT, Dai DQ, Dayarathne MC, Devadatha B, Dissanayake AJ, Dissanayake LS, Doilom M, Dong W, Fan XL, Goonasekara ID, Hongsan S, Huang SK, Jayawardena RS, Jeewon R, Karunarathna A, Konta S, Kumar V, Lin CG, Liu JK, Liu NG, Luangsa-ard J, Lumyong S, Luo ZL, Marasinghe DS, McKenzie EHC, Niego AGT, Niranjan M, Perera RH, Phukhamsakda C, Rathnayaka AR, Samarakoon MC, Samarakoon SMBC, Sarma VV, Senanayake IC, Shang QJ, Stadler M, Tibpromma S, Wanasinghe DN, Wei DP, Wijayawardene NN, Xiao YP, Yang J, Zeng XY, Zhang SN, Xiang MM (2020a) Refined families of Sordariomycetes. *Mycosphere* 11(1): 305–1059. <https://doi.org/10.5943/mycosphere/11/1/7>
- Hyde KD, de Silva NI, Jeewon R, Bhat DJ, Phookamsak R, Doilom M, Boonmee S, Jayawardena RS, Maharachchikumbura SSN, Senanayake IC, Manawasinghe IS, Liu NG, Abeywickrama PD, Chaiwan N, Karunarathna A, Pem D, Lin CG, Sysouphanthong P, Luo ZL, Wei DP, Wanasinghe DN, Norphanphoun C, Tennakoon DS, Samarakoon MC, Jayasiri SC, Jiang HB, Zeng XY, Li JF, Wijesinghe SN, Devadatha B, Goonasekara ID, Brahmanage RS, Yang EF, Aluthmuhandiram JVS, Dayarathne MC, Marasinghe DS, Li WJ, Dissanayake LS, Dong W, Huanraluek N, Lumyong S, Liu JK, Karunarathna SC, Jones EBG, Al-Sadi AM, Xu JC, Harishchandra D, Sarma VV (2020b) AJOM new records and collections of fungi: 1–100. *Asian Journal of Mycology* 3(1): 22–294. <https://doi.org/10.5943/ajom/3/1/3>
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsaard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang KL, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The Faces of Fungi database: Fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74(1): 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jeewon R, Liew ECY, Hyde KD (2003) Molecular systematics of the Amphisphaeriaceae based on cladistic analyses of partial LSU rDNA gene sequences. *Mycological Research* 107(12): 1392–1402. <https://doi.org/10.1017/S095375620300875X>
- Ju YM, Rogers JD (1996) A Revision of the Genus *Hypoxylon*. *Mycologia Memoir* 20: 353–363.

- Ju YM, Hsieh HM, Suwannasai N (2023) *Xylaria furcata* reconsidered and nine resembling species. *Botanical Studies* 64(1): 21. <https://doi.org/10.1186/s40529-023-00392-x>
- Kang JC, Hyde KD, Kong RYC (1999) Studies on Amphisphaeriales. The genera excluded from the Amphisphaeriaceae, Cainiaceae and Clypeosphaeriaceae. *Fungal Diversity* 2: 135–151. <https://doi.org/10.1007/BF02464294>
- Karimi O, Chethana KWT, Farias ARG, Li QR (2023) Two new records of Xylariales species from Northern Thailand. *Phytotaxa* 598(3): 187–205. <https://doi.org/10.11646/phytotaxa.598.3.1>
- Konta S, Hongsanan S, Tibpromma S, Thongbai B, Maharachchikumbura SSN, Bahkali AH, Hyde KD, Boonmee S (2016) An advance in the endophyte story: Oxydothidaceae fam. nov. with six new species of *Oxydothis*. *Mycosphere* 7(9): 1425–1446. <https://doi.org/10.5943/mycosphere/7/9/15>
- Konta S, Hyde KD, Phookamsak R, Xu JC, Maharachchikumbura SSN, Daranagama DA, McKenzie EH, Boonmee S, Tibpromma S, Eungwanichayapant PD, Samarakoon MC (2020) Polyphyletic genera in Xylariaceae (Xylariales): *Neoxylaria* gen. nov. and *Stilbohypoxydon*. *Mycosphere* 11(1): 2629–2651. <https://doi.org/10.5943/mycosphere/11/1/17>
- Koyani RD, Patel HR, Vasava AM, Rajput KS (2016) Xylariaceae: Overview and addition to fungal diversity of Gujarat state. *Studies in Fungi* 1(1): 69–79. <https://doi.org/10.5943/sif/1/1/6>
- Kuhnert E, Surup F, Sir EB, Lambert C, Hyde KD, Hladki AI, Romero AI, Stadler M (2015) *Lenormandins* A–G, new azaphilones from *Hypoxydon lenormandii* and *Hypoxydon jaklitschii* sp. nov., recognised by chemotaxonomic data. *Fungal Diversity* 71(1): 165–184. <https://doi.org/10.1007/s13225-014-0318-1>
- Kuhnert E, Surup F, Halecker S, Stadler M (2017) *Minutellins* A – D, azaphilones from the stromata of *Annulohypoxydon minutellum* (Xylariaceae). *Phytochemistry* 137: 66–77. <https://doi.org/10.1016/j.phytochem.2017.02.014>
- Lambert C, Pourmoghadam MJ, Cedeño-Sánchez M, Surup F, Khodaparast SA, Kri-sai-Greilhuber I, Voglmayr H, Stradal TEB, Stadler M (2021) Resolution of the *Hypoxydon fuscum* complex (Hypoxyloaceae, Xylariales) and discovery and biological characterization of two of its prominent secondary metabolites. *Journal of Fungi (Basel, Switzerland)* 7(2): 131. <https://doi.org/10.3390/jof7020131>
- Li QR, Habib K, Long SH, Wu YP, Zhang X, Hu HM, Wu QZ, Liu LL, Zhou SX, Shen XC, Kang JC (2024a) Unveiling fungal diversity in China: New species and records within the Xylariaceae Family. *Mycosphere* 15(1): 275–364.
- Li QR, Habib K, Wu YP, Long SH, Zhang X, Hu HM, Wu QZ, Liu LL, Lin Y, Shen XC, Kang JC (2024b) Revisiting *Xylaria* diversity in Southern China: Descriptions of 40 new species. *Journal of Systematics and Evolution*, 1–75. <https://doi.org/10.1111/jse.13058>
- Liu Y, Whelen S, Hall B (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Liu JK, Hyde KD, Jones EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SSN, McKenzie EHC, Phookamsak R, Phukhamsakda C, Shenoy BD, Abdel-Wahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dissanayake AJ, Doilom M, D'souza MJ, Fan XL, Goonasekara ID, Hirayama K, Hongsanan S, Jayasiri SC, Jayawardena RS, Karunarathna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Peršoh D, Pinruan U, Senanayake IC, Somrithipol S, Suetrong S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D,

- Wijayawardene NN, Wanasinghe D, Wisitrassameewong K, Zeng XY, Abdel-Aziz FA, Adamčík S, Bahkali AH, Boonyuen N, Bulgakov T, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Lewis D, Li XH, Liu XZ, Liu ZY, Matsumura M, Mortimer PE, Rambold G, Randrianjohany E, Sato G, Sri-Indrasutdhi V, Tian CM, Verbeke A, von Brackel W, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi E (2015) Fungal diversity notes 1–110: Taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72(1): 1–197. <https://doi.org/10.1007/s13225-015-0324-y>
- Ma H, Song Z, Pan X, Li Y, Yang Z, Qu Z (2022) Multi-Gene Phylogeny and Taxonomy of *Hypoxyton* (Hypoxyloaceae, Ascomycota) from China. *Diversity* 14(1): 37. <https://doi.org/10.3390/d14010037>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Gateway Computing Environments Workshop, GCE 2010, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Montagne JPFC (1840) Plantes cellulaires exotiques. *Annales des Sciences Naturelles Botanique* 13: 339–359.
- O'Donnell K, Cigelnik E, Nirenberg HI (1998) Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* 90(3): 465–493. <https://doi.org/10.1080/00275514.1998.12026933>
- Okane I, Toyama K, Nakagiri A, Suzuki KI, Srikulchai P, Sivichai S, Hywel-Jones N, Potacharoen W, Læssøe T (2008) Study of endophytic Xylariaceae in Thailand: Diversity and taxonomy inferred from rDNA sequence analyses with saprobes forming fruit bodies in the field. *Mycoscience* 49(6): 359–372. <https://doi.org/10.1007/S10267-008-0440-6>
- Palapathwala PLES, Daranagama DA, Abeywickrama K, Kannangara BTSDP (2019) Two new records of *Hypoxyton hypomiltum* and *Hypomontagnella monticulosa* from Pilikuththuwa lowland wet zone forest, Sri Lanka. *Studies in Fungi* 4(1): 142–152. <https://doi.org/10.5943/sif/4/1/17>
- Pan XY, Song ZK, Qu Z, Liu TD, Ma HX (2022) Three new *Xylaria* species (Xylariaceae, Xylariales) on fallen leaves from Hainan Tropical Rainforest National Park. *Mycoskeys* 86: 47–63. <https://doi.org/10.3897/mycokeys.86.71623>
- Pažoutová S, Follert S, Bitzer J, Keck M, Surup F, Šrůtka P, Holuša J, Stadler M (2013) A new endophytic insect-associated *Daldinia* species, recognised from a comparison of secondary metabolite profiles and molecular phylogeny. *Fungal Diversity* 60(1): 107–123. <https://doi.org/10.1007/s13225-013-0238-5>
- Penzig AJO, Saccardo PA (1897) Diagnoses fungorum novorum in insula Java collectorum. Ser. II. *Malpighia* 11: 491–530. <https://doi.org/10.5962/bhl.title.4921>
- Peršoh D, Melcher M, Graf K, Fournie J, Stadler M, Rambold G (2009) Molecular and morphological evidence for the delimitation of *Xylaria hypoxyton*. *Mycologia* 101(2): 256–268. <https://doi.org/10.3852/08-108>
- Pourmoghaddam MJ, Ekiz G, Lambert C, Surup F, Primahana G, Wittstein K, Khodaparast SA, Voglmayr H, Krisai-Greilhuber I, Stradal TE, Stadler M (2022) Studies on the secondary metabolism of *Rosellinia* and *Dematophora* strains (Xylariaceae) from Iran. *Mycological Progress* 21(8): 65. <https://doi.org/10.1007/s11557-022-01816-x>
- Rambaut A (2015) FigTree v1.4: Tree figure drawing tool. 2014. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/>
- Rehner SA, Samuels GJ (1994) Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. *Mycological Research* 98(6): 625–634. [https://doi.org/10.1016/S0953-7562\(09\)80409-7](https://doi.org/10.1016/S0953-7562(09)80409-7)
- Rogers JD (2000) Thoughts and musings on tropical Xylariaceae. *Mycological Research* 104(12): 1412–1420. <https://doi.org/10.1017/S0953756200003464>

- Sarbhoy AK, Lal G, Varshney JL (1971) Fungi of India. Navyug Traders, 148 pp.
- Samarakoon MC, Hyde KD, Maharachchikumbura SS, Stadler M, Gareth Jones EB, Promputtha I, Suwannarach N, Camporesi E, Bulgakov TS, Liu JK (2022) Taxonomy, phylogeny, molecular dating and ancestral state reconstruction of *Xylariomycetidae* (Sordariomycetes). *Fungal Diversity* 112(1): 1–88. <https://doi.org/10.1007/s13225-021-00495-5>
- Scherlach K, Boettger D, Remme N, Hertweck C (2010) The chemistry and biology of cytochalasans. *Natural Product Reports* 27(6): 869–886. <https://doi.org/10.1039/b903913a>
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere : Journal of Fungal Biology* 11(1): 2678–2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Senanayake IC, Rossi W, Leonardi M, Weir A, McHugh M, Rajeshkumar KC, Verma RK, Karunarathna SC, Tibpromma S, Ashtekar N, Ashtamoorthy SK, Raveendran S, Kour G, Singh A, De la Peña-Lastra S, Mateos A, Kolařík M, Antonín V, Ševčíková H, Esteve-Raventós F, Larsson E, Pancorbo F, Moreno G, Altés A, Turégano Y, Du T-Y, Lu L, Li Q-R, Kang J-C, Gunaseelan S, Kezo K, Kaliyaperumal M, Fu J, Samarakoon MC, Gafforov Y, Teshaboeva S, Kunjan PC, Chamaparambath A, Flakus A, Etayo J, Rodriguez-Flakus P, Zhurbenko MP, de Silva NI, Tennakoon DS, Latha KPD, Manimohan P, Raj KNA, Calabon MS, Ahmadpour A, Heidarian Z, Alavi Z, Alavi F, Ghosta Y, Azizi R, Luo M, Zhao M-P, Kularathnage ND, Hua L, Yang Y-H, Liao C-F, Zhao H-J, Lestari AS, Jayasiri SC, Yu F-M, Lei L, Liu J-W, Karimi O, Tang S-M, Sun Y-R, Wang Y, Zeng M, Htet ZH, Linaldeddu BT, Alves A, Phillips AJL, Bregant C, Montecchio L, De Kesel A, Hustad VP, Miller AN, Fedosova AG, Kučera V, Raza M, Hussain M, Chen Y-P, Thiyagaraja V, Gomdola D, Rathnayaka AR, Dissanayake AJ, Suwannarach N, Hongsanan S, Maharachchikumbura SSN, Dissanayake LS, Wijayawardene NN, Phookamsak R, Lumyong S, Jones EBG, Yapa N, Wanasinghe DN, Xie N, Doilom M, Manawasinghe IS, Liu J-K, Zhao Q, Xu B, Hyde KD, Song J (2023) Fungal diversity notes 1611–1716: Taxonomic and phylogenetic contributions on fungal genera and species emphasis in south China. *Fungal Diversity* 122(1): 161–403. <https://doi.org/10.1007/s13225-023-00523-6>
- Shenoy BD, Jeewon R, Hyde KD (2005) *Oxydothis bambusicola*, a new ascomycete with a huge subapical ascus ring found on bamboo in Hong Kong. *Nova Hedwigia* 80(3–4): 511–518. <https://doi.org/10.1127/0029-5035/2005/0080-0511>
- Song ZK, Zhu AH, Liu ZD, Qu Z, Li Y, Ma HX (2022) Three New Species of *Hypoxylon* (Xylariales, Ascomycota) on a Multigene Phylogeny from Medog in Southwest China. *Journal of Fungi* 11: 8(5): 500. <https://doi.org/10.3390/jof8050500>
- Stadler M, Kuhnert E, Persöhl D, Fournier J (2013) The Xylariaceae as model example for a unified nomenclature following the “One fungus-one name” (1F1N) concept. *Mycology* 4(1): 5–21. <https://doi.org/10.1080/21501203.2013.782478>
- Stamatakis A (2014) RAXML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Standley PC (1933) The flora of Barro Colorado Island, Panama. Harvard University, Contributions from the Arnold Arboretum 5, 178 pp. <https://doi.org/10.5962/bhl.title.153450>
- Sugita R, Hirayama K, Shirouzu T, Tanaka K (2022) *Spirodecosporaceae* fam. nov. (Xylariales, Sordariomycetes) and two new species of *Spirodecospora*. *Fungal Systematics and Evolution* 10(1): 217–229. <https://doi.org/10.3114/fuse.2022.10.09>

- Sugita R, Yoshioka R, Tanaka K (2024) Anthostomella-like fungi on bamboo: Four new genera belonging to a new family Pallidoperidiaceae (Xylariales). *Mycoscience* 65(1): 28–46. <https://doi.org/10.47371/mycosci.2023.11.005>
- Sun YR, Liu NG, Samarakoon MC, Jayawardena RS, Hyde KD, Wang Y (2021) Morphology and phylogeny reveal Vamsapriyaceae fam. nov. (Xylariales, Sordariomycetes) with two novel *Vamsapriya* species. *Journal of Fungi* (Basel, Switzerland) 7(11): 891. <https://doi.org/10.3390/jof7110891>
- Tang AM, Jeewon R, Hyde KD (2009) A re-evaluation of the evolutionary relationships within the Xylariaceae based on ribosomal and protein-coding gene sequences. *Fungal Diversity* 34(1): 127–155.
- Taylor JE, Hyde KD (2003) Microfungi of tropical and temperate palms. Fungal Diversity Press, Hong Kong, 1–459.
- Taylor JE, Hyde KD, Jones EBG (1999) Endophytic fungi associated with the temperate palm, *Trachycarpus fortunei*, within and outside its natural geographic range. *The New Phytologist* 142(2): 335–346. <https://doi.org/10.1046/j.1469-8137.1999.00391.x>
- Tibpromma S, Hyde KD, McKenzie EHC, Bhat DJ, Phillips AJL, Wanasinghe DN, Samarakoon MC, Jayawardena RS, Dissanayake AJ, Tennakoon DS, Doilom M, Phookamsak R, Tang AMC, Xu J, Mortimer PE, Promputtha I, Maharachchikumbura SSN, Khan S, Karunarathna SC (2018) – Fungal diversity notes 840–928: Micro-fungi associated with Pandanaceae. *Fungal Diversity* 93(1): 1–160. <https://doi.org/10.1007/s13225-018-0408-6>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Voglmayr H, Aguirre-Hudson MB, Wagner HG, Tello S, Jaklitsch WM (2019) Lichens or endophytes? The enigmatic genus *Leptosillia* in the Leptosilliaceae fam. nov. (Xylariales), and *Furfurella* gen. nov. (Delonicicolaceae). *Persoonia* 42(1): 228–260. <https://doi.org/10.3767/persoonia.2019.42.09>
- Wang YZ, Hyde KD (1999) *Hyponectria buxi* with notes on the Hyponectriaceae. *Fungal Diversity* 3: 159–172.
- Wangsawat N, Ju YM, Phosri C, Whalley AJ, Suwannasai N (2021) Twelve new taxa of *Xylaria* associated with termite nests and soil from northeast Thailand. *Biology* (Basel) 10(7): 575. <https://doi.org/10.3390/biology10070575>
- Wendt L, Sir EB, Kuhnert E, Heitkamp S, Lambert C, Hladki AI, Romero AI, Luangsa-ard JJ, Srikitkulchai P, Peršoh D, Stadler M (2018) Resurrection and emendation of the Hypoxylaceae, recognised from a multigene phylogeny of the Xylariales. *Mycological Progress* 17(1–2): 115–154. <https://doi.org/10.1007/s11557-017-1311-3>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols: a guide to methods and applications. Academic Press, San Diego, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğdu M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkálčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro

- I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa – 2021. *Mycosphere* 13(1): 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Wong MKM, Hyde KD (2001) Fungi on grasses: New species of *Ascotaiwania*, *Diaporthe* and *Oxydothis* (Ascomycetes). *Cryptogamie. Mycologie* 22(1): 19–28. [https://doi.org/10.1016/S0181-1584\(01\)01055-7](https://doi.org/10.1016/S0181-1584(01)01055-7)
- Yang EF, Tibpromma S, Karunarathna SC, Phookamsak R, Xu JC, Zhao ZX, Karunanayake C, Promputtha I (2022) Taxonomy and phylogeny of novel and extant taxa in Pleosporales associated with *Mangifera indica* from Yunnan, China (Series I). *Journal of Fungi* (Basel, Switzerland) 8(2): 152. <https://doi.org/10.3390/jof8020152>
- Yurnaliza Y, Jamilah I, Hartanto A, Lutfia A (2021) Screening of endophytic fungi from oil palm (*Elaeis guineensis*) in producing exopolysaccharides. *Biodiversitas* (Surakarta) 22(3). <https://doi.org/10.13057/biodiv/d220350>
- Zhang JF, Liu JK, Hyde KD, Chen YY, Ran HY, Liu ZY (2023) Ascomycetes from karst landscapes of Guizhou Province, China. *Fungal Diversity* 122(1): 1–160. <https://doi.org/10.1007/s13225-023-00524-5>

Supplementary material 1

Supplementary information

Authors: Omid Karimi, Naghmeh Afshari, Raheleh Asghari, Qirui Li, K. W. Thilini Chethana, Kevin D. Hyde, Fatimah O. Alotibi

Data type: docx

Explanation note: **table S1**. List of taxa used for the phylogenetic reconstruction (Xylariaceae). GenBank accession numbers, specimen number, origin, status and reference. **table S2**. List of taxa used for the phylogenetic reconstruction (Hypoxylaceae). **table S3**. List of taxa used for the phylogenetic reconstruction (Oxydothidaceae).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.107.127749.suppl1>