

# An overview of Melanommataceae (Pleosporales, Dothideomycetes): Current insight into the host associations and geographical distribution with some interesting novel additions from plant litter

Danushka S. Tennakoon<sup>1,2,3</sup> , Kasun M. Thambugala<sup>4,5,6</sup> , Nimali I. de Silva<sup>7,8</sup> , Hai-Yan Song<sup>1,2</sup>, Nakarin Suwannarach<sup>7,8</sup> , Fu-Sheng Chen<sup>3</sup>, Dian-Ming Hu<sup>1,2</sup> 

<sup>1</sup> Bioengineering and Technological Research Centre for Edible and Medicinal Fungi, Jiangxi Agricultural University, Nanchang 330045, China

<sup>2</sup> Nanchang Key Laboratory of Edible and Medicinal Fungi, Jiangxi Agricultural University, Nanchang 330045, China

<sup>3</sup> Jiangxi Provincial Key Laboratory of Subtropical Forest Resource Cultivation, Jiangxi Agricultural University, Nanchang 330045, China

<sup>4</sup> Genetics and Molecular Biology Unit, Faculty of Applied Sciences, University of Sri Jayewardenepura, Gangodawila, Nugegoda 10250, Sri Lanka

<sup>5</sup> Center for Biotechnology, Department of Zoology, University of Sri Jayewardenepura, Nugegoda 10250, Sri Lanka

<sup>6</sup> Center for Plant Materials and Herbal Product Research, Department of Botany, Faculty of Applied Sciences, University of Sri Jayewardenepura, Nugegoda 10250, Sri Lanka

<sup>7</sup> Center of Excellence in Microbial Diversity and Sustainable Utilization, Chiang Mai University, Chiang Mai 50200, Thailand

<sup>8</sup> Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai, 50200, Thailand

Corresponding author: Dian-Ming Hu (hudianming1@163.com)



Academic editor: Nalin Wijayawardene

Received: 9 April 2024

Accepted: 19 May 2024

Published: 17 June 2024

**Citation:** Tennakoon DS, Thambugala KM, de Silva NI, Song H-Y, Suwannarach N, Chen F-S, Hu D-M (2024) An overview of Melanommataceae (Pleosporales, Dothideomycetes): Current insight into the host associations and geographical distribution with some interesting novel additions from plant litter. MycoKeys 106: 43–96. <https://doi.org/10.3897/mycokeys.106.125044>

**Copyright:** © Danushka S. Tennakoon et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

## Abstract

Melanommataceous species exhibit high diversity with a cosmopolitan distribution worldwide and show a prominent saprobic lifestyle. In this study, we explored five saprobic species collected from plant litter substrates from terrestrial habitats in China and Thailand. A combination of morphological characteristics and multi-locus phylogenetic analyses was used to determine their taxonomic classifications. Maximum Likelihood and Bayesian Inference analyses of combined LSU, SSU, ITS and *tef1-α* sequence data were used to clarify the phylogenetic affinities of the species. *Byssosphaeria poaceicola* and *Herpotrichia zingiberacearum* are introduced as new species, while three new host records, *Bertiella fici*, *By. siamensis* and *Melanomma populicola* are also reported from litter of *Cinnamomum verum*, *Citrus trifoliata* and *Fagus sylvatica*, respectively. Yet, despite the rising interest in the melanommataceous species, there is a considerable gap in knowledge on their host associations and geographical distributions. Consequently, we compiled the host-species associations and geographical distributions of all the so far known melanommataceous species.

**Key words:** Biodiversity, multi-gene phylogeny, new host records, Pleosporales, saprobes, systematics

## Introduction

Fungi occur in a wide range of ecosystems (Hawksworth 2001; De Silva et al. 2016; Hernández-Restrepo et al. 2017). With over 98,334 extant species, Ascomycota is the largest phylum of fungi and is widely distributed in terrestrial, freshwater and marine environments (Hill et al. 2021; Senanayake et

al. 2022; Bánki et al. 2023). The class Dothideomycetes is estimated to have 32,365 species and is one of the most ecologically diverse group of ascomycetes (Hongsanan et al. 2020; Bánki et al. 2023). Pleosporales is considered the largest and most diverse order in the class Dothideomycetes (Zhang et al. 2012; Hongsanan et al. 2020). They exhibit a wide range of lifestyles (e.g. biotrophs, endophytes, epiphytes, hemibiotrophs, pathogens, saprobes) and can be found worldwide, including terrestrial, marine and freshwater environments (Jones et al. 2019; Hongsanan et al. 2020; Calabon et al. 2022; Gao et al. 2023). In addition, they are extremely adaptable to various ecological niches and can exist in anaerobic, aquatic, mutualistic, terrestrial and even in severe habitats, such as deserts (Zhang et al. 2012; Hyde et al. 2013; Hongsanan et al. 2020). Pleosporales species play significant functional roles from an agricultural, ecological and economic perspectives (Zhang et al. 2012; Raja et al. 2017; Wen et al. 2020; Pimenta et al. 2021). According to the recent outline of Wijayawardene et al. (2022), Pleosporales consists of 91 families.

Melanommataceae is one of the species-rich families in Pleosporales, Dothideomycetes. It was introduced by Winter (1885) to include *Melanomma* as the type genus and species have globose or depressed ascomata, fissitunicate asci, pigmented and phragmosporous ascospores (Tian et al. 2015; Hongsanan et al. 2020). The ordinal level placement of Melanommataceae was controversial for a long time and Barr (1983) introduced Melanommatales to accommodate all taxa which have trabeculate pseudoparaphyses. The nature of trabeculate pseudoparaphyses was broadly discussed by Liew et al. (2000) and illustrated that they are generally 1 µm or thinner, clearly anastomose between the asci and are embedded in a gelatinous matrix. Subsequently, Barr (1990) introduced five genera in Melanommataceae (*Byssosphaeria*, *Keissleriella*, *Melanomma*, *Ostropella* and *Strickeria*) based on erumpent to superficial ascomata and thick-walled peridium. Over time, Melanommataceae has been transferred from Melanommatales to Pleosporales with the revolution of DNA-based molecular phylogenetic studies (Eriksson 2006; Wang et al. 2007; Zhang et al. 2012; Hyde et al. 2013; Tian et al. 2015; Hongsanan et al. 2020).

The species of Melanommataceae have cosmopolitan distribution worldwide in temperate, subtropical and tropical regions (Hyde et al. 2013; Tian et al. 2015; Jaklitsch and Voglmayr 2017; Kularathnage et al. 2022). They can play a vital role as saprobes, endophytes or hyperparasites and occur on twigs or bark of various woody plants in terrestrial, marine or freshwater habitats (Tian et al. 2015; Hashimoto et al. 2017; Tennakoon et al. 2018; Hongsanan et al. 2020; Gao et al. 2023). In addition, some species (e.g. *Seifertia alpina*) have been recorded as plant pathogens and cause bud blight or bud blast disease of *Rhododendron* species (Glawe and Hummel 2006; Li et al. 2016a). Interestingly, several species have also been reported from soil (e.g. *Herpotrichia gelasinoporoides*, *H. striatispora* and *Pleotrichocladium opacum*), on lichen species (e.g. *Aposphaeria ramalinae*) and on mushroom species (e.g. *Exosporiella fungorum* on *Thelephora* sp.) (Karsten 1892; Pitard and Harmand 1911).

The cosmopolitan nature of Melanommataceae is further supported by the numerous novel genera and species that have been introduced in the past years. Based on the year of introduction, we compiled the data and revealed that nine genera were introduced between 1800 and 1899 and nine more genera between 1900 and 1999. In addition, 18 genera were introduced between 2000 and 2024




**Figure 1.** **a** The number of melanommataceous genera introduced in different time periods **b** the number of melanommataceous species introduced in different time periods (Source – MycoBank Database).




**Figure 2.** The melanommataceous genera introduced in different time periods (Source – MycoBank Database).

(Fig. 1). This rapid increase may be primarily due to the revolution in fungal taxonomical studies with DNA sequence-based molecular phylogenies in the last two decades. Interestingly, six genera were introduced in the year 2018, such as *Marjia*, *Melanocucurbitaria*, *Melanodiplodia*, *Monoseptella*, *Pseudobyssosphaeria* and *Uzbekistanica* (Fig. 2). Consequently, 36 genera are currently accepted in Melanommataceae, viz. *Alpinaria*, *Aposphaeria*, *Asymmetricospora*, *Bertiella*, *Bicrouania*, *Byssosphaeria*, *Calyptronectria*, *Camposporium*, *Dematiomelanomma*, *Exosporiella*, *Fusiconidium*, *Herpotrichia*, *Mamillisphaeria*, *Marjia*, *Melanocamarosporioides*, *Melanocamarosporium*, *Melanocucurbitaria*, *Melanodiplodia*, *Melanomma*, *Monoseptella*, *Muriformistrickeria*, *Navicella*, *Neobyssosphaeria*, *Petrakia*, *Phragmocephala*, *Phragmotrichum*, *Pleotrichocladium*, *Praetumpfia*, *Pseudobyssosphaeria*, *Pseudodidymella*, *Pseudostrickeria*, *Sarimanas*, *Seifertia*,

*Tumularia*, *Uzbekistanica* and *Xenostigmina* (Hongsan et al. 2020; Gao et al. 2023). Conversely, the species discoveries are much higher during the 1900 and 1999 period (226 species) and thirty-nine species have been introduced between 1800 and 1899. However, despite having introduced 18 genera during the 2000–2024 period, only 76 species have been introduced (Fig. 1). Nevertheless, it could be much more in future with the extensive taxon samplings, particularly in poorly studied countries/regions, substrates and hosts.

We are exploring the fungal diversity of plant litter substrates with the aim of clarifying their taxonomy, based on morphology coupled with multi-gene phylogeny (Thambugala et al. 2017; Tennakoon et al. 2018, 2021, 2023; Wanasinghe et al. 2018). Thus, we have collected five taxa from China and Thailand which belong to the family Melanommataceae. The objectives of this study are to identify the melanommataceous taxa associated with plant litter using both morphological and phylogenetic approaches and to provide an updated checklist of species in Melanommataceae. This study provides a database on melanommataceous species for future studies, increases knowledge of fungal diversity and helps to understand their global distribution and host associations.

## Materials and methods

### Sample collection and examination

Fresh fungal specimens were collected from plant litter (dead wood and leaves) from Chiang Mai, Thailand and Kunming, China. The collected specimens were taken back to the laboratory in zip lock bags and paper envelopes. All the samples were subjected for an incubation period (one day) in plastic boxes lined with wet tissue paper. The micro- and macro-morphological characteristics were observed as described by Tennakoon et al. (2023). Sections of ascomata were taken manually and were mounted in distilled water. A stereomicroscope (AXIOSKOP 2 PLUS Series, Göttingen, Germany) was used to examine the surface morphological characteristics of fungal fruiting bodies. Micro-morphological characteristics, such as asci, ascospores and pseudoparaphyses were examined using Axioskop 2 Plus (Göttingen, Germany) compound microscope. Images were taken using Canon Axiocam 506 color digital camera (Hanover, Germany) fitted to a Axioskop 2 Plus (Göttingen, Germany) compound microscope. The micro morphological characteristics, such as colour, shape, height and diameter of ascomata, asci, ascospores, peridium and pseudoparaphyses were recorded. Indian ink was used to inspect the existence of the mucilaginous sheath in ascospores. The prepared slides were permanently preserved using lactoglycerol and sealed by applying nail-polish around the margins of cover slips. All measurements were obtained using Tarosoft (R) Image Framework application. Adobe Photoshop CS3 Extended version 10.0 software (Adobe Systems, USA) was used to construct the photo plates. Specimens were deposited in the Herbarium of Department of Biology (CMUB) and Sustainable Development of Biological Resources Laboratory (SDBR), Faculty of Science, Chiang Mai University and Herbarium of Fungi, Jiangxi Agricultural University (HFJAU), Nanchang, China. The Faces of Fungi (FOF) and Index Fungorum (IF) numbers were obtained for new species (*Byssosphaeria poaceicola* and *Herpotrichia zingiberacearum*) as mentioned in Jayasiri et al. (2015) and Index Fungorum (2024).

## DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted from fungal fruiting bodies on the natural substrate by using a DNA extraction kit (E.Z.N.A.® Forensic DNA Kit, D3591-01, Omega BIO-TEK) following the manufacturer's protocol. DNA products were intended for use as a template for PCR and stored at 4 °C and the duplicates were kept at -20 °C for long-term storage. Four genomic regions were amplified, the internal transcribed spacer (ITS) region (ITS1-5.8S-ITS2), the 28S large subunit rDNA (LSU), 18S small subunit rDNA (SSU) and the translation elongation factor 1-alpha gene (*tef1-α*). The primers ITS4 and ITS5 were used to amplify the ITS (White et al. 1990), LR0R and LR5 primers for LSU (Vilgalys and Hester 1990), NS1 and NS4 for SSU (White et al. 1990) and EF1-983F and EF1-2218R primers for *tef1-α* (Rehner 2001). The amplification reactions were performed in a total reaction volume of 25 µl, which contained 9.5 µl of sterilised distilled water, 12.5 µl of 2 × Power Taq PCR MasterMix (a premix and ready to use solution, including 0.1 Units/µl Taq DNA Polymerase, 500 µM dNTP Mixture each (dATP, dCTP, dGTP, dTTP) (Bioteke Co., China), 1 µl of each forward and reverse primers (stock concentration 10 pM) and 1 µl of DNA template. The polymerase chain reaction (PCR) thermal cycle programmes for ITS, LSU, SSU and *tef1-α* genes amplification were adjusted as described in Tennakoon et al. (2022). To check the quality of PCR products, agarose gel electrophoresis (1%) was conducted. The purified PCR products were subjected for sequencing at Sangon Biotech (Shanghai) Co., Ltd, China. Generated sequences were deposited in GenBank and accession numbers were listed (Table 1).

## Phylogenetic analyses

The obtained sequences were initially assembled (forward and reverse sequences) using SeqMan v. 7.0.0 (DNASTAR, Madison, WI). Assembled sequences were subjected to BLAST search in GenBank to obtain strains which have high similarities (<https://blast.ncbi.nlm.nih.gov/>). In addition, some other sequences for taxa in Melanommataceae were obtained using recent publications (Kularathnage et al. 2022; Gao et al. 2023). In total, 85 isolates were used for phylogenetic analyses including *Hysterium angustatum* (MFLU 16-1179) as the outgroup taxon. The combined dataset comprised four genes, ITS, LSU, SSU and *tef1-α*. Single gene sequences were aligned with MAFFT v.7.490 online application (Katoh and Standley 2013) (<https://mafft.cbrc.jp/alignment/software/>) and manually improved in necessary places. Aligned sequences were combined using BioEdit v.7.2.5 (Hall 1999).

The concatenated aligned dataset was analysed separately using Maximum Likelihood (ML) and Bayesian Inference (BI). Maximum Likelihood analysis was performed using the online portal CIPRES Science Gateway v. 3.3 (Miller et al. 2010), with RAxML-HPC v.8 on XSEDE (8.2.12) tool (Stamatakis et al. 2008; Stamatakis 2014) using the GTR+I+G model of nucleotide evolution. Evolutionary models for each barcode were determined using MrModelTest v. 2.3 (Nylander 2004) under the Akaike Information Criterion (AIC).

MrBayes 3.2.1 (Ronquist et al. 2012) was used to analyse Bayesian Inference phylogenies and was run with four chains of 3,000,000 generations and trees were sampled every 100<sup>th</sup> generation. The initial 20% of sampled data

**Table 1.** GenBank and culture collection accession numbers of species included in the phylogenetic study. The newly-generated sequences are shown in bold face.

| Fungal Species                      | Strain/Voucher No. | GenBank Accession Number |                 |                 |                 |
|-------------------------------------|--------------------|--------------------------|-----------------|-----------------|-----------------|
|                                     |                    | ITS                      | LSU             | SSU             | <i>tef-1α</i>   |
| <i>Alpinaria rhododendri</i>        | KT 2520            | LC203335                 | LC203360        | LC203314        | LC203388        |
| <i>Aposphaeria corallinolutea</i>   | MFLU 15-2752       | KY554202                 | KY554197        | KY554200        | KY554205        |
| <i>A. corallinolutea</i>            | GLMC 1355          | MT153708                 | MT156159        | –               | –               |
| <i>Bertiella ellipsoidea</i>        | MFLU 16-0583       | KX765261                 | KX765262        | –               | –               |
| <i>B. ellipsoidea</i>               | MFLUCC 17-2015     | MG543922                 | MG543913        | –               | –               |
| <i>B. fici</i>                      | MFLUCC 20-0229     | –                        | MW063223        | MW079351        | MW183786        |
| <i>B. fici</i>                      | NCYUCC 19-0260     | –                        | MW063224        | MW079352        | MW183787        |
| <i>B. fici</i>                      | NCYUCC 19-0290     | –                        | MW063225        | MW079353        | MW183788        |
| <b><i>B. fici</i></b>               | <b>CMUB 40045</b>  | –                        | <b>PP460772</b> | <b>PP460764</b> | <b>PP475453</b> |
| <i>B. macrospora</i>                | IL 5005            | –                        | GU385150        | –               | –               |
| <i>B. macrospora</i>                | SMH 3953           | –                        | –               | –               | GU327744        |
| <i>Beverwykella pulmonaria</i>      | CBS 283.53         | KY189974                 | GU301804        | –               | –               |
| <i>Byssosphaeria diffusa</i>        | AFTOL ID 1588      | –                        | DQ678071        | DQ678019        | DQ677915        |
| <i>By. diffusa</i>                  | CBS 250.62         | –                        | –               | GU205239        | –               |
| <i>By. jamaicana</i>                | SMH 1403           | –                        | GU385152        | –               | GU327746        |
| <i>By. jamaicana</i>                | SMH 3085           | –                        | GU385154        | –               | –               |
| <i>By. jamaicana</i>                | SMH 3464           | –                        | GU385153        | –               | –               |
| <i>By. macarangae</i>               | MFLUCC 17-2655     | MH389782                 | MH389778        | MH389780        | MH389784        |
| <i>By. musae</i>                    | MFLUCC 11-0146     | KP744435                 | KP744477        | KP753947        | MH581149        |
| <i>By. phoenicis</i>                | ZHKUCC 21-0122     | ON180685                 | ON180683        | ON180691        | ON243583        |
| <i>By. phoenicis</i>                | ZHKUCC 21-0123     | ON180686                 | ON180684        | ON180692        | ON243584        |
| <i>By. rhodomphala</i>              | SMH3086            | –                        | GU385155        | –               | –               |
| <i>By. rhodomphala</i>              | ANM 942            | –                        | GU385160        | –               | –               |
| <i>By. rhodomphala</i>              | SMH 3402           | –                        | GU385170        | –               | –               |
| <i>By. rhodomphala</i>              | GKM L153N          | –                        | GU385157        | –               | GU327747        |
| <i>By. salebrosa</i>                | SMH 2387           | –                        | GU385162        | –               | GU327748        |
| <i>By. schiedermayeriana</i>        | SMH 1816           | –                        | GU385159        | –               | –               |
| <i>By. schiedermayeriana</i>        | SMH 1269           | –                        | GU385158        | –               | –               |
| <i>By. schiedermayeriana</i>        | SMH 3157           | –                        | GU385163        | –               | GU327745        |
| <i>By. siamensis</i>                | MFLUCC 10-0099     | –                        | KT289895        | KT289897        | –               |
| <i>By. siamensis</i>                | MFLUCC 17-1800     | MG543923                 | MG543914        | MG543917        | –               |
| <b><i>By. siamensis</i></b>         | <b>HFJAU10336</b>  | <b>PP460780</b>          | <b>PP460773</b> | <b>PP460765</b> | <b>PP475454</b> |
| <i>By. taiwanense</i>               | MFLUCC 17-2643     | MH389783                 | MH389779        | MH389781        | MH389785        |
| <i>By. villosa</i>                  | GKM 204 N          | –                        | GU385151        | –               | GU327751        |
| <b><i>By. poaceicola</i></b>        | <b>HFJAU10337</b>  | <b>PP460781</b>          | <b>PP460774</b> | <b>PP460766</b> | <b>PP475455</b> |
| <b><i>By. poaceicola</i></b>        | <b>HFJAU10338</b>  | <b>PP460782</b>          | <b>PP460775</b> | <b>PP460767</b> | <b>PP475456</b> |
| <i>Fusiconidium aquaticum</i>       | KUMCC 15-0300      | –                        | KX641894        | KX641895        | KX641896        |
| <i>F. mackenziei</i>                | MFLUCC 14-0434     | –                        | KX611112        | KX611114        | KX611118        |
| <i>Herpotrichia herpotrichoides</i> | GKM 212N           | –                        | GU385169        | –               | –               |
| <i>H. herpotrichoides</i>           | SMH 5167           | –                        | GU385175        | –               | –               |
| <i>H. macrotricha</i>               | GKM 196N           | –                        | GU385176        | –               | GU327755        |
| <i>H. macrotricha</i>               | SMH 269            | –                        | GU385177        | –               | –               |
| <i>H. macrotricha</i>               | SMH 269            | –                        | GU385177        | –               | GU327756        |
| <i>H. vaginatispora</i>             | MFLUCC 13-0865     | –                        | KT934252        | KT934256        | KT934260        |

| Fungal Species                          | Strain/Voucher No. | GenBank Accession Number |                 |                 |                 |
|-----------------------------------------|--------------------|--------------------------|-----------------|-----------------|-----------------|
|                                         |                    | ITS                      | LSU             | SSU             | <i>tef-1α</i>   |
| <i>H. xiaokongense</i>                  | KUMCC 21-0004      | –                        | MZ408889        | MZ408891        | MZ394066        |
| <b><i>H. zingiberacearum</i></b>        | <b>HFJAU10332</b>  | <b>PP460783</b>          | <b>PP460776</b> | <b>PP460768</b> | <b>PP475457</b> |
| <b><i>H. zingiberacearum</i></b>        | <b>HFJAU10333</b>  | <b>PP460784</b>          | <b>PP460777</b> | <b>PP460769</b> | <b>PP475458</b> |
| <b><i>H. zingiberacearum</i></b>        | <b>HFJAU10334</b>  | <b>PP460785</b>          | <b>PP460778</b> | <b>PP460770</b> | <b>PP475459</b> |
| <i>Hysterium angustatum</i>             | MFLU 16-1179       | KX611363                 | KX611364        | KX611365        | –               |
| <i>Marjia tianschanica</i>              | TASM 6120          | MG828909                 | MG829019        | MG829126        | MG829206        |
| <i>M. tianschanica</i>                  | TASM 6121          | MG828910                 | MG829020        | MG829127        | MG829206        |
| <i>Marjia uzbekistanica</i>             | TASM 6122          | MG828911                 | MG829021        | MG829128        | MG829208        |
| <i>Melanocucurbitaria uzbekistanica</i> | MFLUCC 17-0829     | MG828912                 | MG829022        | MG829129        | MG829209        |
| <i>Melanodiplodia tianschanica</i>      | TASM 6111          | MG828914                 | MG829023        | MG829130        | MG829210        |
| <i>Me. tianschanica</i>                 | TASM 6112          | MG828915                 | MG829024        | MG829131        | MG829211        |
| <i>Me. tianschanica</i>                 | MFLUCC 17-0805     | MG828913                 | MG829025        | MG829132        | MG829212        |
| <i>Melanomma japonicum</i>              | KT 3425            | LC203320                 | LC203338        | LC203292        | LC203367        |
| <i>Mel. populicola</i>                  | CBS 543.70         | NR_170056                | NG_075164       | NG_070237       | –               |
| <i>Mel. populicola</i>                  | CPC 27203          | MT223817                 | MT223910        | –               | –               |
| <i>Mel. populicola</i>                  | CBS 350.82         | MT223815                 | JF740265        | –               | –               |
| <i>Mel. populicola</i>                  | CBS 130330         | –                        | JF740328        | –               | –               |
| <b><i>Mel. populicola</i></b>           | <b>HFJAU10335</b>  | <b>PP460786</b>          | <b>PP460779</b> | <b>PP460771</b> | <b>PP475460</b> |
| <i>Mel. pulvis pyrius</i>               | KT 2110            | LC203322                 | LC203340        | LC203294        | LC203368        |
| <i>Mel. pulvis pyrius</i>               | KT 2113            | LC203323                 | LC203341        | LC203295        | LC203369        |
| <i>Mel. pulvis pyrius</i>               | AH 375             | LC203324                 | LC203342        | LC203296        | LC203370        |
| <i>Mel. pulvis pyrius</i>               | KH 27              | LC203325                 | LC203343        | LC203297        | LC203371        |
| <i>Mel. pulvis-pyrius</i>               | CBS 124080         | MH863349                 | GU456323        | GU456302        | GU456265        |
| <i>Monoseptella rosae</i>               | MFLUCC 17-0815     | MG828916                 | MG829026        | MG829133        | MG829213        |
| <i>Mo. rosae</i>                        | TASM 6114          | MG828917                 | MG829027        | MG829134        | MG829214        |
| <i>Mo. tuberculata</i>                  | CBS 256.84         | –                        | GU301851        | –               | GU349006        |
| <i>Muriformistrickeria rubi</i>         | MFLUCC 15-0681     | –                        | KT934253        | KT934257        | KT934261        |
| <i>Petrakia irregularis</i>             | CBS 306.67         | NR_164281                | MH870670        | –               | –               |
| <i>Phragmocephala atra</i>              | MFLUCC 15-0021     | KP698721                 | KP698725        | KP698729        | –               |
| <i>P. garethjonesii</i>                 | MFLUCC 15-0018     | KP698722                 | KP698726        | KP698730        | –               |
| <i>Pleotrichocladium opacum</i>         | CBS 450.70         | MH859791                 | KY853524        | –               | –               |
| <i>Pl. opacum</i>                       | CBS 709.92         | KY853464                 | KY853526        | –               | –               |
| <i>Pseudostrickeria rosae</i>           | MFLUCC 17-0643     | MG828954                 | MG829065        | MG829169        | MG829234        |
| <i>Ps. mutabilis</i>                    | SMH 1541           | –                        | GU385209        | –               | –               |
| <i>Sarimanas pseudofluviatile</i>       | KT 760             | LC001717                 | LC001714        | LC001711        | –               |
| <i>S. shirakamiense</i>                 | KT 3000            | –                        | LC001715        | LC001712        | –               |
| <i>Seifertia azaleae</i>                | DAOM 239136        | –                        | EU030276        | –               | –               |
| <i>Se. shangrilaensis</i>               | MFLUCC 16-0238     | –                        | KU954100        | KU954101        | KU954102        |
| <i>Uzbekistanica rosae-hissaricae</i>   | MFLUCC 17-0819     | MG828976                 | MG829087        | MG829187        | MG829242        |
| <i>U. rosae-hissaricae</i>              | MFLUCC 17-0820     | GU269840                 | MG829088        | MG829188        | MG829243        |
| <i>Xenostigmia zilleri</i>              | CBS 115685         | GU269840                 | GU253857        | LC203316        | GU384553        |

Abbreviations: ANM, A.N. Miller; AFTOL, Assembling the Fungal Tree of Life project; CBS, Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CPC, Working collection of Pedro Crous housed at CBS; DAOM, Plant Research Institute, Department of Agriculture (Mycology), Ottawa, Canada; GLMC, culture collections of the Senckenberg Museum of Natural History Görlitz, Germany; GKM, G.K. Mugambi; IL, I. Lopez; KUMCC, Culture Collection of Chinese Academy of Sciences, Kunming, China; KH, K. Hirayama; KT, K. Tanaka; MFLU, Mae Fah Luang University; MFLUCC, Mae Fah Luang University Culture Collection; NCYUCC, National Chiayi University Culture Collection, China; SMH, S.M. Huhndorf; ZHKUCC, Zhongkai University of Agriculture and Engineering Culture Collection, China.

were discarded as burn-in. The phylograms were visualised using the FigTree v.1.4.0 tool (Rambaut 2012) and reorganised in Adobe Illustrator® CS3 (Version 15.0.0, Adobe®, San Jose, CA). The alignments and sequences were deposited in TreeBASE, submission ID: 31212 (<http://www.treebase.org/>) and GenBank (<https://www.ncbi.nlm.nih.gov/>), respectively.

## Geographical distribution and host associations of melanommataceous species

We enumerated 340 species of Melanommataceae, grouped into 36 genera, along with their geographic distribution and host associations. The necessary data were obtained from published books, publications in reputable journals, Species Fungorum (<https://www.speciesfungorum.org>), MycoBank (<https://www.mycobank.org/>), the U.S. National Fungus Collections Fungus-Host Database (Farr and Rossman 2024) and online sources (Catalogue of Life Checklist). The gathered data were mentioned with appropriate references (Table 2). In total, 296 articles were studied for information on the 340 melanommataceous species that are currently legitimate (Table 2) and accessed through Google Scholar searches. MycoBank (<https://www.mycobank.org/>) was used to illustrate the nomenclature validity of the taxa. However, numerous species have not been verified using molecular data, so most species should be correctly identified, based on modern taxonomic concepts. Therefore, the distribution may vary slightly.

The distribution map was prepared using MapChart programme (<https://www.mapchart.net/index.html>), a platform from which a personalised map of the world using different colours can be generated. The Sankey diagram was created to show the species distribution through plant host families using the free online tool SankeyMATIC by Steve Bogart ([www.sankeymatic.com](http://www.sankeymatic.com)).

## Results

### Phylogenetic analyses

Phylogenetic analyses of combined LSU, SSU, ITS and *tef1-α* sequences comprised 3480 characters including gaps. *Hysterium angustatum* (MFLU 16-1179) was used as the outgroup taxon. The RAxML analysis of the combined dataset yielded a best scoring tree (Fig. 3). The final ML optimisation likelihood value was -16620.192034. There were 35.32% undetermined characters or gaps and 1071 distinct alignment patterns. Estimated base frequencies were A = 0.247534, C = 0.235205, G = 0.272353, T = 0.244907; substitution rates AC = 1.775315, AG = 3.161447, AT = 1.766474, CG = 1.002405, CT = 11.511066, GT = 1.000; proportion of invariable sites I = 0.535605; gamma distribution shape parameter  $\alpha$  = 0.513218. The Bayesian analysis has resulted in 30,000 trees after 3,000,000 generations. Bootstrap support values for ML higher than 70% and BYPP greater than 0.90 are given above each branch respectively (Fig. 3). All analyses (ML and BYPP) showed similar topologies and concurred with previous studies (Pem et al. 2019; Kularathnage et al. 2022; Gao et al. 2023).

According to the phylogeny, our collection HFJAU10336, HFJAU10337 and HFJAU10338 cluster within *Byssosphaeria* species. HFJAU10337 and




**Figure 3.** Phylogram generated from Maximum Likelihood analysis is based on combined LSU, SSU, ITS and *tef1-α* sequence data. The tree is rooted with *Hysterium angustatum* (MFLU 16-1179). The new isolates are in red and ex-type strains are indicated in bold face. Bootstrap support values  $\geq 70\%$  from the Maximum Likelihood (ML) and Bayesian Posterior Probabilities (BYPP) values  $\geq 0.90$  are given above the nodes, respectively.




Figure 3. Continued.

HFJAU10338 cluster together and show an independent lineage, sister to *B. phoenicis* isolates (ZHKUCC 21-0122 and ZHKUCC 21-0123) with 83% ML and 0.99 BYPP statistical support. The isolate HFJAU10336 clusters with *B. siamensis* isolates (MFLUCC 10-0099, MFLUCC 17-1800 and KUMCC 21-0339) in a monophyletic clade. In addition, CBUB 40045 groups with *Bertiella fici* in a 100% ML and 1.00 BYPP supported clade. Three isolates (HFJAU10332, HFJAU10333 and HFJAU10334) cluster with *Herpotrichia* species, and show a close phylogenetic relationship with *H. melanotricha* (GKM 196N, SMH 269 and JCM 14419). As well as HFJAU10335 groups with *Melanomma populicola* (CBS 130330, CBS 350.82, CBS 543.70 and CPC 272203) in a 99% ML and 0.99 BYPP supported clade.

**Table 2.** Host association and geographical distribution of reported melanommataceous species.

| Species                            | Host                                                 | Host family             | Locality                    | References                                                         |
|------------------------------------|------------------------------------------------------|-------------------------|-----------------------------|--------------------------------------------------------------------|
| <i>Alpinaria rhododendri</i>       | <i>Rhododendron</i> sp.                              | Ericaceae               | Austria and Japan           | Hashimoto et al. (2017); Jaklitsch and Voglmayr (2017)             |
| <i>Aposphaeria allantella</i>      | <i>Solanum tuberosum</i>                             | Solanaceae              | Germany                     | Farr and Rossman (2024)                                            |
| <i>Aposphaeria anomala</i>         | Unidentified host                                    | –                       | Italy                       | Farr and Rossman (2024)                                            |
| <i>Aposphaeria arachidis</i>       | <i>Arachis hypogaea</i>                              | Fabaceae                | India                       | Kulkarni (1974); Mathur (1979); Farr and Rossman (2024)            |
| <i>Aposphaeria bambusae</i>        | <i>Bambusa</i> sp.                                   | Poaceae                 | Brazil                      | Bresadola (1926); Farr and Rossman (2024)                          |
| <i>Aposphaeria bombacis</i>        | <i>Bombax macrocarpum</i>                            | Malvaceae               | Germany                     | Diedicke (1912)                                                    |
| <i>Aposphaeria brunneotincta</i>   | <i>Castanea vesca</i>                                | Fagaceae                | The United States           | Thaxter (1922)                                                     |
| <i>Aposphaeria buddlejae</i>       | <i>Buddleja davidii</i>                              | Buddlejaceae            | Ukraine                     | Farr and Rossman (2024)                                            |
| <i>Aposphaeria calligoni</i>       | <i>Calligonum aphyllum</i>                           | Polygonaceae            | Kazakhstan                  | Farr and Rossman (2024)                                            |
| <i>Aposphaeria canavaliae</i>      | <i>Canavalia</i> sp.                                 | Fabaceae                | Fiji                        | Massee (1906); Dingley et al. (1981); Farr and Rossman (2024)      |
| <i>Aposphaeria caraganae</i>       | <i>Caragana arborescens</i>                          | Fabaceae                | Russia                      | Farr and Rossman (2024)                                            |
| <i>Aposphaeria caricicola</i>      | <i>Carex rossii</i>                                  | Cyperaceae              | The United States           | Farr and Rossman (2024)                                            |
| <i>Aposphaeria caulina</i>         | <i>Cerefolium sylvestre</i>                          | Apiaceae                | Finland                     | Karsten (1905)                                                     |
| <i>Aposphaeria charticola</i>      | –                                                    | –                       | The United States           | Saccardo (1911)                                                    |
| <i>Aposphaeria cladoniae</i>       | <i>Cladonia fimbriata</i>                            | Cladoniaceae            | Germany                     | Allescher (1896)                                                   |
| <i>Aposphaeria conica</i>          | <i>Quercus</i> sp.                                   | Fagaceae                | Italy                       | Saccardo and Traverso (1910)                                       |
| <i>Aposphaeria corallinolutea</i>  | <i>Fraxinus excelsior</i> and <i>Kerria japonica</i> | Oleaceae and Rosaceae   | The Netherlands             | De Gruyter et al. (2013)                                           |
| <i>Aposphaeria dendrophomoides</i> | <i>Corylus avellana</i>                              | Betulaceae              | Italy                       | Saccardo (1921)                                                    |
| <i>Aposphaeria denudata</i>        | <i>Cydonia vulgaris</i>                              | Rosaceae                | Hungary                     | Saccardo and Traverso (1910)                                       |
| <i>Aposphaeria desertorum</i>      | <i>Haloxylon aphyllum</i>                            | Amaranthaceae           | Kazakhstan                  | Farr and Rossman (2024)                                            |
| <i>Aposphaeria elymi</i>           | <i>Elymus arenarius</i>                              | Poaceae                 | Germany                     | Diedicke (1912)                                                    |
| <i>Aposphaeria ephedrae</i>        | <i>Ephedra</i> sp.                                   | Ephedraceae             | Kazakhstan and Ukraine      | Hayova and Minter (2009); Farr and Rossman (2024)                  |
| <i>Aposphaeria epicorticalis</i>   | <i>Corylus avellana</i>                              | Betulaceae              | Italy                       | Saccardo (1921)                                                    |
| <i>Aposphaeria eragrostidis</i>    | <i>Eragrostis</i> sp.                                | Poaceae                 | Eritrea, Ethiopia and Iraq  | Castellani and Ciferri (1950); Farr and Rossman (2024)             |
| <i>Aposphaeria eurotiae</i>        | <i>Eurotia eversmanniana</i>                         | Amaranthaceae           | Kazakhstan                  | Farr and Rossman (2024)                                            |
| <i>Aposphaeria ferrum-equinum</i>  | Unidentified host                                    | –                       | Italy                       | Tassi (1900)                                                       |
| <i>Aposphaeria freticola</i>       | <i>Fagus</i> sp. and <i>Nothofagus antarctica</i>    | Fagaceae, Nothofagaceae | Argentina, Chile and Poland | Farr (1973); Mulencko et al. (2008); Farr and Rossman (2024)       |
| <i>Aposphaeria gallicola</i>       | Unidentified host                                    | –                       | Italy                       | Farr and Rossman (2024)                                            |
| <i>Aposphaeria gregaria</i>        | <i>Salix</i> sp.                                     | Salicaceae              | Germany                     | Diedicke (1912)                                                    |
| <i>Aposphaeria halimodendri</i>    | <i>Halimodendron halodendron</i>                     | Fabaceae                | Ukraine                     | Farr and Rossman (2024)                                            |
| <i>Aposphaeria haloxyli</i>        | <i>Haloxylon aphyllum</i>                            | Amaranthaceae           | Kazakhstan                  | Farr and Rossman (2024)                                            |
| <i>Aposphaeria hapalophragmii</i>  | <i>Hapalophragmium acaciae</i>                       | Fabaceae                | Somalia                     | Trotter (1931); Mujica and Vergara (1945); Farr and Rossman (2024) |

| Species                             | Host                                        | Host family             | Locality                                                     | References                                                                         |
|-------------------------------------|---------------------------------------------|-------------------------|--------------------------------------------------------------|------------------------------------------------------------------------------------|
| <i>Aposphaeria henryana</i>         | <i>Salix alba</i>                           | Salicaceae              | Italy                                                        | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria heveae</i>           | <i>Hevea brasiliensis</i>                   | Euphorbiaceae           | Sri Lanka                                                    | Petch (1917)                                                                       |
| <i>Aposphaeria hippuridis</i>       | <i>Hippuris vulgaris</i>                    | Plantaginaceae          | Germany                                                      | Ade (1923)                                                                         |
| <i>Aposphaeria hospitae</i>         | <i>Kleinhovia</i> sp.                       | Malvaceae               | Sri Lanka                                                    | Tassi (1900)                                                                       |
| <i>Aposphaeria humicola</i>         | Unidentified host                           | –                       | The Netherlands                                              | Oudemans (1902)                                                                    |
| <i>Aposphaeria ilicis</i>           | <i>Ilex aquifolium</i>                      | Aquifoliaceae           | Germany                                                      | Diedicke (1912)                                                                    |
| <i>Aposphaeria iliensis</i>         | <i>Halimodendron halodendron</i>            | Fabaceae                | Kazakhstan                                                   | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria jubaeae</i>          | <i>Jubaea spectabilis</i>                   | Arecaceae               | Chile                                                        | Spegazzini (1921); Farr (1973); Mujica and Vergara (1945); Farr and Rossman (2024) |
| <i>Aposphaeria kiefferiana</i>      | <i>Quercus</i> sp.                          | Fagaceae                | Italy                                                        | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria kravtzevii</i>       | <i>Eurotia eversmanniana</i>                | Amaranthaceae           | Kazakhstan                                                   | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria lentisci</i>         | <i>Pistacia</i> sp.                         | Anacardiaceae           | Greece and Spain                                             | Urries (1957); Pantidou (1973); Farr and Rossman (2024)                            |
| <i>Aposphaeria lignicola</i>        | <i>Acacia arabica</i>                       | Fabaceae                | Pakistan                                                     | Ahmad (1964)                                                                       |
| <i>Aposphaeria major</i>            | <i>Rubus parviflorus</i>                    | Rosaceae                | The United States                                            | Sydow and Sydow (1907)                                                             |
| <i>Aposphaeria majuscula</i>        | <i>Vitis vinifera</i>                       | Vitaceae                | France                                                       | Saccardo and Trotter (1913)                                                        |
| <i>Aposphaeria martinii</i>         | <i>Sabal</i> sp.                            | Arecaceae               | United States                                                | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria mediella</i>         | <i>Pinus</i> sp.                            | Pinaceae                | Greece, Finland and Poland                                   | Karsten (1884); Pantidou (1973); Mullenko et al. (2008); Farr and Rossman (2024)   |
| <i>Aposphaeria melaleucae</i>       | <i>Melaleuca leucadendra</i>                | Myrtaceae               | Australia                                                    | Hennings (1903); Shivas and Alcorn (1996); Farr and Rossman (2024)                 |
| <i>Aposphaeria mesembryanthemi</i>  | <i>Mesembryanthemum</i> sp.                 | Aizoaceae               | Portugal                                                     | Costa and Camara (1952)                                                            |
| <i>Aposphaeria mojunkumica</i>      | <i>Haloxylon aphyllum</i>                   | Amaranthaceae           | Kazakhstan                                                   | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria montbretiae</i>      | <i>Crocasmia</i> sp.                        | Iridaceae               | Azerbaijan and Georgia                                       | Siemaszko (1923)                                                                   |
| <i>Aposphaeria musarum</i>          | <i>Musa sapientum</i>                       | Musaceae                | Argentina                                                    | Farr (1973); Farr and Rossman (2024)                                               |
| <i>Aposphaeria nigra</i>            | <i>Betula alba</i>                          | Betulaceae              | Germany                                                      | Diedicke (1912)                                                                    |
| <i>Aposphaeria oxalidis</i>         | <i>Oxalis tuberosa</i>                      | Oxalidaceae             | Bolivia                                                      | Farr and Stevenson (1963); Farr and Rossman (2024)                                 |
| <i>Aposphaeria pakistanica</i>      | Unidentified host                           | –                       | Pakistan                                                     | Ahmad (1971)                                                                       |
| <i>Aposphaeria phellodendri</i>     | <i>Phellodendron amurense</i>               | Rutaceae                | Ukraine                                                      | Farr and Rossman (2024)                                                            |
| <i>Aposphaeria pinea</i>            | <i>Pinus sylvestris</i>                     | Pinaceae                | France and Germany                                           | Saccardo (1884); Mullenko et al. (2008); Farr and Rossman (2024)                   |
| <i>Aposphaeria pini-densiflorae</i> | <i>Pinus densiflora</i>                     | Pinaceae                | Japan                                                        | Sawada (1950); Farr and Rossman (2024)                                             |
| <i>Aposphaeria polonica</i>         | <i>Tilia platyphyllos</i>                   | Pinaceae                | Poland                                                       | Mullenko et al. (2008); Farr and Rossman (2024)                                    |
| <i>Aposphaeria populea</i>          | <i>Populus</i> sp.                          | Salicaceae              | The United Kingdom                                           | Smith and Ramsbottom (1914); Dennis (1986)                                         |
| <i>Aposphaeria pulviscula</i>       | <i>Fagus sylvatica</i> and <i>Salix</i> sp. | Fagaceae and Salicaceae | Austria, France, Iceland, Italy, the Netherlands and Ukraine | Saccardo (1880); Spaulding (1961); Sutton (1980); Farr and Rossman (2024)          |

| Species                            | Host                                                             | Host family                                | Locality                  | References                                                           |
|------------------------------------|------------------------------------------------------------------|--------------------------------------------|---------------------------|----------------------------------------------------------------------|
| <i>Aposphaeria punicina</i>        | <i>Punica granatum</i>                                           | Lythraceae                                 | China and Malta           | Saccardo (1915); Tai (1979)                                          |
| <i>Aposphaeria purpurascens</i>    | <i>Acer pseudoplatanus</i>                                       | Sapindaceae                                | Italy                     | Peyronel (1915)                                                      |
| <i>Aposphaeria ramalinae</i>       | <i>Ramalina</i> sp. (lichen)                                     | –                                          | France                    | Pitard and Harmand (1911)                                            |
| <i>Aposphaeria reaumuriae</i>      | <i>Reaumuria</i> sp.                                             | Tamaricaceae                               | Azerbaijan                | Farr and Rossman (2024)                                              |
| <i>Aposphaeria rhois</i>           | <i>Rhus oxyacantha</i>                                           | Anacardiaceae                              | Libya                     | El-Buni and Rattan (1981); Trotter (1912)                            |
| <i>Aposphaeria rostrata</i>        | Unidentified host                                                | –                                          | The Netherlands           | Oudemans (1902)                                                      |
| <i>Aposphaeria rubefaciens</i>     | <i>Salix</i> sp.                                                 | Salicaceae                                 | Italy and the Netherlands | Bubák and Kabát (1905)                                               |
| <i>Aposphaeria rudis</i>           | <i>Picea excelsa</i>                                             | Pinaceae                                   | Finland                   | Karsten (1905)                                                       |
| <i>Aposphaeria salicis</i>         | <i>Salix</i> sp.                                                 | Salicaceae                                 | Germany and India         | Diedicke (1912); Mathur (1979)                                       |
| <i>Aposphaeria salicum</i>         | <i>Salix viminalis</i>                                           | Salicaceae                                 | Germany                   | Sydow and Sydow (1903)                                               |
| <i>Aposphaeria santolinae</i>      | <i>Santolina chamaecyparissus</i>                                | Asteraceae                                 | Ukraine                   | Farr and Rossman (2024)                                              |
| <i>Aposphaeria sepulta</i>         | <i>Citrus aurantium</i>                                          | Rutaceae                                   | Italy                     | Saccardo (1884)                                                      |
| <i>Aposphaeria sequoiae</i>        | <i>Sequoia</i> sp.                                               | Cupressaceae                               | Denmark                   | Lind (1913)                                                          |
| <i>Aposphaeria silenes</i>         | <i>Silene otites</i>                                             | Caryophyllaceae                            | Russia                    | Farr and Rossman (2024)                                              |
| <i>Aposphaeria sphaerospora</i>    | <i>Betula alba</i>                                               | Betulaceae                                 | Italy                     | Peyronel (1918)                                                      |
| <i>Aposphaeria striolata</i>       | <i>Populus deltoides</i>                                         | Salicaceae                                 | The United States         | Saccardo (1916)                                                      |
| <i>Aposphaeria taquarae</i>        | Bambusoideae sp.                                                 | Poaceae                                    | Brazil                    | Viégas (1945)                                                        |
| <i>Aposphaeria tiliana</i>         | <i>Tilia cordata</i>                                             | Malvaceae                                  | Ukraine                   | Gucevic (1977)                                                       |
| <i>Aposphaeria tragopogonis</i>    | <i>Tragopogon dubius</i>                                         | Asteraceae                                 | Romania                   | Sandu-Ville and Mititiuc (1971)                                      |
| <i>Aposphaeria turmalis</i>        | <i>Diospyros virginiana</i>                                      | Ebenaceae                                  | The United States         | Ellis and Everhart (1902); Cash (1952)                               |
| <i>Aposphaeria ulmicola</i>        | <i>Ulmus</i> sp.                                                 | Ulmaceae                                   | The United Kingdom        | Saccardo (1884)                                                      |
| <i>Aposphaeria zaeae</i>           | <i>Zea mays</i>                                                  | Poaceae                                    | Azerbaijan and Georgia    | Farr and Rossman (2024)                                              |
| <i>Asymmetricospora calamicola</i> | <i>Calamus caryotoides</i>                                       | Arecaceae                                  | Australia                 | Fröhlich and Hyde (1998); Zhang et al. (2012)                        |
| <i>Bertiella botryosa</i>          | <i>Ulmus</i> sp.                                                 | Ulmaceae                                   | The United States         | Morgan (1904)                                                        |
| <i>Bertiella ellipsoidea</i>       | Unidentified host                                                | –                                          | Thailand                  | Hyde et al. (2016)                                                   |
| <i>Bertiella fici</i>              | <i>Cinnamomum verum</i> ,<br><i>Ficus septica</i>                | Lauraceae,<br>Moraceae                     | China and Thailand        | Tennakoon et al. (2021); this study                                  |
| <i>Bertiella gelatinosa</i>        | Unidentified host                                                | –                                          | Brazil                    | Almeida et al. (2017)                                                |
| <i>Bertiella rhodospila</i>        | <i>Cyrilla</i> sp., <i>Populus</i> sp.<br>and <i>Quercus</i> sp. | Cyrillaceae,<br>Fagaceae and<br>Salicaceae | The United States         | Barr et al. (1986)                                                   |
| <i>Bertiella striatispora</i>      | Unidentified host                                                | –                                          | India                     | Niranjan and Sarma (2019)                                            |
| <i>Bicrouania maritima</i>         | <i>Atraphaxis spinosa</i><br>and <i>Halimione portulacoides</i>  | Amaranthaceae,<br>Polygonaceae             | France and Uzbekistan     | Kohlmeyer and Volkmann-Kohlmeyer (1990); Gafforov (2017)             |
| <i>Byssosphaeria alnea</i>         | <i>Alnus</i> sp.                                                 | Betulaceae                                 | The United States         | Barr (1984); Farr and Rossman (2024)                                 |
| <i>Byssosphaeria erumpens</i>      | <i>Litsea</i> sp.                                                | Lauraceae                                  | China                     | Chen and Hsieh (2004); Li and Zhuang (2008); Farr and Rossman (2024) |

| Species                               | Host                                                                                                                                                                                                                                                                                                                                                                                                                 | Host family                                                                 | Locality                                            | References                                                                                            |
|---------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------|
| <i>Byssosphaeria erythrinae</i>       | <i>Erythrina indica</i>                                                                                                                                                                                                                                                                                                                                                                                              | Fabaceae                                                                    | France                                              | Barr (1984)                                                                                           |
| <i>Byssosphaeria guangdongense</i>    | <i>Phoenix roebelenii</i>                                                                                                                                                                                                                                                                                                                                                                                            | Arecaceae                                                                   | China                                               | Xiong et al. (2023)                                                                                   |
| <i>Byssosphaeria hainanensis</i>      | Unidentified host                                                                                                                                                                                                                                                                                                                                                                                                    | –                                                                           | China                                               | Li and Zhuang (2008)                                                                                  |
| <i>Byssosphaeria jamaicana</i>        | <i>Bambusa</i> sp., <i>Quercus</i> sp. and <i>Salix</i> sp.                                                                                                                                                                                                                                                                                                                                                          | Fagaceae and Poaceae, Salicaceae                                            | China, Czech Republic, Jamaica and Mexico           | Barr (1984); Sivanesan and Hsieh (1989); Wang et al. (2004); Medel (2007); Farr and Rossman (2024)    |
| <i>Byssosphaeria juniperi</i>         | <i>Juniperus</i> sp.                                                                                                                                                                                                                                                                                                                                                                                                 | Cupressaceae                                                                | The United States                                   | Wang et al. (2004); Farr and Rossman (2024)                                                           |
| <i>Byssosphaeria macarangae</i>       | <i>Macaranga tanarius</i>                                                                                                                                                                                                                                                                                                                                                                                            | Euphorbiaceae                                                               | China                                               | Tennakoon et al. (2018)                                                                               |
| <i>Byssosphaeria musae</i>            | <i>Musa</i> sp.                                                                                                                                                                                                                                                                                                                                                                                                      | Musaceae                                                                    | Thailand                                            | Liu et al. (2015); Farr and Rossman (2024)                                                            |
| <i>Byssosphaeria oviformis</i>        | <i>Saccharum arundinaceum</i>                                                                                                                                                                                                                                                                                                                                                                                        | Poaceae                                                                     | China and Jamaica                                   | Barr (1984); Lu et al. (2000); Wong and Hyde (2001); Farr and Rossman (2024)                          |
| <i>Byssosphaeria phoenicis</i>        | <i>Phoenix roebelenii</i>                                                                                                                                                                                                                                                                                                                                                                                            | Arecaceae                                                                   | China                                               | Kularathnage et al. (2022)                                                                            |
| <i>Byssosphaeria poaeicola</i>        | <i>Arundo pliniana</i>                                                                                                                                                                                                                                                                                                                                                                                               | Poaceae                                                                     | China                                               | This study                                                                                            |
| <i>Byssosphaeria rhodomphala</i>      | <i>Acer pseudoplatanus</i> and <i>Populus</i> sp.                                                                                                                                                                                                                                                                                                                                                                    | Salicaceae and Sapindaceae                                                  | Brazil, China, Poland and the United States         | Cooke (1887); Scheuer and Chlebicki (1997); Reblova (1997); Chen and Hsieh (2004); Wang et al. (2004) |
| <i>Byssosphaeria salebrosa</i>        | <i>Vaccinium</i> sp.                                                                                                                                                                                                                                                                                                                                                                                                 | Ericaceae                                                                   | The United States                                   | Barr (1984); Barr et al. (1986); Farr and Rossman (2024)                                              |
| <i>Byssosphaeria schiedermayriana</i> | <i>Sambucus nigra</i>                                                                                                                                                                                                                                                                                                                                                                                                | Adoxaceae                                                                   | Austria                                             | Barr (1984)                                                                                           |
| <i>Byssosphaeria semen</i>            | <i>Pyrus americana</i>                                                                                                                                                                                                                                                                                                                                                                                               | Rosaceae                                                                    | The United States                                   | Barr (1984)                                                                                           |
| <i>Byssosphaeria siamensis</i>        | <i>Pandanus</i> sp. and <i>Citrus trifoliata</i>                                                                                                                                                                                                                                                                                                                                                                     | Pandanaceae and Rutaceae                                                    | China and Thailand                                  | Tian et al. (2015); Hyde et al. (2018); Farr and Rossman (2024); this study                           |
| <i>Byssosphaeria taiwanense</i>       | <i>Macaranga tanarius</i>                                                                                                                                                                                                                                                                                                                                                                                            | Euphorbiaceae                                                               | China                                               | Tennakoon et al. (2018)                                                                               |
| <i>Byssosphaeria xestothele</i>       | <i>Cornus florida</i> and <i>Robinia pseudoacacia</i>                                                                                                                                                                                                                                                                                                                                                                | Cornaceae and Fabaceae                                                      | Sweden and the United States                        | Barr (1984); Eriksson (2014); Farr and Rossman (2024)                                                 |
| <i>Calyptronectria argentinensis</i>  | <i>Foeniculum piperitum</i> and <i>Manihot carthaginensis</i>                                                                                                                                                                                                                                                                                                                                                        | Apiaceae, Euphorbiaceae                                                     | Argentina                                           | Spegazzini (1909); Farr (1973); Farr and Rossman (2024)                                               |
| <i>Calyptronectria indica</i>         | <i>Annona squamosa</i>                                                                                                                                                                                                                                                                                                                                                                                               | Annonaceae                                                                  | India                                               | Pande (2008)                                                                                          |
| <i>Calyptronectria platensis</i>      | <i>Manihot carthagenensis</i>                                                                                                                                                                                                                                                                                                                                                                                        | Euphorbiaceae                                                               | Argentina                                           | Spegazzini (1909); Farr (1973); Farr and Rossman (2024)                                               |
| <i>Camposporium antennatum</i>        | <i>Acacia aulacocarpa</i> , <i>Caesalpinia echinata</i> , <i>Cinnamomum japonicum</i> , <i>Cocos nucifera</i> , <i>Drymophloeus pachycladus</i> , <i>Eucalyptus globulus</i> , <i>Ficus erecta</i> , <i>Laurus nobilis</i> , <i>Machilus</i> sp., <i>Mucuna ferruginea</i> , <i>Neolitsea scrobiculata</i> , <i>Phoenix hanceana</i> , <i>Pinus massoniana</i> , <i>Quercus</i> sp. and <i>Trachycarpus fortunei</i> | Arecaceae, Fabaceae, Fagaceae, Lauraceae, Moraceae, Myrtaceae, and Pinaceae | California, China, the United Kingdom and Venezuela | Harkness (1884); Lu et al. (2000); Castaneda-Ruiz et al. (2003); Farr and Rossman (2024)              |
| <i>Camposporium appendiculatum</i>    | Unidentified host                                                                                                                                                                                                                                                                                                                                                                                                    | –                                                                           | China                                               | Hyde et al. (2020)                                                                                    |

| Species                            | Host                                                                                                                                                                                                                                            | Host family                                                                                    | Locality                                                          | References                                                                            |
|------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|-------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| <i>Camposporium atypicum</i>       | <i>Mesua ferrea</i>                                                                                                                                                                                                                             | Calophyllaceae                                                                                 | India                                                             | Koukol and Delgado (2021)                                                             |
| <i>Camposporium cambrense</i>      | <i>Alnus</i> sp., <i>Carpinus betulus</i> , <i>Fagus</i> sp., <i>Freycinetia anksia</i> , <i>Laurus nobilis</i> and <i>Quercus</i> sp.                                                                                                          | Betulaceae, Fagaceae, Lauraceae, and Pandanaceae                                               | China, New Zealand, Poland, Russia and the United Kingdom         | Hughes (1951); Whitton et al. (2002); Mullenko et al. (2008); Farr and Rossman (2024) |
| <i>Camposporium chinense</i>       | Unidentified host                                                                                                                                                                                                                               | –                                                                                              | China                                                             | Xu et al. (2021)                                                                      |
| <i>Camposporium dulciaquae</i>     | Unidentified host                                                                                                                                                                                                                               | –                                                                                              | Thailand                                                          | Calabon et al. (2021)                                                                 |
| <i>Camposporium fusisporum</i>     | <i>Pandanus</i> sp.                                                                                                                                                                                                                             | Pandanaceae                                                                                    | Brunei                                                            | Whitton et al. (2002)                                                                 |
| <i>Camposporium himalayanum</i>    | <i>Phoenix</i> sp.                                                                                                                                                                                                                              | Arecaceae                                                                                      | India                                                             | Adamcik et al. (2015)                                                                 |
| <i>Camposporium hyalinum</i>       | <i>Elaeagnus macrophylla</i> , and <i>Fagus</i> sp.                                                                                                                                                                                             | Elaeagnaceae and Fagaceae                                                                      | The United Kingdom                                                | Abdullah (1980); Kirk and Spooner (1984)                                              |
| <i>Camposporium hyderabadense</i>  | <i>Borassus flabellifer</i> , <i>Machilus thunbergii</i> and <i>Mucuna ferruginea</i>                                                                                                                                                           | Arecaceae, Fabaceae, and Lauraceae                                                             | China, India and Japan                                            | Rao and Rao (1964); Ichinoe (1971); Farr and Rossman (2024)                           |
| <i>Camposporium indicum</i>        | <i>Borassus flabellifer</i>                                                                                                                                                                                                                     | Arecaceae                                                                                      | India                                                             | Rao and Rao (1964)                                                                    |
| <i>Camposporium japonicum</i>      | <i>Acacia confusa</i> , <i>Betula pendula</i> , <i>Cinnamomum japonicum</i> , <i>Freycinetia arborea</i> , <i>Litchi chinensis</i> , <i>Machilus thunbergii</i> , <i>Mucuna ferruginea</i> , <i>Paulownia kawakamii</i> and <i>Quercus</i> sp.  | Betulaceae, Lauraceae, Pandanaceae, Paulowniaceae, and Sapindaceae                             | China and Japan                                                   | Ichinoe (1971); Matsushima (1980); Farr and Rossman (2024)                            |
| <i>Camposporium laundonii</i>      | <i>Aralia elata</i> , <i>Pasania edulis</i> and <i>Rosa</i> sp.                                                                                                                                                                                 | Araliaceae, Fagaceae, and Rosaceae                                                             | Japan and New Zealand                                             | Ellis (1976); Watanabe (1993); Farr and Rossman (2024)                                |
| <i>Camposporium lycopodiellae</i>  | <i>Lycopodiella inundata</i>                                                                                                                                                                                                                    | Lycopodiaceae                                                                                  | Germany                                                           | Hyde et al. (2020)                                                                    |
| <i>Camposporium marylandicum</i>   | Unidentified host                                                                                                                                                                                                                               | –                                                                                              | The United States                                                 | Shearer (1974)                                                                        |
| <i>Camposporium microsporum</i>    | <i>Borassus flabellifer</i>                                                                                                                                                                                                                     | Arecaceae                                                                                      | India                                                             | Rao and Rao (1964)                                                                    |
| <i>Camposporium multiseptatum</i>  | Unidentified host                                                                                                                                                                                                                               | –                                                                                              | China                                                             | Hyde et al. (2020)                                                                    |
| <i>Camposporium ontariense</i>     | <i>Acer saccharum</i> and <i>Freycinetia arborea</i>                                                                                                                                                                                            | Sapindaceae and Pandanaceae                                                                    | Canada and the United States                                      | Matsushima (1983); Whitton et al. (2002)                                              |
| <i>Camposporium pellucidum</i>     | <i>Betula pendula</i> , <i>Caesalpinia echinata</i> , <i>Carpodetus serratus</i> , <i>Elaeagnus</i> sp., <i>Fagus</i> sp., <i>Laurus</i> sp., <i>Pasania glabra</i> , <i>Picea abies</i> , <i>Rhopalostylis</i> sp. and <i>Sorbus aucuparia</i> | Arecaceae, Betulaceae, Elaeagnaceae, Fabaceae, Lauraceae, Pinaceae, Rosaceae, and Rousseeaceae | Brazil, Japan, Poland, New Zealand, Russia and the United Kingdom | Hughes (1951); Matsushima (1983); Mullenko et al. (2008); Farr and Rossman (2024)     |
| <i>Camposporium quercicola</i>     | <i>Quercus germana</i>                                                                                                                                                                                                                          | Fagaceae                                                                                       | Mexico                                                            | Mercado Sierra et al. (1995)                                                          |
| <i>Camposporium ramosum</i>        | <i>Freycinetia</i> sp.                                                                                                                                                                                                                          |                                                                                                | Australia and the United States                                   | Whitton et al. (2002)                                                                 |
| <i>Camposporium scolecosporium</i> | Unidentified host                                                                                                                                                                                                                               | –                                                                                              | Papua New Guinea                                                  | Kobayasi (1971)                                                                       |
| <i>Camposporium septatum</i>       | Unidentified host                                                                                                                                                                                                                               | –                                                                                              | Thailand                                                          | Hyde et al. (2020)                                                                    |
| <i>Camposporium valdivianum</i>    | <i>Sophora microphylla</i>                                                                                                                                                                                                                      | Fabaceae                                                                                       | The United States                                                 | Koukol and Delgado (2021)                                                             |

| Species                              | Host                                                                                     | Host family                              | Locality                        | References                                                                                           |
|--------------------------------------|------------------------------------------------------------------------------------------|------------------------------------------|---------------------------------|------------------------------------------------------------------------------------------------------|
| <i>Camposporium verruculosum</i>     | <i>Clematis vitalba</i>                                                                  | Ranunculaceae                            | Italy                           | Koukol and Delgado (2021)                                                                            |
| <i>Dematiomelanomma yunnanense</i>   | <i>Hypericum monogynum</i> and <i>Rubus parvifolius</i>                                  | Hypericaceae and Rosaceae                | China                           | Gao et al. (2023)                                                                                    |
| <i>Exosporiella fungorum</i>         | <i>Thelephora</i> sp. (leathery earthfans/ mushroom sp.)                                 | –                                        | Sweden                          | Karsten (1892)                                                                                       |
| <i>Fusiconidium mackenziei</i>       | <i>Clematis vitalba</i>                                                                  | Ranunculaceae                            | Italy                           | Li et al. (2017)                                                                                     |
| <i>Herpotrichia alligata</i>         | <i>Opuntia</i> sp.                                                                       | Cactaceae                                | Sweden and the United States    | Barr (1992); Farr and Rossman (2024)                                                                 |
| <i>Herpotrichia alpincola</i>        | <i>Aconitum</i> sp.                                                                      | Ranunculaceae                            | Hungary and Slovakia            | Rehm (1906)                                                                                          |
| <i>Herpotrichia arizonica</i>        | <i>Carnegiea gigantea</i>                                                                | Cactaceae                                | The United States               | Barr (1992); Farr and Rossman (2024)                                                                 |
| <i>Herpotrichia australis</i>        | <i>Sclerocarya caffra</i>                                                                | Anacardiaceae                            | South Africa                    | Bose (1961)                                                                                          |
| <i>Herpotrichia bakeri</i>           | <i>Sambucus javanica</i>                                                                 | Viburnaceae                              | Philippines                     | Teodoro (1937); Farr and Rossman (2024)                                                              |
| <i>Herpotrichia bambusana</i>        | <i>Bambusa vulgaris</i>                                                                  | Poaceae                                  | Brazil                          | Hennings (1908); Eriksson and Yue (1998)                                                             |
| <i>Herpotrichia boldoae</i>          | <i>Boldea fragrans</i> and <i>Peumus boldus</i>                                          | Monimiaceae                              | Chile                           | Spegazzini (1910); Mujica and Vergara (1945); Spaulding (1961); Farr (1973); Farr and Rossman (2024) |
| <i>Herpotrichia brasiliensis</i>     | Unidentified host                                                                        | –                                        | Brazil                          | Rick (1933)                                                                                          |
| <i>Herpotrichia brenckleana</i>      | <i>Urtica gracilis</i>                                                                   | Urticaceae                               | Sweden                          | Eriksson (2014); Farr and Rossman (2024)                                                             |
| <i>Herpotrichia caesalpiniae</i>     | <i>Caesalpinia sepiaria</i>                                                              | Fabaceae                                 | South Africa                    | Doidge (1948); Sivanesan (1971)                                                                      |
| <i>Herpotrichia calamicola</i>       | <i>Calamus caryotoides</i>                                                               | Arecaceae                                | Australia                       | Fröhlich and Hyde (2000); Farr and Rossman (2024)                                                    |
| <i>Herpotrichia callimorpha</i>      | <i>Chamaenerion angustifolium</i> , <i>Salix</i> sp. and <i>Xanthophyllum flavescens</i> | Onagraceae, Polygalaceae, and Salicaceae | Denmark and India               | Munk (1957); Rabenhorst (1869); Pande (2008); Farr and Rossman (2024)                                |
| <i>Herpotrichia caulogena</i>        | <i>Silene nutans</i>                                                                     | Caryophyllaceae                          | Luxembourg                      | Feltgen (1903)                                                                                       |
| <i>Herpotrichia chilensis</i>        | <i>Proustia pungens</i>                                                                  | Asteraceae                               | Chile                           | Feltgen (1903); Farr (1973); Farr and Rossman (2024)                                                 |
| <i>Herpotrichia cirrhostoma</i>      | Unidentified host                                                                        | –                                        | Sri Lanka                       | Berkeley and Broome (1873); Petch (1912)                                                             |
| <i>Herpotrichia dalisayi</i>         | Unidentified host                                                                        | –                                        | The Philippines                 | Hyde and Aptroot (1998)                                                                              |
| <i>Herpotrichia decidua</i>          | Unidentified host                                                                        | –                                        | The United States               | Barr (1992); Farr and Rossman (2024)                                                                 |
| <i>Herpotrichia detzneriae</i>       | <i>Detzneria tubata</i>                                                                  | Plantaginaceae                           | Papua New Guinea                | Kobayasi (1971); Shaw (1984); Farr and Rossman (2024)                                                |
| <i>Herpotrichia diffusa</i>          | <i>Juglans cinerea</i> and <i>Populus</i> sp.                                            | Juglandaceae                             | Palestine and the United States | Ellis and Everhart (1892); Ellis (1895)                                                              |
| <i>Herpotrichia ellisii</i>          | <i>Abies</i> sp.                                                                         | Pinaceae                                 | Canada                          | Barr (1992); Farr and Rossman (2024)                                                                 |
| <i>Herpotrichia ephedrae</i>         | <i>Ephedra distachya</i>                                                                 | Ephedraceae                              | France                          | Kuhnholz-Lordat and Barry (1949)                                                                     |
| <i>Herpotrichia fusispora</i>        | Unidentified host                                                                        | –                                        | China                           | Chen and Hsieh (2004); Farr and Rossman (2024)                                                       |
| <i>Herpotrichia gelasinoporoides</i> | Soil                                                                                     | –                                        | India                           | Von Arx (1981)                                                                                       |
| <i>Herpotrichia henkeliana</i>       | <i>Phragmites communis</i>                                                               | Poaceae                                  | Germany                         | Sydow and Sydow (1921)                                                                               |

| Species                              | Host                                                                                                                                                                                                                                                            | Host family                                                                                             | Locality                                                                                                    | References                                                                                                                                 |
|--------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Herpotrichia herbarum</i>         | <i>Achillea millefolium</i>                                                                                                                                                                                                                                     | Asteraceae                                                                                              |                                                                                                             | Wehmeyer (1952); Barr (1992)                                                                                                               |
| <i>Herpotrichia herpotrichoides</i>  | <i>Carya</i> sp., <i>Epilobium</i> sp., <i>Rhododendron hirsutum</i> , <i>Ribes</i> sp. and <i>Rubus</i> sp.                                                                                                                                                    | Ericaceae, Juglandaceae, Grossulariaceae, Onagraceae, and Rosaceae                                      | Austria, Denmark, Germany, Poland, the United States and the United Kingdom                                 | Cannon (1982); Barr (1984); Cannon et al. (1985); Mullenko et al. (2008); Zhang et al. (2012); Tian et al. (2015); Farr and Rossman (2024) |
| <i>Herpotrichia hippocrateae</i>     | <i>Hippocratea grahamii</i>                                                                                                                                                                                                                                     | Celastraceae                                                                                            | India                                                                                                       | Tilak and Talde (1975); Pande (2008)                                                                                                       |
| <i>Herpotrichia indica</i>           | <i>Duranta plumieri</i>                                                                                                                                                                                                                                         | Verbenaceae                                                                                             | India                                                                                                       | Anahosur (1970); Pande (2008); Farr and Rossman (2024)                                                                                     |
| <i>Herpotrichia laricina</i>         | <i>Larix decidua</i>                                                                                                                                                                                                                                            | Pinaceae                                                                                                | Luxembourg                                                                                                  | Feltgen (1901)                                                                                                                             |
| <i>Herpotrichia leptospora</i>       | Unidentified host                                                                                                                                                                                                                                               | –                                                                                                       | –                                                                                                           | Kirschstein (1911)                                                                                                                         |
| <i>Herpotrichia lignicola</i>        | Unidentified host                                                                                                                                                                                                                                               | –                                                                                                       | Belgium                                                                                                     | Bose (1961)                                                                                                                                |
| <i>Herpotrichia macrotricha</i>      | <i>Acer spicatum</i> , <i>Agropyron repens</i> , <i>Agrostis alba</i> , <i>Arundinaria</i> sp., <i>Carex</i> sp., <i>Cocos nucifera</i> , <i>Eupatorium formosanum</i> , <i>Fagus sylvatica</i> , <i>Fraxinus</i> sp., <i>Rubus</i> sp. and <i>Solidago</i> sp. | Arecaceae, Asteraceae, Cupressaceae, Cyperaceae, Fagaceae, Oleaceae, Poaceae, Rosaceae, and Sapindaceae | China, India, the United Kingdom and the United States                                                      | Dennis (1978); Saccardo (1883); Barr (1968, 1984); Pande (2008); Chen and Hsieh (2004); Farr and Rossman (2024)                            |
| <i>Herpotrichia mangrovei</i>        | Unidentified host                                                                                                                                                                                                                                               | –                                                                                                       | China                                                                                                       | Jones and Vrijmoed (2003)                                                                                                                  |
| <i>Herpotrichia melanotricha</i>     | <i>Hevea brasiliensis</i>                                                                                                                                                                                                                                       | Euphorbiaceae                                                                                           | Congo                                                                                                       | Saccas (1954)                                                                                                                              |
| <i>Herpotrichia millettiae</i>       | <i>Millettia</i> sp.                                                                                                                                                                                                                                            | Fabaceae                                                                                                | Malaysia                                                                                                    | Sivanesan (1971)                                                                                                                           |
| <i>Herpotrichia monospermatis</i>    | <i>Butea monosperma</i>                                                                                                                                                                                                                                         | Fabaceae                                                                                                | India                                                                                                       | Pande (2008); Farr and Rossman (2024)                                                                                                      |
| <i>Herpotrichia mulleri</i>          | <i>Artemisia nilagirica</i> and <i>Butea monosperma</i>                                                                                                                                                                                                         | Asteraceae, Fabaceae                                                                                    | India                                                                                                       | Pande (2008); Farr and Rossman (2024)                                                                                                      |
| <i>Herpotrichia myriangii</i>        | <i>Carica papaya</i>                                                                                                                                                                                                                                            | Caricaceae                                                                                              | Java                                                                                                        | Raciborski (1909)                                                                                                                          |
| <i>Herpotrichia nectrioides</i>      | Melastomataceae sp.                                                                                                                                                                                                                                             | Melastomataceae                                                                                         | Brazil                                                                                                      | Rehm (1901)                                                                                                                                |
| <i>Herpotrichia nigra</i>            | <i>Abies</i> sp., <i>Calocedrus decurrens</i> , <i>Cedrus libani</i> , <i>Chamaecyparis nootkatensis</i> , <i>Juniperus</i> sp., <i>Phyllodoce</i> sp., <i>Picea</i> sp., <i>Pinus</i> sp., <i>Pseudotsuga menziesii</i> and <i>Rhododendron</i> sp.            | Cupressaceae, Ericaceae, and Pinaceae                                                                   | Austria, Canada, France, Germany, Italy, Norway, Poland, Switzerland, Turkey, Ukraine and the United States | Shaw (1973); Ginns (1986); French (1989); Farr and Rossman (2024)                                                                          |
| <i>Herpotrichia nigrotuberculata</i> | <i>Elaeis guineensis</i> and <i>Phyllostachys reticulata</i>                                                                                                                                                                                                    | Arecaceae and Poaceae                                                                                   | Japan and Tanzania                                                                                          | Pirozynski (1972); Farr and Rossman (2024)                                                                                                 |
| <i>Herpotrichia nypicola</i>         | <i>Nypa fruticans</i>                                                                                                                                                                                                                                           | Arecaceae                                                                                               | Malaysia                                                                                                    | Hyde et al. (1999); Farr and Rossman (2024)                                                                                                |
| <i>Herpotrichia occulta</i>          | <i>Eucalyptus</i> sp.                                                                                                                                                                                                                                           | Myrtaceae                                                                                               | Brazil                                                                                                      | Rick (1933)                                                                                                                                |
| <i>Herpotrichia ochrostoma</i>       | <i>Fraxinus excelsior</i>                                                                                                                                                                                                                                       | Oleaceae                                                                                                | Luxembourg                                                                                                  | Feltgen (1903)                                                                                                                             |
| <i>Herpotrichia palmicola</i>        | <i>Calamus caryotoides</i> , <i>Daemonorops</i> sp. and <i>Licuala ramsayi</i>                                                                                                                                                                                  | Arecaceae                                                                                               | Australia and China                                                                                         | Hyde et al. (1999); Fröhlich and Hyde 2000; Zhuang (2001); Farr and Rossman (2024)                                                         |
| <i>Herpotrichia pandei</i>           | <i>Saccharum spontaneum</i>                                                                                                                                                                                                                                     | Poaceae                                                                                                 | India                                                                                                       | Bose (1961)                                                                                                                                |
| <i>Herpotrichia petrakiana</i>       | <i>Fagus sylvatica</i>                                                                                                                                                                                                                                          | Fagaceae                                                                                                | –                                                                                                           | Bose (1961)                                                                                                                                |

| Species                                 | Host                                                                                          | Host family                                | Locality                                                      | References                                                                         |
|-----------------------------------------|-----------------------------------------------------------------------------------------------|--------------------------------------------|---------------------------------------------------------------|------------------------------------------------------------------------------------|
| <i>Herpotrichia philippinensis</i>      | <i>Alstonia scholaris</i>                                                                     | Apocynaceae                                | The Philippines                                               | Rehm (1914); Teodoro (1937); Farr and Rossman (2024)                               |
| <i>Herpotrichia pinetorum</i>           | <i>Herpotrichia pinetorum</i>                                                                 | Pinaceae                                   | Austria and India                                             | Winter (1885); Mueller (1958); Petrak (1962); Farr and Rossman (2024)              |
| <i>Herpotrichia quinqueseptata</i>      | <i>Abies lasiocarpa</i> , <i>Larix europaea</i> , <i>Picea</i> sp. and <i>Populus tremula</i> | Pinaceae and Salicaceae                    | Canada, Czech Republic, Germany, Sweden and the United States | Shaw (1973); Minter (1981); Ginns (1986); Eriksson (2014); Farr and Rossman (2024) |
| <i>Herpotrichia rara</i>                | <i>Tanacetum vulgare</i>                                                                      | Asteraceae                                 | Germany                                                       | Kirschstein (1935)                                                                 |
| <i>Herpotrichia rhenana</i>             | <i>Epilobium angustifolium</i> , <i>Rhododendron hirsutum</i> and <i>Ribes</i> sp.            | Ericaceae, Grossulariaceae, and Onagraceae | Austria and the United States                                 | Fuckel (1870); Remler (1979); Farr and Rossman (2024)                              |
| <i>Herpotrichia rhodospiloides</i>      | <i>Populus deltoides</i>                                                                      | Salicaceae                                 | The United States                                             | Peck (1909)                                                                        |
| <i>Herpotrichia rhodosticta</i>         | <i>Carex paniculata</i> and <i>Populus</i> sp.                                                | Cyperaceae and Salicaceae                  | Germany and the United States                                 | Saccardo (1882); Samuels (1976); Farr and Rossman (2024)                           |
| <i>Herpotrichia setosa</i>              | <i>Betula glandulosa</i> and <i>Myrica gale</i>                                               | Betulaceae and Myricaceae                  | Canada and Ireland                                            | Barr (1992); Farr and Rossman (2024)                                               |
| <i>Herpotrichia striatispora</i>        | Soil                                                                                          | –                                          | South Africa                                                  | Papendorf and Arx (1966)                                                           |
| <i>Herpotrichia symphoricarpi</i>       | <i>Symphoricarpos</i> sp.                                                                     | Caprifoliaceae                             | The United States                                             | Barr (1984); Farr and Rossman (2024)                                               |
| <i>Herpotrichia tenuispora</i>          | <i>Urtica dioica</i>                                                                          | Urticaceae                                 | Germany                                                       | Kirschstein (1906)                                                                 |
| <i>Herpotrichia vaginatispora</i>       | <i>Trifolium</i> sp.                                                                          | Fabaceae                                   | Italy                                                         | Tian et al. (2015)                                                                 |
| <i>Herpotrichia villosa</i>             | Unidentified host                                                                             | –                                          | Brazil                                                        | Samuels and Müller (1978)                                                          |
| <i>Herpotrichia xiaokongensis</i>       | <i>Prunus</i> sp.                                                                             | Rosaceae                                   | China                                                         | Hyde et al. (2021)                                                                 |
| <i>Herpotrichia zingiberacearum</i>     | <i>Hedychium coronarium</i>                                                                   | Zingiberaceae                              | China                                                         | This study                                                                         |
| <i>Mamillisphaeria dimorphospora</i>    | Unidentified host                                                                             | –                                          | Australia                                                     | Hyde et al. (1996)                                                                 |
| <i>Marjia tianshanica</i>               | <i>Cerasus tianshanica</i>                                                                    | Rosaceae                                   | Uzbekistan                                                    | Wanasinghe et al. (2018)                                                           |
| <i>Marjia uzbekistanica</i>             | <i>Rosa</i> sp.                                                                               | Rosaceae                                   | Uzbekistan                                                    | Wanasinghe et al. (2018)                                                           |
| <i>Melanocamarosporioides ugamica</i>   | <i>Lonicera altmannii</i>                                                                     | Caprifoliaceae                             | Uzbekistan                                                    | Pem et al. (2019)                                                                  |
| <i>Melanocamarosporium galiicola</i>    | <i>Galium</i> sp.                                                                             | Rubiaceae                                  | Italy                                                         | Wijayawardene et al. (2016)                                                        |
| <i>Melanocucurbitaria uzbekistanica</i> | <i>Acer pubescens</i>                                                                         | Sapindaceae                                | Uzbekistan                                                    | Wanasinghe et al. (2018)                                                           |
| <i>Melanodiplodia tianschanica</i>      | <i>Rosa</i> sp.                                                                               | Rosaceae                                   | Uzbekistan                                                    | Wanasinghe et al. (2018)                                                           |
| <i>Melanomma acanthophilum</i>          | <i>Cereus quisco</i>                                                                          | Cactaceae                                  | Chile                                                         | Spegazzini (1923); Farr (1973); Farr and Rossman (2024)                            |
| <i>Melanomma afflatum</i>               | Unidentified host                                                                             | –                                          | The United States                                             | Shear (1941)                                                                       |
| <i>Melanomma anceps</i>                 | Unidentified host                                                                             | –                                          | Jawa                                                          | Höhnel (1909)                                                                      |
| <i>Melanomma andinum</i>                | <i>Bulnesia retamo</i>                                                                        | Zygophyllaceae                             | Argentina                                                     | Spegazzini (1909); Farr (1973); Farr and Rossman (2024)                            |
| <i>Melanomma artemisiae-maritimae</i>   | <i>Artemisia maritima</i>                                                                     | Asteraceae                                 | Russia                                                        | Lobik (1927); Farr and Rossman (2024)                                              |
| <i>Melanomma aspegrenii</i>             | <i>Carpinus betulus</i> , <i>Cornus</i> sp. and <i>Fagus sylvatica</i>                        | Betulaceae, Cornaceae, and Fagaceae        | Poland                                                        | Fuckel (1870); Mulenko et al. (2008); Farr and Rossman (2024)                      |

| Species                            | Host                                    | Host family   | Locality                                | References                                                          |
|------------------------------------|-----------------------------------------|---------------|-----------------------------------------|---------------------------------------------------------------------|
| <i>Melanomma aurantiicola</i>      | <i>Citrus</i> sp.                       | Rutaceae      | Paraguay                                | Spegazzini (1920); Farr (1973); Farr and Rossman (2024)             |
| <i>Melanomma aurantiiphila</i>     | <i>Citrus</i> sp.                       | Rutaceae      | Paraguay                                | Spegazzini (1920); Farr (1973); Farr and Rossman (2024)             |
| <i>Melanomma australiense</i>      | Unidentified host                       | –             | Australia                               | Hyde and Goh (1999)                                                 |
| <i>Melanomma brachythele</i>       | <i>Sambucus</i> sp.                     | Adoxaceae     | The United Kingdom                      | Saccardo (1878)                                                     |
| <i>Melanomma bubakii</i>           | <i>Campanula stricta</i>                | Campanulaceae | Turkey                                  | Bubák (1914)                                                        |
| <i>Melanomma cacheutense</i>       | <i>Baccharis glutinosa</i>              | Asteraceae    | Argentina                               | Spegazzini (1909); Farr (1973); Farr and Rossman (2024)             |
| <i>Melanomma caesalpiniae</i>      | <i>Caesalpinia cearense</i>             | Fabaceae      | Brazil                                  | Hennings (1908)                                                     |
| <i>Melanomma caryophagum</i>       | <i>Carya</i> sp. and <i>Juglans</i> sp. | Juglandaceae  | The United States                       | Fairman (1921)                                                      |
| <i>Melanomma castillejae</i>       | <i>Castilleja pallida</i>               | Orobanchaceae | Siberia                                 | Farr and Rossman (2024)                                             |
| <i>Melanomma ceratoniae</i>        | <i>Ceratonia siliqua</i>                | Fabaceae      | Spain                                   | Farr and Rossman (2024)                                             |
| <i>Melanomma chilense</i>          | <i>Proustia pungens</i>                 | Asteraceae    | Chile                                   | Spegazzini (1910); Farr (1973); Farr and Rossman (2024)             |
| <i>Melanomma citricola</i>         | <i>Citrus medica</i>                    | Rutaceae      | Bangladesh and India                    | Sydow et al. (1911); Rao (1969); Farr and Rossman (2024)            |
| <i>Melanomma conjunctum</i>        | <i>Thuja plicata</i>                    | Cupressaceae  | The United States                       | Farr and Rossman (2024)                                             |
| <i>Melanomma cryptostegiae</i>     | <i>Cryptostegia grandiflora</i>         | Apocynaceae   | India                                   | Pande (2008); Farr and Rossman (2024)                               |
| <i>Melanomma cucurbitarioideum</i> | <i>Pentaphylloides fruticosa</i>        | Rosaceae      | China                                   | Yuan and Barr (1994); Farr and Rossman (2024)                       |
| <i>Melanomma dactylosporum</i>     | Unidentified host                       | –             | Brazil                                  | Rehm (1901)                                                         |
| <i>Melanomma dinghuense</i>        | Unidentified host                       | –             | China                                   | Inderbitzin and Huang (2001)                                        |
| <i>Melanomma distinctum</i>        | <i>Pentaphylloides fruticosa</i>        | Rosaceae      | Russia                                  | Vasilyeva (1987)                                                    |
| <i>Melanomma drimydis</i>          | <i>Drymis</i> sp.                       | Winteraceae   | Brazil                                  | Rehm (1901)                                                         |
| <i>Melanomma dryinum</i>           | <i>Quercus</i> sp.                      | Fagaceae      | Belgium                                 | Mouton (1900)                                                       |
| <i>Melanomma dzungaricum</i>       | <i>Eurotia eversmanniana</i>            | Amaranthaceae | Kazakhstan                              | Vasilyeva (1987)                                                    |
| <i>Melanomma ebeni</i>             | <i>Ebenus stellata</i>                  | Fabaceae      | Iran                                    | González Fragoso (1918)                                             |
| <i>Melanomma epiphytica</i>        | <i>Bambusa</i> sp.                      | Poaceae       | Indonesia                               | Raciborski (1909); Eriksson and Yue (1998); Farr and Rossman (2024) |
| <i>Melanomma gigantea</i>          | Unidentified host                       | –             | India                                   | Pande (1979)                                                        |
| <i>Melanomma glumarum</i>          | <i>Oryza sativa</i>                     | Poaceae       | China, India, Japan and the Philippines | Watson (1971); Reinking (1919); Farr and Rossman (2024)             |
| <i>Melanomma gregarium</i>         | <i>Populus</i> sp.                      | Salicaceae    | The United States                       | Cash (1953); Farr and Rossman (2024)                                |
| <i>Melanomma halimodendri</i>      | <i>Halimodendron halodendron</i>        | Fabaceae      | Kazakhstan                              | Farr and Rossman (2024)                                             |
| <i>Melanomma haloxyli</i>          | <i>Haloxylon aphyllum</i>               | Amaranthaceae | Kazakhstan                              | Farr and Rossman (2024)                                             |
| <i>Melanomma helianthemii</i>      | <i>Helianthemum rupifragum</i>          | Cistaceae     | Ukraine                                 | Gucevic (1969)                                                      |
| <i>Melanomma heraclei</i>          | <i>Heracleum pubescens</i>              | Apiaceae      | Ukraine                                 | Dudka et al. (2004); Farr and Rossman (2024)                        |
| <i>Melanomma herpotrichum</i>      | <i>Populus</i> sp.                      | Salicaceae    | Luxembourg                              | Feltgen (1903)                                                      |
| <i>Melanomma japonicum</i>         | <i>Fagus crenata</i>                    | Fagaceae      | Japan                                   | Hashimoto et al. (2017); Farr and Rossman (2024)                    |

| Species                          | Host                                                                                                                                                                                                                                                                                                        | Host family                                                                                         | Locality                                                                    | References                                                                                   |
|----------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|
| <i>Melanomma jenynsii</i>        | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | The United Kingdom                                                          | Saccardo (1883)                                                                              |
| <i>Melanomma juniperi</i>        | <i>Juniperus virginiana</i>                                                                                                                                                                                                                                                                                 | Cupressaceae                                                                                        | The United States                                                           | Fairman (1905); Cash (1953); Farr and Rossman (2024)                                         |
| <i>Melanomma langloisii</i>      | <i>Salix nigra</i>                                                                                                                                                                                                                                                                                          | Salicaceae                                                                                          | The United States                                                           | Farr and Rossman (2024)                                                                      |
| <i>Melanomma lithophilae</i>     | <i>Sobolewsia lithophila</i>                                                                                                                                                                                                                                                                                | Apocynaceae                                                                                         | Ukraine                                                                     | Dudka et al. (2004); Farr and Rossman (2024)                                                 |
| <i>Melanomma longicolle</i>      | <i>Acer</i> sp. and <i>Citrus limon</i>                                                                                                                                                                                                                                                                     | Aceraceae and Rutaceae                                                                              | Italy and the United Kingdom                                                | Saccardo (1875); Farr and Rossman (2024)                                                     |
| <i>Melanomma marathawadense</i>  | On paper                                                                                                                                                                                                                                                                                                    | –                                                                                                   | India                                                                       | Tilak and Kale (1970)                                                                        |
| <i>Melanomma margaretae</i>      | <i>Dryas octopetala</i>                                                                                                                                                                                                                                                                                     | Rosaceae                                                                                            | Poland                                                                      | Mulenko et al. (2008); Farr and Rossman (2024)                                               |
| <i>Melanomma martinianum</i>     | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | New Zealand                                                                 | Saccardo (1883)                                                                              |
| <i>Melanomma mate</i>            | <i>Ilex paraguensis</i>                                                                                                                                                                                                                                                                                     | Aquifoliaceae                                                                                       | Argentina                                                                   | Spegazzini (1908); Farr (1973); Farr and Rossman (2024)                                      |
| <i>Melanomma medium</i>          | <i>Acer negundo</i> , <i>Calligonum</i> sp. and <i>Tamarix</i> sp.                                                                                                                                                                                                                                          | Sapindaceae and Tamaricaceae                                                                        | Canada, Italy and the United Kingdom                                        | Saccardo (1878); Farr (1973); Farr and Rossman (2024)                                        |
| <i>Melanomma mindorenses</i>     | <i>Arenga mindorensis</i>                                                                                                                                                                                                                                                                                   | Arecaceae                                                                                           | The Philippines                                                             | Rehm (1913)                                                                                  |
| <i>Melanomma mojunkumica</i>     | <i>Haloxylon aphyllum</i>                                                                                                                                                                                                                                                                                   | Amaranthaceae                                                                                       | Kazakhstan                                                                  | Farr and Rossman (2024)                                                                      |
| <i>Melanomma moravicum</i>       | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | Slovakia                                                                    | Farr and Rossman (2024)                                                                      |
| <i>Melanomma mutabile</i>        | <i>Solanum dulcamara</i>                                                                                                                                                                                                                                                                                    | Solanaceae                                                                                          | Luxembourg                                                                  | Feltgen (1901)                                                                               |
| <i>Melanomma myricae</i>         | <i>Myrica gale</i>                                                                                                                                                                                                                                                                                          | Myricaceae                                                                                          | Sweden                                                                      | Eriksson (2014); Farr and Rossman (2024)                                                     |
| <i>Melanomma nigriseda</i>       | <i>Fagus</i> sp.                                                                                                                                                                                                                                                                                            | Fagaceae                                                                                            | The United States                                                           | Fairman (1922)                                                                               |
| <i>Melanomma obliterans</i>      | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | The United Kingdom                                                          | Saccardo (1883)                                                                              |
| <i>Melanomma obtusissimum</i>    | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | Cuba                                                                        | Farr and Rossman (2024)                                                                      |
| <i>Melanomma oryzae</i>          | <i>Oryza sativa</i>                                                                                                                                                                                                                                                                                         | Poaceae                                                                                             | Japan                                                                       | Farr and Rossman (2024)                                                                      |
| <i>Melanomma oxysporum</i>       | <i>Quercus</i> sp.                                                                                                                                                                                                                                                                                          | Fagaceae                                                                                            | The United States                                                           | Hawksworth (1985)                                                                            |
| <i>Melanomma panici-miliacei</i> | <i>Panicum miliaceum</i>                                                                                                                                                                                                                                                                                    | Poaceae                                                                                             | Siberia                                                                     | Farr and Rossman (2024)                                                                      |
| <i>Melanomma philippinense</i>   | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | The Philippines                                                             | Sydow and Sydow (1914)                                                                       |
| <i>Melanomma populicola</i>      | <i>Cornus</i> sp., <i>Fagus sylvatica</i> , <i>Picea abies</i> , <i>Populus</i> sp., <i>Quercus</i> sp. and <i>Sorbus aucuparia</i>                                                                                                                                                                         | Cornaceae, Fagaceae, Pinaceae, Rosaceae, and Salicaceae                                             | China, Croatia, Germany and the Netherlands                                 | Crous et al. (2020); Farr and Rossman (2024); this study                                     |
| <i>Melanomma praeandinum</i>     | <i>Salvia gilliesii</i>                                                                                                                                                                                                                                                                                     | Lamiaceae                                                                                           | Argentina                                                                   | Spegazzini (1909); Farr (1973); Farr and Rossman (2024)                                      |
| <i>Melanomma pulveracea</i>      | Unidentified host                                                                                                                                                                                                                                                                                           | –                                                                                                   | China                                                                       | Teng (1936)                                                                                  |
| <i>Melanomma pulvis-pyrius</i>   | <i>Acer</i> sp., <i>Albizia julibrissin</i> , <i>Alhagi</i> sp., <i>Alnus</i> sp., <i>Berberis</i> sp., <i>Betula</i> sp., <i>Bupleurum fruticosum</i> , <i>Campsis radicans</i> , <i>Carpinus betulus</i> , <i>Celtis australis</i> , <i>Corylus</i> sp., <i>Larix decidua</i> and <i>Pinus sylvestris</i> | Apiaceae, Betulaceae, Berberidaceae, Bignoniaceae, Cannabaceae, Fabaceae, Pinaceae, and Sapindaceae | Canada, Czech Republic, Japan, Poland, Russia, Scotland, Sweden and Ukraine | Mulenko et al. (2008); Hashimoto et al. (2017); Crous et al. (2020); Farr and Rossman (2024) |

| Species                            | Host                                                                                 | Host family                                   | Locality                                                                                                             | References                                                             |
|------------------------------------|--------------------------------------------------------------------------------------|-----------------------------------------------|----------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------|
| <i>Melanomma pyriostictum</i>      | Unidentified host                                                                    | –                                             | The United Kingdom                                                                                                   | Cooke (1887)                                                           |
| <i>Melanomma rhododendri</i>       | <i>Rhododendron</i> sp.                                                              | Ericaceae                                     | Austria, Belgium, Germany, Italy, Luxembourg, the Netherlands, Switzerland, the United Kingdom and the United States | Rehm (1881, 1906); Bommer and Rousseau (1890); Farr and Rossman (2024) |
| <i>Melanomma ribis</i>             | <i>Ribes</i> sp.                                                                     | Grossulariaceae                               | The United States                                                                                                    | Barr (1990)                                                            |
| <i>Melanomma rubicundum</i>        | <i>Lythrum</i> sp.                                                                   | Lythraceae                                    | Sweden                                                                                                               | Bommer and Rousseau (1890); Farr and Rossman (2024)                    |
| <i>Melanomma sanguinarium</i>      | Unidentified host                                                                    | –                                             | –                                                                                                                    | Saccardo (1878)                                                        |
| <i>Melanomma saviczii</i>          | <i>Thymus pseudohumillimus</i>                                                       | Lamiaceae                                     | Ukraine                                                                                                              | Dudka et al. (2004); Farr and Rossman (2024)                           |
| <i>Melanomma scrophulariae</i>     | <i>Scrophularia rupestris</i>                                                        | Scrophulariaceae                              | Ukraine                                                                                                              | Gucevic (1967)                                                         |
| <i>Melanomma sordidissimum</i>     | <i>Eriobotrya japonica</i>                                                           | Rosaceae                                      | Argentina                                                                                                            | Spegazzini (1909); Farr (1973); Farr and Rossman (2024)                |
| <i>Melanomma sparsum</i>           | <i>Abies</i> sp.                                                                     | Pinaceae                                      | Switzerland                                                                                                          | Fuckel (1873)                                                          |
| <i>Melanomma spiniferum</i>        | <i>Morus alba</i>                                                                    | Moraceae                                      | The United States                                                                                                    | Cash (1953); Farr and Rossman (2024)                                   |
| <i>Melanomma subandinum</i>        | <i>Atriplex pamparum</i>                                                             | Amaranthaceae                                 | Argentina                                                                                                            | Cash (1953); Farr and Rossman (2024)                                   |
| <i>Melanomma subdispersum</i>      | <i>Betula</i> sp., and <i>Fagus</i> sp.                                              | Betulaceae and Fagaceae                       | Canada, Ireland, Poland, and the United Kingdom                                                                      | Connors (1967); Mulencko et al. (2008); Farr and Rossman (2024)        |
| <i>Melanomma submojunkumica</i>    | <i>Haloxylon aphyllum</i>                                                            | Amaranthaceae                                 | Kazakhstan                                                                                                           | Farr and Rossman (2024)                                                |
| <i>Melanomma thespesiae</i>        | <i>Thespesia</i> sp.                                                                 | Malvaceae                                     | India                                                                                                                | Farr and Rossman (2024)                                                |
| <i>Melanomma trevoae</i>           | <i>Trevoa trinervia</i>                                                              | Rhamnaceae                                    | Chile                                                                                                                | Farr (1973); Farr and Rossman (2024)                                   |
| <i>Melanomma vile</i>              | <i>Quercus</i> sp.                                                                   | Fagaceae                                      | Sweden                                                                                                               | Fuckel (1870)                                                          |
| <i>Melanomma xylariae</i>          | Unidentified host                                                                    | –                                             | Brazil                                                                                                               | Höhnelt (1907)                                                         |
| <i>Monoseptella rosae</i>          | <i>Rosa</i> sp.                                                                      | Rosaceae                                      | Uzbekistan                                                                                                           | Wanasinghe et al. (2018)                                               |
| <i>Muriformistrickeria rosae</i>   | <i>Rosa</i> sp.                                                                      | Rosaceae                                      | Italy                                                                                                                | Wanasinghe et al. (2018)                                               |
| <i>Muriformistrickeria rubi</i>    | <i>Rubus</i> sp.                                                                     | Rosaceae                                      | Italy                                                                                                                | Tian et al. (2015)                                                     |
| <i>Navicella costaricensis</i>     | Unidentified host                                                                    | –                                             | The United States                                                                                                    | El-Shafie et al. (2005)                                                |
| <i>Navicella diabola</i>           | <i>Castanopsis</i> sp.                                                               | Fagaceae                                      | China                                                                                                                | Aptroot (2003)                                                         |
| <i>Navicella pallida</i>           | <i>Elaeocarpus</i> sp.                                                               | Elaeocarpaceae                                | Papua New Guinea                                                                                                     | Aptroot and Iperen (1998)                                              |
| <i>Navicella pileata</i>           | <i>Fraxinus</i> sp., <i>Salix fragilis</i> , <i>Tilia</i> sp. and <i>Quercus</i> sp. | Fagaceae, Malvaceae, Oleaceae, and Salicaceae | Finland, Lithuania and Norway                                                                                        | Fabre (1879); Holm and Holm (1988); Farr and Rossman (2024)            |
| <i>Navicella xinjiangensis</i>     | <i>Lonicera hispida</i> , <i>Haloxylon ammodendron</i>                               | Amaranthaceae and Caprifoliaceae              | China                                                                                                                | Yuan and Zhao (1994); Farr and Rossman (2024)                          |
| <i>Neobyssosphaeria clematidis</i> | <i>Clematis vitalba</i>                                                              | Ranunculaceae                                 | The United Kingdom                                                                                                   | Phukhamsakda et al. (2020)                                             |

| Species                              | Host                                                                                                               | Host family                                                | Locality                                           | References                                                                              |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|----------------------------------------------------|-----------------------------------------------------------------------------------------|
| <i>Petrakia aesculi</i>              | <i>Aesculus turbinata</i>                                                                                          | Sapindaceae                                                | Japan                                              | Jaklitsch and Voglmayr (2017)                                                           |
| <i>Petrakia deviata</i>              | <i>Acer campestre</i>                                                                                              | Sapindaceae                                                | Georgia and Switzerland                            | Watzl (1937); Gross et al. (2017)                                                       |
| <i>Petrakia echinata</i>             | <i>Acer</i> sp.                                                                                                    | Sapindaceae                                                | Italy, Slovakia, Switzerland and the United States | Sydow and Sydow (1913); Li et al. (2016a); Gross et al. (2017); Farr and Rossman (2024) |
| <i>Petrakia fagi</i>                 | <i>Fagus crenata</i>                                                                                               | Fagaceae                                                   | Japan                                              | Beenken et al. (2020)                                                                   |
| <i>Petrakia greenei</i>              | <i>Acer saccharinum</i>                                                                                            | Sapindaceae                                                | The United States                                  | Beenken et al. (2020)                                                                   |
| <i>Petrakia irregularis</i>          | <i>Acer pseudoplatanus</i>                                                                                         | Sapindaceae                                                | The Netherlands and Poland                         | Mulenko et al. (2008); Farr and Rossman (2024)                                          |
| <i>Petrakia juniperi</i>             | <i>Juniperus</i> sp.                                                                                               | Cupressaceae                                               | Germany                                            | Bedlan (2017)                                                                           |
| <i>Petrakia liobae</i>               | <i>Fagus sylvatica</i>                                                                                             | Fagaceae                                                   | Switzerland                                        | Beenken et al. (2020)                                                                   |
| <i>Petrakia minima</i>               | <i>Fagus japonica</i>                                                                                              | Fagaceae                                                   | Japan                                              | Hashimoto et al. (2017); Beenken et al. (2020)                                          |
| <i>Petrakia paracochinensis</i>      | <i>Miscanthus floridulus</i>                                                                                       | Poaceae                                                    | China                                              | Wong et al. (2002)                                                                      |
| <i>Phragmocephala atra</i>           | <i>Rhopalostylis sapida</i> and <i>Urtica</i> sp.                                                                  | Arecaceae, Urticaceae                                      | New Zealand and the United Kingdom                 | Mason and Hughes (1951); McKenzie et al. (1992)                                         |
| <i>Phragmocephala elegans</i>        | <i>Gymnanthes lucida</i>                                                                                           | Euphorbiaceae                                              | Brazil and Cuba                                    | Castañeda (1985)                                                                        |
| <i>Phragmocephala elliptica</i>      | <i>Elaeagnus</i> sp., <i>Filipendula denudata</i> , <i>Laurus</i> sp. <i>Quercus robur</i> and <i>Sambucus</i> sp. | Adoxaceae, Elaeagnaceae, Fagaceae, Lauraceae, and Rosaceae | Canada, Russia, Ukraine and the United Kingdom     | Hughes (1979); Dennis (1986); Holubtsova and Andrianova (2008); Farr and Rossman (2024) |
| <i>Phragmocephala garethjonesii</i>  | Unidentified host                                                                                                  | –                                                          | China                                              | Su et al. (2015)                                                                        |
| <i>Phragmocephala glanduliformis</i> | <i>Corticium coronatum</i> , <i>Picea obovate</i> and <i>Quercus</i> sp.                                           | Corticaceae, Fagaceae, and Pinaceae                        | Austria                                            | Hughes (1955); Farr and Rossman (2024)                                                  |
| <i>Phragmocephala hughesii</i>       | Unidentified host                                                                                                  | –                                                          | China                                              | Wu and Zhuang (2005)                                                                    |
| <i>Phragmocephala minima</i>         | <i>Abies balsamea</i> and <i>Fagus sylvatica</i>                                                                   | Fagaceae and Pinaceae                                      | Canada and the United Kingdom                      | Mason and Hughes (1951); Connors (1967)                                                 |
| <i>Phragmocephala prolifera</i>      | <i>Populus tremuloides</i> and <i>Urtica dioica</i>                                                                | Salicaceae and Urticaceae                                  | Belgium and Canada                                 | Hughes (1979); Farr and Rossman (2024)                                                  |
| <i>Phragmocephala stemphylioides</i> | <i>Carya</i> sp., <i>Cistus</i> sp. and <i>Pistacia lentiscus</i>                                                  | Anacardiaceae, Cistaceae, and Juglandaceae                 | Brazil, Canada, China and Italy                    | Hughes (1958); Zhuang (2005); Lunghini et al. (2013); Farr and Rossman (2024)           |
| <i>Phragmotrichum andamanense</i>    | <i>Strobilanthes</i> sp.                                                                                           | Acanthaceae                                                | India                                              | Bhat and Kendrick (1993)                                                                |
| <i>Phragmotrichum chailletii</i>     | <i>Abies</i> sp. and <i>Picea</i> sp.                                                                              | Pinaceae                                                   | Canada, Switzerland, Romania and the United States | Kunze and Schmidt (1823); Ginns (1986); Farr and Rossman (2024)                         |
| <i>Phragmotrichum karstenii</i>      | <i>Acer platanoides</i>                                                                                            | Sapindaceae                                                | Finland                                            | Sutton and Pirozynski (1965)                                                            |
| <i>Phragmotrichum rivoclarinum</i>   | <i>Acer</i> sp., <i>Alnus</i> sp. and <i>Salix</i> sp.                                                             | Betulaceae, Salicaceae, and Sapindaceae                    | Canada, Italy and the United Kingdom               | Sutton and Pirozynski (1966); Ginns (1986); Farr and Rossman (2024)                     |
| <i>Phragmotrichum vassiljevae</i>    | <i>Alnus kamtschatica</i>                                                                                          | Betulaceae                                                 | Russia                                             | Melnik (1984)                                                                           |

| Species                               | Host                                                                                      | Host family                             | Locality                                                                                                                                     | References                                                                            |
|---------------------------------------|-------------------------------------------------------------------------------------------|-----------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| <i>Pleotrichocladium opacum</i>       | Soil                                                                                      | –                                       | Spain                                                                                                                                        | Hernández-Restrepo et al. (2017)                                                      |
| <i>Praetumpfia obducens</i>           | <i>Fraxinus excelsior</i>                                                                 | Oleaceae                                | Austria and Sweden                                                                                                                           | Jaklitsch and Voglmayr (2017)                                                         |
| <i>Pseudobyssosphaeria bambusae</i>   | <i>Bamboo</i> sp.                                                                         | Poaceae                                 | Thailand                                                                                                                                     | Hyde et al. (2018)                                                                    |
| <i>Pseudostrickeria muriformis</i>    | <i>Origanum vulgare</i>                                                                   | Lamiaceae                               | Italy                                                                                                                                        | Tian et al. (2015)                                                                    |
| <i>Pseudostrickeria ononidis</i>      | <i>Ononis spinosa</i>                                                                     | Fabaceae                                | Italy                                                                                                                                        | Tian et al. (2015)                                                                    |
| <i>Pseudostrickeria rosae</i>         | <i>Rosa</i> sp.                                                                           | Rosaceae                                | Italy                                                                                                                                        | Wanasinghe et al. (2018)                                                              |
| <i>Sarimanas pseudofluviatile</i>     | Unidentified host                                                                         | –                                       | Japan                                                                                                                                        | Liu et al. (2015)                                                                     |
| <i>Sarimanas shirakamiense</i>        | <i>Swida controversa</i>                                                                  | Cornaceae                               | Japan                                                                                                                                        | Liu et al. (2015)                                                                     |
| <i>Seifertia alpina</i>               | <i>Rhododendron ponticum</i>                                                              | Ericaceae                               | Austria                                                                                                                                      | Beenken et al. (2020)                                                                 |
| <i>Seifertia azaleae</i>              | <i>Ledum groenlandicum</i> ,<br><i>Leucopogon costatus</i><br>and <i>Rhododendron</i> sp. | Ericaceae                               | Australia, Canada, China, Germany, Italy, Japan, the Netherlands, New Zealand, Panama, Switzerland, the United Kingdom and the United States | White and Hamilton (1935); Partridge and Morgan-Jones (2002); Farr and Rossman (2024) |
| <i>Seifertia shangrilaensis</i>       | <i>Rhododendron decorum</i>                                                               | Ericaceae                               | China                                                                                                                                        | Li et al. (2016b)                                                                     |
| <i>Tumularia aquatica</i>             | <i>Alnus glutinosa</i> ,<br><i>Phragmites</i> sp. and<br><i>Quercus</i> sp.               | Betulaceae,<br>Fagaceae, and<br>Poaceae | South Africa and the United Kingdom                                                                                                          | Sivanesan (1984); Marvanová and Descals (1987); Farr and Rossman (2024)               |
| <i>Tumularia tuberculata</i>          | <i>Fagus sylvatica</i> ,<br><i>Quercus</i> sp.                                            | Fagaceae                                | Hungary                                                                                                                                      | Gönczöl (1976)                                                                        |
| <i>Uzbekistanica pruni</i>            | <i>Prunus armeniaca</i>                                                                   | Rosaceae                                | Russia                                                                                                                                       | Hyde et al. (2020)                                                                    |
| <i>Uzbekistanica rosae-hissaricae</i> | <i>Rosa</i> sp.                                                                           | Rosaceae                                | Uzbekistan                                                                                                                                   | Wanasinghe et al. (2018)                                                              |
| <i>Uzbekistanica vitis-viniferae</i>  | <i>Vitis vinifera</i>                                                                     | Vitaceae                                | Ukraine                                                                                                                                      | Crous et al. (2020)                                                                   |
| <i>Uzbekistanica yakutkhanika</i>     | <i>Rosa</i> sp.                                                                           | Rosaceae                                | Uzbekistan                                                                                                                                   | Wanasinghe et al. (2018)                                                              |
| <i>Xenostigmina aceris</i>            | <i>Acer macrophyllum</i>                                                                  | Sapindaceae                             | The United States                                                                                                                            | Hashimoto et al. (2017)                                                               |

## Taxonomy

### *Bertiella* (Sacc.) Sacc

**Notes.** *Bertiella* was established by Saccardo and Sydow (1899) to include *B. macrospora* as the type species. The species have superficial ascomata, cylindrical-clavate asci and hyaline, 1-septate (when immature) and pale brown, 3-septate (when mature) ascospores (Tian et al. 2015; Hongsanan et al. 2020). To the present time, there are six *Bertiella* species in Species Fungorum (2024). Of them, molecular data are available only for three species. *Bertiella* species have been reported from six different plant families, Cyrillaceae, Fagaceae, Lauraceae, Moraceae, Salicaceae and Ulmaceae (Fig. 10).

***Bertiella fici* Tennakoon, C.H. Kuo & K.D. Hyde, Fungal Diversity 108: 29 (2021)**

Index Fungorum: IF555314

Facesoffungi Number: FoF09317

Fig. 4

**Description.** *Saprobic* on dead leaves of *Cinnamomum verum* J. Presl (Lauraceae). **Sexual morph:** **Ascomata** 160–220 × 230–280 µm ( $\bar{x}$  = 180 × 240 µm, n = 15), solitary or scattered, semi-immersed to superficial, appeared as black dots on host surface, globose to subglobose, glabrous, unilocular, ostiolate. **Peridium** 12–20 µm wide, thick-walled, carbonaceous, composed of several layers of brown to dark brown pseudoparenchymatous cells, cells towards the inside hyaline, arranged in a **textura angularis**, fusing at the outside indistinguishable from the host tissues. **Hamathecium** comprising numerous, 1–2 µm wide, hyaline, septate, cellular pseudoparaphyses. **Asci** 50–60 × 7.5–8.5 µm ( $\bar{x}$  = 52 × 7.8 µm, n = 20), 8-spored, bitunicate, fissitunicate, cylindrical to cylindrical-clavate, short pedicellate, apically rounded, with a distinct ocular chamber. **Ascospores** 14–18 × 4–5 µm ( $\bar{x}$  = 15 × 4.2 µm, n = 40), overlapping, 1–2-seriate, fusi-




**Figure 4.** *Bertiella fici* (CMUB 40045, new host record) **a** appearance of ascomata on the host **b, c** close-up of ascomata **d** vertical section of an ascoma **e** peridium **f** pseudoparaphyses **g, h** asci **i–n** ascospores. Scale bars: 75 µm (**d**); 10 µm (**e**); 20 µm (**f–h**); 7 µm (**i–n**).

form, initially hyaline, becoming yellowish-brown at maturity, 1-septate, slightly curved, slightly constricted at the septum, guttulate, smooth-walled. **Asexual morph:** Undetermined.

**Material examined.** THAILAND, Chiang Rai, Doi Mae Salong Mountain, on a dead leaf of *Cinnamomum verum* (Lauraceae), 15 June 2020, D. S. Tennakoon, DMS002 (CMUB 40045).

**Known hosts.** *Cinnamomum verum* and *Ficus septica* (Tennakoon et al. 2021; this study).

**Known distribution.** China and Thailand (Tennakoon et al. 2021; this study)

**Notes.** *Bertiella fici* was introduced by Tennakoon et al. (2021) from dead leaves of *Ficus septica* in China. The morphological characteristics of our collection (CMUB 40045) tally well with the *B. fici* in having solitary or scattered, semi-immersed to superficial ascomata, cylindrical to cylindrical-clavate asci and yellowish-brown, 1-septate ascospores with overlapping size ranges (Tennakoon et al. 2021). Multi-gene phylogeny (LSU, SSU, ITS and *tef1-α*) also indicates that our collection nested with *B. fici* isolates in a 100% ML and 1.00 BYPP supported clade. This was further confirmed by having only two nucleotide differences in the LSU and SSU genes between our collection and the type of *Bertiella fici*. Thus, we conclude our collection as a new host record of *Bertiella fici* from *Cinnamomum verum*. In addition, this is the first *Bertiella fici* record from Thailand.

### ***Byssosphaeria* Cooke**

**Notes.** Cooke and Plowright (1879) established *Byssosphaeria* to accommodate *B. keithii* as the type species. *Byssosphaeria* species have superficial ascomata with bright yellow, orange or red flat apices around the ostiole, with dependent hyphal appendages that merge with the subiculum below and hyaline ascospores (Tian et al. 2015; Tennakoon et al. 2018). Species have cosmopolitan distribution as saprobes in various plant substrates (e.g. dead leaves, wood). As well, *Byssosphaeria* species have been reported from 15 plant families (Fig. 10). Currently, there are 18 accepted species in Species Fungorum (2024).

### ***Byssosphaeria poaceicola* Tennakoon & D.M. Hu, sp. nov.**

Index Fungorum: IF901733

Facesoffungi Number: FoF15542

Fig. 5

**Etymology.** Named after the host family (Poaceae) where this fungus was collected.

**Holotype.** HFJAU10338.

**Description.** **Saprobic** on dead stem of *Arundo pliniana* Turra (Poaceae). **Sexual morph: Ascomata** 550–650 × 600–800 µm ( $\bar{x}$  = 610 × 715 µm,  $n$  = 10), solitary to gregarious, superficial, dark brown to black, setose, coriaceous, unilocular, globose to subglobose, non-papillate, apex rounded with an orange to yellow ostiole, ostiole central, with pore-like opening, periphysate. **Peridium** 30–45 µm wide, thick-walled, composed of 6–7 layers of dark brown cells, orange to yellow near ostiole, arranged in **textura angularis**. **Hamathecium** 1–2.5 µm wide, com-




**Figure 5.** *Byssosphaeria poaceicola* (HFJAU10338, holotype) **a, b** appearance of ascomata on the host **c** vertical section of an ascoma **d** ostiole **e** peridium **f** pseudoparaphyses **g–i** asci **j–o** ascospores. Scale bars: 200  $\mu\text{m}$  (**c**); 20  $\mu\text{m}$  (**d, e**); 50  $\mu\text{m}$  (**f–i**); 10  $\mu\text{m}$  (**j–o**).

prising dense, filiform, anastomosing, septate, trabeculate pseudoparaphyses, embedded in a gelatinous matrix. **Asci** 165–180  $\times$  12–15  $\mu\text{m}$  ( $\bar{x}$  = 171  $\times$  13  $\mu\text{m}$ ,  $n$  = 20), 8-spored, bitunicate, fissitunicate, cylindrical clavate, apically rounded, long pedicellate (30–40  $\mu\text{m}$ ), with an indistinct ocular chamber. **Ascospores** 32–40  $\times$  7–8  $\mu\text{m}$  ( $\bar{x}$  = 36  $\times$  7.5  $\mu\text{m}$ ,  $n$  = 30), overlapping, 1–2-seriate, ellipsoid to fusiform, initially hyaline, pale brown when mature, 1-septate, constricted at the septum, slightly curved, guttulate, smooth-walled. **Asexual morph**: Undetermined.

**Material examined.** CHINA, Yunnan Province, Kunming, on a dead stem of *Arundo pliniana* (Poaceae), 22 July 2016, D. S. Tennakoon, KDS30 (HFJAU10338, **holotype**); *ibid.* 28 August 2016, KDS29 (HFJAU10337, **paratype**).

**Notes.** In the combined LSU, SSU, ITS and *tef1-a* phylogenetic analysis, two strains of *Byssosphaeria poaceicola* (HFJAU10337 and HFJAU10338) formed a monophyletic clade sister to *By. phoenicis* strains (ZHKUCC 21-0122 and ZHKUCC 21-012) with 83% ML and 0.99 BYPP statistical support. Morphologically,

they share similarities in having superficial, dark brown to black, coriaceous, non-papillate ascomata, cylindrical clavate asci and ellipsoid to fusiform, 1-septate, pale brown ascospores (Kularathnage et al. 2022). Although *Byssosphaeria poaceicola* can be distinguished from *By. phoenicis* in their size differences of asci ( $165\text{--}180 \times 12\text{--}15 \mu\text{m}$  vs.  $100\text{--}160 \times 10\text{--}15 \mu\text{m}$ ) and ascospores ( $32\text{--}40 \times 7\text{--}8 \mu\text{m}$  vs.  $25\text{--}30 \times 5\text{--}7 \mu\text{m}$ ) (Kularathnage et al. 2022). In addition, a comparison of the 497 nucleotides across the ITS (+5.8S) gene region of *By. poaceicola* and *By. phoenicis* shows 16 base pair differences (3.21%). It is interesting to notice that the *Byssosphaeria* species have not been collected much from Poaceae hosts, except for bamboo species (Jiang et al. 2022). Based on this finding, it appears that *Byssosphaeria* species can adapt to a variety of habitats, although there are limited studies to investigate their diversity on various hosts and regions.

***Byssosphaeria siamensis* Boonmee, Q. Tian & K.D. Hyde, Fungal Diversity 74: 283 (2015)**

Index Fungorum: IF551430

Facesoffungi Number: FoF01026

Fig. 6

**Description.** *Saprobic* on dead stem of *Citrus trifoliata* L. (Rutaceae). **Sexual morph:** **Ascomata**  $250\text{--}400 \times 300\text{--}500 \mu\text{m}$  ( $\bar{x} = 320 \times 410 \mu\text{m}$ ,  $n = 10$ ), solitary to gregarious, superficial, dark brown to black, setose, coriaceous, unilocular, globose to subglobose, non-papillate, apex rounded with an orange to yellow ostiole, ostiole central, with pore-like opening. **Peridium**  $20\text{--}35 \mu\text{m}$  wide, thick-walled, composed of several layers of dark brown cells, orange to yellow near ostiole, arranged in *textura angularis* to *textura prismatica*. **Hamathecium**  $1\text{--}2.5 \mu\text{m}$  wide, comprising dense, filiform, anastomosing, septate, pseudoparaphyses, embedded in a gelatinous matrix. **Asci**  $110\text{--}130 \times 11\text{--}13 \mu\text{m}$  ( $\bar{x} = 120 \times 12 \mu\text{m}$ ,  $n = 20$ ), 8-spored, bitunicate, fissitunicate, cylindrical clavate, apically rounded, long pedicellate ( $20\text{--}35 \mu\text{m}$ ), with an ocular chamber. **Ascospores**  $30\text{--}40 \times 6.5\text{--}8 \mu\text{m}$  ( $\bar{x} = 31 \times 7 \mu\text{m}$ ,  $n = 30$ ), overlapping, 1–2-seriate, ellipsoid to fusiform, initially hyaline, pale brown when mature, 1-septate, constricted at the septum, slightly curved, smooth-walled or verrucose. **Asexual morph:** Undetermined.

**Material examined.** CHINA, Yunnan Province, Kunming, on a dead stem of *Citrus trifoliata* (Rutaceae), 27 July 2016, D. S. Tennakoon, KDS27 (HFJAU10336).

**Known hosts.** *Citrus trifoliata* (this study).

**Known distribution.** China, Thailand (Tian et al. 2015; Hyde et al. 2018; this study).

**Notes.** The morphological characteristics of our collection (HFJAU10336) similar to the type of *Byssosphaeria siamensis* in their superficial, dark brown to black, setose, non-papillate ascomata, cylindrical clavate asci and ellipsoid to fusiform, pale brown, 1-septate ascospores (Tian et al. 2015). In addition, both share overlapping size ranges of asci ( $110\text{--}130 \times 11\text{--}13 \mu\text{m}$  vs.  $112\text{--}148 \times 10\text{--}16 \mu\text{m}$ ) and ascospores ( $30\text{--}40 \times 6.5\text{--}8 \mu\text{m}$  vs.  $40.5\text{--}50 \times 7\text{--}11 \mu\text{m}$ ) (Tian et al. 2015). However, our collection is lacking a mucilaginous sheath which is present in the type species (Tian et al. 2015). According to the multi-gene phylogeny, our collection nested with *By. siamensis* isolates in 86% ML and 0.98 BYPP supported




**Figure 6.** *Byssosphaeria siamensis* (HFJAU10336, new host record) **a** appearance of ascomata on the host **b** close-up of ascomata **c** vertical section of an ascoma **d** ostiole **e** peridium **f** pseudoparaphyses **g, h** asci **i–m** ascospores. Scale bars: 200  $\mu\text{m}$  (**c**); 20  $\mu\text{m}$  (**d, e**); 50  $\mu\text{m}$  (**f–h**); 10  $\mu\text{m}$  (**i–m**).

clade and close to the isolate MFLUCC 17-1800 with 88% ML, 0.99 BYPP support. Therefore, based on both morphology and phylogeny evidence, we introduce our collection as a new host record of *B. siamensis* from *Citrus trifoliata* in China.

### *Herpotrichia* Fuckel

**Notes.** The diverse genus *Herpotrichia* was established by Fuckel (1868) to include two species, *H. rhenana* and *H. rubi*, but without designating a type species. Thus, Bose (1961) assigned *H. rhenana* as the lectotype. Subsequently, Holm (1979) assigned *H. herpotrichoides* (synonymous with *H. rubi*) as the

generic type (Cannon 1982). The species have erumpent to superficial ascomata, clavate to cylindrical asci with hyaline to pale brown, 1-septate ascospores (Sivanesan 1984; Tian et al. 2015). The asexual morph is pyrenochaeta-like with or without setae on the surface of the pycnidia (Sivanesan 1984; Hongsanant et al. 2020). *Herpotrichia* species have been reported as saprobes, mostly in dead wood substrates in 34 plant families (Fig. 10).

***Herpotrichia zingiberacearum* Tennakoon & D.M. Hu, sp. nov.**

Index Fungorum: IF901734

Facesoffungi Number: FoF15543

Fig. 7

**Etymology.** Named after the host family (Zingiberaceae) where this fungus was collected.

**Holotype.** HFJAU10332.

**Description.** **Saprobic** on dead stem of *Hedychium coronarium* J. Koenig (Zingiberaceae). **Sexual morph: Ascomata** 250–350 × 240–320 µm ( $\bar{x}$  = 298 × 262 µm,  $n$  = 10), solitary to clustered, superficial, dark brown to black, setose, coriaceous, unilocular, globose to subglobose, rounded apex broadly cap-like, ostiolate. **Peridium** 15–25 µm wide, thick-walled, composed of 4–5 layers of dark brown to black cells, arranged in **textura angularis**. **Hamathecium** 1–3 µm wide, comprising dense, filiform, anastomosing, septate, branched pseudoparaphyses, embedded in a gelatinous matrix. **Asci** 85–98 × 10–14 µm ( $\bar{x}$  = 94 × 12 µm,  $n$  = 20), 8-spored, bitunicate, fissitunicate, cylindrical clavate, apically rounded, short pedicellate, with an ocular chamber. **Ascospores** 25–30 × 5–6 µm ( $\bar{x}$  = 28 × 5.2 µm,  $n$  = 30), overlapping, 1–2-seriate, fusoid with narrowly rounded ends, hyaline, 1-septate, straight to slightly curved, constricted at the septum, with large guttules, surrounded by an expanded gelatinous sheath pointed at both ends, 2.5–4 µm wide, smooth-walled. **Asexual morph:** Undetermined.

**Material examined.** CHINA, Yunnan Province, Kunming, on a dead stem of *Hedychium coronarium* (Zingiberaceae), 20 July 2016, D. S. Tennakoon, KDS12 (HFJAU10332, **holotype**); *ibid.* 21 August 2016, KDS13 (HFJAU10333, **paratype**); *ibid.* 25 August 2016, KDS18 (HFJAU10334, **paratype**).

**Notes.** *Herpotrichia zingiberacearum* is isolated from the dead stem of *Hedychium coronarium* (Zingiberaceae). The newly-generated sequences *H. zingiberacearum* (LSU, SSU, ITS and *tef1-α*) formed a monophyletic clade closely related to *H. macrotricha* with 90% ML and 0.99 BYPP statistical support. Morphologically, they share similarities in having dark brown to black, setose, coriaceous ascomata, cylindrical clavate asci and hyaline, 1-septate ascospores (Mugambi and Huhndorf 2009). However, *H. zingiberacearum* can be distinguished from *H. macrotricha* in their smaller asci (85–98 × 10–14 µm vs. 115–145 × 11–13 µm) and ascospores (25–30 × 5–6 µm vs. 30–35 × 4–6 µm) (Tanaka and Hosoya 2006). On the other hand, *H. zingiberacearum* (HFJAU10332) differs from *H. macrotricha* (GKM 196N) by a comparison of the 694 nucleotides across the *tef1-α* gene region which shows 20 base pair differences (3.02%). This finding addresses future studies to explore the diversity of *Herpotrichia* species in different geographic regions and host plants, as these species have not previously been described from Zingiberaceae hosts (Table 2).




**Figure 7.** *Herpotrichia zingiberacearum* (HFJAU10332, holotype) **a, b** appearance of ascomata on the host **c** vertical section of an ascoma **d** peridium **e** pseudoparaphyses **f–h** asci **i–l** ascospores **m** ascospore stained in Indian ink showing a mucilaginous sheath. Scale bars: 100 µm (c); 20 µm (d); 50 µm (e–h); 15 µm (i–m).

### ***Melanomma* Nitschke ex Fuckel**

**Notes.** *Melanomma* was validly established by Fuckel (1870) with *M. pulvis-pyrius* as the type species. These species have cosmopolitan distribution worldwide and characterised in having carbonaceous ascomata and hyaline or brown, 2–3-septate ascospores (Tian et al. 2015; Crous et al. 2020). The asexual morph can be either coelomycetes or hyphomycetes (Hyde et al. 2011). Currently, 94 species are accepted in this genus (Species Fungorum 2024) and have been reported from 39 plant families (Fig. 10).

### ***Melanomma populicola* Crous & R.K. Schumach, Fungal Systematics and Evolution 6: 201 (2020)**

Index Fungorum: IF552757

Facesoffungi Number: FoF2887

Fig. 8

**Basionym.** *Aposphaeria populina* Died., Krypt.-Fl. Brandenburg (Leipzig) 9: 206 (1912).

**Synonym.** *Melanomma populinum* (Died.) Phukhams. & K.D. Hyde [as '*populina*'], Fungal Diversity 83: 49. 2017.




**Figure 8.** *Melanomma populicola* (HFJAU10335, new host record) **a** appearance of ascomata on the host **b** close-up of ascoma **c** vertical section of an ascoma **d** peridium **e** pseudoparaphyses **f–h** asci **i–n** ascospores. Scale bars: 75  $\mu\text{m}$  (**c**); 10  $\mu\text{m}$  (**d**); 40  $\mu\text{m}$  (**e–h**); 10  $\mu\text{m}$  (**i–n**).

**Description.** *Saprobic* on dead stem of *Fagus sylvatica* L. (Fagaceae). **Sexual morph:** **Ascomata** 130–200  $\times$  200–300  $\mu\text{m}$  ( $\bar{x}$  = 150  $\times$  255  $\mu\text{m}$ ,  $n$  = 15), solitary or scattered, immersed, erumpent through host surface, black, multi-loculate, globose to subglobose, ostiolate. **Peridium** 10–15  $\mu\text{m}$  wide, thick-walled, carbonaceous, composed of several layers of light brown to dark brown pseudoparenchymatous cells, cells towards the inside hyaline, arranged in a *textura angularis*, fusing at the outside indistinguishable from the host tissues. **Hamathecium** comprising numerous, 1–2  $\mu\text{m}$  wide, hyaline, septate, filiform pseudoparaphyses. **Asci** 80–110  $\times$  6–8  $\mu\text{m}$  ( $\bar{x}$  = 96  $\times$  7  $\mu\text{m}$ ,  $n$  = 20), 8-spored, bitunicate, fissitunicate, cylindrical, apically rounded, short pedicellate with furcate end, with an indistinct ocular chamber. **Ascospores** 13–17  $\times$  4–5.2  $\mu\text{m}$  ( $\bar{x}$  = 15  $\times$  4.8  $\mu\text{m}$ ,  $n$  = 35), overlapping, 1–2-seriate, ellipsoid, initially hyaline, becoming light brown at maturity, 3-septate, straight to slightly curved, slightly constricted at the septa, guttulate, smooth-walled. **Asexual morph:** See Crous et al. (2020).

**Material examined.** CHINA, Kunming, on a dead stem of *Fagus sylvatica* (Fagaceae), 25 July 2016, D. S. Tennakoon, KDS25 (HFJAU10335).

**Known hosts.** *Populus canadensis*, *Picea abies*, *Quercus* sp., *Sorbus aucuparia* (Crous et al. 2020; this study).

**Known distribution.** China, Germany, The Netherlands (Crous et al. 2020; this study).

**Notes.** *Melanomma populicola* was introduced by Diedicke (1912) as *Aposphaeria populina*, which was a phoma-like species collected from *Populus canadensis*. An epitype for *A. populina* was established by De Gruyter et al. (2013). Tibpromma et al. (2017) synonymised *A. populina* under *Melanomma* and erected as *M. populina*. Subsequently, this was validated by Crous et al. (2020) and erected as *M. populicola*. Morphological characteristics of our collection (HFJAU10335) fit well with the *M. populicola* in having black, globose to subglobose ascomata, cylindrical, apically rounded asci and ellipsoid, 3-septate, light brown ascospores. In addition, there are overlapping size ranges of asci (80–110 × 6–8 µm vs. 91–106 × 6.5–7.5 µm) and ascospores (13–17 × 4–5.2 µm vs. 15.1 × 5 µm) (Crous et al. 2020). Multi-gene phylogeny also shows that our collection groups with *M. populicola* isolates in a 99% ML and 0.99 BYPP supported clade. Therefore, we introduce our collections as a new host record of *M. populicola* from *Fagus sylvatica*.

### Geographical distribution and host associations of melanommataceous species

Based on the data collected, it appears that the members of Melanommataceae are widely distributed around the world, comprising subtropical, tropical and temperate regions, such as Austria, Australia, Brazil, Canada, Finland, Germany, Japan, India, Thailand, Papua New Guinea, South Africa, Ukraine and the United States). The highest number of species have been reported from the United States (48 species). This is followed by China (35 species), Italy (30 species), India (25 species), Germany (23 species) and the United Kingdom (22 species) (indicated by red areas, Fig. 9). Moreover, Japan (17 species), Brazil (16 species), Ukraine (16 species), Canada (15 species), Kazakhstan (13 species), Poland (13 species) and Austria (11 species) are indicated by green areas (Fig. 9). Twenty-four countries have the species number range between 3 and 10, for instance Argentina, Russia and Sweden (10 species), Uzbekistan (9 species), Australia, France and Switzerland (8 species), Chile (7 species), Luxembourg, New Zealand, the Philippines and Thailand (6 species), Finland (5 species), Belgium and South Africa (4 species) and three species in nine countries (Azerbaijan, Czechia, Denmark, Georgia, Hungary, Papua New Guinea, Slovakia, Spain and Sri Lanka) (indicated by blue areas, Fig. 9). In addition, 36 countries reported two or a single species (indicated by yellow areas, Fig. 9). This may be because those countries have very limited taxonomic investigations on melanommataceous species. Consequently, it would be important to conduct more comprehensive collections, carry out more taxonomic studies and identify the species in those poorly-explored countries (e.g. African continent). In Asia, the highest number of melanommataceous species have reported from China, India and Thailand (Fig. 9), while Austria, Germany, Italy, Poland and the United Kingdom in Europe also have high numbers.




**Figure 9.** Distribution of so far reported species in Melanommataceae worldwide. Colour gradient shows the number of recorded species from lowest (yellow) to highest (red) and no records (grey).

Some species exist over multiple continents, while others appear to have limited distribution and are currently reported from one or a few countries (Fig. 9). Nine species of Melanommataceae are intercontinental and occur on more than five countries. Some highly diverse species include *Aposphaeria pulviscula*, *Camposporium cambrense*, *C. pellucidum*, *Herpotrichia herpotrichoides*, *H. nigra*, *H. quinquesepata*, *Melanomma pulvis-pyrius*, *M. rhododendri* and *Seifertia azaleae* (Table 2). As well, nine species have been reported from four countries (*Byssosphaeria jamaicana*, *By. rhodomphala*, *Camposporium antennatum*, *Herpotrichia macrotricha*, *Melanomma glumarum*, *M. subdispersum*, *Petrakia echinata*, *Phragmocephala elliptica* and *Phragmotrichum chailetii*) and nine species from three countries (*Aposphaeria eragrostidis*, *A. freticola*, *A. mediella*, *Camposporium hyderabadense*, *Melanomma medium*, *M. populicola*, *Navicella pileata*, *Phragmocephala stemphylioides* and *Phragmotrichum rivoclarinum*). In contrast, the distribution of 313 species has been limited to two or to one country, based on the current information. We believe that the distribution of those species may be much higher with future collections and taxonomical investigations.

When focusing on the host association of melanommataceous species, most of have been discovered in decaying wood or submerged woody substrates (e.g. *Aposphaeria corallinolutea*, *A. rudis*, *Asymmetricospora calamicola*, *Bertiella ellipsoidea*, *Byssosphaeria juniperi*, *Calyptronectria argentinensis*, *Camposporium chinense*, *C. marylandicum*, *Herpotrichia alpincola*, *Marjia uzbekistanica*, *Melanomma dinghuense*, *Phragmocephala garethjonesii* and *Sarimanas pseudofluviatile*) (Table 2). Some species have been recorded from decaying leaves (e.g. *Bertiella fici*, *Byssosphaeria musae*, *Phragmocephala elegans*). As

well, it is noteworthy to mention that some species have been recorded from soil (e.g. *Herpotrichia gelasinosporoides*, *H. striatispora* and *Pleotrichocladium opacum*). In addition, *Aposphaeria ramalinae* has been collected from a lichen species (*Ramalina implectens*) in France (Pitard and Harmand 1911) and *Exosporiella fungorum* from a mushroom species (*Thelephora* sp. fibre vase or earth fan mushroom) in Sweden (Karsten 1892).

According to the host associations, some species of Melanommataceae are highly diverse and have been collected from more than 10 host species (e.g. *Camposporium antennatum*, *C. japonicum*, *C. pellucidum*, *Herpotrichia macrotricha*, *H. nigra* and *Melanomma pulvis-pyrius*), while some are from more than five host species (e.g. *Camposporium cambrense*, *Herpotrichia herpotrichoides*, *Melanomma populicola* and *Phragmocephala elliptica*). The highest number of melanommataceous species have reported from the plant family Fagaceae (32 species). The most common Fagaceae hosts are *Fagus* spp. (e.g. *F. crenata*, *F. sylvatica*) and *Quercus* species (e.g. *Q. germana*). This is followed by Fabaceae (24 species), Rosaceae (23 species), Salicaceae (22 species), Poaceae (19 species), Pinaceae and Arecaceae (18 species), Sapindaceae (16 species), Betulaceae (15 species), Amaranthaceae (12 species), Asteraceae (10 species), Ericaceae (9 species), Cupressaceae, Euphorbiaceae and Lauraceae (7 species), Rutaceae (6 species) and Malvaceae, Oleaceae and Pandanaceae (5 species) (Fig. 10). In addition, a range of 2–5 species have recorded from 30 plant families and single melanommataceous species have reported from 35 plant families. Thus, up to date, the species of the family have been reported from 82 plant families. However, some species have been collected from unidentified hosts and, thus, their host association remained unresolved (e.g. *Aposphaeria anomala*, *A. pakistanica*, *Bertiella ellipsoidea*, *B. gelatinosa*, *Camposporium appendiculatum* and *C. chinense*). The host specificity for most of the species has not yet been clarified, as they have been recorded from various plant families. Though, it was noted by Jaklitsch and Voglmayr (2017) that *Praetumpfia obducens* may be host specific to *Fraxinus* species.

## Discussion

The members of Melanommataceae have been well-studied in last two decades, leading to many exciting discoveries (17 new genera and 76 new species), which may be mostly linked to the progress made in the DNA sequence data. The initial classification of these species was primarily based on morphological characteristics (e.g. globose or depressed ascomata, trabeculate pseudoparaphyses, fissitunicate asci, pigmented ascospores) associated with hand drawings (Saccardo 1883; Winter 1885; Morgan 1904; Spegazzini 1909; Shear 1941). However, morphology-based identification is challenging since most species can share similar characteristics, which may lead to misinterpretations. Thus, a combination of DNA-based molecular phylogeny and morphological traits has become a widely accepted tool for current fungal classifications (Tian et al. 2015; Hashimoto et al. 2017; Jaklitsch and Voglmayr 2017; Purahong et al. 2017; Kularathnage et al. 2022). Mostly, the large subunit (28S, LSU), small subunit (18S, SSU), internal transcribed spacers (ITS1–5.8S–ITS2), translation elongation factor 1 gene (*tef1-α*) and RNA polymerase second largest subunit (*rpb2*) molecular markers have been used in current phylogenetic




**Figure 10.** The melanommataceous species distribution through plant host families.

analyses of Melanommataceae (Tian et al. 2015; Tennakoon et al. 2018; Pem et al. 2019; Kularathnage et al. 2022).

Melanommataceae genera are highly varied, both morphologically and phylogenetically. Of them, some are highly diverse with numerous species (e.g. *Aposphaeria*: 83 species, *Melanomma*: 81 species, *Herpotrichia*: 61 species, *Camposporium*: 25 species, *Byssosphaeria*: 18 species and *Petrakia*: 10 species), while some have fewer species (*Phragmocephala*: 9 species, *Bertiella*: 6 species, *Navicella*: 5 species, *Phragmotrichum*: 5 species, *Uzbekistanica*: 4 species, *Calyptronectria*: 3 species, *Pseudostrickeria*: 3 species, *Seifertia*: 3 species, *Marjia*: 2 species, *Muriformistrickeria*: 2 species, *Sarimanas*: 2 species, and *Tumularia*: 2 species). In addition, some genera are monotypic and need more collections for their expansion (e.g. *Alpinaria*, *Asymmetricospora*, *Bicrouania*, *Dematiomelanomma*, *Exosporiella*, *Fusiconidium*, *Mamillisphaeria*, *Melanocamarosporioides*, *Melanocamarosporium*, *Melanocucurbitaria*, *Melanodiplodia*, *Monoseptella*, *Neobysosphaeria*, *Pleotrichocladium*, *Praetumpfia*, *Pseudobysosphaeria* and *Xenostigmina*). However, it is noteworthy to mention that some genera currently lack molecular data and, thus, their phylogenetic position is uncertain (e.g. *Asymmetricospora*, *Bicrouania*, *Calyptronectria*, *Exosporiella*, *Mamillisphaeria* and *Navicella*) (Hongsanan et al. 2020). As well, only two species of *Aposphaeria* have molecular data out of the 84 species listed in the Species Fungorum (2024). Consequently, to resolve their phylogenetic placements, further collections and investigations are essentially required.

In this study, we introduced two new species and three new host records collected from China and Thailand. The new species, *Byssosphaeria poaceicola* and *Herpotrichia zingiberacearum* can be distinguished from related species in their morphology and DNA molecular data. The morphological characteristics of the new host records strongly tally with their type species and phylogeny analyses also provide evidence for their placements. These new host records also demonstrate their adaptability to a broad range of habitats and there could be many more. Thus, many fungal species and host associations are waiting for us and we should undertake further explorations.

## Acknowledgements

Dr. Yang Gao is thanked for his support in depositing the herbarium specimens. The authors would like to thank Shaun Pennycook (Manaaki Whenua Landcare Research, New Zealand) for his guidance on the fungal nomenclature.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

No funding was reported.

## Author contributions

Conceptualisation, D.S.T. N.I.d.S. and D.M.H; methodology, D.S.T. and N.I.d.S; software, D.S.T. and N.I.S.; validation, K.M.T., N.I.d.S., H.Y.S., N.S., F.S.C. and D.M.H; formal analysis, D.S.T. and N.I.d.S; investigation, K.M.T., N.I.d.S., H.Y.S., N.S., F.S.C. and D.M.H; resources, D.S.T; data curation, D.S.T. and N.I.d.S; writing – original draft preparation, D.S.T., K.M.T., N.I.d.S., N.S. and D.M.H; writing – review and editing, K.M.T., N.I.d.S., H.Y.S., N.S., F.S.C. and D.M.H; supervision, D.M.H; project administration, D.M.H; funding acquisition, D.M.H. All authors have read and agreed to the published version of the manuscript.

## Author ORCIDs

Danushka S. Tennakoon  <https://orcid.org/0000-0003-2306-1255>

Kasun M. Thambugala  <https://orcid.org/0000-0002-6210-0504>

Nimali I. de Silva  <https://orcid.org/0000-0002-1577-280X>

Nakarin Suwannarach  <https://orcid.org/0000-0002-2653-1913>

Dian-Ming Hu  <https://orcid.org/0000-0002-4750-2871>

## Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Abdullah S (1980) Two hyphomycetes on litter in stagnant water from Britain. *Transactions of the British Mycological Society* 75(3): 514–517. [https://doi.org/10.1016/S0007-1536\(80\)80140-9](https://doi.org/10.1016/S0007-1536(80)80140-9)
- Adamcik S, Cai L, Chakraborty D, Chen XH, Cotter HVT, Dai DQ, Dai YC, Das K, Deng C, Ghobad-Nejhad M, Hyde KD, Langer E, Latha KPD, Liu F, Liu SL, Liu T, Lv W, Lv SX, Machado AR, Pinho DB, Pereira OL, Prasher IB, Rosado AWC, Qin J, Qin WM, Verma RK, Wang Q, Yang ZL, Yu XD, Zhou LW, Buyck B (2015) Fungal biodiversity profiles 1–10. *Cryptogamie. Mycologie* 36(2): 121–166. <https://doi.org/10.7872/crym/v36.iss2.2015.121>
- Ade A (1923) *Mykologische Beiträge*. *Hedwigia* 64: 286–320. <https://doi.org/10.2307/911161>
- Ahmad S (1964) Contributions to the fungi of West Pakistan. IV. *Biologia Lahore* 10: 1–6.
- Ahmad S (1971) Contributions to the fungi of West Pakistan. X. *Biologia Lahore* 17: 1–26.
- Allescher A (1896) Diagnosen einiger neuer, im Jahre 1895 gesammelter Arten Bayerischer Pilze aus der Abtheilung der Fungi Imperfecti. *Berichte der Bayerischen Botanischen Gesellschaft zur Erforschung der Heimischen Flora* 4: 31–40.
- Almeida DAC, Gusmão LFP, Miller AN (2017) A new species of *Bertiella* (Melanommataceae) from Brazil and a key to accepted species. *Mycosphere: Journal of Fungal Biology* 8(4): 392–396. <https://doi.org/10.5943/mycosphere/8/4/1>
- Anahosur KH (1970) Ascomycetes of Coorg (India). III. *Sydowia* 24: 177–182.
- Aptroot A (2003) Pyrenocarpous lichens and related non-lichenized ascomycetes from Taiwan. *The Journal of the Hattori Botanical Laboratory* 93: 155–173. [https://doi.org/10.18968/jhbl.93.0\\_155](https://doi.org/10.18968/jhbl.93.0_155)
- Aptroot A, Iperen A (1998) New ascomycetes and ascomycete records from Papua New Guinea. *Nova Hedwigia* 67(3–4): 481–497. <https://doi.org/10.1127/nova.hedwigia/67/1998/481>

- Bánki O, Roskov Y, Döring M, Ower G, Vandepitte L, Hobern D, Remsen D, Schalk P, De Walt RE, Keping M (2023) Catalogue of Life Checklist (Version 2023-01-12). Catalogue of Life. <https://doi.org/10.48580/dfqz>
- Barr ME (1968) The Venturiaceae in North America. *Canadian Journal of Botany* 46(6): 799–864. <https://doi.org/10.1139/b68-111>
- Barr ME (1983) Muriform ascospores in class ascomycetes. *Mycotaxon* 18: 149–157.
- Barr ME (1984) *Herpotrichia* and its segregates. *Mycotaxon* 20: 1–38.
- Barr ME (1990) Melanommatales (Loculoascomycetes). *North American Flora* 13: 1–129.
- Barr ME (1992) Notes on the Lophiostomataceae (Pleosporales). *Mycotaxon* 45: 191–221.
- Barr ME, Rogerson CT, Smith SJ, Haines JH (1986) An annotated catalog of the pyrenomycetes described by Charles H. Peck. *Bull. New York State Museum of Natural History* 459: 1–74. <https://doi.org/10.5962/bhl.title.135547>
- Bedlan G (2017) *Petrakia juniperi* sp. nov., ein neuer Pilz an *Juniperus* sp. *Journal für Kulturpflanzen* 69: 173–174.
- Beenken L, Gross A, Queloz V (2020) Phylogenetic revision of *Petrakia* and *Seifertia* (Melanommataceae, Pleosporales): New and rediscovered species from Europe and North America. *Mycological Progress* 19(5): 417–440. <https://doi.org/10.1007/s11557-020-01567-7>
- Berkeley MJ, Broome CE (1873) Enumeration of the fungi of Ceylon. Part II. *Journal of the Linnean Society. Botany* 14: 29–140. <https://doi.org/10.1111/j.1095-8339.1873.tb00301.x>
- Bhat DJ, Kendrick B (1993) Twenty-five new conidial fungi from the Western Ghats and the Andaman Islands (India). *Mycotaxon* 49: 19–90.
- Bommer E, Rousseau M (1890) Contributions a la Flore Mycologique de Belgique III. *Bulletin de la Société royale de Botanique de Belgique* 29: 205–302.
- Bose SK (1961) Studies on *Massarina* Sacc. and related genera. *Journal of Phytopathology* 41(2): 151–213. <https://doi.org/10.1111/j.1439-0434.1961.tb04048.x>
- Bresadola G (1926) *Selecta mycologica* II. *Student Trent* 7: 51–81.
- Bubák F (1914) Fungi in 'Wissenschaftliche Ergebnisse der Expedition nach Mesopotamien'. *Annalen des Naturhistorischen Museums in Wien* 28: 189–218.
- Bubák F, Kabát JE (1905) Vierter Beitrag zur Pilzflora von Tirol. *Österreichische. Botanisches Zhurnal* 55: 181–186. <https://doi.org/10.1007/BF01793320>
- Calabon MS, Jones EBG, Boonmee S, Doilom M, Lumyong S, Hyde KD (2021) Five novel freshwater ascomycetes indicate high undiscovered diversity in lotic habitats in Thailand. *Journal of Fungi (Basel, Switzerland)* 7(2): 1–27. <https://doi.org/10.3390/jof7020117>
- Calabon MS, Hyde KD, Jones EG, Luo ZL, Dong W, Hurdeal VG, Gentekaki E, Rossi W, Leonardi M, Thiyagaraja V, Lestari AS, Shen H-W, Bao D-F, Boonyuen N, Zeng M (2022) Freshwater fungal numbers. *Fungal Diversity* 114(1): 3–235. <https://doi.org/10.1007/s13225-022-00503-2>
- Cannon PF (1982) A note on the nomenclature of *Herpotrichia*. *Transactions of the British Mycological Society* 79(2): 338–339. [https://doi.org/10.1016/S0007-1536\(82\)80123-X](https://doi.org/10.1016/S0007-1536(82)80123-X)
- Cannon PF, Hawksworth DL, Sherwood-Pike MA (1985) *The British Ascomycotina. An Annotated Checklist*. Commonwealth Mycological Institute, Kew, Surrey, England: 302.
- Cash EK (1952) A record of the fungi named by JB Ellis (No. 2). *Division of Mycology and Disease Survey, Bureau of Plant Industry, Soils, and Agricultural Engineering*,

- Agricultural Research Administration, US Department of Agriculture. <https://doi.org/10.5962/bhl.title.149755>
- Cash EK (1953) A record of the fungi named by J.B. Ellis (Part 2). U.S.D.A. Special Publ. 2: 167–345.
- Castañeda RF (1985) Deuteromycotina de Cuba. Hyphomycetes III: 1–42.
- Castaneda-Ruiz RF, Iturriaga T, Minter DW, Saikawa M, Vidal G, Velazquez-Noa S (2003) Microfungi from Venezuela, A new species of *Brachydesmiella*, a new combination, and new records. Mycotaxon 85: 211–229.
- Castellani E, Ciferri R (1950) Mycoflora Erythraea, Somalia et Aethiopia Suppl. 1. Atti dell'Istituto Botanico e del Laboratorio Crittogamico dell'Università di Pavia, 52.
- Chen CY, Hsieh WH (2004) *Byssosphaeria* and *Herpotrichia* from Taiwan, with notes on the taxonomic relationship between these two genera. Sydowia 56: 24–38.
- Connors IL (1967) An Annotated Index of Plant Diseases in Canada and Fungi Recorded on Plants in Alaska, Canada and Greenland. Research Branch Canadian Department of Agriculture 1251: 1–381.
- Cooke MC (1887) Synopsis pyrenomycetum. Grevillea 15: 80–86. [https://doi.org/10.1175/1520-0493\(1887\)15\[86:V\]2.0.CO;2](https://doi.org/10.1175/1520-0493(1887)15[86:V]2.0.CO;2)
- Cooke MC, Plowright CB (1879) British Sphaeriacei. Grevillea 7: 77–89. <https://doi.org/10.1093/nq/s5-XII.292.89b>
- Costa MEA, Camara MS (1952) Species Aliquae Mycologicae Lusitaniae. Portugaliae Acta Biologica Series A 3: 294–307.
- Crous PW, Wingfield MJ, Schumacher RK, Akulov A, Bulgakov TS, Carnegie AJ, Jurjević Ž, De-cock C, Denman S, Lombard L, Lawrence DP, Stack AJ, Gordon TR, Bostock RM, Burgess T, Summerell BA, Taylor PWJ, Edwards J, Hou LW, Cai L, Rossman AY, Wöhner T, Allen WC, Castlebury LA, Visagie CM, Groenewald JZ (2020) New and Interesting Fungi. 3. Fungal Systematics and Evolution 6(1): 157–231. <https://doi.org/10.3114/fuse.2020.06.09>
- De Gruyter J, Woudenberg JHC, Aveskamp MM, Verkley GJM, Groenewald JZ, Crous PW (2013) Redisposition of Phoma-like anamorphs in Pleosporales. Studies in Mycology 75: 1–36. <https://doi.org/10.3114/sim0004>
- De Silva NI, Lumyong S, Hyde KD, Bulgakov T, Phillips AJL, Yan JY (2016) Mycosphere essays 9: Defining biotrophs and hemibiotrophs. Mycosphere: Journal of Fungal Biology 7(5): 545–559. <https://doi.org/10.5943/mycosphere/7/5/2>
- Dennis RWG (1978) British Ascomycetes. J. Cramer, Vaduz. 585.
- Dennis RWG (1986) Fungi of the Hebrides. Royal Botanic Gardens, Kew: 383.
- Diedicke H (1912) Sphaeropsidaeae. Kryptogamen–Flora der Mark Brandenburg. 9(1): 1–240.
- Dingley JM, Fullerton RA, McKenzie EHC (1981) Survey of Agricultural Pests and Diseases. Technical Report Volume 2. Records of Fungi, Bacteria, Algae, and Angiosperms Pathogenic on Plants in Cook Islands, Fiji, Kiribati, Niue, Tonga, Tuvalu, and Western Samoa. Food and Agriculture Organization, 485.
- Doidge EM (1948) South African Ascomycetes in the National Herbarium. Part VI. Botalia 4(4): 837–880. <https://doi.org/10.4102/abc.v4i4.1734>
- Dudka IO, Heluta VP, Tykhonenko YY, Andrianova TV, Hayova VP, Prydiuk MP, Dzhagan VV, Isikov VP (2004) Fungi of the Crimean Peninsula. M.G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine, 452.
- El-Buni AM, Rattan SS (1981) Check List of Libyan Fungi. Al Faateh University, Faculty of Science, Department of Botany, Tripoli, 169.
- El-Shafie AE, Aptroot A, Al-Bahry SN, Al-Kindi AY (2005) A new species of *Navicella* from Costa Rica. Sydowia 57: 19–22.

- Ellis JB (1895) Notes on some specimens of Pyrenomycetes in the Schweinitz herbarium of the Academy. *Proceedings. Academy of Natural Sciences of Philadelphia* 47: 20–31.
- Ellis MB (1976) Dematiaceous Hyphomycetes; Commonwealth Mycological Institute, Kew, UK, 344–353. <https://doi.org/10.1079/9780851983653.0000>
- Ellis JB, Everhart BM (1892) The North American Pyrenomycetes. Johnson Reprint Corporation. <https://doi.org/10.5962/bhl.title.55690>
- Ellis JB, Everhart BM (1902) New Alabama fungi. *Journal of Mycology* 8(2): 62–73. <https://doi.org/10.2307/3752297>
- Eriksson OE (2006) Outline of Ascomycota. *Myconet* 12: 1–82.
- Eriksson OE (2014) Checklist of the non-lichenized ascomycetes of Sweden. *Acta Universitatis Upsaliensis. Symbolae Botanicae Upsalienses* 36: 499.
- Eriksson OE, Yue JZ (1998) Bambusicolous pyrenomycetes, an annotated checklist. *Myconet* 1: 25–78.
- Fabre JH (1879) Essai sur les Sphériacées du Département de Vaucluse. *Annales des Sciences Naturelles Botanique Series* 6: 66–118.
- Fairman CE (1905) The Pyrenomyceteae of Orleans County, New York. *Proceedings of the Rochester Academy of Science*. 4: 165–191.
- Fairman CE (1921) The fungi of our common nuts and pits. *Proceedings of the Rochester Academy of Science*. 6: 73–115.
- Fairman CE (1922) New or rare fungi from various localities. *Proceedings of the Rochester Academy of Science* 6: 117–139.
- Farr ML (1973) An annotated list of Spegazzini's fungus taxa, Vol. 1. *Bibliotheca Mycologica* 35: 1–823.
- Farr DF, Rossman AY (2024) Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.ars-grin.gov/fungaldatabases> [accessed on 1 February 2024]
- Farr ML, Stevenson JA (1963) Eine ergänzungsliste bolivianischer pilze. *Sydowia* 17: 37–69.
- Feltgen J (1901) Vorstudien zu einer Pilzflora des Grossherzogthums Luxemburg. I. Theil. – Ascomycetes. *Nachträge II*: 1–243.
- Feltgen J (1903) Vorstudien zu einer Pilzflora des Grossherzogthums Luxemburg. I. Theil. – Ascomycetes. *Nachträge III*: 1–328.
- French AM (1989) California Plant Disease Host Index. Calif. Dept. Food Agric., Sacramento, 394.
- Fröhlich J, Hyde KD (1998) Fungi from palms. XXXIX. *Asymmetricospora* gen. et sp. nov. (Melanommataceae). *Sydowia* 50: 182–186.
- Fröhlich J, Hyde KD (2000) Palm Microfungi. *Fungal Diversity Research Series* 3: 1–393.
- Fuckel L (1868) Fungi rhenani exsic. 7: no. 2171.
- Fuckel L (1870) *Symbolae mycologicae*. Beiträge zur Kenntniss der Rheinischen Pilze. *Jahrbücher des Nassauischen Vereins für Naturkunde*. 23–24: 1–459. <https://doi.org/10.5962/bhl.title.47117>
- Fuckel L (1873) *Symbolae mycologicae*. Beiträge zur Kenntniss der rheinischen Pilze. Zweiter Nachtrag, 1–99.
- Gafforov YS (2017) A preliminary checklist of Ascomycetous microfungi from southern Uzbekistan. *Mycosphere: Journal of Fungal Biology* 8(4): 660–696. <https://doi.org/10.5943/mycosphere/8/4/12>
- Gao Y, Zhong T, Bhat JD, de Farias ARG, Dawoud TM, Hyde KD, Xiong W, Li Y, Gui H, Yang X, Wu S (2023) Pleomorphic *Dematiomelanomma yunnanense* gen. et sp. nov. (Asco-

- mycota, Melanommataceae) from grassland vegetation in Yunnan, China. *MycKeys* 98: 273–297. <https://doi.org/10.3897/mycokeys.98.107093>
- Ginns JH (1986) Compendium of plant disease and decay fungi in Canada 1960–1980. Research Branch Canadian Department of Agriculture 1813: 416. <https://doi.org/10.5962/bhl.title.58888>
- Glawe DA, Hummel RL (2006) North American records for *Seifertia azaleae*, cause of *Rhododendron* bud blight disease. *Pacific Northwest Fungi* 1: 1–6. <https://doi.org/10.2509/pnwf.2006.001.005>
- Gönczöl J (1976) *Monotosporella tuberculata*, a new species of aquatic Hyphomycetes from Hungary. *Nova Hedwigia* 27: 493–500.
- González Fragoso R (1918) *Pugillus secundus* mycetorum Persiae. *Boletín de la Real Sociedad Española de Historia Natural* 18: 78–85.
- Gross A, Beenken L, Dubach V, Queloz V, Tanaka K, Hashimoto A, Holdenrieder O (2017) *Pseudodidymella fagi* and *Petrakia deviata*: Two closely related tree pathogens new to central Europe. *Forest Pathology* 47(5): e12351. <https://doi.org/10.1111/efp.12351>
- Gucevic SA (1967) Species fungorum novae in plantis Tauriae endemicis collectae. *Novosti Sistematiki Nizshikh Rastenii* 4: 212–217.
- Gucevic SA (1969) De speciebus fungorum novis notula. *Novosti Sistematiki Nizshikh Rastenii* 6: 135–142.
- Gucevic SA (1977) De speciebus novis fungorum Pycnidialium e Tauria. *Novosti Sistematiki Nizshikh Rastenii* 14: 71–80.
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Harkness HW (1884) New species of California fungi. *Bulletin of the California Academy of Sciences* 1: 29–47.
- Hashimoto A, Matsumura M, Hirayama K, Fujimoto R, Tanaka K (2017) Pseudodidymellaceae fam. nov.: Phylogenetic affiliations of mycopappus-like genera in Dothideomycetes. *Studies in Mycology* 87(1): 187–206. <https://doi.org/10.1016/j.simyco.2017.07.002>
- Hawksworth DL (1985) A redistribution of the species referred to the ascomycete genus *Microthelia*. *Bulletin of the British Museum for Natural History* 14: 43–181.
- Hawksworth DL (2001) The magnitude of fungal diversity: The 1.5 million species estimate revisited. *Mycological Research* 105(12): 1422–1432. <https://doi.org/10.1017/S0953756201004725>
- Hayova VP, Minter DW (2009) *Aposphaeria ephedrae*. *Descriptions of Fungi and Bacteria Set* 180, 1792: 1–3. <https://doi.org/10.1079/DFB/20093210586>
- Hennings P (1903) Fungi Australienses. *Beiblatt. Hedwigia* 42: 73–88. <https://doi.org/10.1038/scientificamerican01311903-73>
- Hennings P (1908) Fungi paraënses III. *Hedwigia* 48: 101–117.
- Hernández-Restrepo M, Gené J, Castañeda-Ruiz RF, Mena-Portales J, Crous PW, Guarro J (2017) Phylogeny of saprobic microfungi from Southern Europe. *Studies in Mycology* 86: 53–97. <https://doi.org/10.1016/j.simyco.2017.05.002>
- Hill R, Leitch IJ, Gaya E (2021) Targeting Ascomycota genomes: What and how big? *Fungal Biology Reviews* 36: 52–59. <https://doi.org/10.1016/j.fbr.2021.03.003>
- Höhnelt F (1907) Ergebnisse der Botanischen Expedition der Kaiserlichen Akademie der Wissenschaften nach Süd-Brasilien 1901. II. Band. Thallophyta und Bryophyta. *Eumycetes et Myxomycetes*, 1–45.
- Höhnelt F (1909) Fragmente zur Mykologie. (VI. Mitteilung, Nr. 182 bis 288). *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften Math. naturw. Klasse Abt. I*. 118: 275–452.

- Holm L (1979) Index nominum genericorum (plantarum). In: Farr ER, Leussink JA, Staf-leu FA (Eds) The Hague, W. Junk, 189 pp.
- Holm L, Holm K (1988) Studies in the Lophiostomataceae with emphasis on the Swed-ish species. *Symbolae Botanicae Upsalienses* 28: 1–50.
- Holubtsova YI, Andrianova TV (2008) New Species of Anamorphic Fungi from Novgorod-Severskoe Poles'e (Ukraine). *Mikologiya i Fitopatologiya* 42: 28–34.
- Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie EHC, Sarma VV, Boonmee S, Lücking R, Pem D, Bhat DJ, Liu N, Tennakoon DS, Karunarathna A, Jiang SH, Jones EBG, Phillips AJL, Manawasinghe IS, Tibpromma S, Jayasiri SC, Sandamali D, Jayawardena RS, Wijayawardene NN, Ekanayaka AH, Jeewon R, Lu YZ, Dissanay-ake AJ, Zeng XY, Luo Z, Tian Q, Phukhamsakda C, Thambugala KM, Dai D, Chethana KWT, Ertz D, Doilom M, Liu JK, Pérez-Ortega S, Suija A, Senwanna C, Wijesinghe SN, Konta S, Niranjana M, Zhang SN, Ariyawansa HA, Jiang HB, Zhang JF, de Silva NI, Thi-yagaraja V, Zhang H, Bezerra JDP, Miranda-González R, Aptroot A, Kashiwadani H, Harishchandra D, Aluthmuhandiram JVS, Abeywickrama PD, Bao DF, Devadatha B, Wu HX, Moon KH, Gueidan C, Schumm F, Bundhun D, Mapook A, Monkai J, Chomnunti P, Samarakoon MC, Suetrong S, Chaiwan N, Dayarathne MC, Yang J, Rathnayaka AR, Bhunjun CS, Xu JC, Zheng J, Liu G, Feng Y, Xie N (2020) Refined families of Dothideo-mycetes: Dothideomycetidae and Pleosporomycetidae. *Mycosphere: Journal of Fun-gal Biology* 11(1): 1553–2107. <https://doi.org/10.5943/mycosphere/11/1/13>
- Hughes SJ (1951) Studies on Micro-fungi. III. *Mastigosporium*, *Camposporium*, and *Cer-atosporium*. *Mycological Papers* 36: 1–43.
- Hughes SJ (1955) Microfungi. I. Cordana, Brachysporium, Phragmocephala. *Canadian Journal of Botany* 33(3): 259–268. <https://doi.org/10.1139/b55-019>
- Hughes SJ (1958) Revisiones Hyphomycetum aliquot cum appendice de nominibus re-jiciendis. *Canadian Journal of Botany* 36(6): 727–836. <https://doi.org/10.1139/b58-067>
- Hughes SJ (1979) Relocation of species of Endophragmia auct. with notes on relevant generic names. *New Zealand Journal of Botany* 17(2): 139–188. <https://doi.org/10.1080/0028825X.1979.10426887>
- Hyde KD, Aptroot A (1998) A new species of *Herpotrichia* from the tropics. *Nova Hedwigia* 66(1–2): 247–249. <https://doi.org/10.1127/nova.hedwigia/66/1998/247>
- Hyde KD, Goh TK (1999) Some new melannommataceous fungi from woody substrata and a key to genera of lignicolous loculoascomycetes in freshwater. *Nova Hedwigia* 68: 251–272. <https://doi.org/10.1127/nova.hedwigia/68/1999/251>
- Hyde KD, McKenzie EHC, KoKo TW (2011) Towards incorporating anamorphic fungi in a natural classification checklist and notes for 2010. *Mycosphere: Journal of Fungal Biology* 2: 1–88.
- Hyde KD, Wong SW, Jones EBG (1996) Tropical Australian freshwater fungi. 11. *Mamilli-sphaeria dimorphospora* gen. et sp. nov. and notes on freshwater ascomycetes with dimoprphic ascospores. *Nova Hedwigia* 62: 513–520.
- Hyde KD, Goh TK, Lu BS, Alias SA (1999) Eleven new intertidal fungi from *Nypa fru-ticans*. *Mycological Research* 103(11): 1409–1422. <https://doi.org/10.1017/S0953756299008667>
- Hyde KD, Jones EBG, Liu JK, Ariyawansa HA, Boehm E, Boonmee S, Braun U, Chomnunti P, Crous PW, Dai DQ, Diederich P, Dissanayake AJ, Doilom M, Doveri F, Hongsanan S, Jayawardena RS, Lawrey JD, Li YM, Liu YX, Lücking R, Monkai J, Muggia L, Nelsen MP, Pang KL, Phookamsak R, Senanayake IC, Shearer CA, Suetrong S, Tanaka K, Thambu-gala KM, Wijayawardene NN, Wikee S, Wu HX, Zhang Y, Begoña AH, Alias SA, Aptroot

- A, Bahkali AH, Bezerra JL, Bhat DJ, Camporesi E, Chuksa E, Gueidan C, Hawksworth DL, Hirayama K, Hoog SD, Kang JK, Knudsen K, Li WJ, Li XH, Liu ZY, Mapook A, McKenzie EHC, Miller AN, Mortimer PE, Phillips AJL, Raja HA, Scheuer C, Schumm F, Taylor JE, Tian Q, Tibpromma S, Wanasinghe DN, Wang Y, Xu JC, Yacharoen S, Yan JY, Zang M (2013) Families of Dothideomycetes. *Fungal Diversity* 63(1): 1–313. <https://doi.org/10.1007/s13225-013-0263-4>
- Hyde KD, Hongsanan S, Jeewon R, Bhat DJ, McKenzie EHC, Jones EBG, Phookamsak R, Ariyawansa HA, Boonmee S, Zhao Q, Abdel-Aziz FA, Abdel-Wahab MA, Banmai S, Chomnunti P, Cui BK, Daranagama DA, Das K, Dayarathne MC, de Silva NI, Dissanayake AJ, Doilom M, Ekanayaka AH, Gibertoni TB, Go'es-Neto A, Huang SK, Jayasiri SC, Jayawardena RS, Konta S, Lee HB, Li WJ, Lin CG, Liu JK, Lu YZ, Luo ZL, Manawasinghe IS, Manimohan P, Mapook A, Niskanen T, Norphanphoun C, Papizadeh M, Perera RH, Phukhamsakda C, Richter C, de Santiago ALCMA, Drechsler-Santos ER, Senanayake IC, Tanaka K, Tennakoon TMD, Thambugala KM, Tian Q, Tibpromma S, Thongbai B, Vizzini A, Wanasinghe DN, Wijayawardene NN, Wu HX, Yang J, Zeng XY, Zhang H, Zhang JF, Bulgakov TS, Camporesi E, Bahkali AH, Amoozegar AM, Araujo-Neta LS, Ammirati JF, Baghela A, Bhatt RP, Bojantchev S, Buyck B, da Silva GA, de Lima CLF, de Oliveira RJV, de Souza CAF, Dai YC, Dima B, Duong TT, Ercole E, Mafalda-Freire F, Ghosh A, Hashimoto A, Kamolhan S, Kang JC, Karunarathna SC, Kirk PM, Kyto-vuori I, Lantieri A, Liimatainen K, Liu ZY, Liu XZ, Lu'cking R, Medardi G, Mortimer PE, Nguyen TTT, Promputtha I, Raj KNA, Reck MA, Lumyong S, Shahzadeh-Fazeli SA, Stadler M, Soudi MR, Su HY, Takahashi T, Tangthirasun N, Uniyal P, Wang Y, Wen TC, Xu JC, Zhang ZK, Zhao YC, Zhou JZ, Zhu L (2016) Fungal diversity notes 367–490: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 80: 1–270. <https://doi.org/10.1007/s13225-016-0373-x>
- Hyde KD, Chaiwan N, Norphanphoun C, Boonmee S, Camporesi E, Chethana KWT, Dayarathne MC, de Silva NI, Dissanayake AJ, Ekanayaka AH, Hongsanan S, Huang SK, Jayasiri SC, Jayawardena RS, Jiang HB, Karunarathna A, Lin CG, Liu JK, Liu NG, Lu YZ, Luo ZL, Maharachchimbura SSN, Manawasinghe IS, Pem D, Perera RH, Phukhamsakda C, Samarakoon MC, Senwana C, Shang QJ, Tennakoon DS, Thambugala KM, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Zhang JF, Zhang SN, Bulgakov TS, Bhat DJ, Cheewangkoon R, Goh TK, Jones EBG, Kang JC, Jeewon R, Liu ZY, Lumyong S, Kuo CH, McKenzie EHC, Wen TC, Yan JY, Zhao Q (2018) Mycosphere notes 169–224. *Mycosphere: Journal of Fungal Biology* 9(2): 271–430. <https://doi.org/10.5943/mycosphere/9/2/8>
- Hyde KD, Dong Y, Phookamsak R, Jeewon R, Bhat DJ, Jones EBG, Liu NG, Abeywickrama PD, Mapook A, Wei D, Perera RH, Manawasinghe IS, Pem D, Bundhun D, Karunarathna A, Ekanayaka AH, Bao DF, Li J, Samarakoon MC, Chaiwan N, Lin CG, Phutthacharoen K, Zhang SN, Senanayake IC, Goonasekara ID, Thambugala KM, Phukhamsakda C, Tennakoon DS, Jiang HB, Yang J, Zeng M, Huanraluek N, Liu JK, Wijesinghe SN, Tian Q, Tibpromma S, Brahmanage RS, Boonmee S, Huang SK, Thiagaraja V, Lu YZ, Jayawardena RS, Dong W, Yang EF, Singh SK, Singh SM, Rana S, Lad SS, Anand G, Devadatha B, Niranjana M, Sarma VV, Liimatainen K, Aguirre-Hudson B, Niskanen T, Overall A, Alvarenga RLM, Gibertoni TB, Pfliegler WP, Horváth E, Imre A, Alves AL, da Silva Santos AC, Tiago PV, Bulgakov TS, Wanasinghe DN, Bahkali AH, Doilom M, Elgorban AM, Maharachchikumbura SSN, Rajeshkumar KC, Haelewaters D, Mortimer PE, Zhao Q, Lumyong S, Xu J, Sheng J (2020) Fungal diversity notes 1151–1276: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 100(1): 5–277. <https://doi.org/10.1007/s13225-020-00439-5>

- Hyde KD, Suwannarach N, Jayawardena RS, Manawasinghe IS, Liao CF, Doilom M, Cai L, Zhao P, Buyck B, Phukhamsakda C, Su WX, Fu YP, Li Y, Zhao RL, He MQ, Li JX, Tibpromma S, Lu L, Tang X, Kang JC, Ren GC, Gui H, Hofstetter V, Ryoo R, Antonín V, Hurdeal VG, Gentikaki E, Zhang JY, Lu YZ, Senanayake IC, Yu FM, Zhao Q, Bao DF (2021) Mycosphere notes 325–344–Novel species and records of fungal taxa from around the world. *Mycosphere: Journal of Fungal Biology* 12(1): 1101–1156. <https://doi.org/10.5943/mycosphere/12/1/14>
- Ichinoe M (1971) *Camposporium* species from Japan. *Transactions of the British Mycological Society Japan* 12: 79–88.
- Inderbitzin P, Huang ZL (2001) *Melanomma dinghuense*, a new loculoascomycete with Munk pore-like perforations from Dinghushan Biosphere Reserve in Southern China. *Mycoscience* 42(2): 187–191. <https://doi.org/10.1007/BF02464136>
- Index Fungorum (2024) Index Fungorum. <http://www.indexfungorum.org/names/Names.asp> [accessed 6 January 2024]
- Jaklitsch WM, Voglmayr H (2017) Three former taxa of *Cucurbitaria* and considerations on *Petrakia* in the Melanommataceae. *Sydowia* 69: 81–95. <https://doi.org/10.12905/0380.sydowia69-2017-0081>
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat DJ, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang KA, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsanan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The faces of fungi database: Fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74(1): 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jiang H, Phookamsak R, Hongsanan S, Bhat DJ, Mortimer PE, Suwannarach N, Karkumyan P, Xu J (2022) A review of bambusicolous Ascomycota in China with an emphasis on species richness in southwest China. *Studies in Fungi* 7(1): 1–33. <https://doi.org/10.48130/SIF-2022-0020>
- Jones EG, Vrijmoed LL (2003) Biodiversity of marine fungi in Hong Kong coastal waters. *Perspectives on Marine Environment Change in Hong Kong and Southern China*. Hong Kong University Press, Hong Kong, 75–92.
- Jones EG, Pang KL, Abdel-Wahab MA, Scholz B, Hyde KD, Boekhout T, Ebel R, Rateb ME, Henderson L, Sakayaroj J, Suetrong S, Dayarathne MC, Kumar V, Raghukumar S, Sridhar KR, Bahkali AHA, Gleason FH, Norphanphoun C (2019) An online resource for marine fungi. *Fungal Diversity* 96(1): 347–433. <https://doi.org/10.1007/s13225-019-00426-5>
- Karsten PA (1884) *Fragmenta mycologica XIV*. *Hedwigia* 23: 57–59.
- Karsten PA (1892) *Finlands mögelsvampar (Hyphomyctes fennici)*. *Bidrag till Kännedom av Finlands Natur och. Folk (Kobenhavn)* 51: 343–534.
- Karsten PA (1905) *Fungi novi nonnullis exceptis in Fennia lecti*. *Acta Societatis pro Fauna et Flora Fennica* 27: 1–16.
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kirk PM, Spooner BM (1984) An account of the fungi of Arran, Gigha and Kintyre. *Kew Bulletin* 38(4): 503–597. <https://doi.org/10.2307/4108573>

- Kirschstein W (1906) Neue märkische Ascomyceten. Verhandlungen des Botanischen Vereins der Provinz Brandenburg. 48: 39–61.
- Kirschstein W (1911) Sphaeriales. Kryptogamen–Flora der Mark Brandenburg 7: 164–304.
- Kirschstein W (1935) Neue und seltene Ascomyceten. Annales Mycologici 33: 202–229.
- Kobayasi Y (1971) Mycological reports from New Guinea and the Solomon Islands (1–11). Bulletin of the National Science Museum, Tokyo 14: 367–551.
- Kohlmeyer J, Volkmann-Kohlmeyer B (1990) Revision of marine species of *Didymosphaeria* (Ascomycotina). Mycological Research 94(5): 685–690. [https://doi.org/10.1016/S0953-7562\(09\)80669-2](https://doi.org/10.1016/S0953-7562(09)80669-2)
- Koukol O, Delgado G (2021) Why morphology matters: The negative consequences of hasty descriptions of putative novelties in asexual ascomycetes. IMA Fungus 12(1): 1–8. <https://doi.org/10.1186/s43008-021-00073-z>
- Kuhnholz-Lordat G, Barry JP (1949) Quelques micromycètes parasites intéressants du sud de la France. Bulletin de la Société Mycologique de France 65: 121–131.
- Kularathnage ND, Tennakoon DS, Zhu X, Zhou J, Su B, Xie Y, Chen Q, Calabon MS, Kirk PM, Senanayake IC, Doilom M, Xu B, Dong W, Song J (2022) Reinstating *Dyfrlomycetes* and introducing *Melomastia pyriformis* sp. nov. (Pleurotremataceae, Dyfrlomycetales) from Guangdong Province, China. Current Research in Environmental & Applied Mycology 13(1): 426–438. <https://doi.org/10.5943/cream/13/1/16>
- Kulkarni RL (1974) Three fungi from groundnut leaf surface. Current Science 43: 561–562. [https://doi.org/10.1016/0039-6028\(74\)90276-3](https://doi.org/10.1016/0039-6028(74)90276-3)
- Kunze G, Schmidt JC (1823). Mykologische (2): 1–176. [https://doi.org/10.1016/S0140-6736\(02\)95514-5](https://doi.org/10.1016/S0140-6736(02)95514-5)
- Li WY, Zhuang WY (2008) Notes on the genus *Byssosphaeria* (Melanommataceae) from China. Junwu Xuebao 27: 48–53.
- Li GJ, Hyde KD, Zhao RL, Hongsanan S, Abdel-Aziz FA, AbdelWahab MA, Alvarado P, Alves-Silva G, Ammirati JF, Ariyawansa HA, Baghela A, Bahkali AH, Beug M, Bhat DJ, Bojantchev D, Boonpratuang T, Bulgakov TS, Camporesi E, Boro MC, Ceska O, Chakraborty D, Chen JJ, Chethana KWT, Chomnunti P, Consiglio G, Cui BK, Dai DQ, Dai YC, Daranagama DA, Das K, Dayarathne MC, De Crop E, De Oliveira RJV, de Souza CAF, de Souza JI, Dentinger BTM, Dissanayake AJ, Doilom M, Drechsler-Santos ER, Gho-bad-Nejhad M, Gilmore SP, Góes-Neto A, Gorczak M, Haitjema CH, Hapuarachchi KK, Hashimoto A, He MQ, Henske JK, Hirayama K, Iribarren MJ, Jayasiri SC, Jayawardena RS, Jeon SJ, Jerônimo GH, Jesus AL, Jones EBG, Kang JC, Karunarathna SC, Kirk PM, Konta S, Kuhnert E, Langer E, Lee HS, Lee HB, Li WJ, Li XH, Liimatainen K, Lima DX, Lin CG, Liu JK, Liu XZ, Liu ZY, Luangsa-ard JJ, Lücking R, Lumbsch HT, Lumyong S, Leaño EM, Marano AV, Matsumura M, McKenzie EHC, Mongkolsamrit S, Mortimer PE, Nguyen TTT, Niskanen T, Norphanphoun C, O'Malley MA, Parnmen S, Pawłowska J, Perera RH, Phookamsak R, Phukhamsakda C, Pires-Zottarelli CLA, Raspé O, Reck MA, Rocha SCO, de Santiago ALCMA, Senanayake IC, Setti L, Shang QJ, Singh SK, Sir EB, Solomon KV, Song J, Srikitkulchai P, Stadler M, Suetrong S, Takahashi H, Takahashi T, Tanaka K, Tang LP, Thambugala KM, Thanakitpipattana D, Theodorou MK, Thongbai B, Thummarukcharoen T, Tian Q, Tibpromma S, Verbeken A, Vizzini A, Vlasák J, Voigt K, Wanasinghe DN, Wang Y, Weerakoon G, Wen HA, Wen TC, Wijayawardene NN, Wongkanoun S, Wrzosek M, Xiao YP, Xu JC, Yan JY, Yang J, Da Yang S, Hu Y, Zhang JF, Zhao J, Zhou LW, Peršoh D, Phillips AJL, Maharachchikumbura SSN (2016a) Fungal diversity notes 253–366: Taxonomic and phylogenetic contributions to fungal taxa. Fungal Diversity 78(1): 1–237. <https://doi.org/10.1007/s13225-016-0366-9>

- Li JF, Phookamsak R, Mapook A, Boonmee S, Bhat JD, Hyde KD, Lumyong S (2016b) *Seifertia shangrilaensis* sp. nov. (Melanommataceae), a new species from Southwest China. *Phytotaxa* 273(1): 34–42. <https://doi.org/10.11646/phytotaxa.273.1.3>
- Li JF, Jeewon R, Luo ZL, Phookamsak R, Bhat DJ, Mapook A, Phukhamsakda C, Camporesi E, Lumyong S, Hyde KD (2017) Morphological characterization and DNA based taxonomy of *Fusiconidium* gen. nov. with two novel taxa within Melanommataceae (Pleosporales). *Phytotaxa* 308(2): 206–218. <https://doi.org/10.11646/phytotaxa.308.2.2>
- Liew ECY, Aptroot A, Hyde KD (2000) Phylogenetic significance of the pseudoparaphyses in Loculoascomycete taxonomy. *Molecular Phylogenetics and Evolution* 16(3): 392–402. <https://doi.org/10.1006/mpev.2000.0801>
- Lind J (1913) Danish fungi as represented in the herbarium of E. Rostrup, 1–648. <https://doi.org/10.5962/bhl.title.54591>
- Liu JK, Hyde KD, Jones EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SSN, McKenzie EHC, Phookamsak R, Phukhamsakda C, Shenoy BD, Abdel-Wahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dissanayake AJ, Doilom M, Dsouza MJ, Fan XL, Goonasekara ID, Hirayama K, Hongsanan S, Jayasiri SC, Jayawardena RS, Karunarathna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Peršoh D, Pinruan U, Senanayake IC, Somrithipol S, Suetrong S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D, Wijayawardene NN, Wanasinghe D, Wisitrassameewong K, Zeng XY, Abdel-Aziz FA, Adamčík S, Bahkali AH, Boonyuen N, Bulgakov T, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Lewis J, Li XH, Liu XZ, Liu ZY, Matsumura M, Mortimer PE, Rambold G, Randrianjohany E, Sato G, Sri-Indrasudthi V, Tian CM, Verbeken A, von Brackel W, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi E (2015) Fungal diversity notes 1–110: Taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72(1): 1–197. <https://doi.org/10.1007/s13225-015-0324-y>
- Lobik A (1927) Materialien zur Mykoflora des Terski Kreises. *Bolezni Rast.* 16: 157–199.
- Lu B, Hyde KD, Ho WH, Tsui KM, Taylor JE, Wong KM, Yanna K, Zhou D (2000) Checklist of Hong Kong Fungi. Fungal Diversity Press, Hong Kong, 207.
- Lunghini D, Granito VM, Di Lonardo DP, Maggi O, Persiani AM (2013) Fungal diversity of saprotrophic litter fungi in a Mediterranean maquis environment. *Mycologia* 105(6): 1499–1515. <https://doi.org/10.3852/13-103>
- Marvanová L, Descals E (1987) New taxa and new combinations of ‘aquatic hyphomycetes’. *Transactions of the British Mycological Society* 89(4): 499–507. [https://doi.org/10.1016/S0007-1536\(87\)80084-0](https://doi.org/10.1016/S0007-1536(87)80084-0)
- Mason EW, Hughes SJ (1951) *Phragmocephala* gen. nov. Hyphomycetorum. *The Naturalist London*, 97–105.
- Massee GE (1906) Fungi exotici, V. *Bulletin of Miscellaneous Informations of the Royal Botanical Gardens Kew*, 256–258. <https://doi.org/10.2307/4118220>
- Mathur RS (1979) The Coelomycetes of India. Bishen Singh Mahendra Pal Singh, Delhi, India, 460.
- Matsushima T (1980) Matsushima Mycological Memoirs No. 1. Saprophytic Microfungi from Taiwan, Part 1. Hyphomycetes. Matsushima Fungus Collection, Kobe, Japan, 82.
- Matsushima T (1983) Matsushima Mycological Memoirs 3. Matsushima Mycological Memoirs 3: 1–90. <https://doi.org/10.2307/3793127>
- McKenzie EHC, O’Sullivan PJ, Wilkie JP (1992) A list of type specimens of New Zealand fungi held in DSIR Plant Protection Herbarium (PDD). *Mycotaxon* 43: 77–156.

- Medel R (2007) First record of the genus *Byssosphaeria* (Pleosporales) in Mexico and Venezuela. *Mycotaxon* 100: 247–250.
- Melnik VA (1984) Coelomycetes of the SSSR. II. *Nagrajomyces dictyosporus* Melnik gen. et sp. nov. and *Phragmotrichum vassiljevae* Melnik sp.nov. *Mikologiâ i Fitopatologiâ* 18: 9–14.
- Mercado Sierra A, Heredia G, Mena Portales J (1995) New species of dematiaceous hyphomycetes from Veracruz, Mexico. *Mycotaxon* 55: 491–499.
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for Inference of Large Phylogenetic Trees. SC10 Workshop on Gateway Computing Environments (GCE10). <https://doi.org/10.1109/GCE.2010.5676129>
- Minter DW (1981) Microfungi on needles, twigs and cones of pines in Czechoslovakia. *Česká Mykologie* 35: 90–101.
- Morgan AP (1904) New species of Pyrenomycetes. *Journal of Mycology* 10(4): 161–162. <https://doi.org/10.2307/3752680>
- Mouton V (1900) Quatrième notice sur des ascomycètes nouveaux ou peu connus. *Bulletin de la Société Royale de Botanique de Belgique*. 39: 37–53.
- Mueller E (1958) Pilze aus dem Himalaya II. *Sydowia* 12: 160–184.
- Mugambi GK, Huhndorf SM (2009) Molecular phylogenetics of pleosporales: Melanommataceae and Lophiostomataceae re-circumscribed (Pleosporomycetidae, Dothideomycetes, Ascomycota). *Studies in Mycology* 64: 103–121. <https://doi.org/10.3114/sim.2009.64.05>
- Mujica F, Vergara C (1945) Flora fungosa Chilena. Índice preliminar de los huéspedes de los hongos chilenos y sus referencias bibliográficas. Imprenta Stanley, 199.
- Mulenko W, Majewski T, Ruszkiewicz-Michalska M (2008) A preliminary checklist of Micromycetes in Poland. *W. Szafer Institute of Botany. Polish Academy of Sciences* 9: 752.
- Munk A (1957) Danish Pyrenomycetes. A Preliminary Flora. *Dansk Botanisk Arkiv* 17: 1–491.
- Niranjan M, Sarma VV (2019) New species and new records of Melanommataceae (Pleosporales) from Andaman Islands. *Kavaka* 52: 103–108. <https://doi.org/10.36460/Kavaka/54/2020/38-42>
- Nylander JAA (2004) MrModeltest 2.0. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Oudemans CAJA (1902) Contributions á la flore mycologique des Pays-Bas. XVIII. *Nederlandsch Kruidkundig Archief* 2: 633–781.
- Pande A (1979) Ascomycetes of Western India VI. *Acta Botanica Indica* 7: 181–183.
- Pande A (2008) Ascomycetes of Peninsular India. Scientific Publishers (India), Jodhpur, 584.
- Pantidou ME (1973) Fungus-host index for Greece. *Benaki Phytopathol. Institute of Kiphissia, Athens*, 382.
- Papendorf MC, von Arx JA (1966) *Herpotrichia striatispora* a new ascomycete from South Africa. *Nova Hedwigia* 12: 395–397.
- Partridge EC, Morgan-Jones G (2002) Notes on hyphomycetes LXXXVIII: New genera in which to classify *Alysidium resinae* and *Pycnostysanus azaleae*, with a consideration of *Sorocybe*. *Mycotaxon* 83: 335–352.
- Peck CH (1909) New species of fungi. *Bulletin of the Torrey Botanical Club* 36(3): 153–157. <https://doi.org/10.2307/2479189>
- Pem D, Jeewon R, Gafforov Y, Hongsanan S, Phukhamsakda C, Promputtha I, Doilom M, Hyde KD (2019) *Melanocamarosporioides ugamica* gen. et sp. nov., a novel mem-

- ber of the family Melanommataceae from Uzbekistan. *Mycological Progress* 18(3): 471–481. <https://doi.org/10.1007/s11557-018-1448-8>
- Petch T (1912) Revisions of Ceylon fungi (Part III). *Annals of the Royal Botanic Gardens Peradeniya* 5: 265–301.
- Petch T (1917) Additions to Ceylon fungi. *Annals of the Royal Botanic Gardens Peradeniya* 6: 195–256. <https://doi.org/10.2307/4111564>
- Petrak F (1962) *Mykologische Beitrage zur osterreichischen Flora*. *Sydowia* 16: 155–198.
- Peyronel B (1915) Primo Elenco di Funghi di val San Martino o valle della Germanasca. *Memorie della Reale Accademia delle Scienze di Torino Series* 66: 1–58.
- Peyronel B (1918) Micromiceti di Val Germanasca. *Nuovo Giornale Botanico Italiano*. 25: 405–464.
- Phukhamsakda C, McKenzie EHC, Phillips AJL, Jones EBG, Bhat DJ, Stadler M, Bhunjun CS, Wanasinghe DN, Thongbai B, Camporesi E, Ertz D, Jayawardena RS, Perera RH, Ekanayake AH, Tibpromma S, Doilom M, Xu JC, Hyde KD (2020) Microfungi associated with *Clematis* (Ranunculaceae) with an integrated approach to delimiting species boundaries. *Fungal Diversity* 102(1): 1–203. <https://doi.org/10.1007/s13225-020-00448-4>
- Pimenta LP, Gomes DC, Cardoso PG, Takahashi JA (2021) Recent findings in azaphilone pigments. *Journal of Fungi* (Basel, Switzerland) 7(7): e541. <https://doi.org/10.3390/jof7070541>
- Pirozynski KA (1972) Microfungi of Tanzania. I. Miscellaneous fungi on oil palm. *Mycological Paper* 129: 1–39.
- Pitard CJ, Harmand J (1911) Contribution à l'étude des lichens des Îles Canaries. *Bulletin de la Société Botanique de France* 58 (Mém. 22): 1–72. <https://doi.org/10.1080/00378941.1911.10829212>
- Purahong W, Pietsch KA, Lentendu G, Schöps R, Bruelheide H, Wirth C, Buscot F, Wubet T (2017) Characterization of unexplored deadwood mycobiome in highly diverse subtropical forests using culture-independent molecular technique. *Frontiers in Microbiology* 8: e574. <https://doi.org/10.3389/fmicb.2017.00574>
- Rabenhorst GL (1869) *Fungi Europaei exsiccati, Klotzschii herbarii vivi mycologici continuatio*. Editio nov. Series secunda. Cent. 13(1201–no): 1300.
- Raciborski M (1909) Parasitische und epiphytische Pilze Javas. *Bulletin International de l'Academie des Sciences de Cracovie Classe des Sciences Mathematiques et Naturelles*. 3: 346–394.
- Raja HA, Paguigan ND, Fournier J, Oberlies NH (2017) Additions to *Lindgomyces* (Lindgomycetaceae, Pleosporales, Dothideomycetes), including two new species occurring on submerged wood from North Carolina, USA, with notes on secondary metabolite profiles. *Mycological Progress* 16(5): 535–552. <https://doi.org/10.1007/s11557-017-1282-4>
- Rambaut A (2012) FigTree Version 1.4.0. Edinburgh: Institute of Evolutionary Biology; University of Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/> [accessed January 10, 2024]
- Rao VG (1969) Fungi on Citrus from India. *Sydowia* 23: 215–224. <https://doi.org/10.3406/scrip.1969.3366>
- Rao RP, Rao D (1964) Some species of *Camposporium* Harkn. from India. *Antonie van Leeuwenhoek* 30(1): 60–64. <https://doi.org/10.1007/BF02046701>
- Reblova M (1997) Revision and reclassification of some *Chaetosphaeria* species. *Czech Mycology* 50(2): 73–83. <https://doi.org/10.33585/cmy.50201>

- Rehm H (1881) Ascomyceten in getrockneten Exemplaren herausgegeben. Berichte des Naturhistorischen Vereins Augsburg. 26: 1–132.
- Rehm H (1901) Beiträge zur Pilzflora von Südamerika. XII. Sphaeriales. Hedwigia 40: 100–124.
- Rehm H (1906) Beiträge zur Ascomycetenflora der Voralpen und Alpen. Plant Systematics and Evolution 56(8): 291–348. <https://doi.org/10.1007/BF01672393>
- Rehm H (1913) Ascomycetes philippinenses – III. Philippine Journal of Science 8: 391–406.
- Rehm H (1914) Ascomycetes philippinenses – V. Leaflets of Philippine Botany. 6: 2191–2237.
- Rehner S (2001) Primers for Elongation Factor 1- $\alpha$  (EF1- $\alpha$ ). <http://ocid.NACSE.ORG/research/deephyphae/EF1primer.pdf>
- Reinking OA (1919) Host Index of Diseases of Economic Plants in the Philippines. Philippine Agriculturist 8: 38–54.
- Remler P (1979) Ascomyceten auf Ericaceen in den Ostalpen. Bibliotheca Mycologica 68: 1–321.
- Rick JE (1933) Monographia *Sphaerialium astromaticorum* Riograndensium: II. Brotéria Série Trimestral. Ciências Naturais. 2: 169–201.
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Saccardo PA (1875) Fungi Veneti novi vel critici. Series III. Hedwigia 14: 68–76.
- Saccardo PA (1878) Fungi Italici autographice delineati a Prof. P.A. Saccardo. Patavii 1878. – Fascicoli V.–VIII. sistentes tab. 161–320. Michelia 1: 326–350.
- Saccardo PA (1880) Conspectus generum fungorum Italiae inferiorum nempe ad Sphaeropsideas, Melanconieas et Hyphomyceteas pertinentium systemate sporologico dispositum. Michelia 2: 1–38.
- Saccardo PA (1882) Sylloge Pyrenomycetum. Vol. I. Sylloge Fungorum 1: 1–768.
- Saccardo PA (1883) Sylloge Pyrenomycetum, Vol. II. Sylloge Fungorum 2: 1–813.
- Saccardo PA (1884) Sylloge Sphaeropsidearum et Melanconiearum Omnium Hucusque Cognitarum. Sylloge Fungorum 3: 1–840.
- Saccardo PA (1911) Notae mycologicae. Series XIII. Annales Mycologici 9: 249–257.
- Saccardo PA (1915) Fungi ex insula Melita (Malta) lecti a Doct. Caruana–Gatto et Doct. G. Borg annis MCMXIII et MCMIV. Nuovo Giornale Botanico Italiano 22: 24–76.
- Saccardo PA (1916) Notae mycologicae. Series XX. Nuovo Giornale Botanico Italiano. 23: 185–234.
- Saccardo PA (1921) Notae mycologicae. Series XXIV. I. Fungi Singaporenses Barbesiani. Bolletino dell’Orto Botanico Regia dell’Università de Napoli 6: 39–73.
- Saccardo PA, Sydow P (1899) Sylloge Fungorum 14: 1–724.
- Saccardo PA, Traverso JB (1910) Index Iconum Fungorum. A–L. Sylloge Fungorum 19: 1–1158.
- Saccardo PA, Trotter A (1913) Supplementum Universale, Pars IX. Sylloge Fungorum 22: 1–1612.
- Saccas AM (1954) Les champignons de l’hévéa en Afrique Equatoriale Française. Journal d’Agriculture Tropicale et de Botanique Appliquée 1(7): 333–355. <https://doi.org/10.3406/jatba.1954.2165>
- Samuels GJ (1976) A revision of the fungi formerly classified as *Nectria* subgenus *Hyphonectria*. Memoirs of the New York Botanical Garden 26: 1–126.

- Samuels GJ, Müller E (1978) Life-history studies of Brazilian Ascomycetes. 4. Three species of *Herpotrichia* and their Pyrenochaeta-like anamorphs. *Sydowia* 31: 157–168.
- Sandu-Ville C, Mititiuc M (1971) Für die Wissenschaft unbekannte Pilze aus dem Gebiete des zukünftigen Stausees am Eisernen Tor. *Feddes Repertorium* 81(8–9): 625–627. <https://doi.org/10.1002/fedr.19710810810>
- Sawada K (1950) Represent Forest Experiment Station, Tokyo 46: 111–150. [Fungi inhabiting on conifers in the Tohoku district. II. Fungi on various conifers except “Sugi”]
- Scheuer C, Chlebicki A (1997) Recent collections of miscellaneous microfungi from South Poland. *Acta Mycologica* 32(2): 147–172. <https://doi.org/10.5586/am.1997.014>
- Senanayake IC, Pem D, Rathnayaka AR, Wijesinghe SN, Tibpromma S, Wanasinghe DN, Phookamsak R, Kularathnage ND, Gomdola D, Harishchandra D, Dissanayake LS, Xiang M, Ekanayaka AH, McKenzie EHC, Hyde KD, Zhang H, Xie N (2022) Predicting global numbers of teleomorphic ascomycetes. *Fungal Diversity* 114(1): 237–278. <https://doi.org/10.1007/s13225-022-00498-w>
- Shaw CG (1973) Host fungus index for the Pacific Northwest – I. Hosts. Washington State University, Agricultural Experiment Station Bulletin 765: 1–121.
- Shaw DE (1984) Microorganisms in Papua New Guinea. Department of Primary Industries and Research Bulletin 33: 1–344.
- Shear CL (1941) Mycological notes. V. *Mycologia* 33(3): 318–332. <https://doi.org/10.1080/00275514.1941.12020821>
- Shearer CA (1974) Fungi of the Chesapeake Bay and its Tributaries IV. Three new species from the Patuxent River. *Mycologia* 66(1): 16–24. <https://doi.org/10.1080/00275514.1974.12019568>
- Shivas RG, Alcorn JL (1996) A checklist of plant pathogenic and other microfungi in the rainforests of the wet tropics of northern Queensland. *Australasian Mycological Society* 25: 158–173. <https://doi.org/10.1071/AP96029>
- Siemaszko W (1923) Fungi caucasici novi vel minus cogniti. II. Diagnoses specierum novarum ex Abchazia Adzariaque provenientium. *Acta Societatis Botanicorum Poloniae* 1(1): 1–10. <https://doi.org/10.5586/asbp.1923.003>
- Sivanesan A (1971) The genus *Herpotrichia* Fuckel. *Mycological Papers* 127: 1–37.
- Sivanesan A (1984) The bitunicate ascomycetes and their anamorphs. *J. Cramer, Vaduz*, 701.
- Sivanesan A, Hsieh WH (1989) New species and new records of ascomycetes from Taiwan. *Mycological Research* 93(3): 340–351. [https://doi.org/10.1016/S0953-7562\(89\)80161-3](https://doi.org/10.1016/S0953-7562(89)80161-3)
- Smith AL, Ramsbottom J (1914) New or rare microfungi. *Transactions of the British Mycological Society* 4: 318–330. [https://doi.org/10.1016/S0007-1536\(12\)80035-0](https://doi.org/10.1016/S0007-1536(12)80035-0)
- Spaulding P (1961) Foreign Diseases of Forest Trees of the World. U.S.D.A. Agric. Handb. 197: 1–361.
- Species Fungorum (2024) Species Fungorum. <http://www.indexfungorum.org> [accessed on 15 January 2024]
- Spegazzini C (1908) Hongos de la yerba mate. *Anales del Museo Nacional de Historia Natural Buenos Aires* 17: 111–141.
- Spegazzini C (1909) Mycetes Argentinenses. Series IV. *Anales del Museo Nacional de Historia Natural Buenos Aires Series 3* 12: 257–458.
- Spegazzini C (1910) Fungi Chilenses. Contribución al Estudio de los Hongos Chilenos. *Revista de la Facultad de Agronomía Universidade Nacional de La Plata* 6: 1–205.

- Spegazzini C (1920) Sobre algunas enfermedades y hongos que afectan las plantas de 'Agrios' en el Paraguay. *Anales de la Sociedad Científica Argentina* 90: 155–188.
- Spegazzini C (1921) Mycetes chilenses. *Boletín de la Academia Nacional de Ciencias en Córdoba* 25: 1–124.
- Spegazzini C (1923) Honguillos exóticos. *Boletín de la Academia Nacional de Ciencias en Córdoba* 26: 369–403.
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Stamatakis A, Hoover P, Rougemont J (2008) A rapid bootstrap algorithm for the RAxML web servers. *Systematic Biology* 57(5): 758–771. <https://doi.org/10.1080/10635150802429642>
- Su HY, Udayanga D, Luo ZL, Manamgoda DS, Zhao YC, Yang J, Liu XY, McKenzie EHC, Zhou DQ, Hyde KD (2015) Hyphomycetes from aquatic habitats in Southern China: Species of *Curvularia* (Pleosporaceae) and *Phragmocephala* (Melanommataceae). *Phytotaxa* 226(3): 201–216. <https://doi.org/10.11646/phytotaxa.226.3.1>
- Sutton BC (1980) The Coelomycetes – fungi imperfecti with pycnidia, acervuli and stromata. Commonwealth Mycological Institute, Kew, Surrey, England, 696.
- Sutton BC, Pirozynski KA (1965) Notes on microfungi. II. *Transactions of the British Mycological Society* 48(3): 349–366. [https://doi.org/10.1016/S0007-1536\(65\)80055-9](https://doi.org/10.1016/S0007-1536(65)80055-9)
- Sutton BC, Pirozynski KA (1966) *Phragmotrichum karstenii*. *Transactions of the British Mycological Society* 49(3): 522. [https://doi.org/10.1016/S0007-1536\(66\)80099-2](https://doi.org/10.1016/S0007-1536(66)80099-2)
- Sydow H, Sydow P (1903) *Mycotheca Germanica Fasc. II* (no. 51–100). *Annales Mycologici* 1: 536–539.
- Sydow H, Sydow P (1907) *Novae fungorum species. IV. Annales Mycologici* 5: 338–340.
- Sydow H, Sydow P (1913) *Novae fungorum species XI. Annales Mycologici* 11: 402–408.
- Sydow H, Sydow P (1914) Fungi from northern Palawan. *Philippine Journal of Science, Section C: Botany* 9: 157–189.
- Sydow P, Sydow H (1921) *Novae fungorum species. XVI. Annales Mycologici* 18: 154–160.
- Sydow H, Sydow P, Butler EJ (1911) *Fungi Indiae orientalis pars III. Annales Mycologici* 9: 372–421.
- Tai FL (1979) *Sylloge Fungorum Sinicorum*. Sci. Press, Acad. Sin., Peking, 1527.
- Tanaka K, Hosoya T (2006) Some new records of Loculoascomycetes from Yakushima Island, southern Japan. *Bulletin of the National Science Museum, Series B (Botany)* 32: 151–160.
- Tassi F (1900) *Novae Micromycetum species de scripta e et iconibus illustratae. Bollettino del Laboratorio de Orto Botanico Reale Universita Siena* 3: 52–57.
- Teng SC (1936) Additional fungi from China II. *Sinensia* 7: 490–527.
- Tennakoon DS, Jeewon R, Kuo CH, Hyde KD (2018) Phylogenetic and morphological characterization of *Byssosphaeria macarangae* sp. nov., and *B. taiwanense* sp. nov. from *Macaranga tanarius*. *Phytotaxa* 364(3): 211–226. <https://doi.org/10.11646/phytotaxa.364.3.1>
- Tennakoon DS, Kuo CH, Maharachchikumbura SSN, Thambugala KM, Gentekaki E, Phillips AJ, Bhat DJ, Wanasinghe DN, de Silva NI, Promputtha I, Hyde KD (2021) Taxonomic and phylogenetic contributions to *Celtis formosana*, *Ficus ampelas*, *F. septica*, *Macaranga tanarius* and *Morus australis* leaf litter inhabiting microfungi. *Fungal Diversity* 108(1): 1–215. <https://doi.org/10.1007/s13225-021-00474-w>

- Tennakoon DS, Thambugala KM, Silva NID, Suwannarach N, Lumyong S (2022) A taxonomic assessment of novel and remarkable fungal species in Didymosphaeriaceae (Pleosporales, Dothideomycetes) from plant litter. *Frontiers in Microbiology* 13: e1016285. <https://doi.org/10.3389/fmicb.2022.1016285>
- Tennakoon DS, de Silva NI, Maharachchikumbura SS, Bhat DJ, Kumla J, Suwannarach N, Lumyong S (2023) Exploring More on Dictyosporiaceae: The Species Geographical Distribution and Intriguing Novel Additions from Plant Litter. *Diversity* 15(3): e410. <https://doi.org/10.3390/d15030410>
- Teodoro NG (1937) An Enumeration of Philippine Fungi. Tech. Bull. Dept. Agric. Comm. Manila 4: 1–585.
- Thambugala KM, Wanasinghe DN, Phillips AJL, Camporesi E, Bulgakov TS, Phukhamsakda C, Ariyawansa HA, Goonasekara ID, Phookamsak R, Dissanayake A, Tennakoon DS (2017) Mycosphere notes 1–50: Grass (Poaceae) inhabiting Dothideomycetes. *Mycosphere: Journal of Fungal Biology* 8(4): 697–796. <https://doi.org/10.5943/mycosphere/8/4/13>
- Thaxter R (1922) Reliquiae Farlowianae distributed from the Farlow Herbarium of Harvard University. *Mycologia* 14(3): 99–103. <https://doi.org/10.1080/00275514.1922.12020377>
- Tian Q, Liu JK, Hyde KD, Wanasinghe DN, Boonmee S, Jayasiri SC, Luo ZL, Taylor JE, Phillips AJL, Bhat DJ, Li WJ, Ariyawansa H, Thambugala KM, Gareth Jones EB, Chomnunti P, Bahkali AH, Xu JC, Camporesi E (2015) Phylogenetic relationships and morphological reappraisal of Melanommataceae (Pleosporales). *Fungal Diversity* 74(1): 267–324. <https://doi.org/10.1007/s13225-015-0350-9>
- Tibpromma S, Hyde KD, Jeewon R, Maharachchikumbura SS, Liu JK, Bhat DJ, Jones EBG, McKenzie E, Camporesi E, Bulgakov TS, Doilom M, Santiago AM, Das K, Manimohan P, Gibertoni TB, Lim YW, Ekanayaka AH, Thongbai B, Lee HB, Yang J, Kirk PM, Sysouphanthong P, Singh SK, Boonmee S, Dong W, Raj KN, Latha KP, Phookamsak R, Phukhamsakda C, Konta S, Jayasiri SC, Norphanphoun C, Tennakoon D, Li J, Dayarathne MC, Perera RH, Xiao Y, Wanasinghe DN, Senanayake IC, Goonasekara ID, Silva NI, Mapook A, Jayawardena RS, Dissanayake AJ, Manawasinghe IS, Chethana KW, Luo Z, Hapuarachchi KK, Baghela A, Soares AM, Vizzini A, Meiras-Ottoni A, Mešić A, Dutta AK, Souza CA, Richter C, Lin C, Chakrabarty D, Daranagama DA, Lima DX, Chakraborty D, Ercole E, Wu F, Simonini G, Vasquez G, Silva GA, Plautz HL, Ariyawansa HA, Lee HS, Kušan I, Song J, Sun J, Karmakar J, Hu K, Semwal KC, Thambugala KM, Voigt K, Acharya K, Rajeshkumar KC, Ryvarden L, Jadan M, Hosen MI, Mikšić M, Samarakoon MA, Wijayawardene NN, Kim NK, Matočec N, Singh PN, Tian Q, Bhatt RP, Oliveira RJ, Tulloss RE, Aamir S, Kaewchai S, Marathe SD, Khan S, Hongsanan S, Adhikari S, Mehmood T, Bandyopadhyay TK, Svetasheva TY, Nguyen TT, Antonín V, Li W, Wang Y, Indoliya Y, Tkálčec Z, Elgorban AM, Bahkali AH, Tang A, Su H, Zhang H, Promputtha I, Luangsa-Ard J, Xu J, Yan J, Kang JC, Stadler M, Mortimer PE, Chomnunti P, Zhao Q, Phillips AJ, Nontachaiyapoom S, Wen T, Karunarathna SC (2017) Fungal diversity notes 491–602: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 83(1): 1–261. <https://doi.org/10.1007/s13225-017-0378-0>
- Tilak ST, Kale SB (1970) Contribution to our knowledge of Ascomycetes of India XXIV. *Sydowia* 24: 81–88.
- Tilak ST, Talde UK (1975) Contribution of our knowledge of Ascomycetes of India XXXV. *Maharashtra Vidnyan Mandir Patrika*. 10: 15–18.
- Trotter A (1912) Mycetum Tripolitanorum pugillus. *Annales Mycologici* 10: 509–514.
- Trotter A (1931) Supplementum Universale, Pars X. Myxomycetae, Myxobacteriaceae, Deuteromycetae, Mycelia sterilia. *Sylloge Fungorum*. 25: 1–1093.

- Urries MJ (1957) Hongos Microscopicos De Canarias. El Museo Canario, 140.
- Vasilyeva LN (1987) Pirenomitsety i Lokuloaskomitsety severa Dalbnego Vostoka, 1–255.
- Viégas AP (1945) Alguns fungos do Brasil XI. Fungi Imperfecti (Sphaeropsidales). *Bragantia* 5: 717–779. <https://doi.org/10.1590/S0006-87051945000900002>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Von Arx JA (1981) The genera of fungi sporulating in pure culture, 3<sup>rd</sup> edn. Stechert-Hafner, New York, 288.
- Wanasinghe DN, Phukhamsakda C, Hyde KD, Jeewon R, Lee HB, Gareth Jones EB, Tibpromma S, Tennakoon DS, Dissanayake AJ, Jayasiri SC, Gafforov Y, Camporesi E, Bulgakov TS, Ekanayake AH, Perera RH, Samarakoon MC, Goonasekara ID, Mapook A, Li WJ, Senanayake IC, Li J, Norphanphoun C, Doilom M, Bahkali AH, Xu J, Mortimer PE, Tibell L, Tibell S, Karunarathna SC (2018) Fungal diversity notes 709–839: Taxonomic and phylogenetic contributions to fungal taxa with an emphasis on fungi on Rosaceae. *Fungal Diversity* 89(1): 1–236. <https://doi.org/10.1007/s13225-018-0395-7>
- Wang YZ, Aptroot A, Hyde KD (2004) Revision of the ascomycete genus *Amphisphaeria*. Fungal Diversity Press, Hong Kong, 168.
- Wang HK, Aptroot A, Crous PW, Jeewon R, Hyde KD (2007) The polyphyletic nature of Pleosporales: An example from *Massariosphaeria* based on rDNA and RBP2 gene phylogenies. *Mycological Research* 111(11): 1268–1276. <https://doi.org/10.1016/j.mycres.2007.08.014>
- Watanabe T (1993) *Camposporium laundonii* on root tissue of *Aralia elata* from Japan. *Transactions of the British Mycological Society, Japan* 34: 71–76.
- Watson AJ (1971) Foreign bacterial and fungus diseases of food, forage, and fiber crops. USDA Agricultural Research Service, 111.
- Watzl O (1937) On the trees and shrubs of the region of the Khodja group drained by the Baramba and their leaf diseases. (A contribution to the flora of the south-western central Caucasus.) II. Fungi on living leaves. *Beihefte zum Botanischen Centralblatt* 57: 431–440.
- Wehmeyer LE (1952) Some pyrenomycetous fungi from Mt. Rainier National Park. *Sydowia* 6: 412–428.
- Wen S, Fan W, Guo H, Huang C, Yan Z, Long Y (2020) Two new secondary metabolites from the mangrove endophytic fungus Pleosporales sp. SK7. *Natural Product Research* 34(20): 2919–2925. <https://doi.org/10.1080/14786419.2019.1598993>
- White RP, Hamilton CC (1935) Diseases and insect pests of *Rhododendron* and *Azalea*. New Jersey Agricultural Experiment Station 350: 1–22.
- White TJ, Bruns T, Lee SJWT, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications* 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Whitton SR, McKenzie EHC, Hyde KD (2002) Microfungi on the Pandanaceae: Two new species of *Camposporium* and key to the genus. *Fungal Diversity* 11: 177–187.
- Wijayawardene NN, Hyde KD, Wanasinghe DN, Papizadeh M, Goonasekara ID, Camporesi E, Bhat DJ, McKenzie EHC, Phillips AJL, Diederich P, Tanaka K, Li WJ, Tangthirasun N, Phookamsak R, Dai DQ, Dissanayake AJ, Weerakoon G, Maharachchikumbura SSN, Hashimoto A, Matsumura M, Bahkali AH, Wang Y (2016) Taxonomy and phylogeny of dematiaceous coelomycetes. *Fungal Diversity* 77(1): 1–316. <https://doi.org/10.1007/s13225-016-0360-2>

- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğan M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkálčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalasekera SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa (2021) *Mycosphere* 13: 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Winter G (1885) Pilze–Ascomyceten. GL Rabenhorst's Kryptogamen–Flora von Deutschland. Oesterreich und der Schweiz 1: 65–528.
- Wong MKM, Hyde KD (2001) Diversity of fungi on six species of Gramineae and one species of Cyperaceae in Hong Kong. *Mycological Research* 105(12): 1485–1491. <https://doi.org/10.1017/S0953756201004695>
- Wong MKM, Goh TK, McKenzie EHC, Hyde KD (2002) Fungi on grasses and sedges: *Paratetraploa exappendiculata* gen. et sp. nov., *Petrakia paracochinensis* sp. nov. and *Spadicoides versiseptatis* sp. nov. (dematiaceous hyphomycetes). *Cryptogamie. Mycologie* 23: 195–203.
- Wu WP, Zhuang WY (2005) *Sporidesmium*, *Endophragmiella* and related genera from China. *Fungal Diversity Research Series* 15: 1–531.
- Xiong YR, Manawasinghe IS, Wanasinghe DN, Hongsanan S, Hyde KD, Biao X, Dong ZY (2023) Two new species and a new host record of Pleosporales (Dothideomycetes) from palm (Arecaceae) in Guangdong Province, China. *New Zealand Journal of Botany*: 1–27. <https://doi.org/10.1080/0028825X.2023.2258583>
- Xu ZH, Sun WX, Shi XG, Zhang XG, Xia JW, Castañeda Ruiz RF, Ma J (2021) *Camposporium chinense* sp. nov. from Jiangxi, China. *Mycotaxon* 136(1): 73–78. <https://doi.org/10.5248/136.73>
- Yuan ZQ, Barr ME (1994) New ascomycetous fungi on bush cinquefoil from Xinjiang, China. *Sydowia* 46: 329–337.
- Yuan ZQ, Zhao ZY (1994) Studies on lophiostomataceous fungi from Xinjiang, China. *Sydowia* 46: 162–184.
- Zhang Y, Crous PW, Schoch CL, Hyde KD (2012) Pleosporales. *Fungal Diversity* 53(1): 1–221. <https://doi.org/10.1007/s13225-011-0117-x>
- Zhuang WY (2001) Higher Fungi of Tropical China. *Mycotaxon, Ltd., Ithaca, NY*, 485.
- Zhuang WY (2005) Fungi of northwestern China. *Mycotaxon, Ltd., Ithaca, NY*, 430.