

Research Article

Diversity and distribution patterns of invasive alien plant species in mainland Portugal

Raquel Fernandes¹, Andry Castro¹, Hélia Marchante^{2,3}, Elizabete Marchante^{3,4}, César Capinha^{1,3}

¹ Centre of Geographical Studies, Institute of Geography and Spatial Planning, University of Lisbon, Rua Branca Edmée Marques, 1600-276 Lisboa, Portugal

² Research Center for Natural Resources, Environment and Society (CERNAS), Polytechnic University of Coimbra, Coimbra Agriculture School, Bencanta, 3045-601 Coimbra, Portugal

³ Associate Laboratory Terra, Lisboa, Portugal

⁴ Centre for Functional Ecology - Science for People & the Planet, Department of Life Sciences, University of Coimbra, Calçada Martim de Freitas, 3000-456 Coimbra, Portugal

Corresponding author: Raquel Fernandes (raquel.fernandes@edu.ulisboa.pt)

Abstract

Understanding the distribution and diversity of invasive alien species is increasingly crucial to meet legal requirements and guide effective management. Portugal currently hosts a large number of invasive alien plant species, with significant environmental and socio-economic impacts. However, the distribution of most of these species across the territory and the factors driving their spread and diversity remain largely unexplored. To address this, we present the first atlas of invasive alien plants in mainland Portugal. A total of 96 terrestrial and aquatic invasive alien plants are presented, encompassing all species listed under Portuguese national legislation and the European Union's list of invasive species of concern. Occurrence data were collected from a broad array of sources, including national and international biodiversity observation databases, citizen science data, literature, and data collections owned by managers and researchers—totaling approximately 85,000 records, with a mean of 879 records per species (range: 1–9,190). Each species was characterized based on multiple distributional parameters, and a k-means analysis was used to group species with similar distribution patterns. The richness of invasive plants was mapped at the municipality level. The drivers of their spatial variation were investigated using a comprehensive set of 30 variables representing multiple environmental and human factors. Results show that invasive plants are present in all mainland Portuguese municipalities but with high variability. Four main patterns of distribution were identified: species primarily located along the coast (e.g., *Acacia saligna*) and a gradient of species with narrow (e.g., *Reynoutria japonica*), moderate (e.g., *Ipomoea indica*), and wide distribution ranges (e.g., *Cortaderia selloana*). Invasive species richness was significantly higher in coastal and larger municipalities, particularly those closer to major urban centers and with a higher density of power lines. Our results provide the first comprehensive assessment of the distribution of invasive alien plants in mainland Portugal, establishing a much-needed baseline for future invasion prevention and management efforts.

Key words: Baseline distribution, drivers of invasion, invasion atlas, invasive alien plants, invasive species richness, occupancy



Academic editor: Inderjit

Received: 26 June 2025

Accepted: 21 October 2025

Published: 19 November 2025

Citation: Fernandes R, Castro A, Marchante H, Marchante E, Capinha C (2025) Diversity and distribution patterns of invasive alien plant species in mainland Portugal. NeoBiota 104: 139–162. <https://doi.org/10.3897/neobiota.104.163291>

Copyright: © Raquel Fernandes et al. This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Human activities are facilitating the global movement and introduction of species beyond their native ranges (Pyšek et al. 2020; Lopez et al. 2022; Capinha et al. 2023). Currently, over 37,000 established alien species have been documented worldwide

(IPBES 2023), with approximately 200 new alien species (i.e., “exotic” or “non-native”) recorded annually. Among these, alien plants constitute a significant proportion, with more than 13,000 species established outside their natural ranges (Seebens et al. 2023).

A minor portion of alien plants become invasive, causing significant and diverse negative impacts in regions of introduction. These impacts include reductions in native species diversity and abundance (Powell et al. 2011; Hansen et al. 2020; Beaury et al. 2023), disruption of key ecosystem services (Vieites-Blanco and González-Prieto 2020; Ferreira et al. 2021; Ren et al. 2021; Gallardo et al. 2024), and major economic losses affecting multiple activities, including the loss of crops and productive land (Richardson and Van Wilgen 2004; Le Maitre et al. 2011; Milanović et al. 2020; Reynolds et al. 2020; Sirbu et al. 2022). Additionally, several invasive plants pose environmental risks to human health (Rodríguez et al. 2021). Most of these impactful species have been reported in the American continent, the Asia–Pacific region, Central Asia, and Europe (IPBES 2023). In Europe, at least 7,335 alien vascular plant species have been recorded, of which 1,037 (about 14%) are considered invasive in at least one European territory (Kalusová et al. 2024).

Given the frequent severity of the observed and potential impacts, many national and regional governments now prioritize managing biological invasions, including such species in dedicated legislation. Portugal was one of the first European countries to adopt legislation on invasive alien species (hereafter invasive species), introducing in 1999 a list of species subject to specific restrictions, including bans on commercial use and possession (Ministério do Ambiente 1999). This list included over 40 plant species, with several listed as invasive, such as *Acacia dealbata*, *Carpobrotus edulis*, and *Oxalis pes-caprae*, and others considered to have known ecological risk (either in the early stages of invasion or having a high potential to become invasive in the future), such as *Ludwigia peploides*, *R. japonica*, and *Senecio inaequidens*. This legislation and associated species list were updated by Decree-Law No. 92/2019 (Ministério do Ambiente e Transição Energética 2019), which now includes more than 100 plant species listed as invasive for mainland Portugal. The updated list includes confirmed and potential invasive species known to occur in the territory, as well as others that are presumed absent but expected to become invasive if introduced. Additionally, the European Union Regulation No. 1143/2014 (Official Journal of the European Union 2014) includes the “Union list,” which specifically lists the invasive species of Union concern. This regulation aims to prevent and mitigate negative environmental impacts on European Union countries. Species listed are automatically part of the Portuguese national list, being subject to Decree-Law No. 92/2019. As of the third update of the Union list in 2022, 41 plant species are listed (Official Journal of the European Union 2022).

Over the years, a relevant number of studies have investigated plant invasions in Portugal (Sousa et al. 2018). For example, several have listed the diversity of alien plants (Almeida and Freitas 2000; Almeida and Freitas 2012; Marchante et al. 2014), distribution, or potential invasion risk (Rodríguez-Merino 2023). Others have analyzed individual species such as *Baccharis spicata* (Verloove et al. 2017), *C. edulis* (Maltez-Mouro et al. 2009; Chefaoui and Chozas 2019), *Cotula coronopifolia* (Costa et al. 2009), or *Pontederia crassipes* (Pádua et al. 2022a, 2022b; Mouta et al. 2023). Studies have also focused on *Acacia* species (Marchante et al. 2003; Martins et al. 2016; Vicente et al. 2019; Raposo et al. 2021; Große-Stoltenberg et al. 2023), a highly problematic group of taxa due to their invasive behavior, with nine species already established in the territory (Marchante et al. 2023).

Specific regions and ecosystems have also received particular attention, including protected areas (Gutierrez et al. 2011; Duarte et al. 2023), dune habitats (Duarte et al. 2023), forests (Ferreira et al. 2021), riparian systems (Bernez et al. 2005; Aguiar and Ferreira 2013; Pabst et al. 2022), and inland waters (Oficialdegui et al. 2023).

Despite the numerous valuable contributions and the growing number of citizen science distribution records on different platforms (e.g., the project “Invasoras.pt” at iNaturalist), the overall distribution of invasive plants across mainland Portugal remains largely unknown. This gap persists despite national inventories of invasive species being essential tools for monitoring invasions, guiding policy, and formulating management plans (Latombe et al. 2017; Pabst et al. 2022; Sirbu et al. 2022). The lack of comprehensive distribution data is likely due, in part, to information on individual taxa being dispersed across numerous sources, making collation and mapping a resource-intensive and challenging task. Additionally, for many species, information about distribution at local scales is incomplete or missing.

Here, we present a comprehensive atlas of the distribution of invasive plant richness in mainland Portugal. This atlas was created by compiling and harmonizing data from a wide range of sources, including national and international biodiversity observation databases, scientific literature, and data collections from managers and researchers. In Portugal, municipalities play a central role in territorial management, environmental planning, and decision-making regarding field interventions. This is also the case for local- and national-level resource allocation for invasive species prevention and management, citizen science initiatives, and public awareness campaigns. Hence, municipalities were selected as reference sampling units. With the information collected, we aimed to: i) map the recorded distribution of each invasive alien plant species in the territory; ii) identify species sharing similar distribution patterns; iii) assess spatial patterns of invasive species richness; and iv) evaluate the environmental and socio-economic drivers of spatial variation in invasive species richness values. Our results are expected to provide substantial support for decision-making in local-to-national-scale prevention and management efforts of alien plant invasions in Portugal.

Materials and methods

Study area

Mainland Portugal is located in the western Iberian Peninsula, at the southwestern edge of Europe (Fig. 1). Its territory spans both the Atlantic and Mediterranean biogeographical regions (Roekaerts 2002). The central and the southern regions (Alentejo and Algarve) are characterized by a Mediterranean climate with dry, hot summers, while the northern regions experience a temperate oceanic climate with dry, mild summers (Beck et al. 2023). The southern and coastal areas are generally low-lying, whereas the northern and central regions are more mountainous, with several peaks exceeding 1,000 m in elevation (Ramos and Ramos-Pereira 2020). The territory is divided into 278 municipalities. The population is primarily concentrated in coastal municipalities, particularly in the metropolitan areas of Lisbon and Oporto (Instituto Nacional de Estatística 2022). Forests occupy large areas in all Nomenclature of Territorial Units for Statistics II (NUTS II) regions, especially in the Centre region, while agricultural areas are most represented in Alentejo and the Lisbon Metropolitan Area (Alves et al. 2022).




Figure 1. Location of mainland Portugal showing its administrative divisions, including NUTS II regions (outlined in black) and the 278 municipalities (outlined in grey), used as units of analysis in this study. Population density (persons/km²) is highest in the Lisbon and Oporto Metropolitan Areas, represented by dark red regions.

List of invasive plants

The mapping effort focused on the plant species listed in Portuguese Decree-Law No. 92/2019 (Ministério do Ambiente e Transição Energética 2019) and European Regulation No. 1143/2014 (Official Journal of the European Union 2014). Since our focus is on mainland Portugal, species mentioned in national legislation that pertain only to the insular regions, such as the Madeira archipelago, were excluded. Portuguese Decree-Law No. 92/2019 lists the genus *Acacia* collectively, unlike other taxa, which are listed at the species level. Therefore, we consulted relevant literature (Marchante et al. 2014, 2023) to identify the *Acacia* species known to occur in the territory. Additionally, we included *Watsonia meriana* in our mapping, as the research community recognizes its invasive behavior in several areas of mainland Portugal (Marchante et al. 2014; Morais et al. 2017). Distribution data were collected for a total of 96 species from 64 different genera (Suppl. material 1: table S1.1).

Occurrence database

We gathered occurrence records for each invasive plant species from multiple sources. The online databases consulted included the Global Biodiversity Information Facility (GBIF) (GBIF.org 2024); Flora-On (Flora de Portugal Interactiva 2024), an expert-led project aimed at systematizing information for all vascular plant species in Portugal, including their distribution; and the citizen-science platforms iNaturalist (iNaturalist 2024) (only research-grade records were considered) and Pl@ntNet (Pl@ntNet 2024). Records were also collected from numerous research publications identified in Web of Science and Scopus using the search terms “Portugal” and each species name, as well as from master’s and doctoral theses identified in RENATES, a Portuguese platform containing official information on dissertations produced in Portugal (DGEEC 2024). Additionally, we incorporated unpublished data from the authors, collaborators, and companies involved in invasive plant management actions (see Acknowledgments section). Data up to May 2024 were considered.

The taxonomy of species was defined and harmonized according to the GBIF Backbone Taxonomy (GBIF Secretariat 2023). Occurrence records were collected for all available years and included subspecies (if present). Records identified as having coordinates of stored specimens (e.g., herbarium locations) rather than the actual place of observation in the wild were removed. Additionally, records with a spatial precision of less than 100 m were excluded.

An exception was made for data from Flora-On. This source is crucial for the comprehensiveness of the mapping but provides data as centroids of grid squares at a 10×10 km resolution. Despite its contribution to spatial comprehensiveness, the lower geographical precision may introduce some spatial errors in attributing occurrences to municipalities—the spatial units of mapping and analysis used (see below). Therefore, we performed two mapping exercises and associated analyses: one including Flora-On data and another without it. The results from both datasets were highly congruent (see Results section). Thus, we primarily describe the results, including the entire set of occurrence data, while those excluding Flora-On data are provided in Suppl. material 3.

Given the breadth of consulted sources, we expect to have obtained a representative identification of the species occurring in each municipality. However, as with any similar exercise, our data may still exhibit some gaps and biases, particularly toward regions with higher recording efforts and species that are more conspicuous and easier to identify (see Discussion section for details).

Invasive plant richness mapping and occupancy

To analyze species distribution patterns, we aggregated occurrence records at the municipality level, following the boundaries defined by the Carta Administrativa Oficial de Portugal (CAOP) (Direção-Geral do Território 2024). For each municipality, duplicate records of the same species were removed, retaining only one occurrence per species. Invasive species richness was then calculated as the total number of unique species recorded per municipality. Species occupancy patterns were assessed by quantifying, within each municipality, the number of 1×1 km grid cells in which the species was recorded. The mean percentage of occupied grid cells across municipalities was subsequently calculated to represent overall species occupancy. Analyses were performed in R v4.3.2 (R Core Team 2022).

Assessing patterns of species distributions

To identify similarities in species-level patterns of distribution across the territory, we first characterized the distribution of each species using five variables: the number of municipalities where occurs; the median and interquartile range of distance to the coast (from the centroid of each municipality, in kilometers); and the median and interquartile range of latitude (in decimal degrees).

We then performed a cluster analysis based on these variables. To account for multicollinearity among the variables, we used principal component analysis (PCA), which explained 72.3% of the variance with the first two components (Suppl. material 1: fig. S1.1). Next, we performed a k-means clustering analysis based on the scores of each species in the two-dimensional PCA space. The PCA was performed using the “prcomp” function from the *stats* R package. The k-means procedure employed the “kmeans” function from the same package, and the elbow method (Bholowalia and Kumar 2014) was used to identify the optimal number of clusters (Suppl. material 1: fig. S1.2).

We also examined whether the number of municipalities where a species occurs reflects its overall occupancy across the territory. To do this, we fitted a linear model relating the mean occupancy of species (i.e., the average percentage of occupied 1 × 1 km grid cells per municipality) to the total number of municipalities where it was recorded.

Spatial drivers of invasive plant richness

Explanatory variables

To identify the factors driving the spatial variation in the richness of invasive plants across the territory, we selected a comprehensive set of 30 variables (Table 1) representing distinct environmental and human-related factors, including proxies for recording effort. These variables were mapped at the municipality level and provide a broad representation of the hypotheses on driving factors considered in previous studies (Vicente et al. 2010; Santos et al. 2011; Vicente et al. 2019; Vieites-Blanco and González-Prieto 2020; Pabst et al. 2022).

The variables representing climatic conditions, topography, continentality, socioeconomy (gross domestic product [GDP] and travel time to cities, i.e., the time it takes to travel to the nearest urban center via surface transport) (Weiss et al. 2018), and soil groups consisted of mean values of raster cell data within the boundaries of each municipality. The land-use variables used vector-type base data and were obtained by dividing the total area occupied by each land-use class by the municipality’s total area. Additionally, variables representing dispersal corridors were calculated by summing the extent of these entities (hydrographic network, road infrastructure, railways, and power lines) within each municipality and dividing by its area. The minimum distance to the coastline was determined as the shortest distance from the centroid of each municipality.

The availability of species distribution data can be strongly influenced by varying levels of recording effort (Tiago et al. 2017). Therefore, we considered a set of variables expected to represent spatial patterns in this effort across the territory. These variables were the average number of higher education students per municipality, the number of higher education institutions per

Table 1. Explanatory variables considered as predictors of spatial variation patterns of invasive plant richness. The final set of variables (in bold) was determined after assessing multicollinearity.

Group	Variables	Transformation	Source	VIF
Climate	Annual precipitation amount (kg m ⁻²)		Chelsa (1981–2010) (Karger et al. 2017, 2018)	>5
	Precipitation amount of the wettest month (kg m ⁻²)			>5
	Precipitation amount of the driest month (kg m ⁻²)			>5
	Annual mean temperature (°C)			>5
	Daily mean air temperatures of the driest quarter (°C)			>5
	Daily mean air temperatures of the wettest quarter (°C)			3.44
	Daily mean air temperatures of the coldest quarter (°C)			>5
	Mean daily maximum air temperature of the warmest month (°C)			2.38
	Mean daily minimum air temperature of the coldest month (°C)			>5
Topography	Slope (degrees)		Earth Resources Observation and Science (EROS) Center (2018)	>5
	Elevation (meters)			>5
	Ruggedness Index			3.06
Geography	Minimum distance to coastline (km)			2.58
	Municipality area (km²)	Sqrt	Direção-Geral do Território (2024)	2.55
Land use	Density of artificialized territories (%)	Log(x+1)	COS, Version 2018 (Direção Geral do Território (DGT), 2020)	>5
	Density of agricultural areas (%)			2.02
	Density of forest areas (%)			2.70
	Density of protected areas (%)	Log(x+1)	Instituto da Conservação da Natureza e das Florestas (ICNF) (2022, 2023a, 2023b, 2023c, 2023d)	1.89
	Burnt area (2007–2022) (%)		Instituto da Conservação da Natureza e das Florestas (ICNF) (2024)	2.65
Soil	Average soil pH (15–30 cm layer)	Log	Poggio et al. (2021)	2.32
Density of dispersal corridors	Density of local hydrographic network (km/km²) (2024)		OpenStreetMap. Geofabrik GmbH, and OpenStreetMap Contributors (2018)	2.13
	Density of roads (km/km ²) (2024)	Log(x+1)		>5
	Density of railways (km/km²) (2024)	Log(x+1)		1.92
	Density of power lines (km/km²) (2024)	Log(x+1)	REN—Redes Energéticas Nacionais, SA (see acknowledgments)	1.26
Socioeconomic	Mean population density according to the national census (mean value of 2001, 2011, and 2021; number of persons/km ²)	Log	INE, PORDATA (2024)	>5
	GDP (1990–2015; 2011 international US dollars)	Log	Kummu et al. (2015)	>5
	Travel time to cities (minutes)		Weiss et al. (2018)	2.97
	Sampling effort (academic and citizen-science sampling effort)		DGEEC/ME-MCTE, PORDATA (2021, 2024); INE, PORDATA (2024)	3.21
	Number of iNaturalist projects with IAP records	Log(x+1)	iNaturalist (2024)	2.51
	Number of plant nurseries and aquarium stores	Log(x+1)	Nunes et al. (unpublished)	1.69

municipality, and the number of iNaturalist projects with invasive plant records per municipality. To avoid redundancy in their spatial variation, we projected these variables into a PCA. The first principal component, explaining around 75% of the variance, was retained, representing mainly the joint variation of the number of higher education students and institutions. The scores of this first composite were extracted and used as a new variable named “academic sampling effort” (Table 1). Because the number of iNaturalist projects contributed little to this component, we used the original values of this variable as a proxy for “citizen-science sampling effort”.

Testing for drivers of invasive plant richness

To assess the relationship between the richness of invasive plants across the territory (dependent variable) and the explanatory variables (Table 1), we used a generalized least squares (GLS) regression model (Dormann et al. 2007). Prior to model fitting, square root and logarithmic transformations were applied to several explanatory variables (Table 1). The response variable was also log-transformed to meet model expectations of a Gaussian distribution of error (Ives 2015).

Potential multicollinearity among the explanatory variables was assessed using the variance inflation factor (VIF) (Alin 2010), calculated with the “vifstep” function from the *usdm* R package (Naimi 2022). Only the set of explanatory variables with a VIF < 5 was considered for the GLS model (Table 1).

This model was fitted using the “gl” function from the *nlme* R package (Pinheiro et al. 2023), assuming an exponential decay in the spatial autocorrelation of richness values (Dormann et al. 2007). Significant relationships between dependent and explanatory variables were determined at $\alpha = 0.05$, and the goodness of fit of the model was measured using the pseudo- R^2 of Nakagawa et al. (2017).

Results

Invasive plants over Portuguese municipalities

Of the 96 invasive plants with distribution records in mainland Portugal (Suppl. material 1: table S1.1), 86% are terrestrial, with a smaller portion being aquatic or semi-aquatic (14%), including plants such as *Alternanthera philoxeroides* and *Ludwigia* spp. They include grasses and herbs (47%), trees and shrubs (29%), and only a few climbing plants (5%) and other growth forms (Suppl. material 1: fig. S1.3). These species belong to 36 plant families, mainly Asteraceae (16%), Amaranthaceae (14%), Fabaceae (14%), and Poaceae (9%) (Suppl. material 1: fig. S1.4).

Of the 10 species with records in the greatest number of municipalities (Table 2), the majority are grasses and herbs (50%; e.g., *Datura stramonium*, *C. selloana*) and trees and shrubs (40%; e.g., *A. dealbata*, *Ailanthus altissima*) (Table 2). Six are already present in more than 200 municipalities (Table 2). The species recorded in only a few municipalities (up to 10; Suppl. material 1: table S1.2) are mostly aquatic or semi-aquatic (40%; e.g., *Nymphaea mexicana* or *Ludwigia grandiflora*).

Table 2. The 10 invasive plants recorded in the greatest number of Portuguese municipalities.

Species	Number of municipalities with records of the species
<i>Acacia dealbata</i>	249
<i>Datura stramonium</i>	233
<i>Arundo donax</i>	232
<i>Phytolacca americana</i>	224
<i>Oxalis pes-caprae</i>	213
<i>Cortaderia selloana</i>	208
<i>Ailanthus altissima</i>	198
<i>Acacia melanoxylon</i>	190
<i>Agave americana</i>	187
<i>Robinia pseudoacacia</i>	186

Four species have been recorded in a single municipality (*A. philoxeroides*, *Amaranthus hypochondriacus*, *Amaranthus muricatus*, and *Baccharis halimifolia*; Suppl. material 1: table S1.2). The distribution of each species per municipality is provided in Suppl. material 4.

The three species with the highest mean percentage of occupancy in mainland Portugal—*O. pes-caprae* (7.53%), *Arundo donax* (6.90%), and *C. selloana* (6.87%)—are also among the most widely distributed, occurring in more than 200 municipalities (Table 2, Suppl. material 1: table S1.3). Their broad geographic range reflects both high local presence and widespread dispersal across the country. In contrast, *Arctotheca calendula* (5.88%) and *Acacia longifolia* (5.58%) rank fourth and fifth in mean occupancy but have a more restricted national distribution. Their presence is concentrated in coastal regions, suggesting high local abundance but limited expansion at a broader scale (Suppl. material 1: table S1.3).

Similarly, the regression model assessing the relationship between the number of municipalities in which a species was recorded and its mean occupancy revealed a statistically significant positive association ($p < 0.05$), explaining 42% of the observed variation. This suggests that species occurring in a greater number of municipalities also tend to occupy larger areas within those municipalities. However, the substantial unexplained variation indicates that additional factors influence species occupancy patterns (Fig. 2).




Figure 2. Linear regression model showing the relationship between the total number of municipalities where each species was recorded and its average occupancy (%). Each dot represents a species. The dashed line indicates the fitted linear relationship.

Distribution patterns of invasive plant richness

Overall, the species are grouped into four distinct clusters (Fig. 3, Suppl. materials 1, 3: table S1.4, fig. S3.1). The first cluster (Fig. 3) includes 26 species predominantly recorded in coastal areas, such as *A. saligna* (Fig. 4a) and *C. coronopifolia*.




Figure 3. Clusters of invasive plants according to their distribution patterns in mainland Portugal. Cluster 1 (green): species primarily located along the coast; Cluster 2 (blue): mostly species with narrow ranges; Cluster 3 (red): moderately widespread species; Cluster 4 (yellow): mostly widespread species. The two PCA axes represent variation in the total number of municipalities of occurrence (N_{muni}), the median distance to the coast ($Dist_coast_median$), the interquartile range of distances to the coast ($Dist_coast_rangeQ$), the median latitude (Lat_median), and the interquartile range of latitudes (Lat_rangeQ).

The remaining three groups are differentiated mainly by the breadth of their distribution ranges. The second cluster includes 14 species characterized by narrow distribution areas (up to 33 municipalities). These species are also generally found in central and northern municipalities (e.g., *R. japonica*, Fig. 4b, and *B. halimifolia*). The third cluster includes 29 mostly moderately widespread species (e.g., *A. longifolia* and *I. indica*, Fig. 4c). Although there are 10 species with narrow distributions (up to 33 municipalities; Suppl. material 1: table S1.4), they were grouped in the third cluster because of their wider latitudinal range than those in the second cluster. The final group consists of 27 species, mostly with very wide ranges (mean number of municipalities = 145 ± 67). Examples of species included in this last group are *A. dealbata* and *C. selleana* (Fig. 4d).




Figure 4. Distribution of example species within each cluster, grouped according to the k-means clustering (Fig. 3) across the Portuguese territory: **a.** Cluster 1 (green)—*Acacia saligna*; **b.** Cluster 2 (blue)—*Reynoutria japonica*; **c.** Cluster 3 (dark red)—*Ipomoea indica*; **d.** Cluster 4 (yellow)—*Cortaderia selloana*.

Patterns and drivers of invasive plant richness

At least one invasive plant species has been recorded in each Portuguese municipality (Fig. 5, Suppl. material 3: fig. S3.2). Richness is markedly higher in coastal municipalities, particularly those near major urban centers such as the metropolitan areas of Lisbon and Oporto. Additionally, certain cities of regional significance, including Coimbra, Figueira da Foz, and Viana do Castelo, as well as locations along the Algarve coast in southern Portugal, also exhibit high numbers of invasive plants. The highest numbers were recorded in Lisbon (63 species), Coimbra (62), Sintra (58), and Figueira da Foz (55). Conversely, lower richness was observed in municipalities located primarily in northeastern Portugal and in some central and inland regions of Alentejo.

The GLS model identified drivers of variation in invasive species richness, achieving a pseudo- R^2 value of 0.59, indicating good explanatory capacity. Four variables exhibited significant relationships with richness values ($p < 0.05$; Table 3, Suppl. material 3: table S3.1). Specifically, invasive species richness increased with larger municipality area, higher density of power lines, greater proximity to the coastline, and shorter travel time to cities.

Table 3. Coefficients and p -value results of each explanatory variable considered in the GLS model. Significant relationships ($p < 0.05$) with variation in invasive plant richness by municipality are shown in bold.

Explanatory variable	Coefficient (SE)	p -value
Precipitation amount of the wettest month (kg m ⁻²)	-0.0009 (0.001)	0.103
Mean daily maximum air temperature of the warmest month (°C)	-0.0323 (0.029)	0.259
Ruggedness Index	0.0105 (0.010)	0.272
Minimum distance to coastline	-0.0074 (0.002)	0.003
Municipality area	0.0478 (0.006)	<.001
Density of agricultural areas	0.0014 (0.003)	0.606
Density of forest areas	0.0005 (0.003)	0.86
Density of protected areas	0.0270 (0.015)	0.077
Burnt area in the last 15 years	0.0004 (0.002)	0.868
Soil pH	0.7854 (1.176)	0.505
Density of local hydrographic network	0.0680 (0.090)	0.449
Density of railways	0.1167 (0.154)	0.448
Density of power lines	0.4782 (0.231)	0.039
Travel time to cities (minutes)	-0.0070 (0.002)	0.003
Sampling effort (academic and citizen-science sampling effort)	0.0445 (0.025)	0.071
Number of iNaturalist projects with IAP records	0.1681 (0.189)	0.374
Number of plant nurseries and aquarium stores	0.0778 (0.045)	0.082

Discussion

Our assessment reveals that invasive alien plant species are now established throughout mainland Portugal, yet their patterns of distribution and occupancy are far from uniform. By integrating approximately 85,000 occurrence records from multiple sources, we identified the distribution of 96 invasive plant species across the study area. While previous works estimated the number of invasive alien plants occurring in the country (Almeida and Freitas 2000; Almeida and Freitas 2012; Morais et al. 2017), they did not assess their spatial distribution at the national scale. Other studies examined nationwide distribution patterns but focused on single species (e.g., González et al. 2020). This study provides new insights into the distribution of invasive plant species and their richness in mainland Portugal.




Figure 5. Richness of invasive alien plant species in mainland Portugal, corresponding to the total number of these species recorded in each municipality.

Spatial patterns of invasive species richness

Coastal and urbanized areas, particularly around Lisbon, Oporto, and the Algarve, emerge as hotspots of invasive species richness. In contrast, inland and northeastern regions remain less heavily invaded. Proximity to the coastline and shorter travel times to urban centers were identified as significant predictors of invasive species richness, reinforcing the role of urbanization, human disturbance, and accessibility in driving invasions as favorable conditions for the introduction, establishment, and spread of alien species in many regions (e.g., Pyšek and Richardson 2010; Santos et al. 2011; Dawson et al. 2017; Pyšek et al. 2020; Dimitrakopoulos et al. 2022; Polce et al. 2023; Musarella et al. 2024; Ivison et al. 2025; Montagnani et al. 2025). This reflects the demographic and economic structure of Portugal, which is strongly coast-biased, with population and related activities concentrated along the western and southern coastal regions, particularly in and around the Lisbon and Oporto metropolitan areas (Instituto Nacional de Estatística 2022). Some widespread invasive species in Portugal are known to have been initially brought to major urban centers as ornamentals (e.g., *I. indica* or *Erigeron karvinskianus*; Marchante et al. 2014), while others were introduced specifically to coastal ecosystems in past decades, especially for dune fixation and erosion control (e.g., *A. longifolia* or *C. edulis*; Marchante et al. 2014).

Invasive species richness was also positively associated with municipality size and the density of power-line corridors. The effect of area aligns with the classic species–area relationship, whereby larger spatial units tend to encompass a greater variety of habitats and a broader representation of the regional species pool, supporting more stable and abundant populations and contributing to higher observed species richness (Matthews et al. 2021). On the other hand, linear infrastructures such as power lines may function as ecological corridors. These areas are subject to vegetation cutting interventions, which cause disturbance and eliminate competition—factors known to facilitate the dispersal and establishment of invasive plant species (Dalu et al. 2023). Despite this, the role of such linear infrastructures in plant invasion in Portugal remains underexplored.

Although climate is a known driver of alien species distributions (Dutra Silva et al. 2021), climatic variables were not significant predictors of invasive plant richness in this study. This may reflect the stronger influence of anthropogenic factors, namely human distribution and activities. Moreover, despite its relatively small size globally, mainland Portugal presents relatively diverse climatic conditions, shaped largely by its latitudinal gradient, with warmer, dry Mediterranean-type climates in southern regions and cooler, humid temperate climates in northern coastal areas (Beck et al. 2023). We anticipated a significant relationship between soil pH and invasive species richness, given that calcareous and serpentine soils—known for their high pH and challenging chemical profiles—tend to support fewer invasive plants. However, this expected effect may have been obscured because municipality-level pH values represent averages across diverse soil types, thereby diluting the influence of localized extremes.

Further insights emerged from the cluster analysis (Fig. 3), which revealed four main distribution patterns ranging from narrowly confined inland species to widely distributed taxa spanning both latitudinal and coastal–inland gradients. While most species conformed to expected patterns, notable exceptions emerged (e.g., *C. selloana*, which was classified as widespread, is relatively uncommon in inland regions; González et al. 2020). This highlights how presence-only data at the municipal level may obscure finer-scale spatial variation.

These spatial and ecological patterns carry critical implications for monitoring and management. Species recorded in only a few but geographically dispersed municipalities, such as *L. grandiflora*, may reflect either early-stage invasions or underreporting. Differentiating between these scenarios is crucial, as early detection offers a window of opportunity for effective containment or eradication. Targeted field surveys and systematic monitoring should prioritize such cases to reduce uncertainty and support timely interventions.

Finally, the analysis revealed a moderate positive association ($R^2 = 0.42$) between the number of municipalities in which a species occurs and its mean occupancy, suggesting that widespread species are also more likely to be locally dominant. This is evident for well-established invaders such as *A. dealbata*, *A. donax*, *O. pes-caprae*, and *Phytolacca americana*, whose ecological impacts in mainland Portugal are well documented (Aguar and Ferreira 2013; Castro et al. 2016; Duarte et al. 2020; Marchante et al. 2023). In contrast, some species, such as *A. longifolia*, *A. saligna*, and *C. edulis*, show high local occupancy but limited national spread (mainly in coastal dune areas; Marchante et al. 2014), suggesting environmentally constrained yet intense localized invasions.

Species richness as an indicator: opportunities and limitations

The measure of invasive species richness, derived from municipality-level presence data, provides a valuable and standardized overview of invasive plant distributions across mainland Portugal. While this approach does not capture local abundance—which could offer further insight into ecological dominance and impact—it remains a practical method for identifying general patterns of invasion. As with many large-scale biodiversity assessments, the underlying data are primarily sourced from opportunistic records, including citizen-science contributions. Although these sources provide broad spatial coverage and large volumes of data, they can introduce spatial biases related to observer effort, site accessibility, and species detectability (Isaac et al. 2014; Tiago et al. 2017; Westekemper et al. 2018). As a result, the apparent absence of species in certain areas may reflect lower sampling effort rather than true absence, particularly for less conspicuous or recently introduced taxa.

Moreover, invasive species richness does not differentiate among species in terms of their ecological impact or management urgency. For example, the detection of *Fallopia japonica* in a new location may warrant immediate action due to its known severe ecological consequences (Aguilera et al. 2010), while a new occurrence of *E. karvinskianus* might pose a more localized and manageable concern. While our analysis adopts an equal-weight approach across species, it establishes a foundational framework that can be enhanced through species-specific risk assessments and prioritization strategies, which are vital for informed resource allocation and targeted management.

Despite limitations inherent in opportunistic and citizen-science data (Robinson et al. 2017), these data offer a cost-effective way to collect species records in greater numbers and at more locations. Moreover, using invasive species richness as a pressure indicator is a helpful tool to identify areas where conservation and management efforts should be prioritized (Fleishman et al. 2006; Santos et al. 2011). However, while it is a useful tool for spotting potential invasion hotspots, it should not be interpreted as a direct measure of invasion severity or ecological impact.

Conclusion

This study provides insights into the spatial patterns and environmental and human drivers of invasive plant richness in mainland Portuguese municipalities. Coastal urban areas show the highest invasive species richness, largely driven by greater accessibility. This suggests that municipalities in these regions should prioritize invasive plant management due to their higher levels of impact. At the same time, municipalities with fewer invasive species must act early to prevent their spread, thereby increasing the effectiveness and sustainability of management efforts nationwide. Despite relying on the total number of invasive plants in each municipality, and not their abundance, the results offer a valuable baseline for identifying invasion hotspots and guiding management priorities. Continued monitoring and systematic data collection are essential to support effective prevention and control of plant invasions across the country.

Acknowledgments

We would like to express our thanks to the many experts who contributed by providing useful information on occurrence records: Ascendi, Altri, Infraestruturas de Portugal, S.A., Liliana Duarte, Mónica Almeida, Sílvia Martins, MED–Mediterranean Institute for Agriculture, Environment and Development, and the Botany Lab/Applied Ecology and Conservation Research Group of the University of Évora; the National Electricity Transmission Network shapefile provided by David Almeida (Redes Energéticas Nacionais [REN]); and information on plant nurseries and aquarium store data collected by Ana Sofia Nunes (Nunes et al. unpublished).

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

We used AI to polish the manuscript text and help develop the R code for the methodology.

Funding

RF was supported by a grant (PRT/BD/152100/2021 and <https://doi.org/10.54499/PRT/BD/153505/2021>) funded by the Portuguese Foundation for Science and Technology (FCT) under the MIT Portugal Program. AC was supported by a grant (PRT/BD/152100/2021 and <https://doi.org/10.54499/PRT/BD/152100/2021>) funded by the Portuguese FCT under the MIT Portugal Program. AC, RF, and CC acknowledge support from FCT through funding to the CEG/IGOT Research Unit (UID/295/2025 and <https://doi.org/10.54499/UID/00295/2025>).

Author contributions

RF: Conceptualization, Data curation, Methodology, Writing—original draft, Writing—review and editing. AC: Methodology, Writing—original draft, Writing—review and editing. HM: Conceptualization, Supervision, Writing—review and editing. EM: Conceptualization, Supervision,

Writing—review and editing. CC: Conceptualization, Supervision, Methodology, Writing—original draft, Writing—review and editing.

Author ORCIDs

Raquel Fernandes  <https://orcid.org/0000-0002-3401-115X>

Andry Castro  <https://orcid.org/0000-0001-5635-5271>

Hélia Marchante  <https://orcid.org/0000-0002-3247-5663>

Elizabete Marchante  <https://orcid.org/0000-0003-1303-7489>

César Capinha  <https://orcid.org/0000-0002-0666-9755>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Aguir F, Ferreira MT (2013) Plant invasions in the rivers of the Iberian Peninsula, south-western Europe: A review. *Plant Biosystems* 147(4): 1107–1119. <https://doi.org/10.1080/11263504.2013.861539>
- Aguilera AG, Alpert P, Dukes JS, Harrington R (2010) Impacts of the invasive plant *Fallopia japonica* (Houtt.) on plant communities and ecosystem processes. *Biological Invasions* 12: 1243–1252. <https://doi.org/10.1007/s10530-009-9543-z>
- Alin A (2010) Multicollinearity. *Wiley Interdisciplinary Reviews: Computational Statistics* 2(3): 370–374. <https://doi.org/10.1002/wics.84>
- Almeida J, Freitas H (2000) A flora exótica e invasora de Portugal. *Portugaliae Acta Biol* 19: 159–176.
- Almeida J, Freitas H (2012) Exotic flora of continental Portugal—a new assessment. *Boccone* 24: 231–237. <https://www.researchgate.net/publication/235457718>
- Alves A, Marcelino F, Gomes E, Rocha J, Caetano M (2022) Spatiotemporal Land-Use Dynamics in Continental Portugal 1995–2018. *Sustainability* 14(23). <https://doi.org/10.3390/su142315540>
- Beaury E, Sofaer H, Early R, Pearse I, Blumenthal D, Corbin J, Diez J, Dukes J, Barnett D, Ibáñez I, Petri L, Vilà M, Bradley B (2023) Macroscale analyses suggest invasive plant impacts depend more on the composition of invading plants than on environmental context. *Global Ecology and Biogeography* 32(11): 1964–1976. <https://doi.org/10.1111/geb.13749>
- Beck H, McVicar T, Vergopolan N, Berg A, Lutsko N, Dufour A, Zeng Z, Jiang X, van Dijk A, Miralles D (2023) High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections. *Scientific Data* 10(1): 724. <https://doi.org/10.1038/s41597-023-02549-6>
- Berneiz I, Ferreira M, Albuquerque A, Aguiar F (2005) Relations between river plant richness in the Portuguese floodplains and the widespread water knotgrass (*Paspalum Paspalodes*). *Hydrobiologia* 551(1): 121–130. <https://doi.org/10.1007/s10750-005-4454-1>
- Bholowalia P, Kumar A (2014) EBK-Means: A clustering technique based on elbow method and K-means in WSN. *International Journal of Computer Applications* 105(9).
- Capinha C, Essl F, Porto M, Seebens H (2023) The worldwide networks of spread of recorded alien species. *Proceedings of the National Academy of Sciences of the United States of America* 120(1): e2201911120. <https://doi.org/10.1073/pnas.2201911120>
- Castro S, Castro M, Ferrero V, Costa J, Tavares D, Navarro L, Loureiro J (2016) Invasion fosters change: independent evolutionary shifts in reproductive traits after *oxalis pes-caprae* L. Introduction. *Frontiers in Plant Science* 7: 874. <https://doi.org/10.3389/fpls.2016.00874>
- Chefaoui RM, Chozas S (2019) Abandonment of traditional saltworks facilitates degradation of halophytic plant communities and *Carpobrotus edulis* invasion. *Applied Vegetation Science* 22(3): 444–453. <https://doi.org/10.1111/avsc.12436>

- Costa J, Neto C, Arsénio P, Capelo J (2009) Geographic variation among Iberian communities of the exotic halophyte *Cotula coronopifolia*. *Botanica Helvetica* 119(1): 53–61. <https://doi.org/10.1007/s00035-009-0056-2>
- Dalu T, Stam E, Ligege M, Cuthbert R (2023) Highways to invasion: Powerline servitudes as corridors for alien plant invasions. *African Journal of Ecology* 61(2): 379–388. <https://doi.org/10.1111/aje.13121>
- Dawson W, Moser D, Van Kleunen M, Kreft H, Pergl J, Pyšek P, Weigelt P, Winter M, Lenzner B, Blackburn T, Dyer E, Cassey P, Scrivens S, Economo E, Guénard B, Capinha C, Seebens H, García-Díaz P, Nentwig W, et al. (2017) Global hotspots and correlates of alien species richness across taxonomic groups. *Nature Ecology & Evolution* 1. <https://doi.org/10.1038/s41559-017-0186>
- DGEEC (2024) RENATES - Registo Nacional de Teses e Dissertações. <https://renates2.dgeec.mec.pt/>
- DGEEC/ME-MCTES, PORDATA (2021) Alunos matriculados no ensino superior: total, por subsistema e por tipo de ensino. <https://www.pordata.pt/pt/estatisticas/educacao/ensino-superior/alunos-inscritos-no-ensino-superior-por-subsistema-de-ensino>
- DGEEC/ME-MCTES, PORDATA (2024) Estabelecimentos de ensino superior: total e por tipo de ensino. <https://www.pordata.pt/municipios/estabelecimentos+de+ensino+superior+total+e+por+tipo+de+ensino-336>
- Dimitrakopoulos P, Koukoulas S, Michelaki C, Galanidis A (2022) Anthropogenic and environmental determinants of alien plant species spatial distribution on an island scale. *The Science of the Total Environment* 805: 150314. <https://doi.org/10.1016/j.scitotenv.2021.150314>
- Direção-Geral do Território (2024) Carta Administrativa Oficial de Portugal. <https://www.dgterritorio.gov.pt/cartografia/cartografia-tematica/caop>
- Direção-Geral do Território (2020) Carta de Uso e Ocupação do Solo para 2018. <https://www.dgterritorio.gov.pt/Carta-de-Uso-e-Ocupacao-do-Solo-para-2018?language=en>
- Dormann C, McPherson J, Araújo M, Bivand R, Bolliger J, Carl G, Davies R, Hirzel A, Jetz W, Daniel Kissling W, Kühn I, Ohlemüller R, Peres-Neto P, Reineking B, Schröder B, Schurr F, Wilson R (2007) Methods to account for spatial autocorrelation in the analysis of species distributional data: A review. *Ecography* 30(5): 609–628. <https://doi.org/10.1111/j.2007.0906-7590.05171.x>
- Duarte L, Gomes C, Marchante H, Marchante E (2020) Integrating knowledge of ecological succession into invasive alien plant management: A case study from Portugal. *Applied Vegetation Science* 23(3): 328–339. <https://doi.org/10.1111/avsc.12488>
- Duarte L, Marchante E, Marchante H (2023) Managing an invasive tree in coastal dunes: The importance of follow-up treatments to improve the recovery of protected habitats. *Frontiers in Environmental Science* 11: 1113876. <https://doi.org/10.3389/fenvs.2023.1113876>
- Dutra Silva L, Bento Elias R, Silva L (2021) Modelling invasive alien plant distribution: A literature review of concepts and bibliometric analysis. *Environmental Modelling & Software* 145: 105203. <https://doi.org/10.1016/j.envsoft.2021.105203>
- Earth Resources Observation and Science (EROS) Center (2018) USGS EROS Archive - Digital Elevation - Shuttle Radar Topography Mission (SRTM) 1 Arc-Second Global. <https://www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-shuttle-radar-topography-mission-srtm-1>
- Ferreira V, Figueiredo A, Graça M, Marchante E, Pereira A (2021) Invasion of temperate deciduous broadleaf forests by N-fixing tree species – consequences for stream ecosystems. *Biological Reviews of the Cambridge Philosophical Society* 96(3): 877–902. <https://doi.org/10.1111/brev.12682>
- Fleishman E, Noss R, Noon B (2006) Utility and limitations of species richness metrics for conservation planning. *Ecological Indicators* 6(3): 543–553. <https://doi.org/10.1016/j.ecoind.2005.07.005>
- Flora de Portugal Interactiva (2024) Flora de Portugal Interactiva. <https://flora-on.pt/>
- GBIF Secretariat (2023) GBIF Backbone Taxonomy. Checklist dataset. <https://doi.org/10.15468/39omei>

- González F, Serrano B, Urchaga A, García de Enterría S, Marchante H, Almeida M, Alves HN, Herrero Á, García A (2020) Estratégia transnacional de luta contra a Cortaderia selloana no Arco Atlântico no quadro do projeto LIFE STOP Cortaderia (LIFE17 NAT/ES/000495). Projeto LIFE STOP Cortaderia, Santander.
- Große-Stoltenberg A, Lizarazo I, Brundu G, Paiva Gonçalves V, Prado Osco L, Masemola C, Müllerová J, Werner C, Kotze I, Oldeland J (2023) Remote Sensing of Invasive Australian Acacia Species: State of the Art and Future Perspectives. In: Richardson DM, Le Roux JJ, Marchante E (Eds) Wattles: Australian Acacia Species Around the World. CABI, Boston, 474–495. <https://doi.org/10.1079/9781800622197.0029>
- Gutierrez F, Reis E, Lobo A (2011) Acacia saligna (Labill.) H. Wendl in the Sesimbra County: Invaded habitats and potential distribution modeling. Journal of Coastal Research 64: 403–407. <https://www.researchgate.net/publication/235921262>
- Hansen W, Wollny J, Otte A, Eckstein R, Ludwig K (2020) Invasive legume affects species and functional composition of mountain meadow plant communities. Biological Invasions 23(1): 281–296. <https://doi.org/10.1007/s10530-020-02371-w>
- iNaturalist (2024) iNaturalist. <https://www.inaturalist.org>
- INE, PORDATA (2024) População residente com 15 e mais anos segundo os Censos: total e por nível de escolaridade completo mais elevado. <https://www.pordata.pt/municipios/populacao+residente+com+15+e+mais+anos+segundo+os+censos+total+e+por+nivel+de+escolaridade+completo+mais+elevado-69>
- Instituto da Conservação da Natureza e das Floresta [ICNF] (2022) Reservas da Biosfera - Programa Man and the Biosphere da UNESCO. <https://sig.icnf.pt/portal/home/item.html?id=6d5b9ca-6f0e746a28c1a06170a0a76cc>
- Instituto da Conservação da Natureza e das Floresta [ICNF] (2023a) Rede Nacional de Áreas Protegidas (RNAP). <https://sig.icnf.pt/portal/home/item.html?id=02b7a03f8fbd4dada77f5f3e5f91f186>
- Instituto da Conservação da Natureza e das Floresta [ICNF] (2023b) Sítios RAMSAR - Convenção sobre Zonas Húmidas. <https://sig.icnf.pt/portal/home/item.html?id=8eb4f474eab3491b9a-0b40a11b83d170>
- Instituto da Conservação da Natureza e das Floresta [ICNF] (2023c) Zonas de Proteção Especial da Diretiva Aves (ZPE) - RN2000. <https://sig.icnf.pt/portal/home/item.html?id=a158877a57eb4f-5fbad767d36e261fab>
- Instituto da Conservação da Natureza e das Floresta [ICNF] (2023d) Zonas Especiais de Conservação / Sítios da Diretiva Habitats (ZEC/SIC) - RN2000. <https://sig.icnf.pt/portal/home/item.html?id=a158877a57eb4f5fbad767d36e261fab>
- Instituto da Conservação da Natureza e das Floresta [ICNF] (2024) Territórios ardidos (Área ardida entre 1975 e 2023). <https://sig.icnf.pt/portal/home/item.html?id=6d5b9ca6f0e746a28c1a06170a0a76cc>
- Instituto Nacional de Estatística (2022) Censos 2021. XVI Recenseamento Geral da População. VI Recenseamento Geral da Habitação: Resultados definitivos. <https://www.ine.pt/xurl/pub/65586079>
- IPBES (2023) Summary for Policymakers of the Thematic Assessment Report on Invasive Alien Species and their Control of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES secretariat. <https://doi.org/10.5281/zenodo.7430692>
- Isaac N, van Strien A, August T, de Zeeuw M, Roy DB (2014) Statistics for citizen science: Extracting signals of change from noisy ecological data. Methods in Ecology and Evolution 5(10): 1052–1060. <https://doi.org/10.1111/2041-210X.12254>
- Ives A (2015) For testing the significance of regression coefficients, go ahead and log-transform count data. Methods in Ecology and Evolution 6(7): 828–835. <https://doi.org/10.1111/2041-210X.12386>
- Iverson K, Howard C, Baldini L, Essl F, Pyšek P, Dawson W, Speed J (2025) Horizon scanning of potential invasive alien plant species and their distribution in Norway under a changing climate. Ecography c07604. <https://doi.org/10.1002/ecog.07604>

- Kalusová V, Čeplová N, Danihelka J, Večeřa M, Pyšek P, Albert A, Anastasiu P, Biurrun I, Boch S, Cottaz C, Essl F, Kuzemko A, Maslo S, Mifsud S, Protopopova V, Shevera M, Sirbu C, Svenning J, Welk E, Axmanová I (2024) Alien plants of Europe: An overview of national and regional inventories. *Preslia* 96(2): 149–182. <https://doi.org/10.23855/preslia.2024.149>
- Karger D, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler M (2017) Climatologies at high resolution for the earth's land surface areas. *Scientific Data* 4: 170122. <https://doi.org/10.1038/sdata.2017.122>
- Karger D, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler M (2018) Data from: Climatologies at high resolution for the earth's land surface areas. Dryad Digital Repository.
- Kummu M, Taka M, Guillaume J (2015) Data from: Gridded global datasets for Gross Domestic Product and Human Development Index over 1990–2015. [Dryad, Dataset]
- Latombe G, Pyšek P, Jeschke J, Blackburn T, Bacher S, Capinha C, Costello MJ, Fernández M, Gregory R, Hobern D, Hui C, Jetz W, Kumschick S, McGrannachan C, Pergl J, Roy H, Scalera R, Squires ZE, Wilson J, et al. (2017) A vision for global monitoring of biological invasions. *Biological Conservation* 213: 295–308. <https://doi.org/10.1016/j.biocon.2016.06.013>
- Le Maitre D, Gaertner M, Marchante E, Ens E, Holmes P, Pauchard A, O'Farrell PJ, Rogers AM, Blanchard R, Blignaut J, Richardson D (2011) Impacts of invasive Australian acacias: Implications for management and restoration. *Diversity & Distributions* 17(5): 1015–1029. <https://doi.org/10.1111/j.1472-4642.2011.00816.x>
- Lopez B, Allen J, Dukes J, Lenoir J, Vil M, Blumenthal DM, Beaury E, Fusco E, Laginhas B, Lyn Morelli T, Sorte C, Maceda-Veiga A, Whitlock R, Bradley BA (2022) Global environmental changes more frequently offset than intensify detrimental effects of biological invasions. *Proceedings of the National Academy of Sciences of the United States of America* 119(9): 1–7. <https://doi.org/10.1073/pnas>
- Maltez-Mouro S, Maestre F, Freitas H (2009) Weak effects of the exotic invasive *Carpobrotus edulis* on the structure and composition of Portuguese sand-dune communities. *Biological Invasions* 12(7): 2117–2130. <https://doi.org/10.1007/s10530-009-9613-2>
- Marchante H, Marchante E, Freitas H (2003) Invasion of the Portuguese dune ecosystems by the exotic species *Acacia longifolia* (Andrews) Willd.: effects at the community level. *Plant Invasions: Ecological Threats and Management Solutions*: 75–85. <https://www.researchgate.net/publication/232717067>
- Marchante H, Morais M, Freitas H, Marchante E (2014) Guia prático para a identificação de plantas invasoras em Portugal. Imprensa da Universidade de Coimbra. <https://doi.org/10.14195/978-989-26-0786-3>
- Marchante E, Gouveia AC, Brundu G, Marchante H (2023) Australian Acacia Species in Europe. In: Richardson DM, Le Roux JJ, Marchante E (Eds) *Wattles: Australian Acacia Species Around the World*. CABI, Boston, 148–166. <https://doi.org/10.1079/9781800622197.0010>
- Martins F, Alegria C, Gil A (2016) Mapping invasive alien *Acacia dealbata* Link using ASTER multispectral imagery: A case study in central-eastern of Portugal. *Forest Systems* 25(3). <https://doi.org/10.5424/fs/2016253-09248>
- Matthews T, Triantis K, Whittaker RJ (2021) *The Species–Area Relationship: Theory and Application*. Cambridge University Press, 1–502. <https://doi.org/10.1017/9781108569422>
- Milanović M, Knapp S, Pyšek P, Kühn I (2020) Linking traits of invasive plants with ecosystem services and disservices. *Ecosystem Services* 42: 1–10. <https://doi.org/10.1016/j.ecoser.2020.101072>
- Ministério do Ambiente (1999) Diário da República n.o 295/1999, Série I-A de 1999-12-21, páginas 9100–9115. <https://diariodarepublica.pt/dr/detalhe/decreto-lei/565-1999-661769>
- Ministério do Ambiente e Transição Energética (2019) Diário da República n.o 130/2019, Série I de 2019-07-10, 3428–3442. <https://diariodarepublica.pt/dr/detalhe/decreto-lei/92-2019-123025739>

- Montagnani C, Gentili R, Karrer G, Caronni S, Acosta A, Adorni M, Alessandrini A, Ardenghi NMG, Buffa G, Bartolucci F, Belisario A, Bonali F, Bonini I, Brancaloni L, Buccheri M, Castello M, Conti F, Di Carlo F, Domina G, Fenu G, Forte L, Galasso G, Gubellini L, Iberite M, Labadini A, Luccioli E, Maccioni S, Marcucci R, Martini F, Medagli P, Montanari S, Morelli V, Nepi C, Nicoletta G, Pellizzari M, Enrico Vito Perrino, Piccoli F, Prisco I, Prosser F, Repetto G, Rosati L, Selvaggi A, Tavano M, Vallariello R, Vidali M, Vivona L, Wagensommer RP, Citterio S (2025) The invasion history of *Ambrosia psilostachya* (Asteraceae) in Italy: First record and distribution over more than one century. *Plant Biosystems* 159(3): 522–533. <https://doi.org/10.1080/11263504.2025.2485990>
- Morais M, Marchante E, Marchante H (2017) Big troubles are already here: Risk assessment protocol shows high risk of many alien plants present in Portugal. *Journal for Nature Conservation* 35: 1–12. <https://doi.org/10.1016/j.jnc.2016.11.001>
- Mouta N, Silva R, Pinto E, Vaz A, Alonso J, Gonçalves J, Honrado J, Vicente J (2023) Sentinel-2 Time Series and Classifier Fusion to Map an Aquatic Invasive Plant Species along a River—The Case of Water-Hyacinth. *Remote Sensing* 15(13): 3248. <https://doi.org/10.3390/rs15133248>
- Musarella C, Laface V, Angiolini C, Bacchetta G, Bajona E, Banfi E, Barone G, Biscotti N, Bonsanto D, Calvia G, Cambria S, Capuano A, Caruso G, Crisafulli A, Del Guacchio E, Di Gristina E, Domina G, Fanfarillo E, Fascetti S, Fiaschi T, Galasso G, Mascia F, Mazzacava G, Mei G, Minisale P, Motti R, Perrino EV, Picone R, Pinzani L, Podda L, Potenza G, Rosati L, Stinca A, Tavilla G, Villano C, Wagensommer R, Spampinato G (2024) New alien plant taxa for Italy and Europe: An update. *Plants* 13: 620. <https://doi.org/10.3390/plants13050620>
- Naimi B (2022) Package usdm. Uncertainty Analysis for Species Distribution Models. <http://r-gis.net>
- Nakagawa S, Johnson P, Schielzeth H (2017) The coefficient of determination R² and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *Journal of the Royal Society, Interface* 14(134): 1–11. <https://doi.org/10.1098/rsif.2017.0213>
- Official Journal of the European Union (2014) Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32014R1143&qid=1719589404376>
- Official Journal of the European Union (2022) Commission implementing regulation (EU) 2022/1203 of 12 July 2022 amending Implementing Regulation (EU) 2016/1141 to update the list of invasive alien species of Union concern. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32022R1203>
- Oficialdegui F, Zamora-Marín J, Guareschi S, Anastácio P, García-Murillo P, Ribeiro F, Miranda R, Cobo F, Gallardo B, García-Berthou E, Boix D, Arias A, Cuesta JA, Medina L, Almeida D, Banha F, Barca S, Biurrun I, Cabezas MP, Calero S, Campos JA, Capdevila-Argüelles L, Capinha C, Casals F, Clavero M, Encarnação J, Fernández-Delgado C, Franco J, Guillén A, Hermoso V, Machordom A, Martelo J, Mellado-Díaz A, Morcillo F, Osoz J, Perdices A, Pou-Rovira Q, Rodríguez-Merino A, Ros M, Ruiz-Navarroac A, Sánchez MI, Sánchez-Fernández D, Sánchez-González JR, Sánchez-Gullón E, Teodósio MA, Torralva M, Vieira-Lanero R, Oliva-Paterna FJ (2023) A horizon scan exercise for aquatic invasive alien species in Iberian inland waters. *The Science of the Total Environment* 869: 1–16. <https://doi.org/10.1016/j.scitotenv.2023.161798>
- OpenStreetMap. Geofabrik GmbH, and OpenStreetMap Contributors (2018) Download OpenStreetMap data for this region: Portugal. <https://download.geofabrik.de/europe/portugal.html>
- Pabst R, Dias F, Borda-de-Água L, Rodríguez-González P, Capinha C (2022) Assessing and predicting the distribution of riparian invasive plants in continental Portugal. *Frontiers in Ecology and Evolution* 10: 1–10. <https://doi.org/10.3389/fevo.2022.875578>
- Pádua L, Antão-Geraldes A, Sousa J, Rodrigues M, Oliveira V, Santos D, Miguens M, Castro J (2022a) Water Hyacinth (*Eichhornia crassipes*) Detection using coarse and high resolution multispectral data. *Drones* 6(2): 1–14. <https://doi.org/10.3390/drones6020047>

- Pádua L, Duarte L, Antão-Geraldes A, Sousa J, Castro J (2022b) Spatio-temporal water hyacinth monitoring in the lower Mondego (Portugal) using remote sensing data. *Plants* 11(24): 1–17. <https://doi.org/10.3390/plants11243465>
- Pinheiro J, Bates D, DebRoy S, Sarkar D, Heisterkamp S, Willigen B, Ranke J (2023) Package nlme. Linear and Nonlinear Mixed Effects Models. <https://bugs.r-project.org>
- Poggio L, de Sousa L, Batjes N, Heuvelink G, Kempen B, Ribeiro E, Rossiter D (2021) SoilGrids 2.0: Producing soil information for the globe with quantified spatial uncertainty. *Soil* 7(1): 217–240. <https://doi.org/10.5194/soil-7-217-2021>
- Polce C, Cardoso A, Deriu I, Gervasini E, Tsiamis K, Vigiak O, Zulian G, Maes J (2023) Invasive alien species of policy concerns show widespread patterns of invasion and potential pressure across European ecosystems. *Scientific Reports* 13(1): 8124. <https://doi.org/10.1038/s41598-023-32993-8>
- Powell K, Chase J, Knight T (2011) A synthesis of plant invasion effects on biodiversity across spatial scales. *American Journal of Botany* 98(3): 539–548. <https://doi.org/10.3732/ajb.1000402>
- Pyšek P, Richardson D (2010) Invasive species, environmental change and management, and health. *Annual Review of Environment and Resources* 35: 25–55. <https://doi.org/10.1146/annurev-environ-033009-095548>
- Pyšek P, Hulme P, Simberloff D, Bacher S, Blackburn T, Carlton J, Dawson W, Essl F, Foxcroft L, Genovesi P, Jeschke J, Kühn I, Liebhold A, Mandrak N, Meyerson L, Pauchard A, Pergl J, Roy HE, Seebens H, et al. (2020) Scientists’ warning on invasive alien species. *Biological Reviews of the Cambridge Philosophical Society* 95(6): 1511–1534. <https://doi.org/10.1111/brv.12627>
- R Core Team (2022) The R Project for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Ramos C, Ramos-Pereira A (2020) Landscapes of Portugal: Paleogeographic Evolution, Tectonics and Geomorphology. In: Vieira G, Zêzere JL, Mora C (Eds) *Landscapes and Landforms of Portugal*. Springer International, 3–31. https://doi.org/10.1007/978-3-319-03641-0_1
- Raposo M, Pinto Gomes C, Nunes L (2021) Evaluation of species invasiveness: A case study with *Acacia dealbata* Link. on the slopes of Cabeça (Seia-Portugal). *Sustainability* 13(20): 1–10. <https://doi.org/10.3390/su132011233>
- Ren J, Chen J, Xu C, Van De Koppel J, Thomsen M, Qiu S, Cheng F, Song W, Liu QX, Xu C, Bai J, Zhang Y, Cui B, Bertness M, Silliman B, Li B, He Q (2021) An invasive species erodes the performance of coastal wetland protected areas. *Science Advances* 7: eabi8943. <https://doi.org/10.1126/sciadv.abi8943>
- Reynolds C, Venter N, Cowie B, Marlin D, Mayonde S, Tocco C, Byrne M (2020) Mapping the socio-ecological impacts of invasive plants in South Africa: Are poorer households with high ecosystem service use most at risk? *Ecosystem Services* 42: 101075. <https://doi.org/10.1016/j.ecoser.2020.101075>
- Richardson D, Van Wilgen BW (2004) Invasive alien plants in South Africa: How well do we understand the ecological impacts? *South African Journal of Science* 100: 45–52.
- Robinson O, Ruiz-Gutierrez V, Fink D (2017) Correcting for bias in distribution modelling for rare species using citizen science data. *Diversity & Distributions* 24(4): 460–472. <https://doi.org/10.1111/ddi.12698>
- Rodríguez F, Lombardero-Vega M, San Juan L, Vecillas L, Morchón E, Liendo D, Uranga M, Gandarillas A (2021) Allergenicity to worldwide invasive grass *Cortaderia selloana* as environmental risk to public health. *Scientific Reports* 11: 24426. <https://doi.org/10.1038/s41598-021-03581-5>
- Rodríguez-Merino A (2023) Identifying and managing areas under threat in the Iberian peninsula: An invasion risk atlas for non-native aquatic plant species as a potential tool. *Plants* 12(17). <https://doi.org/10.3390/plants12173069>
- Roekaerts M (2002) The Biogeographical Regions Map of Europe. Basic principles of its creation and overview of its development. European Environment Agency. <https://www.researchgate.net/publication/224840052>

- Santos M, Freitas R, Crespi AL, Hughes S, Cabral JA (2011) Predicting trends of invasive plants richness using local socio-economic data: An application in North Portugal. *Environmental Research* 111(7): 960–966. <https://doi.org/10.1016/j.envres.2011.03.014>
- Seebens H, Meyerson LA, Rahlao SJ, Lenzner B, Tricarico E, Aleksanyan A, Courchamp F, Keskin E, Saeedi H, Tawake A, Pyšek P (2023) Chapter 2: Trends and status of alien and invasive alien species. In: Roy HE, Pauchard A, Stoett P, Renard Truong T (Eds) *Thematic Assessment Report on Invasive Alien Species and their Control of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. IPBES secretariat, Bonn. <https://doi.org/10.5281/zenodo.7430725>
- Sirbu C, Miu I, Gavrilidis A, Gradinaru SR, Niculae I, Preda C, Oprea A, Urziceanu M, Camen-Comanescu P, Nagoda E, Sirbu IM, Memedemin D, Anastasiu P (2022) Distribution and pathways of introduction of invasive alien plant species in Romania. *NeoBiota* 75: 1–21. <https://doi.org/10.3897/neobiota.75.84684>
- Sousa R, Marchante E, Vicente J, Vaz A (2018) *Publicação Científica Sobre Invasões Biológicas*. In: Vicente J, Queiroz A, Silva L, Marchante E, Honrado J (Eds) *As Invasões Biológicas em Portugal: História, Diversidade e Gestão*. Arte e Ciência, Porto.
- Tiago P, Ceia-Hasse A, Marques T, Capinha C, Pereira HM (2017) Spatial distribution of citizen science casuistic observations for different taxonomic groups. *Scientific Reports* 7(1): 1–9. <https://doi.org/10.1038/s41598-017-13130-8>
- Verloove F, Dana E, Alves P (2017) *Baccharis spicata* (Asteraceae), a new potentially invasive species to Europe. *Plant Biosystems* 152(3): 416–426. <https://doi.org/10.1080/11263504.2017.1303001>
- Vicente J, Alves P, Randin C, Guisan A, Honrado J (2010) What drives invasibility? A multi-model inference test and spatial modelling of alien plant species richness patterns in northern Portugal. *Ecography* 33(6): 1081–1092. <https://doi.org/10.1111/j.1600-0587.2010.6380.x>
- Vicente J, Kueffer C, Richardson D, Vaz AS, Cabral J, Hui C, Araújo M, Kühn I, Kull C, Verburg P, Marchante E, Honrado J (2019) Different environmental drivers of alien tree invasion affect different life-stages and operate at different spatial scales. *Forest Ecology and Management* 433: 263–275. <https://doi.org/10.1016/j.foreco.2018.10.065>
- Vieites-Blanco C, González-Prieto S (2020) Invasiveness, ecological impacts and control of acacias in southwestern Europe- A review. *Web Ecology* 20(2): 33–51. <https://doi.org/10.5194/we-20-33-2020>
- Weiss D, Nelson A, Gibson H, Temperley W, Peedell S, Lieber A, Hancher M, Poyart E, Belchior S, Fullman N, Mappin B, Dalrymple U, Rozier J, Lucas T, Howes R, Tusting L, Kang S, Cameron E, Bisanzio D, ... Gething P (2018) A global map of travel time to cities to assess inequalities in accessibility in 2015. *Nature* 553(7688): 333–336. <https://doi.org/10.1038/nature25181>
- Westekemper K, Reinecke H, Signer J, Meißner M, Herzog S, Balkenhol N (2018) Stay on trails – effects of human recreation on the spatiotemporal behavior of red deer *Cervus elaphus* in a German national park. *Wildlife Biology* 2018(1): 1–9. <https://doi.org/10.2981/wlb.00403>

Supplementary material 1

Results with all occurrence data

Authors: Raquel Fernandes, Andry Castro, Hélia Marchante, Elizabete Marchante, César Capinha

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendata-commons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.104.163291.suppl1>

Supplementary material 2

Markdown

Authors: Raquel Fernandes, Andry Castro, Hélia Marchante, Elizabete Marchante, César Capinha

Data type: docx

Explanation note: R code used in our study.

Copyright notice: This dataset is made available under the Open Database License (<http://opendata-commons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.104.163291.suppl2>

Supplementary material 3

Results without FloraOn data

Authors: Raquel Fernandes, Andry Castro, Hélia Marchante, Elizabete Marchante, César Capinha

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendata-commons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.104.163291.suppl3>

Supplementary material 4

Atlas of invasive plant species

Authors: Raquel Fernandes, Andry Castro, Hélia Marchante, Elizabete Marchante, César Capinha

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendata-commons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.104.163291.suppl4>