

Research article

New records of the Bulgarian endemic species *Pilemia serriventris* Holzschuh, 1984 (Cerambycidae: Lamiinae)

Denis Gradinarov¹, Yana Petrova²

(1) Faculty of Biology, Sofia University “St Kliment Ohridski”, 8 Dragan Tzankov Blvd, 1164 Sofia, Bulgaria, dgradinarov@abv.bg ✉; <https://orcid.org/0000-0002-6128-5009> 

(2) National Genetic Laboratory, 2 Zdrave Street, 1431 Sofia, Bulgaria, yanagradinarova@abv.bg ✉; <https://orcid.org/0000-0002-8443-6228> 

Abstract: Bulgarian endemic species *Pilemia serriventris* (Holzschuh, 1984) is reported for the first time with exact data from several localities, including its type locality near Harmanli. The species is distributed in SE Bulgaria along river valleys from the Aegean and the Black Sea drainage basins, inhabiting both riverside and roadside habitats. The host plant of the species appears to be *Anchusa procera* Besser, and the earlier reports of host plants are doubtful.

Keywords: Balkan Peninsula, distribution, host plants, taxonomy

Introduction

The genus *Pilemia* Fairmaire, 1864 includes two subgenera – *Pilemia* Fairmaire, 1864 and *Pseudopilemia* Kasatkin, 2018 (Danilevsky, 2020). Two species of subgenus *Pilemia* s. str. are known from Bulgaria – *P. tigrina* (Mulsant, 1851) and *P. serriventris* Holzschuh, 1984 (Danilevsky, 2020). The first species is monophagous on *Anchusa barrelieri* (All.) Vitman (Boraginaceae) (Tóth et al., 2016) and in Bulgaria it has been reported from a number of localities in Western Stara Planina Mts, Sofia Valley and Northern Bulgaria (Gradinarov, 2016; Georgiev 2020; Gradinarov & Petrova, 2019; 2021).

Pilemia serriventris (Cerambycidae: Lamiinae) is a Bulgarian endemic species (Migliaccio et al., 2007; Danilevsky, 2020), described by Holzschuh (1984) from Harmanli (type locality: “Charmanli”). There are not many other records of the species after its description. The species was reported from “Harmanli-Ljubimec” (Rejzek et al., 2001), Bistrets Vill. (Strandzha Mts) (Georgiev et al., 2005) and from “Lubimec near Harmanli” (Hoskovec et al., 2023). In all cases of finding the species, no geographic coordinates or data on its habitat have been published.

Two species of family Boraginaceae have been reported as host plants of *P. serriventris* – *Cynoglossum officinale* L. (Rejzek et al., 2001) and *A. barrelieri* (Georgiev et al., 2005, repeated by Migliaccio et al., 2007).

In the present work we report several new localities of the species in SE Bulgaria, as well as a new data on the host plants of the species.

Methods

The material for the present study was collected by the authors in the period 2018 – 2023 from Eastern Rhodopes Mts, Maritsa River Valley and Strandzha Mts (Fig. 1). Adult beetles were hand collected from the host plants. The pictures were taken by using digital camera Olympus SP-820UZ (Fig. 1C), Canon PowerShot SX420 IS (Fig. 1A, B, D–F, Fig. 2) and a combination of Canon EOS 2000D digital camera, PRO-CA Camera Adapter, and a microscope Olympus SZ61 (Figs 3–6). The collected specimens are preserved in the Zoological Collection of Sofia University, Faculty of Biology (BFUS).




Fig. 1. Habitats of *Pilemia serriventris* in Bulgaria – (A) Lyubimets locality, 24.v.2021, (B) Harmanly locality, 28.v.2023, (C) Kotlari locality, 27.v.2018 (D) Kotlari locality, 16.iv.2023, (E) Slavyanovo locality, 26.v.2023, (F) Prohod locality, 25.v.2023.

Results and discussion

Pilemia serriventris (Holzschuh, 1984)

Material: Maritsa Riv. Valley, Lyubimets, right bank of Maritsa Riv. (Fig. 1A), 41°51.245'N, 26°05.113'E,

60 m a.s.l., 24.v.2021, roadside verges, 1 ♂, 1 ♀ (copulating pair), D. Gradinarov & Y. Petrova leg.; the same locality, 26.v.2023, 1 ♀, D. Gradinarov & Y. Petrova leg.; Maritsa Riv. Valley, E of Harmanly, right bank of Maritsa Riv. (Fig. 1B), 41°55.564'N, 25°56.000'E, 76 m a.s.l., 28.v.2023, riverine




Fig. 2. Males of *Pilemia serriventris* on host plants – A) Prohod Vill., 25.v.2023; B) Kotlari Vill., 27.v.2023.

vegetation, 1 ♂, 1 ♀ (copulating pair), D. Gradinarov & Y. Petrova leg.; Eastern Rhodopes Mts, 1,5 km SW Kotlari Vill., right bank of Arda Riv. (Fig. 1C, D), 41°36.914'N, 25°43.647'E, 160 m a.s.l., 27.v.2018, riverine vegetation, 2 ♂♂, 1 ♀ (copulating pair including), D. Gradinarov & Y. Petrova leg.; the same locality, 05.vi.2020, 1 ♂, 2 ♀♀, D. Gradinarov & Y. Petrova leg.; the same locality, 27.v.2023, 7 ♂♂, 7 ♀♀, D. Gradinarov & Y. Petrova leg.; Eastern Rhodopes Mts, 1 km W of Slavyanovo Vill., left bank of Biserska Reka Riv. (Fig. 1E), 41°48.433'N, 25°45.064'E, 172 m a.s.l., 27.v.2023, riverine vegetation, 1 ♂, D. Gradinarov & Y. Petrova leg.; Strandzha Mts, NW of Prohod Vill., left bank of Sredetska Reka Riv. (Fig. 1F), 42°19.978'N, 27°03.398'E, 49 m a.s.l., riverine vegetation and roadside verges, 25.v.2023, 3 ♂♂, 1 ♀, D. Gradinarov & Y. Petrova leg.

All specimens were collected from the host plant *Anchusa procera* Besser (Fig. 2). On the same plant species, the process of copulation, as well as feeding on flower petals, were observed. The records of *C. officinale* and *A. barrelieri* as host plants of *P. serriventris*, by Rejzek et al. (2001) and Georgiev et al. (2005) respectively, are rather doubtful. In the localities near Kotlari and Slavyanovo, *A. procera* Besser coexist with other species of genus *Cynoglossum* – *C. hungaricum* Simonk., but we did not observe individuals of *P. serriventris* on this plant species. *Anchusa barrelieri*, which was reported as a host plant of *P. ser-*

riventris from Bistrets (Georgiev et al., 2005) was not found in the vicinity of this village during the present study, as well as nowhere else in the areas visited.

According Holzschuh (1984) and Bense (1995), the longitudinal rows of ocher (yellowish-brown) hairs on the elytra in both sexes, median tooth-like processes on the abdominal sternites 1 and 2 in males as well as the regularly sharpened apex of penis are among the main diagnostic features of *P. serriventris* (Figs 3, 4, 5A, 6A). In some examined specimens of *P. serriventris*, however, the longitudinal rows on the elytra are less pronounced to almost invisible. On the other hand, such rows are well developed in some of the *P. tigrina* specimens in the populations of the Western Stara Planina Mts (Gradinarov, 2016; Gradinarov & Petrova, 2019), as well as in populations of *P. tigrina* in Romania (Crişan et al., 2017). Among the males of *P. tigrina* from the locality of the species in the region of Beledie han (Gradinarov & Petrova, 2021) specimens with well-developed processes on the abdominal sternites 1 and 2 can also be found (Fig. 5B). Thus, the shape of the apex of penis is the most useful character for distinguishing *P. serriventris* (Fig. 6A, B) and *P. tigrina* (Fig. 6C, D). The pygidium in males of *P. serriventris* is significantly wider distally than in *P. tigrina*, which can also be used in distinguishing the two species (Holzschuh, 1984).

Roots of *A. procera* with live Cerambycidae larvae were collected on 9 September 2021, and April




Fig. 3. Males of *Pilemia serriventris* from Bulgaria. A) Harmanli, 28.v.2023; B) Lyubimets, 24.v.2021; C) Kotlari Vill., 27.v.2018; D) Slavyanovo Vill., 27.v.2023; E) Prohod Vill., 25.v.2023. Scale bars: 1 mm.




Fig. 4. Females of *Pilemia serriventris* from Bulgaria. A) Harmanli, 28.v.2023; B) Lyubimets, 24.v.2021; C) Kotlari Vill., 05.vi.2020; D) Prohod Vill., 25.v.2023. Scale bars: 1 mm.




Fig. 5. Abdominal sternites in males of *Pilemia* spp. from Bulgaria. A – *P. serriventris*, Kotlari Vill., 27.v.2018; B – *P. tigrina*, Beledie Han Settlement, 20.v.2020. Scale bar: 1 mm.




Fig. 6. Male genitalia of *Pilemia* spp. from Bulgaria. A, B – apex of penis and parameres of *P. serriventris*, Kotlari Vill., 27.v.2018; C, D – apex of penis and parameres of *P. tigrina*, Beledie Han Settlement, 20.v.2020. Scale bar: 1 mm.

16, 2023, from the locality near Lyubimets and that near Kotlari Vill., respectively. In both cases, single specimens of *Phytoecia* (*Opsilia*) *coerulescens* (Scopoli, 1763) (Cerambycidae: Lamiinae) have emerged from the roots after laboratory rearing. The last species is polyphagous mostly in Boraginaceae (Sama, 2002) and has been reported together with *P.*

tigrina on *A. barrelieri* by earlier authors (Kovács, 2005; Crişan et al., 2017).

In our study, adults of *P. serriventris* were found on the host plants in the period of the second half of May to the beginning of June. Our data is consistent with the known period of activity of the adults of this species (Holzschuh, 1984; Georgiev et al., 2005,

Hoskovec et al., 2023). Adults of *P. serriventris* appear to become active later in spring than those of *P. tigrina*, which in Bulgaria can be found on food plants already in April (Gradinarov 2016; Gradinarov & Petrova, 2019; 2021). The late appearance of the adults of *P. serriventris* is probably related with the late flowering of the host plant *A. procera*, in comparison to *A. barrelieri*.

During the survey we found the species at an altitude of about 50 to 170 m. a.s.l., on alluvial banks of rivers from the Aegean drainage basin (Maritsa Riv., Arda Riv., Biserska Reka Riv.) and from the Black Sea drainage basin (Sredetska Reka Riv.). Near Lyubimets and near the Prohod Vill., it was also found among roadside vegetation at a greater distance (up to about 120 m) from the river. Both habitat types belong to the linear habitats (with linear strips of vegetation) sensu Bennett (2003). The second species of the subgenus *Pilemia* in Bulgaria – *P. tigrina*, has been recorded from linear habitats as well (Gradinarov & Petrova, 2021). In such habitats natural vegetation may have remained intact and they can serve as corridors for the dispersal of plant and animal species (Bennett, 2003). The presence of *P. serriventris* along the rivers Arda and Maritsa is also possible in the neighboring areas of Greece and the European part of Türkiye.

Acknowledgements

The authors would like to thank Asen Asenov (Department of Botany, Sofia University “St. Kliment Ohridski”, Sofia, Bulgaria) for the identification of the plant species of Boraginaceae family. This work has been carried out in the framework of the National Science Program “Environmental Protection and Reduction of Risks of Adverse Events and Natural Disasters”, approved by the Resolution of the Council of Ministers No. 577/17.08.2018 and supported by the Ministry of Education and Science (MES) of Bulgaria (Agreement No. D01-271/09.12.2022).

References

- Bennett A.F. 2003 Linkages in the Landscape: The Role of Corridors and Connectivity in Wildlife Conservation. IUCN, Gland, Switzerland and Cambridge, xiv + 254 pp.
<https://doi.org/10.2305/iucn.ch.2004.fr.1.en>
- Bense U. 1995 Longhorn beetles: illustrated key to the Cerambycidae and Vesperidae of Europe. Margraf Verlag, Weikersheim, 512 pp.
- Crișan A., Manci C.O., Ruicănescu A., Rákossy L. 2017 Information about the biology, ecology and distribution of *Pilemia tigrina* (Mulsant, 1851), in Romania (Coleoptera: Cerambycidae). Entomologica Romanica 21: 9–14.
<https://doi.org/10.24193/entomolrom.21.2>
- Danilevsky M.L. 2020 Taxa from West Europe, and North Africa to countries of former Soviet Union, and Mongolia. In: Danilevsky M.L. (ed.) Catalogue of Palaearctic Coleoptera, vol. 6 (1), Chrysomeloidea I (Vesperidae, Disteniidae, Cerambycidae). Revised and updated edition. Brill, Leiden/Boston, 118–480.
<https://doi.org/10.1163/9789004440333>
- Georgiev G. 2020 New records of longhorn beetles (Coleoptera: Cerambycidae) in entomological collections in Bulgaria. Forest Science 56 (1): 87–99.
- Georgiev G., Simov N., Stojanova A., Doychev D. 2005 New and interesting records of longhorn beetles (Coleoptera: Cerambycidae) in some Bulgarian Mountains. Acta Zoologica Bulgarica 57 (2): 131–138.
- Gradinarov D. 2016 New data on the distribution of *Pilemia tigrina* (Mulsant, 1851) (Cerambycidae: Lamiinae) in Bulgaria. ZooNotes 96: 1–3.
- Gradinarov D., Petrova Y. 2019 Longhorn beetles (Coleoptera: Cerambycidae) from Vrachanska Planina Mountains and Vrachanski Balkan Nature Park. In: Bechev D., Georgiev D. (eds) Faunistic diversity of Vrachanski Balkan Nature Park. Part 2. ZooNotes, Supplement 7, Plovdiv University Press, Plovdiv, 59–80.
- Gradinarov D., Petrova Y. 2021 New records of *Pilemia tigrina* (Mulsant, 1851) (Cerambycidae: Lamiinae) from roadside habitats in Bulgaria. ZooNotes 183: 1–4.
- Hoskovec M., Navrátil D., Jelínek P., Rejzek M. 2023 *Pilemia* (*Pilemia*) *serriventris* (Holzschuh, 1984). In: Cerambycidae. Longhorn beetles (Cerambycidae, Coleoptera) of the West Palaearctic region, neighboring territories and countries of the former Soviet Union.
http://www.cerambyx.uochb.cz/phytoecia_serriventris.php
(accessed 23 November 2023)
- Holzschuh C. 1984 Beschreibung neuer Arten aus der unmittelbaren Verwandtschaft von *Phytoecia*

- (*Pilemia*) *tigrina* (Cerambycidae, Col.). Koleopterologische Rundschau 57: 167–175.
- Migliaccio E., Georgiev G., Gashtarov V. 2007 An annotated list of Bulgarian Cerambycids with special view on the rarest species and endemics (Coleoptera: Cerambycidae). Lambillionea 107 (1), Supplément 1: 1–78.
- Rejzek M., Sama G., Alziar G. 2001 Host Plants of Several Herb-Feeding Cerambycidae Mainly from East Mediterranean Region (Coleoptera: Cerambycidae). Biocosme Mésogéen 17 (4): 263–294.
- Sama G. 2002 Atlas of the Cerambycidae of Europe and the Mediterranean Area. Vol. 1. Northern, Western, Central and Eastern Europe, British Isles and Continental Europe from France (excluding Corsica) to Scandinavia and Urals. Nakladatelstvi Kabourek, Zlin, 173 pp.
- Kovács T. 2005 Adatok a *Pilemia tigrina* (Mulsant, 1851) magyarországi elterjedéséhez és életmódjához (Coleoptera: Cerambycidae). Folia Historico Naturalia Musei Matraensis 29: 145–150.
- Tóth I.Z., Csathó A.I., Buşmachi G., Merkl O. 2016 *Pilemia tigrina*: new and corrected records from the Republic of Moldova, Hungary and Romania (Coleoptera: Cerambycidae). Folia Entomologica Hungarica 77: 33–40.
<https://doi.org/10.17112/foiaenthung.2016.77.33> 
-