

Piper betle L.: Cultural Heritage, Phytochemistry, Pharmacology and Agronomic Significance

Khushboo Rani¹ and Pushpanjali Khare^{2*}

Keywords: *Piper betle* L., Cultural Heritage, Phytochemistry, Pharmacology, Agronomic Significance.

Abstract:

Piper betle L., commonly known as betel leaf or “Paan,” is a perennial evergreen climber belonging to the family Piperaceae that possesses immense cultural, medicinal, and economic importance across South and Southeast Asia. Traditionally revered as “Tambula” in Ayurveda, the plant occupies a central place in religious rituals, social customs, and ethnomedical practices. For centuries, it has been utilized in Ayurveda, Unani, and Siddha systems of medicine for its digestive, carminative, antimicrobial, anti-inflammatory and stimulant properties. Phytochemical investigations have identified a rich spectrum of bioactive compounds, including phenolics, flavonoids, tannins, alkaloids, and essential oils such as eugenol, chavicol, and hydroxychavicol. These constituents contribute to its wide range of pharmacological activities, including antioxidant, antimicrobial, anticancer, anti-diabetic, gastroprotective, hepatoprotective and immunomodulatory effects. Modern scientific validation of traditional claims has enhanced their relevance in phytopharmaceutical research and functional food development. Agronomically, *Piper betle* is an economically valuable tropical climber cultivated under specific shade and humidity conditions, contributing significantly to rural livelihoods. However, sustainable cultivation, disease management and quality improvement remain important research priorities. This study integrates historical, ethnobotanical, phytochemical, pharmacological and agronomic perspectives, highlighting *Piper betle* as a culturally rooted yet scientifically promising medicinal plant resource.

Introduction: Historical, Ethnobotanical and Global Context

Piper betle L., commonly known as the betel leaf or “Paan,” is much more than a simple botanical entity; it is a living bridge between ancient tradition and modern science. A perennial evergreen climber belonging to the Piperaceae family, this plant thrives in the humid, tropical corridors of South and Southeast Asia. For millennia, it has occupied a central role in the lives of millions, sitting at the vibrant intersection of agriculture, medicine, and spiritual ritual. Unlike many medicinal plants that have faded into the background of history, the betel leaf has maintained an unbroken chain of relevance, evolving from a sacred symbol in Vedic texts to a subject of rigorous pharmacological scrutiny in the 21st century.

Khushboo Rani

Patna University, Ashok Rajpath, Patna University Campus, Patna, Bihar, 800005, India

Orcid iD:  <https://orcid.org/0009-0007-9141-2420>

Pushpanjali Khare

Magadh Mahila College, Patna, is North Gandhi Maidan, Patna - 800001, Bihar, India

E-mail:  khare.pushpanjali2@gmail.com; Orcid iD:  <https://orcid.org/0000-0002-1335-4559>

***Corresponding Author:** khare.pushpanjali2@gmail.com

© International Academic Publishing House, 2025

Dr. Saikat Kumar Basu, Dr. Bhanumati Sarkar, Dr. Suparna Sanyal Mukherjee and Dr. Alminda Magbalot-Fernandez (eds.), The Basic Handbook of Indian Ethnobotany and Traditional Medicine [Volume: 4]. ISBN: 978-81-989555-5-5; pp. 85-99; Published online: 24th December, 2025

Historically, the practice of chewing betel leaf is woven into the very DNA of Asian social grace. Archaeological evidence and historical documentation suggest that these customs date back over 2,000 years. In the Indian subcontinent, the leaf is synonymous with hospitality; to offer a prepared "biras" of paan is to offer respect, friendship, and peace. It is an indispensable guest at every major life event, from the sacred fire of weddings to the festive altars of Diwali and Durga Puja, where it symbolizes prosperity and divine blessing.

Beyond the altar and the social gathering, *Piper betle* is a cornerstone of traditional healing. In the hallowed scrolls of Ayurveda, it is celebrated as "Tambula," prized for its ability to clear the breath, stimulate the appetite, and aid digestion. Similarly, the Unani and Siddha systems of medicine have long utilized their warming properties to treat respiratory ailments and skin conditions. Today, as we move into an era of evidence-based phytotherapy, the "Green Gold" of Asia is being rediscovered. Modern science is now confirming what traditional healers have known for generations: that this humble leaf is a chemical powerhouse, packed with bioactive compounds capable of fighting bacteria, reducing inflammation, and perhaps even protecting against cancer.

Taxonomy and Systematics

Classical Taxonomic Classification

Piper betle L. was first formally described by the Swedish botanist Carl Linnaeus in 1753 in his landmark botanical work *Species Plantarum*, which laid the foundation for modern plant nomenclature. The species belongs to the Kingdom Plantae and is classified within the clade Angiosperms, representing flowering plants characterized by enclosed seeds. Within angiosperms, it is grouped under the Magnoliids, an ancient lineage of dicotyledonous plants known for their aromatic compounds and primitive floral structures.

Taxonomically, *Piper betle* falls under the order Piperales and the family Piperaceae. The Piperaceae family is a pantropical group comprising approximately 3,000 species, distributed predominantly in humid tropical and subtropical regions. Members of this family are generally aromatic herbs, shrubs, or climbers, often rich in essential oils and alkaloids. The genus *Piper* is the largest within the family and represents one of the most economically significant genera among tropical plants.

The genus includes several species of global commercial importance, such as *Piper nigrum* (black pepper) and *Piper longum* (long pepper), both widely used as spices and medicinal plants. Within this genus, *Piper betle* is distinguished by its dioecious reproductive nature, cordate leaves, and characteristic pungent aroma resulting from essential oil-rich secretory cells. Unlike wild relatives, it is primarily cultivated and rarely found in a truly wild state, indicating long-term human domestication. The presence of numerous regional cultivars further reflects its agricultural and cultural importance across South and Southeast Asia.

Evolutionary and Phylogenetic Position

Advances in molecular systematics have significantly refined the understanding of phylogenetic relationships within Piperaceae. Early classification relied heavily on morphological traits such as leaf venation, inflorescence structure, and growth habit. However, contemporary studies utilizing chloroplast DNA markers—including *rbcL*, *matK*, and *trnL-F*—along with nuclear ribosomal ITS sequences, have provided more robust evolutionary insights. Research by Jaramillo and Manos (2001) clarified that *Piper betle* belongs to the Asian clade of *Piper*, which is phylogenetically distinct from Neotropical lineages.

Molecular phylogenetic analyses indicate that diversification within Piperaceae began during the early Cretaceous period, corresponding with the radiation of angiosperms. Climatic fluctuations and continental drift likely played major roles in shaping current distribution patterns. The Asian lineage appears to have undergone adaptive radiation influenced by humid tropical ecosystems, which favored the evolution of climbing habits and aromatic secondary metabolites as ecological defense strategies.

Within this evolutionary framework, *Piper betle* occupies a unique position. Although closely related to several Southeast Asian taxa, it demonstrates remarkable cultivar variability. This diversity is primarily attributed to domestication-driven divergence rather than natural speciation. Since the plant is propagated vegetatively through stem cuttings, genetic variability arises from somatic mutations and selective breeding practices maintained over centuries. Such domestication has resulted in region-specific cultivars differing in leaf size, pungency, phytochemical composition, and agronomic performance.

Overall, the taxonomy and systematics of *Piper betle* illustrate the dynamic interplay between evolutionary history and human cultural selection. Its classification within Piperaceae reflects deep phylogenetic roots, while its present-day diversity highlights the profound influence of domestication and ecological adaptation.

Phytochemistry of *Piper betle*

The remarkable pharmacological potential of *Piper betle* is fundamentally rooted in its rich and diverse phytochemical composition. Over the past few decades, advanced analytical techniques such as Gas Chromatography–Mass Spectrometry (GC–MS), Liquid Chromatography–Tandem Mass Spectrometry (LC–MS/MS), High-Performance Liquid Chromatography (HPLC), Nuclear Magnetic Resonance (NMR), and Fourier Transform Infrared Spectroscopy (FTIR) have enabled detailed chemical profiling of the leaf. These investigations have revealed an extensive array of bioactive compounds, including phenolic compounds, phenylpropanoids, flavonoids, alkaloids, terpenoids, sterols and essential oils.

Among these groups, phenolic compounds and phenylpropanoids are particularly dominant and are considered major contributors to the leaf's antioxidant, antimicrobial, and anti-inflammatory properties. Essential oils constitute a significant fraction of the plant's total phytochemical pool and are responsible for its characteristic pungent aroma and therapeutic activity. However, it is important to recognize that the phytochemical profile of *Piper betle* is

not uniform. It varies considerably depending on cultivar type, agro-climatic conditions, soil composition, irrigation practices, leaf maturity stage, and even post-harvest handling and storage conditions.

Comparative phytochemical studies conducted across different regions of India—especially eastern states such as Bihar—have demonstrated notable differences in essential oil composition. Variations are particularly pronounced in the concentrations of hydroxychavicol and eugenol derivatives, two major bioactive constituents. Such regional variation suggests that environmental stressors, soil micronutrients, and climatic factors significantly influence secondary metabolite biosynthesis pathways. This environmental responsiveness enhances the importance of geographic origin in determining both medicinal potency and commercial quality.

Primary Metabolites

While secondary metabolites often receive greater scientific attention due to their pharmacological activities, primary metabolites also play an essential role in the physiological and nutritional value of betel leaves. These compounds are directly involved in growth, metabolism, and cellular function, and they contribute to the plant's overall structural integrity.

Carbohydrates

Betel leaves contain moderate amounts of carbohydrates, including reducing sugars, sucrose, and structural polysaccharides. These carbohydrates are essential for maintaining osmotic balance within plant cells and play a crucial role in determining leaf texture, flexibility, and moisture retention. Since the plant is harvested vegetatively and repeatedly, starch accumulation remains relatively low. Continuous harvesting prevents long-term carbohydrate storage, thereby maintaining a metabolically active state in the foliage. Additionally, soluble sugars may indirectly influence flavor characteristics and microbial stability of fresh leaves.

Proteins and Amino Acids

Proteomic analyses have identified both structural and enzymatic proteins in *Piper betle*. The total protein content typically ranges from 2–4% on a dry-weight basis. Essential amino acids such as lysine, methionine, and tryptophan have been reported, indicating modest nutritional relevance. More importantly, enzymes involved in the phenylpropanoid biosynthetic pathway are abundant in leaf tissues. These enzymes regulate the production of phenolic compounds and essential oils, linking primary metabolic processes directly to secondary metabolite formation. Thus, protein expression plays a critical regulatory role in determining phytochemical richness.

Lipids and Fatty Acids

Although the total lipid content of betel leaves is relatively low, fatty acid profiling reveals the presence of biologically important fatty acids such as linoleic acid, palmitic acid, and oleic

acid. These lipids contribute to cellular membrane stability, maintain structural integrity, and may influence the retention and stabilization of volatile aromatic compounds within leaf tissues. Membrane lipids also support active transport processes that facilitate metabolite synthesis and accumulation.

Secondary Metabolites

Secondary metabolites constitute the biochemical foundation of *Piper betle*'s therapeutic importance. Unlike primary metabolites, which are essential for growth and cellular maintenance, secondary metabolites function primarily in ecological defense, environmental adaptation, and plant–microbe interactions. In *Piper betle*, these compounds are abundant and chemically diverse, contributing significantly to its antimicrobial, antioxidant, anti-inflammatory, and anticancer activities. The major classes include phenylpropanoids, flavonoids, alkaloids, terpenoids, essential oils, tannins and saponins.

Phenylpropanoids

Phenylpropanoids represent the dominant chemical class in betel leaf and are largely responsible for its pungent aroma and pharmacological potency. These compounds are synthesized through the phenylpropanoid biosynthetic pathway, beginning with the amino acid phenylalanine.

Hydroxychavicol (4-allylpyrocatechol) is widely recognized as the principal bioactive constituent. It exhibits strong antioxidant activity by neutralizing reactive oxygen species (ROS) and protecting cells from oxidative stress. Additionally, it possesses potent antimicrobial properties against Gram-positive and Gram-negative bacteria. One of its most studied properties is its anticancer potential, particularly its ability to induce apoptosis in tumor cell lines through ROS-mediated pathways. Experimental studies have demonstrated that hydroxychavicol can trigger mitochondrial dysfunction and activate caspase-dependent cell death mechanisms. Importantly, its concentration varies among cultivars, with eastern Indian varieties—especially those cultivated under humid agro-climatic conditions—often showing higher levels.

Eugenol, another key phenylpropanoid also found in *Syzygium aromaticum* (clove), contributes to the analgesic, antiseptic, and anti-inflammatory properties of betel leaf. It exerts antimicrobial effects by disrupting microbial cell membranes and increasing permeability, leading to leakage of intracellular contents. Moreover, eugenol inhibits inflammatory mediators such as cyclooxygenase-2 (COX-2), thereby reducing inflammatory responses.

Chavibetol and chavicol are structural isomers that significantly influence both the aroma and antimicrobial strength of the leaf. These compounds demonstrate antifungal activity, particularly against *Candida* and *Aspergillus* species, supporting traditional applications in oral hygiene and wound care.

Flavonoids

Flavonoids such as quercetin, catechin, and rutin have been identified through HPLC-DAD analysis. These polyphenolic compounds exhibit powerful free radical scavenging activity and contribute to overall antioxidant capacity. Their ability to chelate metal ions further enhances protection against oxidative damage. Additionally, flavonoids modulate inflammatory cytokines and signaling pathways, contributing to anti-inflammatory and immunomodulatory effects. Geographical studies reveal that total flavonoid content varies significantly, with samples from humid agro-climatic regions such as Bihar showing comparatively higher antioxidant index values in laboratory assays.

Alkaloids

Although present in relatively minor concentrations compared to phenolic compounds, alkaloids contribute to the physiological activity of *Piper betle*. Piperidine derivatives and other nitrogen-containing compounds have been reported. Unlike *Piper nigrum*, which contains substantial amounts of piperine, *Piper betle* lacks high concentrations of this alkaloid, indicating metabolic divergence within the genus. This distinction further emphasizes species-specific phytochemical specialization.

Terpenoids and Essential Oils

Essential oil content in betel leaf ranges between 0.7–2.6%, depending on cultivar, extraction method, and environmental factors. Major constituents include safrole, cineole, caryophyllene, terpinene, and estragole. GC–MS analyses reveal pronounced chemotypic variation across India. Southern cultivars frequently exhibit higher safrole content, whereas eastern cultivars tend to show phenolic dominance. These volatile compounds contribute not only to aroma but also to antimicrobial and anti-inflammatory activities.

Tannins and Saponins

Tannins contribute to the astringent taste and exhibit antimicrobial properties by precipitating microbial proteins. Saponins, on the other hand, possess membrane-permeabilizing properties, which may enhance the bioavailability and synergistic effects of other phytochemicals.

Traditional Medicinal Systems and Classical Documentation

Piper betle holds a distinguished and historically significant position in multiple traditional medicinal systems across South and Southeast Asia. Its therapeutic importance long predates modern pharmacology and is deeply embedded in classical textual traditions, ethnomedical practices, and ritualistic applications. The consistent documentation of betel leaf across Ayurveda, Unani, and Siddha systems reflects cross-cultural validation of its medicinal value.

In **Ayurveda**, betel leaf is referred to as *Tambula*. Classical Ayurvedic texts describe it as *katu* (pungent) and *tikta* (bitter) in taste, with *ushna virya* (hot potency). According to

Ayurvedic pharmacodynamics, its heating nature stimulates *agni* (digestive fire), thereby enhancing digestion, metabolism, and nutrient assimilation. It is traditionally recommended for *Deepana* (appetite stimulation), *Pachana* (digestive enhancement), *Krimighna* (anthelmintic action), *Vranashodhana* (wound cleansing), and *Shothahara* (anti-inflammatory effects). Tambula is commonly included in post-meal digestive preparations due to its carminative properties, which help relieve bloating and flatulence. Its warming action is also believed to improve circulation and respiratory function.

In **Unani medicine**, betel leaf is classified under the temperament *Mizaj-e-Haar Yabis* (warm and dry). It is prescribed for conditions associated with excess phlegm and cold temperament, including bronchitis, flatulence, halitosis, and sexual weakness. Unani scholars describe its ability to strengthen nerves, stimulate circulation, and enhance vitality.

In **Siddha medicine**, betel leaf is used both externally and internally. Externally, it is applied over inflammatory swellings, joint pain, and wounds due to its anti-inflammatory and antiseptic properties. Internally, it functions as an expectorant and digestive stimulant. The convergence of these traditional systems demonstrates pharmacological consistency, reinforcing the therapeutic credibility of *Piper betle* across cultural boundaries.

Ethnobotanical Relevance in Eastern India with Special Reference to Bihar

Eastern India represents one of the most culturally significant regions for betel cultivation and medicinal application. In Bihar, *Piper betle* is deeply integrated into agrarian life, household remedies, and ritual traditions, making it both an economic resource and a socio-cultural symbol.

Household Medicinal Uses in Bihar

Ethnobotanical surveys conducted in districts such as Vaishali, Muzaffarpur, Darbhanga, and Samastipur document extensive traditional uses. A warmed betel leaf is commonly applied to the chest of children and adults to relieve cough and congestion. Leaf paste mixed with mustard oil is applied to joints for pain relief and to reduce swelling. The fresh leaf juice is administered in small quantities for indigestion and gastric discomfort. Crushed leaves are placed over wounds to prevent infection due to their antimicrobial properties. In postpartum care, betel leaf preparations are traditionally used to support lactation and recovery.

Traditional healers, known locally as *Vaidya* or *Ojha*, incorporate betel leaf in polyherbal formulations for fever, respiratory ailments, and digestive disorders. These practices reflect cumulative empirical knowledge transmitted through generations.

Ritual and Therapeutic Symbolism

Beyond medicinal value, betel leaf symbolizes purity, prosperity, and auspiciousness in Bihar. It is indispensable in marriage ceremonies, temple offerings, and festivals such as Chhath Puja. The ritualistic use reinforces its identity not only as a medicinal plant but as a socio-therapeutic symbol linking health, spirituality, and community bonding.

Experimental Pharmacology: Mechanistic Insights

Modern scientific research has validated many traditional claims regarding *Piper betle* through controlled in vitro and in vivo studies.

Antioxidant Mechanisms

Oxidative stress plays a major role in chronic diseases such as diabetes, cancer, and neurodegeneration. Extracts of *Piper betle* demonstrate strong free radical scavenging activity in DPPH, ABTS, and FRAP assays. Bioactive compounds such as hydroxychavicol and eugenol act as electron donors, neutralizing reactive oxygen species (ROS). Mechanistically, they enhance endogenous antioxidant enzymes including superoxide dismutase (SOD), catalase, and glutathione peroxidase. Animal studies report reduced lipid peroxidation in hepatic tissues following administration of aqueous extracts.

Antimicrobial Activity

Betel leaf exhibits broad-spectrum antimicrobial activity. The mechanism includes disruption of microbial cell membrane integrity, leakage of intracellular components, inhibition of biofilm formation, and suppression of quorum sensing pathways. Hydroxychavicol has shown effectiveness against Gram-positive and Gram-negative bacteria, including resistant strains. Antifungal studies demonstrate inhibition of *Candida albicans* growth via mitochondrial dysfunction induction.

Anti-inflammatory and Analgesic Effects

Inflammation underlies many chronic pathologies. Betel leaf extracts reduce inflammatory mediators through COX-2 inhibition, suppression of the NF- κ B pathway, and downregulation of TNF- α and IL-6. Rodent models show significant reduction in carrageenan-induced paw edema. Eugenol contributes to analgesic effects by modulating prostaglandin synthesis.

Antidiabetic Activity

Experimental diabetic models indicate that betel leaf extracts can reduce blood glucose levels and improve insulin sensitivity. The mechanism may involve enhanced glucose uptake and antioxidant-mediated pancreatic protection. These findings suggest potential applications in metabolic disorder management.

Agronomy and Cultivation Systems

Piper betle is a shade-loving perennial climber requiring carefully regulated environmental conditions. Commercial cultivation is labor-intensive and regionally specialized.

Agro-Climatic Requirements

The vine thrives at temperatures between 15–40°C, relative humidity of 60–80%, and annual rainfall of 1500–3000 mm. It grows best in well-drained loamy or alluvial soils. Excess sunlight

causes leaf scorching, while low humidity reduces essential oil concentration. The fertile Gangetic alluvial soils of Bihar and West Bengal provide ideal conditions.

Propagation Techniques

The plant is propagated vegetatively through stem cuttings with 3–5 nodes. Farmers select disease-free mother vines, root cuttings in nursery beds, and transplant them under shade structures. While this ensures genetic uniformity, it reduces overall genetic diversity.

Barouj (Bareja) System

In eastern India, traditional enclosed structures known as Barouj (Bareja in Bihar) regulate microclimatic conditions. Bamboo frameworks covered with straw and jute mats maintain humidity, reduce temperature fluctuations, and protect vines from wind and pests. This indigenous agro-architecture reflects adaptive ecological intelligence.

Soil, Irrigation and Nutrient Management

Successful cultivation of *Piper betle* depends heavily on proper soil management and balanced nutrient supply. The crop performs best in well-drained, fertile loamy or alluvial soils with a pH range of 5.5–7.0. Good drainage is critical because the vine is highly sensitive to waterlogging, which can lead to root rot and fungal infections. In eastern Indian regions such as Bihar, fertile Gangetic alluvial soils provide ideal physical and chemical properties for vine growth.

Organic manures such as farmyard manure (FYM), vermicompost, and green manure are widely incorporated into the soil to enhance organic matter content and microbial activity. Nitrogen promotes vegetative growth and leaf expansion, while potassium plays a significant role in improving essential oil biosynthesis and leaf quality. Phosphorus supports root development and early establishment.

Irrigation management involves frequent light watering rather than heavy flooding. Drip irrigation is increasingly recommended to improve water-use efficiency, reduce fungal disease incidence, and maintain proper soil aeration. During monsoon seasons in Bihar, proper drainage channels are essential to prevent stagnation and protect root systems from excess moisture stress.

Plant Pathology and Pest Management

Betel vine is susceptible to several fungal, bacterial, and viral diseases that significantly affect yield and quality. Among fungal diseases, leaf rot caused by *Phytophthora* species and anthracnose caused by *Colletotrichum* species are the most economically damaging. Powdery mildew also appears under humid conditions. These pathogens reduce leaf marketability and may cause large-scale crop losses.

Integrated Disease Management (IDM) strategies are widely recommended. These include the application of copper-based fungicides, use of biological control agents such as

Trichoderma species, soil solarization, crop rotation, and improved drainage systems. The use of certified disease-free planting material is critical in preventing the spread of infections.

Insect pests such as scale insects, mites, and mealybugs commonly attack betel vines. These pests damage leaf tissues and reduce commercial value. To minimize chemical residues, farmers increasingly adopt neem-based biopesticides and botanical extracts as environmentally safer alternatives. Sustainable pest management is essential for maintaining both productivity and phytochemical integrity.

Post-Harvest Physiology and Storage

Betel leaves are highly perishable due to their high moisture content and rapid respiration rate. Post-harvest handling practices significantly influence shelf life, quality, and market value. Leaves are harvested manually at physiological maturity, and frequent plucking stimulates continued vegetative growth.

Traditional storage methods involve wrapping leaves in banana leaves, sprinkling them lightly with water, and maintaining cool, humid conditions to reduce moisture loss. However, improper storage can lead to fungal growth, discoloration, and loss of essential oil content.

Modern preservation techniques include modified atmosphere packaging, vacuum packaging, and cold storage at temperatures between 10–15°C. These methods slow respiration rates and extend shelf life while preserving aroma and texture. Improved post-harvest management not only reduces economic losses but also ensures maintenance of medicinal and commercial quality.

Economic Importance and Rural Livelihoods

Piper betle cultivation represents a high-value cash crop system in India. The country is among the largest producers and exporters of betel leaves, with major production centers in West Bengal, Odisha, Assam, Andhra Pradesh, Tamil Nadu, and Bihar.

In Bihar, betel cultivation is concentrated in districts such as Vaishali, Muzaffarpur, Darbhanga, Samastipur and Nalanda. The crop provides year-round income due to continuous harvesting cycles. It also generates employment, particularly for women engaged in leaf sorting, grading, bundling, and packaging.

Market linkages connect rural producers to metropolitan cities, ensuring steady demand. Despite its economic significance, the sector faces challenges such as disease outbreaks, climate variability and fluctuating market prices. Strengthening cooperative marketing systems, improving storage facilities, and promoting value-added products could enhance rural livelihoods and ensure long-term sustainability of betel vine cultivation.

Sustainability and Climate Change Challenges

The cultivation of *Piper betle* faces increasing sustainability concerns under changing climatic conditions. As a shade-loving, humidity-dependent crop, betel vine is highly sensitive to temperature fluctuations, irregular rainfall patterns, and extreme weather events. Rising

temperatures may reduce essential oil concentration and increase evapotranspiration stress, while excessive rainfall—particularly in monsoon-dominated regions like Bihar—can lead to waterlogging and higher incidence of fungal diseases such as *Phytophthora* leaf rot.

Climate variability also influences pest dynamics, potentially increasing outbreaks of mites, scale insects, and pathogenic fungi. Traditional enclosed systems such as the Barouj provide some microclimatic buffering; however, prolonged climatic stress may exceed their protective capacity.

Sustainable cultivation practices are therefore critical. These include integrated pest management, organic nutrient supplementation, improved drainage systems, rainwater harvesting, and adoption of drip irrigation. Developing climate-resilient cultivars through selection and biotechnology could further safeguard productivity. Promoting eco-friendly inputs and reducing chemical dependency will ensure environmental sustainability while preserving phytochemical integrity. Addressing climate resilience is essential for protecting both farmer livelihoods and the medicinal value of betel vine.

Future Research Directions

Despite extensive ethnomedicinal documentation and growing pharmacological evidence, several research gaps remain in *Piper betle* studies. Future investigations should prioritize large-scale clinical trials to validate safety, efficacy, and dosage standardization of purified leaf extracts. Standardization of phytochemical markers such as hydroxychavicol and eugenol is essential for developing globally accepted herbal formulations.

Advanced metabolomic and genomic studies could help understand cultivar variation and metabolic regulation pathways. Exploring transcriptomic responses under different agro-climatic conditions may clarify how environmental stress influences phytochemical biosynthesis. Additionally, research into nano-formulations and novel drug delivery systems may enhance bioavailability and therapeutic effectiveness.

From an agronomic perspective, breeding programs focusing on disease resistance and climate tolerance are necessary. Post-harvest technologies aimed at extending shelf life without compromising quality also require further development. Interdisciplinary collaboration between botanists, pharmacologists, agronomists, and policymakers will be crucial for transforming *Piper betle* from a traditional medicinal plant into a scientifically standardized and globally recognized phytotherapeutic resource.

Overall Conclusion

Piper betle L. stands as a remarkable example of the dynamic interplay between traditional knowledge systems and modern scientific validation. Across centuries, it has transitioned from being a sacred cultural symbol and household remedy to an object of rigorous phytochemical and pharmacological investigation. This continuity of relevance reflects not only the plant's deep-rooted presence in South and Southeast Asian societies but also its substantial therapeutic potential grounded in biochemical complexity.

From a taxonomic and evolutionary perspective, *Piper betle* belongs to a lineage with ancient angiosperm origins, yet its present diversity is strongly shaped by domestication and human selection. Vegetative propagation, regional cultivation systems, and environmental influences have produced significant chemotypic variation across cultivars. Such variability underscores the importance of integrating botanical systematics with molecular research to better understand genetic diversity, phytochemical expression, and adaptive potential under changing climates.

Phytochemically, the plant exhibits an intricate profile of primary and secondary metabolites. While primary metabolites sustain cellular metabolism and structural integrity, secondary metabolites—particularly phenylpropanoids such as hydroxychavicol and eugenol—form the pharmacological backbone of its medicinal activity. Flavonoids, terpenoids, alkaloids, tannins, and essential oils collectively contribute to antioxidant, antimicrobial, anti-inflammatory, and metabolic regulatory properties. Advanced analytical techniques have substantiated these bioactive compounds, providing molecular explanations for centuries-old therapeutic practices.

Traditional medicinal systems including Ayurveda, Unani, and Siddha consistently recognize betel leaf as a digestive stimulant, anti-inflammatory agent, antimicrobial remedy, and respiratory aid. The pharmacodynamic concepts of *ushna virya* and *Mizaj-e-Haar Yabis* align closely with experimentally observed thermogenic and circulatory effects. Ethnobotanical evidence from eastern India, particularly Bihar, demonstrates how the plant functions not merely as medicine but as a socio-cultural asset embedded in rituals, ceremonies, and household healthcare. The dual identity of *Piper betle*—as both sacred symbol and therapeutic agent—illustrates the integration of spiritual and physiological health in traditional communities.

Modern pharmacological studies further validate these traditional claims. Antioxidant assays confirm strong free radical scavenging potential, supported by upregulation of endogenous enzymes such as superoxide dismutase and catalase. Antimicrobial mechanisms involve membrane disruption, inhibition of biofilm formation, and suppression of quorum sensing pathways. Anti-inflammatory activity is mediated by COX-2 inhibition and modulation of the NF- κ B pathway, while emerging research suggests promising antidiabetic and anticancer properties through apoptosis induction and metabolic regulation. These findings collectively position *Piper betle* as a potential candidate for evidence-based phytotherapy.

Agronomically, the cultivation of betel vine represents a sophisticated and labor-intensive system. Traditional structures such as the Barouj or Bareja demonstrate indigenous ecological intelligence, effectively regulating humidity, temperature, and microclimatic stability. Soil management, nutrient dynamics, irrigation practices, and integrated pest control significantly influence both yield and phytochemical quality. However, the crop faces increasing challenges from fungal diseases, pest infestations, post-harvest perishability, and climate variability.

Sustainable cultivation strategies—including organic inputs, improved drainage, drip irrigation, and climate-resilient cultivar development—are essential to ensure long-term productivity.

Economically, betel vine serves as a high-value cash crop supporting thousands of rural households across India. In Bihar, districts such as Vaishali, Muzaffarpur, and Darbhanga rely heavily on this crop for year-round income and employment, particularly for women engaged in grading and packaging. Strengthening cooperative marketing networks, post-harvest infrastructure, and value-added product development can enhance rural livelihoods while maintaining quality standards.

Looking forward, the future of *Piper betle* lies in bridging ethnomedicine with translational research. Standardization of bioactive markers, clinical trials for dosage validation, genomic mapping for cultivar improvement, and novel drug delivery systems represent key research priorities. Equally important is the ethical preservation of traditional knowledge and sustainable agro-ecosystems.

In essence, *Piper betle* exemplifies how ancient botanical wisdom can inform contemporary scientific innovation. Its journey from sacred leaf to pharmacological candidate reflects a broader paradigm: that biodiversity, when respected and scientifically explored, holds transformative potential for global health, sustainable agriculture, and cultural continuity. The enduring legacy of this “Green Gold” affirms that the integration of tradition and science is not merely complementary—it is essential for holistic advancement.

References

- Al-Adhroey, A. H., Nor, Z. M., Al-Mekhlafi, H. M., & Mahmud, R. (2011). Antimalarial activity of methanolic leaf extract of *Piper betle* L. *Parasites & Vectors*, 4, 182. <https://doi.org/10.1186/1756-3305-4-182>
- Ali, M., & Khan, T. (2025). Nutraceutical potential of betel leaf. *Frontiers in Nutrition*, 12, 1298456.
- Antioxidant and antimicrobial potential of selected varieties of *Piper betle* L. (2017). PubMed indexed study.
- Arambewela, L. S. R., Arawwawala, L. D. A. M., & Ratnasooriya, W. D. (2005). Antidiabetic activities of aqueous and ethanolic extracts of *Piper betle* leaves in rats. *Journal of Ethnopharmacology*, 102(2), 239–245. <https://doi.org/10.1016/j.jep.2005.06.016>
- Arawwawala, L. D. A. M., Arambewela, L. S. R., & Ratnasooriya, W. D. (2004). Gastroprotective effect of *Piper betle* Linn. leaves grown in Sri Lanka. *Journal of Ayurveda and Integrative Medicine*, 25(2), 120–125.
- Bhadra, S., & Roy, S. (2018). Agronomic practices and cultivation technology of betel vine (*Piper betle* L.). *Indian Journal of Agricultural Sciences*, 88(5), 673–681.
- Chakraborty, D., & Shah, B. (2011). Antimicrobial, antioxidative and antihemolytic activity of *Piper betle* leaf extracts. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3(3), 192–199.
- Chakraborty, S., & Ghosh, P. (2024). Anti-inflammatory and analgesic properties of *Piper betle* extracts. *Natural Product Research*, 38(12), 2150–2158.

- Choudhary, D., & Kale, R. K. (2002). Antioxidant and non-toxic properties of *Piper betle* leaf extract. *Food and Chemical Toxicology*, 40(8), 1177–1183. [https://doi.org/10.1016/S02786915\(02\)00091-0](https://doi.org/10.1016/S02786915(02)00091-0)
- Dasgupta, N., & De, B. (2004). Antioxidant activity of *Piper betle* L. leaf extract in vitro. *Food Chemistry*, 88(2), 219–224. <https://doi.org/10.1016/j.foodchem.2004.01.036>
- Das, A., et al. (2025). Microwave-assisted extraction of bioactive compounds from *Piper betle* leaves. *Journal of Applied Research on Medicinal and Aromatic Plants*, 35, 100482.
- Dwivedi, V., & Tripathi, S. (2014). Review study on potential activity of *Piper betle*. *Journal of Pharmacognosy and Phytochemistry*, 3(4), 93–98.
- Guha, P. (2006). Betel leaf: The neglected green gold of India. *Journal of Human Ecology*, 19(2), 87–93. <https://doi.org/10.1080/09709274.2006.11905992>
- Gupta, P., Maral, A., Dubey, V., Yadav, S., & Rajbhar, S. (2023). Formulation and assessment of *Piper betle* emulgel for antimicrobial application. *Journal of Population Therapeutics & Clinical Pharmacology*.
- Haslan, H., Suhaimi, F. H., Thent, Z. C., & Das, S. (2015). The underlying mechanism of action for various medicinal properties of *Piper betle* (betel). *Clinical Therapeutics*, 166(5), 197–203. <https://doi.org/10.7417/T.2015.1880>
- Hossain, M. S., et al. (2024). Antiviral activity of *Piper betle* leaf phytochemicals. *International Journal of Pharmaceutical Sciences*, 16(3), 255–263.
- Jeng, J. H., Chang, M. C., & Hahn, L. J. (2001). Role of areca nut in betel quid-associated chemical carcinogenesis. *Journal of Oral Pathology & Medicine*, 30(1), 1–9.
- Khanra, R., Dewanjee, S., et al. (2015). Protective role of *Piper betle* against oxidative stress and fibrosis. *Environmental Toxicology and Pharmacology*, 40(1), 60–69.
- Kumar, N., & Singh, R. (2024). Antioxidant potential of *Piper betle* L. and its therapeutic implications. *Journal of Herbal Medicine*, 41, 100705.
- Kumar, N., Misra, P., Dube, A., et al. (2010). *Piper betle* Linn. a maligned pan-Asian plant with potential pharmacological activities. *Journal of Pharmacy Research*, 3(10), 2530–2536.
- Maiti, S., Ghosh, P., & Nag, A. (2020). Bio-actives of betel leaf (*Piper betle* L.): A comprehensive review on extraction, isolation, and biological activity. *Phytotherapy Research*, 34(10), 2609–2627. <https://doi.org/10.1002/ptr.6715>
- Nalina, T., & Rahim, Z. H. A. (2007). The crude aqueous extract of *Piper betle* L. and its antibacterial effect towards *Streptococcus mutans*. *American Journal of Biochemistry and Biotechnology*, 3(1), 10–15. <https://doi.org/10.3844/ajbbsp.2007.10.15>
- Patra, B., & Shukla, S. (2024). Phytochemical diversity and pharmacological properties of *Piper betle* L.: A comprehensive review. *Phytomedicine Plus*, 4(1), 100560.
- Pin, K. Y., & Chuah, A. L. (2010). Antioxidant and antibacterial activity of leaves of *Piper betle* L. *International Journal of Pharmacology*, 6(3), 278–281.

- Pradhan, D., Suri, K. A., Pradhan, D. K., & Biswasroy, P. (2013). Golden heart of the nature: *Piper betle* L. *Journal of Pharmacognosy and Phytochemistry*, 1(6), 147–167.
- Rahman, M. M., et al. (2025). Antimicrobial activity of *Piper betle* leaf extracts against multidrug-resistant bacteria. *BMC Complementary Medicine and Therapies*, 25, 112.
- Rai, M. P., Thilakchand, K. R., et al. (2011). Evaluation of hepatoprotective activity of *Piper betle* leaf extract. *Journal of Pharmacy Research*, 4(3), 756–758.
- Rathee, J. S., Patro, B. S., Mula, S., Gamre, S., & Chattopadhyay, S. (2006). Antioxidant activity of *Piper betle* leaf extract and its constituents. *Journal of Agricultural and Food Chemistry*, 54(24), 9046–9054. <https://doi.org/10.1021/jf061663j>
- Saha, S., et al. (2024). GC-MS analysis of essential oils of *Piper betle* L. *Industrial Crops and Products*, 207, 117628.
- Saini, S. *Piper betel* L.: A Review of Phytochemical and Pharmacological Profile. *International Education and Research Journal*.
- Shah, S. K., & Jhade, D. N. (2018). Antifertility potential of *Piper betle* in female rats. *Biochemical and Biophysical Reports*. <https://doi.org/10.1016/j.bbrep.2018.08.001>
- Sharma, P., & Kaur, G. (2025). Role of phenolic compounds of *Piper betle* in cancer prevention. *Food Bioscience*, 57, 103062.
- Sharma, S., Khan, I. A., Ali, I., et al. (2009). Evaluation of antimicrobial activity of *Piper betle* L. *Indian Journal of Pharmaceutical Sciences*, 71(5), 553–558.
- Shivakumar, S. N., & Hima Bindu, K. (2018). Phytochemistry and Pharmacological Potentiality of Betelvine (*Piper betle* L.). *Journal of Pharmacognosy and Phytochemistry*.
- Tarmale, V. A., Wankhede, S., Sen, A. K., & Uppalwar, S. V. (2024). Pharmacological Properties and Traditional Applications of Betel Leaf. *Journal of Pharma Insights & Research*.
- Wahyuni, T. S., Tumewu, L., & Permanasari, A. A. (2024). Phytochemistry profile of *Piper betle* extract and activity against Hepatitis C virus. *Indonesian Journal of Pharmacy*. <https://doi.org/10.22146/ijp.7071>

HOW TO CITE

Khushboo Rani and Pushpanjali Khare (2025). *Piper betle* L.: Cultural Heritage, Phytochemistry, Pharmacology and Agronomic Significance. © International Academic Publishing House (IAPH), Dr. Saikat Kumar Basu, Dr. Bhanumati Sarkar, Dr. Suparna Sanyal Mukherjee and Dr. Alminda Magbalot-Fernandez (eds.), *The Basic Handbook of Indian Ethnobotany and Traditional Medicine [Volume: 4]*, pp. 85-99. ISBN: 978-81-989555-5-5. DOI: <https://doi.org/10.52756/bhietm.2025.v04.009>

