

Phytochemical Composition, Antimicrobial, Anticancer Properties and Molecular Docking Profiles of *Tragopogon coelesyriacus* Boiss. Extracts

Tuba Unver (✉ tuba.unver@inonu.edu.tr)

Inonu University

Ugur Uzuner

Karadeniz Technical University

Selcen Celik-Uzuner

Karadeniz Technical University

Ismet Gurhan

Inonu University

Nur Sena Sivri

Karadeniz Technical University

Zeynep Ozdemir

Inonu University

Research Article

Keywords: *Tragopogon coelesyriacus*, Phenolic compounds, Antimicrobial activity, Anticancer activity, Molecular docking, In silico analyses

Posted Date: October 13th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3429191/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

Tragopogon coelestriacus is a biennial herbaceous plant of the Asteraceae family that grows mainly in Europe and Asia and is consumed as a vegetable in Turkey. This study aimed to obtain methanol and water extracts from aerial parts (stem, leaf, and flower) of the *T. coelestriacus* to determine its phenolic compounds and subsequently to determine the antimicrobial and anticancer activities of *T. coelestriacus* extracts as well as *in silico* molecular docking profiles for targets. *T. coelestriacus* methanol extract had a higher inhibitory effect against *S. aureus*, *P. aeruginosa*, and *K. pneumonia* (MIC: 0.83, 1.67 and 1.67 mg/mL, respectively) than *E. coli* and *E. aerogenes* (MIC: 53.3 mg/mL). Antifungal activity was observed against all *Candida* species, with the highest inhibition against *C. krusei* (MIC: 0.83 mg/mL). In contrast, *T. coelestriacus* water extract had no inhibitory effect. Furthermore, both extracts of *T. coelestriacus* showed a significant cytotoxic effect on AR42J pancreatic cancer cells. However, HeLa cervical and MDA-MB-231 breast cancer cells were more resistant to the cytotoxic effect of methanol and water extract, respectively. *In silico* analyses showed the inhibitory effects of Keracyanin chloride within the extracts on *S. aureus* MurB protein and the remarkable inhibitory potential of the Naringin molecule on FYN kinase protein regarding AR42J pancreatic cells. This study is the first *in vitro* and *in silico* analysis to show *T. coelestriacus* antimicrobial and anticancer effects along with molecular docking profiles, and it suggests that *T. coelestriacus* can be used as a potent pharmacological and therapeutic-protective agent.

1. Introduction

The use of phytotherapeutic plants in health care is as old as human history and dates back thousands of years [1]. In the last century, considering the severe side effects of chemical antibiotics and anticancer drugs, the popularity of these plants in traditional medicine has increased. The negativities of infectious diseases caused by pathogenic microorganisms and antibiotic resistance to treatment constitute serious problems worldwide [2–5]. Therefore, research on medicinal plants has intensified to discover bioactive plant extracts or compounds against microorganisms. In addition, Cancer has an increased mortality rate and is the second leading cause of death worldwide after cardiovascular diseases [6, 7]. In cancer treatment, resistance to cancer chemotherapy and severe side effects of chemotherapeutic drugs have led people to herbal treatment methods [8]. Anticancer studies with herbal substances have proven that thousands of plant species have anticancer effects worldwide [9–12]. In recent studies with species belonging to the *Tragopogon* genus, it has been stated that this plant may have phytotherapeutic activity [11, 13–15].

Tragopogon (Asteraceae) is represented by 150 species, native to the semi-arid region and growing mainly in Europe and Asia [16]. In Turkey, it is known as a manger or goatee and is consumed as a vegetable by the people of Eastern and Southeastern Anatolia [17]. The members of the *Tragopogon* species are distributed with 26 taxa belonging to approximately 22 species in Turkey [18, 19]. The leaves of the *T. coelestriacus* are grass-like, the stems have few branches, and they secrete a milky sap if the stems or leaves are broken [20]. In Turkish folk medicine, it has been found that some *Tragopogon* species have anthelmintic activities and are used to treat abdominal pain. In addition, the leaves and

stem of *T. coelesyriacus* are used in intestinal disorders [21, 22]. In studies with extracts of *Tragopogon* species, although it has been observed that this plant has antitumor, anti-inflammatory, antimicrobial, wound healing, enzyme inhibitor, antihyperlipidemic, and hepatoprotective effects, there are not enough studies on the toxicity assessment and chemical composition of the extracts [11]. These effects have been associated with compounds in the plant crude extracts. Flavonoids, terpenoids, and several natural products associated with various classes have been determined in numerous taxa of the genus *Tragopogon* [23–26]. Phenolic compounds such as Chlorogenic acid, Quinic acid, Gallic acid, Vitexin, Ferulic Acid, Rosmarinic Acid, and Luteolin were detected in the extracts of *Tragopogon* species [23–25, 27–30]. The plant's antimicrobial and antioxidant properties are related to its phenolic compounds [31]. Phenolic compounds show antimicrobial properties by damaging bacterial cell membranes and DNA during replication and inactivating bacterial enzymes [32]. Moreover, phenolic compounds reduce the redox potential of the environment due to the antioxidant activities of hydroxyl groups and prevent the growth of many aerobic microorganisms [33]. For this purpose, medicinal plants can be used to produce natural compounds with critical therapeutic properties [34].

Various antimicrobial and antitumor studies have been carried out in the literature on different species belonging to the *Tragopogon* genus. In these studies, the inhibition values of different species of *Tragopogon* on various microorganisms varied, and their anticancer activity was evaluated using different parts and extracts of plants on different cell lines [27, 35–38]. However, this study is the first to show the antimicrobial and anticancer properties of *T. coelesyriacus*. The objective of this study was to evaluate the antimicrobial and anticancer activity of water and methanol extracts of *T. coelesyriacus*, along with molecular docking profiles. These *in vitro* and *in silico* results show that *T. coelesyriacus* can be used as a potent herbal pharmacological agent in medicine, pharmaceuticals, the food industry, and even cosmetics products.

2. Materials and Methods

2.1. Collection of *Tragopogon coelesyriacus* and preparation of extracts

T. coelesyriacus (previously known as *Tragopogon porrifolius* subsp. *longirostris* (Sch. Bip.) Greuter, *Tragopogon longirostris* Sch. Bip. var. *longirostris*) aerial parts (stem, leaf, and flower) were collected from Puturge-Taslıpara district of Malatya province (Turkey) in April and May 2022 and brought to Inonu University, Faculty of Pharmacy, Pharmaceutical Botany laboratory, and was identified in this laboratory [39]. The plant aerial parts were dried and then crushed with a grinder before extraction. Two different solvents (water and methanol) were used in the extraction. 10 g of the aerial parts of the plants were weighed, and 100 ml of water or methanol was added to them and left to maceration at room temperature. This process was repeated three times by removing the filtrate and adding solvent. The collected filtrates were combined and evaporated in a rotary evaporator (Heidolph Laborota 4000,

Germany) until the solvents were removed [40]. The remaining extracts were stored in a deep freezer at – 18°C until used in studies.

2.2. LC-HRMS (Liquid Chromatography-High Resolution Mass Spectrometry) Analysis

Liquid Chromatography-High Resolution Mass Spectrometry (LC-HRMS) studies were carried out using an Electrospray Ionization (ESI) Source and a Reversed Phase C18 column on an Agilent 6460 Triple Quad LC-MS/MS with a 1290 Infinity UPLC system. The mobile phases A and B comprised 1% formic acid in water and 1% in acetonitrile, respectively. The gradient elution was applied. The column temperature was set to 30°C, and the mobile phase flow rate was 0.4 mL/min. The compounds were determined and identified by comparing the retention durations of reference compounds with HRMS data from the Eastern Anatolia High Technology Application and Research Center. The mobile phase (1 mL; A: B; 50:50; v/v) was dissolved with 1 mg of dry water and methanol extract. Before being injected into the LC in a volume of 5 µL, the solution was filtered through a 0.45-µm filter.

2.3. Antibacterial and antifungal activity

2.3.1. Strains

The following 9 microorganisms, consisting of 5 bacteria and 4 yeast, were studied: *Staphylococcus aureus* (ATCC 12600), *Escherichia coli* (C2987), *Enterobacter aerogenes* (ATCC 51697), *Pseudomonas aeruginosa* (ATCC 10145), *Klebsiella pneumoniae* (ATCC 13883), *Candida albicans* (ATCC 14053), *Candida tropicalis* (ATCC 13803), *Candida parapsilosis* (ATCC 22019), and *Candida krusei* (ATCC 14243). *E. coli* was purchased from New England Biolabs (NEB, MA, US). Other strains were purchased from the American Type Culture Collection (ATCC, VA, US).

2.3.2. Media

Muller Hinton Broth (MHB), Muller Hinton Agar (MHA), Sabouraud Dextrose Agar (SDA), and Sabouraud Dextrose Broth (SDB) were used for the antimicrobial activity of water and methanol extracts of *T. coelestis*. While MHA and MHB were used for bacterial strains, SDA and SDB were used for yeast strains.

2.3.3. Agar dilution assay

The agar dilution assay was carried out to analyze the antimicrobial properties of *T. coelestis* against nine different microorganisms, namely *S. aureus*, *E. aerogenes*, *E. coli*, *P. aeruginosa*, *K. pneumoniae*, *C. tropicalis*, *C. albicans*, *C. krusei* and *C. parapsilosis*. For two-fold agar dilution method, 640, 320, 160, 80, 40, 20, 10, 5, 2.5, 1.25, 0.6 g (w/v) methanol and water extracts of *T. coelestis* were added to 6 mL agar plates separately to obtain the concentration of 106.6, 53.3, 26.6, 13.3, 6.6, 3.3, 1.67, 0.83, 0.42, 0.21 and 0.1 (mg/mL) respectively. While MHA was used for bacterial strains, SDA was used for yeast strains. Pure MHA and SDA were prepared separately without the extracts for control plates. Direct colony suspensions were prepared in distilled water for each species, and their turbidity was

adjusted to 0.5 McFarland (1×10^8 cfu/mL for bacterial strains, 1×10^6 cfu/mL for yeast strains). Afterward, MHA plates were separated into five distinct regions: *S. aureus* to the 1st region, *E. aerogenes* to the 2nd region, *E. coli* to the 3rd region, *P. aeruginosa* to the 4th region, and *K. pneumoniae* to the 5th region. Furthermore, SDA plates were divided into four regions: *C. albicans* to the 1st region, *C. tropicalis* to the 2nd region, *C. krusei* to the 3rd region, and *C. parapsilosis* to the 4th region, as applied before [41]. 1 μ l of each microorganism from the standard inoculum was inoculated onto the agar plates in each region, to which different *T. coelestis* extracts concentrations were added. In an incubator, inoculated media were incubated at 36°C overnight. Thereafter, the growth of each microorganism strain on the plates was evaluated, and the minimum inhibitory concentrations (MIC) were determined for each strain (Fig. 1). The experiment was performed in triplicate.

2.3.4. Broth microdilution method

In the broth two-fold microdilution assay, the antibacterial and antifungal activity tests of *T. coelestis* were repeated against nine different microorganisms, including *S. aureus*, *E. aerogenes*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, *C. albicans*, *C. tropicalis*, *C. krusei* and *C. parapsilosis*, using a 96-well microplate. 21.33, 10.66, 5.33, 2.66, 1.33, 0.66, 0.33, 0.16, 0.08, and 0.04 mg of methanol and water extracts of *T. coelestis* were added to 200 μ l MHB for bacterial strains, and SDB for yeast strains in each well. For control groups, pure MHB and SDB without extracts were placed in one of the control wells as negative controls, and microorganisms used with MHB and SDB were placed in the other well as positive controls. Each microorganism's direct colony suspension was prepared in distilled water, and their turbidity was adjusted to standard inoculum 0.5 McFarland. Afterward, 1 μ l of *S. aureus*, *E. aerogenes*, *E. coli*, *P. aeruginosa*, *K. pneumoniae*, *C. albicans*, *C. tropicalis*, *C. krusei*, and *C. parapsilosis* were inoculated into the A, B, C, D, E, F, G, H, and I row of the microplate respectively. The microplate with different extracts of *T. coelestis* concentrations was incubated at 36°C for 24 hours, and subsequently, 15 μ l of the resazurin (0.15% w/v) was added to per well. The microplate was incubated at 36°C for 4 hours, and the growth of microorganisms was interpreted by following the change in well colors.

2.4. Anticancer Activity

2.4.1. Cell culture

This study included 3 epithelial cancerous (AR42J, pancreatic cancer; HeLa, cervical cancer and MDA-MB-231 breast cancer) and 2 non-cancerous epithelial human cells (ARPE-19, retina; and MCF10A, breast). AR42J and MDA-MB-231 cells were cultured in RPMI media. HeLa cells were cultured in EMEM media. ARPE-19 and MCF10A cells were cultured in DMEM. Each media consisted of penicillin-streptomycin (1%) and fetal bovine serum (10%). Cells were cultured in a 37°C incubator with 5% CO₂ humidification. Cells were cultured in 96-well plates at 10,000 cells per well. After they reached at 80% confluency, cells were treated with extracts. For this, experiments for each cell included two groups, 1) treated with *T. coelestis* extracts obtained by methanol for 24h, 48h and 72h at 3.125 μ g/ml, 6.25 μ g/ml, 12.5 μ g/ml, 25 μ g/ml, 50 μ g/ml, 100 μ g/ml and 200 μ g/ml, and 2) treated with *T. coelestis* extracts

obtained by water for 24h, 48h and 72h at 3.125 µg/ml, 6.25 µg/ml, 12.5 µg/ml, 25 µg/ml, 59 µg/ml, 100 µg/ml and 200µg/ml. Control wells for each cell were untreated (0 µg/ml). For each dose at each concentration, at least 3 wells were used as intra-experimental repeats. The whole experiments for each cell were repeated as two to three independent replications.

2.4.2. MTT cytotoxicity assay

After treatment with different extracts, media with extracts was removed, and the MTT protocol was applied as before [42]. In this method, 200 µl of media including MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) was added to each well and incubated at 37°C for 2.5 hours. After incubation, media with MTT was removed, and 150 µl of DMSO was added to each well for dissolving formazan crystals. Subsequently, plates were incubated in the dark on a shaker overnight. Absorbances at 570 nm were read the day after. Cell viabilities (as percentages) were calculated according to the viability of untreated cells considered as 100%. If applicable, IC₅₀ values were calculated as before [43].

2.5. Molecular docking

To elucidate the antibacterial and anticancer effects of *T. coelesyriacus* natural compounds, two-way molecular insertion studies were carried out. *In silico* molecular docking studies on *S. aureus* UDP-*N*-acetylenolpyruvylglucosamine reductase (MurB; PDB ID: 1HSK) were performed to investigate antibacterial activity potentials of *T. coelesyriacus* natural compounds. The initial stage of peptidoglycan synthesis in *S. aureus* involves the conversion of phosphoenolpyruvate and UDP-*N*-acetylglucosamine to UDP-*N*-acetylenolpyruvylglucosamine (UDP-GlcNAcEP)₃ by MurA. Afterwards, UDP-GlcNAcEP is reduced to UDP-*N*-acetylmuramic acid by MurB. Then, MurC, MurD, MurE, and MurF enzymes add five amino acid residues (L-Lys, L-Ala, D-Ala-D-Ala, and D-Glu) sequentially to the D-lactyl group of UDP-*N*-acetylmuramic acid in a ribosome-independent manner. Since MurB is a crucial enzyme in the peptidoglycan synthesis of *S. aureus* [44], chemical compounds that hinder MurB activity are considered as potential antimicrobial agents [45].

The FYN protein was investigated to elucidate the anticancer effects of *T. coelesyriacus* natural compounds on pancreatic cells. FYN is a non-receptor tyrosine kinase involved in various cellular processes such as cellular adhesion, cell motility, cell proliferation, apoptosis, cytoskeletal remodeling, integrin-mediated signaling, cell growth and survival regulation, immune response, and axon guidance [46]. When activated upon phosphorylation from its C-terminal tail, it initiates the phosphorylation of PTK2/FAK1 to focal adhesions. FYN eventually regulates cell adhesion and motility through phosphorylation of CTNND1 (delta-catenin) and CTNNB1 (beta-catenin) [47]. It has recently been reported that FYN kinase is overexpressed in many tumors as well as pancreatic cell lines and promotes cancer metastasis and growth [48]. Furthermore, FYN is involved in the regulation of multiple cancer-associated signaling pathways, including interactions with COX-2, STAT5, ERK, AKT, and MET. Therefore, FYN is a manifested therapeutic target for various types of tumors, and suppression of FYN can improve prognosis and prolong patient survival [48].

Molecular docking studies were carried out by employing AutoDock Vina 1.5.7 program [49]. 3D structures of *S. aureus* MurB (PDB ID: 1HSK) and human FYN kinase (PDB ID: 2DQ7) as target protein structures were retrieved from the Protein Data Bank (PDB). Water molecules and any bound ligand were excluded from target protein 3D structures. However, Flavin Adenine Dinucleotide (FAD), the major cofactor of the MurB protein bound to the 1HSK active site, was retained during MurB-related molecular docking studies. Both target protein structures were monomerized, and the protonation states were counted by PROPKA 3.0 at pH = 7.0 [50]. All amino acids were charged accordingly. The pdb files of natural ligands were built using Open Babel [51]. The pdbqt file formats of both target proteins and extracted ligand compounds were prepared by employing AutoDock tools 1.5.7.

Polar hydrogens were added to the 3D structures of both MurB and FYN, and Kollman charges were also calculated. Gasteiger charges were calculated for natural compounds as ligands, and pdbqt files were saved. Each docking assay was applied on both MurB and FYN protein targets via three independent runs using ligand pdbqt files. The grid dimensions for 1HSK were set as $-30\text{\AA} \times 25\text{\AA} \times 30\text{\AA}$ and grid center coordinates were adjusted as $x = 178\text{\AA}$, $y = 155\text{\AA}$, $z = 155\text{\AA}$, respectively. Whereas 2DQ7 docking coordinates were set as $-19\text{\AA} \times 20\text{\AA} \times -15\text{\AA}$ for grid dimensions and $x = 25\text{\AA}$, $y = 35\text{\AA}$, $z = 25\text{\AA}$ for grid center coordinates. The docking outcomes for each compound were investigated in detail, and specific 2D interaction maps were analyzed using the Discovery Studio Accelrys (Discovery Studio Visualizer-v21.1.0.20298) program (Dassault Systems, BIOVIA).

2.6. Statistics

For statistical analyses, version 13 of the SPSS program was used. Cell viabilities (%) calculated from the MTT assay were statistically compared using the UNIANOVA test. p values less than 0.05 were considered as significant. The significance levels used are $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***) and $p < 0.0001$ (****). Experiments included inter and intra-repeats, and standard errors of the mean ($\pm$ s.e.m.) were shown by error bars in the graphs.

3. Results

3.1. Analysis of the extracts using LC-HRMS

The prepared plant extracts were analyzed with LC-HRMS to determine their content. The LC-HRMS analysis of *T. coelestiacus*, which has the highest total phenolic content, was carried out due to the spectrophotometric study. It identified 12 chemicals as being present. Table 1 displays the retention time (Rt), the amount of each chemical, and the discovered molecular ion. In *T. coelestiacus*, 25.12 g/kg of individual phenolic compounds were discovered in methanol extract, and 13.03 g/kg of individual phenolic compounds were discovered in water extract. The most prevalent compound in methanol extract was chlorogenic acid, which had a mass of 13.58 g/kg, followed by quinic acid, 7.81 g/kg, and vitexin, 1.41 g/kg. Additionally, the most prevalent compound in water extract was quinic acid, which had a mass of 8.79 g/kg, followed by epicatechin, 1.36 g/kg.

Table 1
Quantitative phenolic content of water and methanol extracts.

Compounds	Found m/z (molecular ion)	Rt (min)	Water Extract	Methanol Extract
			Amount of phenolic comp. g/kg	Amount of phenolic comp. g/kg
Fumaric acid	114.9	4.032	0.5916	0.5350
Chlorogenic acid	352.9	10.824	0.9763	13.5768
Keracyanin chloride	592.8	9.892	0.0061	0.1100
Quinic acid	190.9	2.477	8.7850	7.8116
Gallic acid	168.9	5.567	0.2541	0.0721
Cyanidin-3- <i>O</i> -glucoside	447.1	10.372	0.0991	0.2877
Epicatechin	289.0	11.508	1.3567	0.6443
Vitexin	430.9	11.548	0.9293	1.4102
Naringin	579.0	11.952	-	0.0037
Ferulic Acid	193.0	12.542	-	0.0210
Rosmarinic Acid	358.8	12.462	0.0286	0.4425
Luteolin	284.9	13.409	-	0.2067

3.2. Antibacterial and Antifungal Activity

3.2.1. Results of agar dilution method

In the agar dilution method, the antibacterial activity results of the methanol extract of *T. coelestriacus* against *S. aureus*, *E. aerogenes*, *E. coli*, *P. aeruginosa*, and *K. pneumonia* are demonstrated in Fig. 1. After incubation at 36 °C 24 hours, the plates of 0.42, 0.21, 0.1 mg/mL and the control (0 mg/mL) methanol extract of *T. coelestriacus* against *S. aureus*; plates, J, K, L, and I (a few colonies) were covered with pale yellow bacterial colonies, respectively. No colonies were detected on plates A, B, C, D, E, F, G, and H treated with 106.6, 53.3, 26.6, 13.3, 6.6, 3.3, 1.67, and 0.83 mg/mL methanol extract of *T. coelestriacus*, respectively. Therefore, the MIC value of the methanol extract of *T. coelestriacus* against *S. aureus* was found to be 0.83 mg/mL. Colonies began to seem from the C-plate, where 26.6 mg/mL methanol extract of *T. coelestriacus* concentration was used for *E. coli* and *E. aerogenes*. Hence, this experiment determined the MIC value of *T. coelestriacus* methanol extract against *E. aerogenes* and *E. coli* to be 53.3 mg/mL. *K. pneumonia* and *P. aeruginosa* colonies were evident from the H-plate, where 0.83 mg/mL methanol extract of *T. coelestriacus* concentration was used. Therefore, the MIC value of the methanol

extract of *T. coelesyriacus* against *K. pneumonia* and *P. aeruginosa* was found to be 1.67 mg/mL. (Fig. 1).

The antifungal activity results of *T. coelesyriacus* methanol extract against *C. albicans*, *C. tropicalis*, *C. krusei*, and *C. parapsilosis* are demonstrated in Fig. 2. After incubation at 36 °C for 24 hours, *C. albicans* colonies began appearing from the C-plate, where 26.6 mg/mL of methanol extract concentration was used. So, the MIC value of the methanol extract of *T. coelesyriacus* against *C. albicans* was found to be 53.3 mg/mL. *C. tropicalis*, *C. krusei*, and *C. parapsilosis* colonies began appearing from the B-plate, I-plate, and D-plate, respectively. Therefore, the MIC values against *C. tropicalis*, *C. krusei*, and *C. parapsilosis* were 106.6, 0.83, and 26.6 mg/mL, respectively. In antibacterial and antifungal studies with water extract of *T. coelesyriacus* against *S. aureus*, *E. aerogenes*, *E. coli*, *K. pneumonia*, *P. aeruginosa*, *C. albicans*, *C. tropicalis*, *C. krusei*, and *C. parapsilosis*, it was observed that water extract of *T. coelesyriacus* did not have any inhibitory effect on the bacterial and yeast strains.

3.2.2. Results of broth microdilution method

The antibacterial and antifungal activity of the methanol extract of *T. coelesyriacus* against *S. aureus*, *E. aerogenes*, *E. coli*, *P. aeruginosa*, *K. pneumonia*, *C. albicans*, *C. tropicalis*, *C. krusei* and *C. parapsilosis* was repeated using the broth dilution assay. Methanol extract of *T. coelesyriacus* was added to wells 1–10 on the microplate at concentrations of 21.33, 10.66, 5.33, 2.66, 1.33, 0.66, 0.33, 0.16, 0.08, and 0.04 mg/mL, respectively. The negative control is the 11th well, which confirms no contamination has occurred during the experiment. The positive control is the 12th well representing the viability of the microbial strains. The MIC value is determined as the methanol extract of *T. coelesyriacus* concentration with the least amount of the extract on the microplate with no change in color. Pink indicates microorganism growth and, thus viability, while blue indicates inhibition of the growth. As a result of this experiment, same as the agar dilution test results, the MIC values of the methanol extract against bacterial strains *S. aureus*, *E. coli*, *E. aerogenes*, *P. aeruginosa*, and *K. pneumonia* were determined as 0.83, 53.3, 53.3, 1.67 and 1.67 mg/mL respectively, in the 9th, 2nd, 2nd, 7th and 7th wells without color change. The MIC values of the methanol extract against yeast strains *C. albicans*, *C. tropicalis*, *C. krusei* and *C. parapsilosis* were found to be 53.3, 106.6, 0.83, and 26.6 mg/mL, respectively, in the 2nd, 1st, 8th and 4th wells with no color change (Fig. 3, Table 2). The antifungal and antibacterial activity of the water extract of *T. coelesyriacus* was repeated using a broth dilution assay. However, it was determined that the water extract of *T. coelesyriacus* had no inhibitory effect on the bacterial and yeast strains.

Table 2
MIC values of the methanol extract of *T. coelestis* against different microorganisms.

Microorganisms	Methanol extract of <i>T. coelestis</i> (mg/mL)
<i>S. aureus</i>	0.83
<i>E. aerogenes</i>	53.3
<i>E. coli</i>	53.3
<i>P. aeruginosa</i>	1.67
<i>K. pneumonia</i>	1.67
<i>C. albicans</i>	53.3
<i>C. tropicalis</i>	106.6
<i>C. krusei</i>	0.83
<i>C. parapsilosis</i>	26.6

3.2.3. Cytotoxic profile in human cancerous and normal cells

The treatment with the maximum dose (200 µg/ml) of methanol extract of *T. coelestis* for 24h and 48h resulted in the death of almost half of AR42J but not at 72h (Fig. 4A, **up panel**), while the viability of MDA-MB-231 cell population decreased at half after the maximum dose for 72h only (Fig. 4C, **up panel**). However, HeLa cells were the most resistant to methanol extract as 70% of cells were still viable after the maximum dose for the extended incubation till 72h (Fig. 4B, **up panel**). AR42J was also more responsive to lower doses (50 µg/ml for 24h and 100 µg/ml for 48h) (Fig. 4A, **up panel**) compared to HeLa and MDA-MB-231 cells.

In the experimental groups for the treatments with water-extracted *T. coelestis*, AR42J was the more responsive cancer cell to the lower doses from 6.25 µg/ml to 200 µg/ml for 72h as half of AR42J died (Fig. 4A, **down panel**). At lower incubations (24h and 48h), the cell death rate in AR42J was higher than other cells at the same doses of water extracts (Fig. 4B and 4C, **down panels**). 50% of HeLa cells died at 200 µg/ml for 72h (Fig. 4B, **down panel**). However, MDA-MB-231 cells were the most resistant cell type to water extract of *T. coelestis* (Fig. 4C, **down panel**).

Statistical analyses for viability were summarized for each cell under each graph in Fig. 4. According to the UNIANOVA test, the dose itself was significant in AR42J, HeLa, and MDA-MB-231 cells with *p* values as 0 (****). Extraction type was significant in only MDA-MB-231 cells (*p* < 0.0001) but not in other cells, as both extracts were cytotoxic in AR42J, and the extraction method did not reveal a significant difference in HeLa cells. However, incubation itself was not significant in MDA-MB-231 cells but significant in HeLa (*p* < 0.0001) and AR42J (*p* < 0.01) cells. The synergistic effect of dose and incubation were significant in

AR42J cells only ($p < 0.001$), while extract and incubation together significantly affected cell viability in AR42J, HeLa, and MDA-MB-231 cells with p values less than 0.0001 (****), 0.5 (*) and 0.001 (***), respectively. The total effect of dose, incubation, and extract was significant in HeLa cells only. Calculated IC50 values representing the cytotoxic dose for half of the cell population are shown in Table 3. The minimum IC50 values were detected for 1) 24h incubation with methanol extract in AR42J cells (Average: 77.31), 2) 72h incubation with water extract in AR42J cells (Average: 75.04), and 3) 72h incubation with methanol extract in MDA-MB-231 cells (Average: 76.69). The less IC50 means, the higher the cytotoxic effect. In normal cells, IC50 values could not be detected because there were no conditions that 50% of cell death occurred.

Table 3
IC50 values for in vitro cytotoxicity of *T. coelestis* in human cancers

Cell	Extract	Incubation	IC50 value (µg/ml)
AR42J	Methanol	24h	77.31 ± 18.58
		48h	86.94 ± 20.04
	Water	48h	92.02 ± 26.38
		72h	75.04 ± 14.07
HeLa	Water	72h	91.30 ± 18.10
MDA-MB-231	Methanol	72h	76.69 ± 2.11

The same experiments were performed for non-cancerous healthy cells including ARPE-19 and MCF10A (Fig. 5A-B). The combined effect of dose, extract type, and incubation was insignificant in ARPE-19 (Fig. 5A) but significant in MCF10A cells (***). Each nominal parameter including dose, incubation, and extract was individually significant in both cells. But the minimum cell viability (around 80%) was detected in normal cells after the treatment with water extraction when compared to cancer cells. In general, the cell viability of normal cells was significantly higher than that of cancerous cells. Therefore, IC50 values for normal cells could not be detectable.

3.3. Molecular docking data

Most of the native compounds of *T. coelestis* were energetically preferred for the *S. aureus* MurB active site. Of these, keracyanin chloride, naringin, and vitexin were determined as the leading MurB inhibitory compounds of *T. coelestis* in the presence of low binding free energy values: -9.4, 9.2 and 8.2 kcal/mol, respectively (Table 4).

Table 4

Molecular docking data and interacting residue profiles of *T. coelestis* native compounds when directed to *S. aureus* MurB enzyme.

Compound Id	Binding Energy 1HSK (kcal/mol)	Interacting Residues	H Bonds
Chlorogenic acid	-7.2	Chain A: ALA154 TYR155 GLY156 TYR187 ARG225 LYS228 GLN229 SER238 PHE240 GLN241 ARG242 PRO243 PHE247 ALA248 GLY249 HIS271 PHE274	4
Cyanidin-3-O-glucoside	-7.9	Chain A: GLY153 ALA154 TYR155 TYR187 ARG188 ARG225 GLN229 SER238 PHE240 GLN241 ARG242 PHE247 ALA248 GLY249 LYS250 HIS271 GLY273 GLU308	5
Epicatechin	-7.1	Chain A: GLY153 ALA154 TYR155 ARG225 GLN229 SER238 ARG242 PHE247 ALA248 GLY249 LYS250 HIS271 ALA272 GLY273 PHE274	3
Ferulic acid	-6.2	Chain A: LEU17 VAL25 ALA37 LYS39 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 ASN86 GLY88 SER89 LEU137 ALA147	4
Fumaric acid	-4.3	Chain A: VAL25 ALA37 ILE38 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 LEU137 ALA147 ASP148 PHE149	4
Gallic acid	-5.8	Chain A: PHE22 LYS39 MET46 SER50 PHE51 LEU52 GLU53 GLU54 ALA55 ASP148 PHE149 GLY150 LEU151 ARG153	2
Keracyanin chloride	-9.4	Chain A: ASN83 MET150 GLY153 ALA154 TYR155 TYR187 ARG188 ARG225 GLN229 GLY237 SER238 PHE240 GLN241 ARG242 PHE247 ALA248 GLY249 LYS250 HIS271 PHE274 GLU308	6
Luteolin	-7.7	Chain A: ASN83 MET150 ALA154 TYR155 TYR187 ARG188 ARG225 GLN229 SER238 PHE240 ARG242 PHE274 GLU308	7
Naringin	-9.2	Chain A: GLY153 ALA154 TYR155 GLY156 GLY157 GLU158 TYR187 ARG188 ARG224 ARG225 LYS228 GLN229 SER238 PHE240 GLN241 ARG242 ALA248 HIS271 PHE274 GLU308	6
Quinic acid	-5.6	Chain A: LEU17 VAL25 ALA37 ILE38 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 LEU137 ALA147 ASP148 PHE149	4
Rosmarinic acid	-7.3	Chain A: ASN83 MET150 TYR187 ARG188 GLY237 SER238 PHE240 GLN241 ARG242 PRO243 HIS246 PHE247 ALA248 GLY249 LYS250 HIS271 GLU308	7
Vitexin	-8.2	Chain A: GLY153 ALA154 TYR155 TYR187 ARG188 ARG225 GLN229 SER238 PHE240 GLN241 ARG242 PHE247 ALA248 GLY249 LYS250 HIS271 PHE274 GLU308	5

In-depth analysis of the keracyanin chloride binding mode to the catalytic core of the *S. aureus* MurB protein revealed that it is capable of forming 6 hydrogen bonds with the 1HSK active site residues

(Fig. 6). Similarly, as an indicator of strong binding, naringin has the potential to form a total of 6 H bonds with 1HSK catalytic residues (Fig. 7). Furthermore, rosmarinic acid and luteolin, other two natural compounds of *T. coelestis*, were able to form 7 H bonds within the catalytic core of MurB, suggesting their potential antibacterial activity against *S. aureus* pathogenic bacteria.

Cytotoxicity analysis revealed that *T. coelestis* natural compounds exhibited selective antitumor activity on pancreatic cancer cells. Pancreatic cancer-specific marker proteins were investigated to elucidate the key players of such anticarcinogenic activity. The pancreatic cancer marker gene product FYN kinase was therefore targeted for *in silico* molecular docking analysis using *T. coelestis* natural compounds. Docking analysis on FYN revealed that most of the native compounds of *T. coelestis* showed a high binding affinity to FYN in the presence of low binding free energy. Among those compounds, naringin, vitexin, keracyanin chloride, luteolin, and cyanidin-3-*O*-glucoside were identified with high binding affinity to FYN protein with low binding free energies, -10.1, -9.8, -9.7, -9.5 and - 9.4 kcal/mol, respectively (Table 5).

Table 5

Molecular docking data and interacting residue profiles of *T. coelestiacus* native compounds when directed to FYN.

Compound Id	Binding Energy 2DQ7 (kcal/mol)	Interacting Residues	H Bonds
Chlorogenic acid	-8.1	Chain X: LEU17 ASN19 VAL25 ALA37 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 ASN86 LYS87 GLY88 SER89 ASN135 LEU137 ALA147 ASP148 PHE149	1
Cyanidin-3-O-glucoside	-9.4	Chain X: LEU17 GLY18 ASN19 VAL25 ALA37 ILE38 LYS39 GLU54 MET58 VAL67 ILE80 THR82 TYR84 MET85 ASN86 LYS87 GLY88 SER89 ALA134 ASN135 ILE136 LEU137 ILE146 ALA147 ASP148	4
Epicatechin	-8.3	Chain X: LEU17 GLY18 VAL25 ALA37 LYS39 GLU54 MET58 VAL67 THR82 GLU83 TYR84 MET85 ASN86 GLY88 SER89 ALA134 ASN135 LEU137 ALA147 ASP148 PHE149	2
Ferulic acid	-6.2	Chain X: LEU17 VAL25 ALA37 LYS39 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 ASN86 GLY88 SER89 LEU137 ALA147	1
Fumaric acid	-4.3	Chain X: VAL25 ALA37 ILE38 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 LEU137 ALA147 ASP148 PHE149	4
Gallic acid	-5.8	Chain X: PHE22 LYS39 MET46 SER50 PHE51 LEU52 GLU53 GLU54 ALA55 ASP148 PHE149 GLY150 LEU151 ARG153	N/A
Keracyanin chloride	-9.7	Chain X: ARG16 LEU17 GLY18 ASN19 VAL25 ALA37 ILE38 LYS39 GLU54 MET58 VAL67 ILE80 VAL81 THR82 TYR84 MET85 ASN86 LYS87 GLY88 SER89 LEU91 ASP92 ARG132 ALA134 ASN135 LEU137 ALA147 ASP148	8
Luteolin	-9.5	Chain X: LEU17 GLY18 VAL25 ALA37 LYS39 GLU54 MET58 VAL67 LEU69 ILE80 THR82 GLU83 TYR84 MET85 ASN86 GLY88 SER89 LEU137 ALA147 ASP148 PHE149	3
Naringin	-10.1	Chain X: LEU17 GLY18 ASN19 VAL25 ALA37 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 ASN86 GLY88 SER89 ASP92 ALA134 ASN135 ILE136 LEU137 ALA147 ASP148 PHE149	4
Quinic acid	-5.6	Chain X: LEU17 VAL25 ALA37 ILE38 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 LEU137 ALA147 ASP148 PHE149	4
Rosmarinic acid	-8.2	Chain X: LEU17 VAL25 ALA37 LYS39 GLU54 MET58 VAL67 ILE80 THR82 GLU83 TYR84 MET85 ASN86 LYS87 GLY88 SER89 ALA134 ASN135 ILE136 LEU137 ILE146 ALA147 ASP148 PHE149	4
Vitexin	-9.8	Chain X: LEU17 GLY18 ASN19 GLY20 GLY23 VAL25 ALA37 LYS39 GLU54 MET58 VAL67 THR82 GLU83 TYR84 MET85 GLY88 SER89 ALA134 ASN135 LEU137 ALA147 ASP148	5

Among the analyzed compounds, *T. coelesyriacus* natural compound naringin was determined to have the highest binding affinity to FYN in the presence of the lowest binding free energy (-10.1 kcal/mol) (Fig. 8).

Furthermore, another ambitious inhibitor candidate for the FYN protein against pancreatic cancer cell lines was identified as keracyanin chloride with its high binding affinity, low binding free energy, and highest (8) H bond-forming ability (Fig. 9).

4. Discussion

Leafy green vegetables are rich in bioactive phenolic compounds and have various biological functions, such as antioxidant, antimicrobial, and antitumor effects resulting from these phenolic compounds. These plants also contain large amounts of polyphenols, such as aromatic compounds, flavonoids, phenolic acids, and the most abundant phytochemicals in the human diet [32]. As a result of our extraction study with *T. coelesyriacus*, 14 phenolic compounds were found by LC-HRMS analysis; chlorogenic acid (13.58 g/kg) was the most prevalent in methanol extract, and quinic acid (8.79 g/kg) was the most prevalent in water extract. The amount of chemicals in the water extract is lower despite more phenolic compounds flowing into the methanol extract. The increase in the number of carbons in the structure of the compounds explains this. The enhanced solubility of compounds like naringin, ferulic acid, and luteolin in methanol, which is somewhat more nonpolar than in water, and the fact that these chemicals are not present in the water extract are due to an increase in the number of nonpolar carbons in the structures. However, even if only slightly, the passage of compounds into water extract can be explained by molecules like chlorogenic acid and rosmarinic acid, whose structure is as large as the others. This is because these molecules have functional groups like ester and carboxylic acid in their structures, increasing water solubility by adding extra hydrogen bonds. In addition, despite its complex structure, the keracyanin chloride molecule was also in the form of salt, although it was in a minimal amount in the water extract (Fig. 10).

In the literature, various results were obtained in antimicrobial studies with different species of the genus *Tragopogon* (*Tragopogon dubius* Scop., *Tragopogon oligolepis* Hartvig & Strid, *Tragopogon porrifolius* L., and *Tragopogon graminifolius* DC.), with varying MIC values. In the *in vitro* study of Moromete et al., inhibitory effects of the two extracts obtained from the root, stem, and leaves of *T. dubius* were analyzed against *S. aureus*, *E. coli*, and *Streptococcus mutans* (for antibacterial), and *C. albicans* (for antifungal). It was determined that these extracts did not have an inhibitory effect on *C. albicans* and *S. aureus*, and the plant root extract had an inhibitory effect against *E. coli* and *S. mutans* at high concentrations (25%, 50%). MIC values could not be determined in this study performed with the disk diffusion method [35]. In the study of Farzaei et al., the antibacterial properties of extracts and the essential oil of *T. graminifolius* aerial parts were examined using disc diffusion and two-fold micro-well dilution methods. In conclusion, in the analysis against *S. aureus*, *K. pneumoniae*, *Proteus vulgaris*, and *Shigella dysenteriae* strains, the highest inhibition was found against *S. dysenteriae* with a MIC value of 2.5 mg/mL [52]. In the antimicrobial activity study of Uysal et al., ethyl acetate, methanol, and water extracts of *T. dubius* leaf

were used. In this study performed by microdilution method, MIC (between 0.06 mg/mL and 0.40 mg/mL) and the minimum bactericidal concentrations (MBC) (between 0.124 mg/mL and 0.45 mg/mL) were determined for bacterial microorganisms. For yeast strains, The MIC and the minimum fungicidal concentrations (MFC) values were found to be between 0.02 and 0.18 mg/mL and between 0.03 to 0.25 mg/mL, respectively [27]. In another *in vitro* study, the antimicrobial activity of different extracts of aerial parts of *T. oligolepis* was investigated using the disc diffusion assay. The best inhibitory activity was seen in the ethyl alcohol extract of the plant against all Gram-negative bacteria with an inhibition zone diameter of 11–21 mm and multi-antibiotic resistant *Staphylococcus*. While the chloroform extract of the plant showed weak inhibition, the ethyl acetate extract did not show any inhibitory effect [22]. In the study using the micro-well dilution assay, *T. porrifolius* antibacterial activity against *S. aureus*, *E. coli*, *Proteus mirabilis*, and *Enterococcus faecalis* was examined, and *T. porrifolius* showed activity only against *E. coli*, and MIC values were not determined [14]. The results above show that most *Tragopogon* species may have antimicrobial properties in studies with extracts from the *Tragopogon* genus. However, the methods used (disk diffusion, etc.) and the lack of MIC values in some studies have reduced their scientific value. Apart from all these studies, antimicrobial and anticancer properties of extracts of *T. coelesyriacus* were studied in our study. In conclusion, MIC values of the methanol extract of *T. coelesyriacus* against *S. aureus*, *E. coli*, *E. aerogenes*, *P. aeruginosa*, *K. pneumonia*, *C. albicans*, *C. tropicalis*, *C. krusei*, and *C. parapsilosis* were determined as 0.83, 53.3, 53.3, 1.67, 1.67, 53.3, 106.6, 0.83, and 26.6 mg/mL respectively. In the antibacterial study, the methanol extract of *T. coelesyriacus* had the highest inhibitory effect against *S. aureus*, followed by *P. aeruginosa*, *K. pneumonia*, *E. coli*, and *E. aerogenes*, respectively. The differences between the antimicrobial properties of plant extracts against Gram-positive and Gram-negative bacteria result from the different cell wall structures of these microorganisms [32]. *S. aureus*, a Gram-positive microorganism, has a single-layer of peptidoglycan. Due to its single-layer structure, it does not show sufficient buffering against the protonation impacts produced by phenolic substances. Thus, bacterial cell cytoplasm may become acidified, and bacterial energy metabolism may be impaired. Nevertheless, lipopolysaccharides (LPS) and small membrane components in the cell membrane of Gram-negative bacteria act as an inhibitory barrier against hydrophobic compounds. Therefore, they are more resistant to the antimicrobial effects of phenolic compounds [53]. In addition, antifungal activity was observed in all *Candida* species, while the highest inhibition was seen in *C. krusei*, with a MIC value of 0.83 mg/mL. In contrast, the water extract of *T. coelesyriacus* had no inhibitory effect on the bacterial and yeast strains. Changes in the antimicrobial effects of *T. coelesyriacus* water and methanol extracts may have resulted from the differences in the phenolic compounds in the extracts.

Plant extracts usually contain antimicrobial compounds, and phenolic compounds form the main component of these antimicrobial compounds [54]. It has been observed that phenolic compounds found in many plant species protect against pathogenic microorganisms [55]. – OH groups in phenolic compounds might be responsible for plants' antimicrobial and antioxidant effects [56]. Phenolic compounds gain antimicrobial properties by causing dysfunction in bacterial cell membranes. The -OH groups in phenolic compounds are highly reactive in aqueous environments. It reacts with several biomolecules and causes the deformation of these molecules, thus preventing bacterial growth and

reproduction. Phenolic compounds also bind to cell walls and proteins, inactivating bacterial enzymes and intercalating with bacterial DNA [57, 58]. In this study, as a result of the LC-HRMS analysis, it was found that both the number and the amount of phenolic compounds in the methanolic extract of *T. coelestyriacus* were higher than in the water extract. Almost 14 times higher chlorogenic acid in methanol extract (13.5768 g/kg) compared to water extract (0.9763 g/kg) may have increased the antimicrobial activity of the plant. Chlorogenic acid causes an important decline in intracellular ATP (adenosine triphosphate) concentration, possibly by affecting cell signaling and energy metabolism. It exerts a bacteriostatic effect by causing an intracellular metabolic imbalance of the glycolysis and tricarboxylic acid (TCA) cycle [59]. Chlorogenic acid increases the plasma and outer membrane permeability, causing loss of barrier function and consequently disrupting bacterial cell membranes [60]. In our study, phenolic compounds such as Keracyanin chloride, Cyanidin-3-*O*-glucoside, Vitexin, and Rosmarinic Acid in methanol extract more than in water extract may have affected the antimicrobial mechanism. Another study found that Cyanidin-3-*O*-glucoside showed outstanding antibacterial activity by showing a bactericidal effect against *S. aureus*. Changes in intracellular PH, intracellular ATP concentration, cellular morphology of *S. aureus*, and decreased membrane integrity caused cell membrane dysfunction [61]. The highest antibacterial activity achieved against *S. aureus* in our results may be associated with cyanidin-3-*O*-glucoside, found in notable amounts in the methanol extract. Rosmarinic acid is a powerful antimicrobial agent that inhibits the growth of many bacterial and yeast strains and can inhibit the biofilm formation of microorganisms. It exhibits antibiofilm properties by destroying established biofilms and moderately reducing exopolysaccharide production. Rosmarinic acid exerts an antifungal effect by reducing mitochondrial activity, altering membrane integrity, and mildly inhibiting protease production in fungal cells [62, 63]. In addition, in our study, three phenolic compounds (Naringin, Ferulic Acid, and Luteolin), which are present in the methanol extract but not in the water extract, may have contributed to the antimicrobial activity of the methanol extract (Table 1). Studies have shown that Naringin, Ferulic Acid, and Luteolin have serious antimicrobial activities by damaging cell membrane integrity or preventing biofilm formation [64–66]. The absence of these three phenolic compounds in the water extract in our study may be associated with the lack of antimicrobial properties of the water extract. Considering that most phenolic compounds contained in *T. coelestyriacus* have bactericidal effects, it is thought that the effect of *T. coelestyriacus* against microorganisms can be in this direction.

Our current molecular insertion studies have provided useful information about the antibacterial potentials of *T. coelestyriacus* leaf extracts that add to previous findings. In addition, molecular docking studies have illuminated for the first time the remarkable inhibitory effects of Keracyanin chloride on *S. aureus* MurB. In addition, the Naringin molecule showed high binding affinity to *S. aureus* MurB in the presence of low binding free energy. Although the antimicrobial activity potential of Naringin has been reported elsewhere [67], we report here for the first time its inhibitory activity on the *S. aureus* MurB enzyme. Our current molecular docking studies have provided information supporting previous findings. In sequence, molecular docking studies on *S. aureus* MurB elucidated its remarkable inhibitory effects of Keracyanin chloride for the first time. Similarly, Naringin molecule showed high binding affinity to *S. aureus* MurB in the presence of low binding free energy. Vitexin has also been identified with promising

inhibitory activity on the *S. aureus* cell wall biosynthesis-related enzyme MurB in the presence of -8.2 kcal/mol binding free energy. Similar inhibitory activity of Vitexin on biofilm formation of *S. aureus* pathogens has also been reported recently [68].

Methanol extract of *T. porrifolius* was found to be cytotoxic in MDA-MB-231 breast cancer cells as well as in Caco-2 colon cancer cells. Cell death was increased in a dose-dependent manner from 5 to 100 µg/ml for 24h and 72h [36]. This study did not cover experimental data on normal cells to compare with cancerous cells. *In vitro*, the cytotoxic effect of ethanol extracts from herbs or roots of *Tragopogon pratensis* L. was shown in leukemic cells, and this effect was found to be correlated with the phenolic content of the extracts. IC₅₀ after 24h incubation value was determined as around 300 µg/ml [38]. Similarly, *T. pratensis* extract up to 100 µg/ml was cytotoxic in HepG2 hepatic cancer cells, but the death of normal cells (HaCaT) was also induced with the extracts [37]. The species with cytotoxic effects shown from *Tragopogon* taxa are *T. porrifolius* and *T. pratensis* only. However, more species (such as *T. graminifolius*, *T. dubius*, *T. oligolepis*, and *T. porrifolius*) were shown to have an antimicrobial effect [11]. To the best of our knowledge, there is no study revealing the anticancer activity of *T. coelestiacus*. The present study suggests that both extracts of *T. coelestiacus* showed a significant cytotoxic effect on AR42J pancreatic cancer cells. However, HeLa cervical and MDA-MB-231 breast cancer cells were found to be more resistant to the cytotoxic effect of methanol and water extract, respectively. Normal epithelial cells were not affected as much as cancerous cells, and their viability was around 80% after both extracts. In the literature, extractions were performed by a range of chemicals, including methanol [69], ethanol [29], hexane [22], chloroform [70], and water [27]. The contents of each extraction method can vary, and this can determine the effect of the plant. In our study, we found that the water extract of *T. coelestiacus* does not have 3 components, including Naringin, Ferulic acid, and Luteloin compared to its methanol extract composition, however, water extract has more amount of 4 components including fumaric acid, quinic acid, gallic acid and epicatechin compared to methanol extract (Table 1). This difference can be suggested not to be deterministic for cytotoxicity in AR42J pancreatic cancer cells as the viability percentages of these cells reduced after both extracts. Nonetheless, the divergence composition may play a specific role in inducing cytotoxicity in HeLa and MDA-MB-231 cells. For instance, Naringin, which is abundant in grapefruit, has been shown to regulate cell cycle arrest [71] and to have a cytotoxic effect in HeLa cells [72]. Epicatechin was found to have anticancer activities on different breast cancer cells, including MDA-MB-231 and MCF7 cells [73]. Our results confirm the previous findings on the specific cytotoxic effect of some metabolites on different cancer cells, however, this is the first study concluding the anticancer effect of *T. coelestiacus* extracts.

Recently, it has been reported that the SRC family FYN kinase is overexpressed in pancreatic cancer cells as a pancreatic cancer-specific marker protein [48]. Experimental findings validated the selective cytotoxic activity of *T. coelestiacus* extracts on AR42J pancreatic cancer cells. The pancreatic cancer-specific FYN kinase was therefore targeted through *in silico* molecular docking analysis using compounds identified from *T. coelestiacus* extracts. Among the tested compounds, naringin, vitexin, keracyanin chloride, luteolin and cyanidin-3-*O*-glucoside were identified with high binding affinity to FYN protein with low binding free energies, -9.8, -9.7, -9.5 and - 9.4 kcal/mol, respectively. Beyond its promising antibacterial

potential, molecular docking data also denoted the remarkable inhibitory potential of the Naringin molecule on FYN kinase protein in the presence of -10.1 kcal/mol binding free energy. Although the anticarcinogenic activity of naringin has been previously reported on various cancers, including breast, lung, stomach, prostate, and colorectal cancers, its effects on pancreatic cancer have been partially reported here for the first time [71]. On the other hand, the detection of vitexin, keracyanin chloride, luteolin, and cyanidin-3-*O*-glycoside compounds with high binding affinity to FYN kinase suggests that their antitumor activities coordinately contribute to the anticarcinogenic activity of Naringin in AR42J pancreatic cancer cell lines.

5. Conclusion

The water and methanol extracts of *T. coelestyriacus* were evaluated for their antibacterial, antifungal, and anticancer properties. The results indicated that the methanol extract of *T. coelestyriacus* has considerable antifungal and antibacterial activity. The bactericidal effects of most of the phenolic compounds contained in *T. coelestyriacus* extracts on microbial cells have been reported in the literature. Phenolic compounds, which are more intense in methanol extract, are thought to affect the antimicrobial activity test result. Furthermore, we can associate the different distributions of phenolic compounds in water and methanol extracts with the variability in the anticancer activity results. Both extracts from *T. coelestyriacus* showed anticancer activity, especially in pancreatic cancer cells, whereas normal cells were more resistant to cell death after treatment with extracts. *In silico* molecular docking studies support experimental data by elucidating inhibition of an antimicrobial protein (MurB involved in cell wall biosynthesis in *S. aureus*) and a cancer-related protein (FYN) specifically increased in pancreatic cancer. Over the last century, the overuse of antibiotics, the resulting antimicrobial resistance, and the severe side effects of chemotherapeutic synthetics and antibiotics have led people to natural compounds, powerful alternatives. Extracts obtained from the edible plant *T. coelestyriacus* have the potential to be used as natural antimicrobial and anticancer therapeutics concerning the phenolic compounds it contains.

Declarations

Author Contributions

T. Unver: Investigation, Methodology, Formal analysis, Writing – original draft, Project administration. U. Uzuner: Software, Investigation, Data curation, Writing - original draft. S. Celik Uzuner: Conceptualization, Investigation, Methodology, Writing - original draft. I. Gurhan: Methodology, Investigation. N.S. Sivri: Methodology, Investigation. Z. Ozdemir: Validation, Data curation, Formal analysis.

Acknowledgments

We would like to thank Assoc. Prof. Ayse Sebnem Erenler, Assist. Prof. Naile Ozturk and Assist. Prof. Hayrettin Tonbul provided technical support for the practical laboratory application. Moreover, we would

like to thank Prof. Dr. Turan Arabacı for helping to identify the plant, and Malatya-Puturge natives, who supported us with their efforts in collecting the studied plant.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Data Availability

The published article incorporates all the generated and analyzed data.

Ethics Approval Not applicable.

Consent to Participate Not applicable.

Consent for Publication Not applicable.

Conflict of Interest The authors affirm that there are no conflicts of interest.

References

1. Petrovska BB (2012) Historical review of medicinal plants' usage. *Pharmacogn. Rev.* 6(11), 1–5. <https://doi.org/10.4103/0973-7847.95849>
2. Dadgostar P (2019) Antimicrobial Resistance: Implications and Costs. *Infect. Drug Resist.* 12:3903–3910. <https://doi.org/10.2147/IDR.S234610>
3. Aslam B, Wang W, Arshad MI, Khurshid M, Muzammil S, Rasool MH, Nisar MA, Alvi RF, Aslam MA, Qamar MU, Salamat MKF, Baloch Z (2018) Antibiotic resistance: a rundown of a global crisis. *Infect. Drug Resist.* 11, 1645–1658. <https://doi.org/10.2147/IDR.S173867>.
4. Simonsen GS (2018) Antimicrobial resistance surveillance in Europe and beyond. *Euro Surveill.* 23(42), 1800560. <https://doi.org/10.2807/1560-7917.ES.2018.23.42.1800560>
5. MacPherson DW, Gushulak BD, Baine WB, Bala S, Gubbins PO, Holtom P, Segarra-Newnham M (2009) Population mobility, globalization, and antimicrobial drug resistance. *Emerg Infect Dis.* 15(11), 1727–1732. <https://doi.org/10.3201/eid1511.090419>
6. GBD (2015) Mortality and Causes of Death Collaborators, Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015 [published correction appears in *Lancet*. 2017 Jan 7;389(10064):e1]. *Lancet* 388(10053):1459-1544
7. Siegel RL, Miller KD, Jemal A (2015) Cancer statistics, 2015. *Cancer J. Clin.* 65(1), 5–29. <https://doi.org/10.3322/caac.21254>
8. Alfarouk KO, Stock CM, Taylor S, Walsh M, Muddathir AK, Verduzco D, Bashir AH, Mohammed OY, Elhassan GO, Harguindey S, Reshkin SJ, Ibrahim ME, Rauch C (2015) Resistance to cancer

- chemotherapy: failure in drug response from ADME to P-gp. *Cancer Cell Int.* 15, 71. <https://doi.org/10.1186/s12935-015-0221-1>.
9. Kumar S, Singh R, Dutta D, Chandel S, Bhattacharya A, Ravichandiran V, Sukla S (2022) In Vitro Anticancer Activity of Methanolic Extract of *Justicia adhatoda* Leaves with Special Emphasis on Human Breast Cancer Cell Line. *Molecules*, 27(23), 8222. <https://doi.org/10.3390/molecules27238222>
 10. Salaroli R, Andreani G, Bernardini C, Zannoni A, La Mantia D, Protti M, Forni M, Mercolini L, Isani G (2022) Anticancer activity of an *Artemisia annua* L. hydroalcoholic extract on canine osteosarcoma cell lines. *Res. Vet. Sci.* 152, 476–484. <https://doi.org/10.1016/j.rvsc.2022.09.012>
 11. Abdalla MA and Zidorn C (2020) The genus *Tragopogon* (Asteraceae): A review of its traditional uses, phytochemistry, and pharmacological properties. *J. Ethnopharmacol.* 250, 112466. <https://doi.org/10.1016/j.jep.2019.112466>.
 12. Graham JG, Quinn ML, Fabricant DS, Farnsworth NR (2000) Plants used against cancer - an extension of the work of Jonathan Hartwell. *J. Ethnopharmacol.* 73(3), 347–377. [https://doi.org/10.1016/s0378-8741\(00\)00341-x](https://doi.org/10.1016/s0378-8741(00)00341-x).
 13. Sarac N (2015) Antioxidant, mutagenic, and antimutagenic activities of *Tragopogon longirostis* var. *longirostis*, an edible wild plant in Turkey. *Indian J. Pharmacol.* 47(4), 414–418. <https://doi.org/10.4103/0253-7613.161267>
 14. Tabaraki R, Nateghi A, Ahmady-Asbchin S (2013) In vitro assessment of antioxidant and antibacterial activities of six edible plants from Iran. *J. Acupunct. Meridian Stud.* 6(3), 159–162. <https://doi.org/10.1016/j.jams.2013.01.016>
 15. Bahadır Acıkara Ö, Saltan HG, Çoban T (2013) Evaluation of antioxidant properties of some *Tragopogon* species growing in Turkey. *Turk J. Pharm. Sci.* 10:3,377-384.
 16. Bell CD, Mavrodiev EV, Soltis PS, Calaminus AK, Albach DC, Cellinese N, Garcia-Jacas N, Soltis DE (2012) Rapid diversification of *Tragopogon* and ecological associates in Eurasia. *J. Evol. Biol.* 25(12), 2470–2480. <https://doi.org/10.1111/j.1420-9101.2012.02616.x>
 17. Baytop T (2000) Anadolu dağlarında 50 yıl: Bir bitki avcısının gözlemleri, Nobel Tıp kitabevi, İstanbul.
 18. Gültepe M, Coşkunçelebi K, Makbul S, Güzel ME (2021) Contribution to the taxonomy of little known *Tragopogon* species endemic to Turkey. *Nord. J. Bot.* 39:1-7. <https://doi.org/10.1111/njb.03242>.
 19. Güner A, Akyıldırım B, Alkayış MF, Çingay B, Kanoğlu SS, Özkan AM, Öztekin M, Tuğ GN (2012) Türkçe bitki adları. In: Güner, A., Aslan, S., Ekim, T., Vural, M., Babaç, M.T. (Eds.) A checklist of the Flora of Turkey (Vascular Plants). Nezahat Gökyiğit Botanik Bahçesi Yayınları Flora Dizisi 1 İstanbul.
 20. Richardson IBK (1976) *Flora Europaea*. Cambridge:University Press Vol. 4
 21. Altundag E and Ozturk M (2011) Ethnomedicinal studies on the plant resources of East Anatolia. Turkey, *Procedia Soc. Behav. Sei.* 19:756-777.
 22. Ugur A, Sarac N, Ceylan O, Emin Duru M, Okmen G, Varol O (2010) Chemical composition of endemic *Tragopogon oligolepis* and studies on the antimicrobial activity against multi-antibiotic-resistant bacteria. *Acta Hort.* (Wagening.) 853, 299–306. <https://doi.org/10.17660/ActaHortic.2010.853.35>.

23. Lusa MG, Martucci MEP, Loeuille BFP, Gobbo-Neto L, Appezzato-da-Gloria B, Da Costa FB (2016) Characterization and evolution of secondary metabolites in Brazilian Vernoniaeae (Asteraceae) assessed by LC-MS fingerprinting. Bot. J. Linn. Soc. 182, 594–611. <https://doi.org/10.1111/boj.12480>
24. Granica S, Piwowarski JP, Randazzo A, Schneider P, Żyżyńska-Granica B, Zidorn C (2015) Novel stilbenoids, including cannabispiradienone glycosides, from *Tragopogon tommasinii* (Asteraceae, Cichorieae) and their potential anti-inflammatory activity. Phytochemistry 117, 254–266. <https://doi.org/10.1016/j.phytochem.2015.06.018>.
25. Sareedenchai V, Ganzera M, Ellmerer EP, Lohwasser U, Zidorn C (2009) Phenolic compounds from *Tragopogon porrifolius* L. Biochem. Syst. Ecol. 37, 234–236. <https://doi.org/10.1016/j.bse.2009.03.004>.
26. Miyase T, Kohsaka H, Ueno A (1992) Tragopogonosides A-I, oleanane saponins from *Tragopogon pratensis*. Phytochemistry 31, 2087–2091. [https://doi.org/10.1016/0031-9422\(92\)80368-O](https://doi.org/10.1016/0031-9422(92)80368-O).
27. Uysal S, Senkardes I, Mollica A, Zengin G, Bulut G, Dogan A, Glamočlija J, Soković M, Lobine D, Mahomoodally MF (2019) Biologically active compounds from two members of the Asteraceae family: *Tragopogon dubius* Scop. and *Tussilago farfara* L. J. Biomol. Struct. Dyn. 37(12), 3269–3281. <https://doi.org/10.1080/07391102.2018.1506361>.
28. Bayrami Z, Hajiaghaee R, Khalighi-Sigaroodi F, Rahimi R, Farzaei MH, Hodjat M, Baeri M, Rahimifard M, Navaei-Nigjeh M, Abdollahi M (2018) Bio-guided fractionation and isolation of active component from *Tragopogon graminifolius* based on its wound healing property. J Ethnopharmacol. 226, 48–55. <https://doi.org/10.1016/j.jep.2018.08.002>
29. Farzaei MH, Khanavi M, Moghaddam G, Dolatshahi F, Rahimi R, Shams-Ardekani MR, Amin G, Hajimahmoodi M (2014) Standardization of *Tragopogon graminifolius* DC. extract based on phenolic compounds and antioxidant activity. J. Chem. 2014:425965
30. Smolarz H, Krzaczek T (1988) Phytochemical studies of the herb *Tragopogon orientalis* L. (Asteraceae). Components of the methanol extract. Acta Soc. Bot. Pol. 57, 93–105. <https://doi.org/10.5586/asbp.1988.009>.
31. Martínez-González MÁ, Hershey MS, Zazpe I, Trichopoulou A (2017) Transferability of the Mediterranean diet to non-Mediterranean countries. What is and what is not the Mediterranean diet. Nutrients, 9(11), 1226. <https://doi.org/10.3390/nu9111226>.
32. Kim SJ, Cho AR, Han J (2013) Antioxidant and antimicrobial activities of leafy green vegetable extracts and their applications to meat product preservation. Food control, 29(1), 112–120. <https://doi.org/10.1016/j.foodcont.2012.05.060>
33. Bayram M and Topuz S (2023) Madımaktan (*Polygonum cognatum* Meissn.) Yanıt Yüzey Yöntemi Kullanılarak Fenolik Bileşik Ekstraksiyonunun Optimizasyonu. Gıda 48(1):118–129.
34. Gündüz M, Çiçek ŞK, Topuz S (2023) Extraction and Optimization of Phenolic Compounds from Butterbur Plant (*Petasites hybridus*) by Ultrasound-Assisted Extraction and Determination of

- Antioxidant and Antimicrobial Activity of Butterbur Extracts. J. Appl. Res. Med. Aromat. Plants 100491. <https://doi.org/10.1016/j.jarmap.2023.100491>
35. Moromete C, Badea V, Arcuş M, Bucur L, Roşoiu B (2016) Study of the antibacterial and antifungal effect of the *Tragopogon dubius* Scop. (Asteraceae) aqueous extracts. Studia Universitatis Vasile Goldis Seria Stiintele Vietii (Life Sciences Series) 26, 473–478.
36. Tenkerian C, El-Sibai M, Daher CF, Mroueh M (2015) Hepatoprotective, antioxidant, and anticancer effects of the *Tragopogon porrifolius* methanolic extract. Evid. Based Complement Altern. Med. 10. <https://doi.org/10.1155/2015/161720>. 2015,
37. Kucekova Z, Mlcek J, Humpolicek P, Rop O (2013) Edible flowers—Antioxidant activity and impact on cell viability. Open Life Sci. 8(10), 1023-1031. <https://doi.org/10.2478/s11535-013-0212-y>.
38. Wegiera M, Smolarz HD, Jedruch M, Korczak M, Koproń K (2012) Cytotoxic effect of some medicinal plants from Asteraceae family on J-45.01 leukemic cell line—pilot study. Acta Pol. Pharm. 69(2), 263–268.
39. Matthews VA, Kupicha FK, Parris BS (1975) *Flora of Turkey and the East Aegean Islands*, (Davis PH, Ed.; Vol. 5). Edinburgh University Press.
40. Radulović N, Stankov-Jovanović V, Stojanović G, Šmelcerović A, Spiteller M, Asakawa Y (2007) Screening of in vitro antimicrobial and antioxidant activity of nine Hypericum species from the Balkans. Food Chem. 103(1), 15-21. <https://doi.org/10.1016/j.foodchem.2006.05.062>
41. Unver T, Erenler AS, Bingul M, Boga M (2023) Comparative Analysis of Antioxidant, Anticholinesterase, and Antibacterial Activity of Microbial Chondroitin Sulfate and Commercial Chondroitin Sulfate. Chem. Biodivers. e202300924. <https://doi.org/10.1002/cbdv.202300924>.
42. Cakmak U, Oz-Tuncay F, Basoglu-Ozdemir S, Ayazoglu-Demir E, Demir I, Colak A, Celik-Uzuner S, Erdem SS, Yildirim N (2021) Synthesis of hydrazine containing piperazine or benzimidazole derivatives and their potential as α -amylase inhibitors by molecular docking, inhibition kinetics and in vitro cytotoxicity activity studies. Med. Chem. Res. 30, 1886–1904. <https://doi.org/10.1007/s00044-021-02785-8>
43. Uzuner SÇ, Birinci E, Tetikoğlu S, Birinci C, Kolaylı S (2021) Distinct epigenetic reprogramming, mitochondrial patterns, cellular morphology, and cytotoxicity after bee venom treatment. Recent Pat. Anticancer Drug Discov. 16(3), 377-392. <https://doi.org/10.2174/1574892816666210422125058>
44. Matsuo M, Kurokawa K, Nishida S, Li Y, Takimura H, Kaito C, Fukuhara N, Maki H, Miura K, Murakami K, Sekimizu K (2003) Isolation and mutation site determination of the temperature-sensitive murB mutants of Staphylococcus aureus. FEMS Microbiol. Lett. 222(1), 107–113. [https://doi.org/10.1016/S0378-1097\(03\)00260-X](https://doi.org/10.1016/S0378-1097(03)00260-X)
45. Nishida S, Kurokawa K, Matsuo M, Sakamoto K, Ueno K, Kita K, Sekimizu K (2006) Identification and characterization of amino acid residues essential for the active site of UDP-N-acetylenolpyruvylglucosamine reductase (MurB) from Staphylococcus aureus. The J. Biol. Chem. 281(3), 1714–1724. <https://doi.org/10.1074/jbc.M509277200>

46. Sen B and Johnson FM (2011) Regulation of SRC family kinases in human cancers. J. Signal Transduct. 2011, 865819. <https://doi.org/10.1155/2011/865819>
47. Posadas EM, Al-Ahmadie H, Robinson VL, Jagadeeswaran R, Otto K, Kasza KE, Tretiakov M, Siddiqui J, Pienta KJ, Stadler WM, Rinker-Schaeffer C, Salgia R (2009) FYN is overexpressed in human prostate cancer. BJU Int. 103(2), 171–177. <https://doi.org/10.1111/j.1464-410X.2008.08009.x>
48. Peng S and Fu Y (2023) FYN: emerging biological roles and potential therapeutic targets in cancer. J. Transl. Med. 21(1), 84. <https://doi.org/10.1186/s12967-023-03930-0>
49. Eberhardt J, Santos-Martins D, Tillack AF, Forli S (2021) AutoDock Vina 1.2.0: New Docking Methods, Expanded Force Field, and Python Bindings. J. Chem. Inf. Model. 61(8), 3891–3898. <https://doi.org/10.1021/acs.jcim.1c00203>
50. Olsson MH, Søndergaard CR, Rostkowski M, Jensen JH (2011) PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions. J. Chem. Theory Comput. 7(2), 525–537. <https://doi.org/10.1021/ct100578z>
51. O'Boyle NM, Banck M, James CA, Morley C, Vandermeersch T, Hutchison GR (2011) Open Babel: An open chemical toolbox. J. Cheminform. 3, 33. <https://doi.org/10.1186/1758-2946-3-33>
52. Farzaei MH, Rahimi R, Attar F, Siavoshi F, Saniee P, Hajimahmoodi M, Mirnezami T, Khanavi M (2014) Chemical composition, antioxidant and antimicrobial activity of essential oil and extracts of *Tragopogon graminifolius*, a medicinal herb from Iran. Nat. Prod. Commun. 9:121–124.
53. Helander LM, Alakomi HL, Latva-Kala K (1998) Characterization of the action of selected essential oil components on Gram-negative bacteria. Journal of Agricultural and Food. Chemistry 46(9):3590e3595. <https://doi.org/10.1021/jf980154m>
54. Lin YT, Labbe RG, Shetty K (2004) Inhibition of *Listeria monocytogenes* in fish and meat systems by use of oregano and cranberry phytochemical synergies. Appl. Environ. Microbiol. 70(9), 5672–5678. <https://doi.org/10.1128/AEM.70.9.5672-5678.2004>
55. Walsh C (2003) Antibiotics: actions, origins, resistance. ASM Press, Washington.
56. Gelssman TA (1963) Flavonoid compounds tannins lignins and related compounds. In Florkin, M., & Stutz, E. H. (Eds.). *Pyrrole pigments, isoprenoid compounds and phenolic plant constituents*, Elsevier, New York, 9:265
57. Brooks GF, Butel JS, Morse SA (1998) Jawetz, Melnick, and Adelberg's Medical Microbiology (21st ed.). Stamford: Appleton & Lange
58. Pelczar MJ, Chan EDS, Krieg NR (1988) The control of microorganisms by physical agents. Microbiology, McGraw-Hill International, New York.
59. Wu Y, Liang S, Zhang M, Wang Z, Wang Z, Ren X (2020) The Effect of Chlorogenic Acid on *Bacillus subtilis* Based on Metabolomics. Molecules. 25(18), 4038. <https://doi.org/10.3390/molecules25184038>
60. Lou Z, Wang H, Zhu S, Ma C, Wang Z (2011) Antibacterial activity and mechanism of action of chlorogenic acid. J. Food Sci. 76(6), M398–M403. <https://doi.org/10.1111/j.1750-3841.2011.02213.x>

61. Li L, Zhou P, Wang Y, Pan Y, Chen M, Tian Y, Zhou H, Yang B, Meng H, Zheng J (2022) Antimicrobial activity of cyanidin-3-O-glucoside-lauric acid ester against *Staphylococcus aureus* and *Escherichia coli*. *Food Chem.* 383, 132410. <https://doi.org/10.1016/j.foodchem.2022.132410>
62. Kernou ON, Azzouz Z, Madani K, Rijo P (2023) Application of Rosmarinic Acid with Its Derivatives in the Treatment of Microbial Pathogens. *Molecules*, 28(10), 4243. <https://doi.org/10.3390/molecules28104243>
63. Ivanov M, Kostić M, Stojković D, Soković M (2022) Rosmarinic acid–Modes of antimicrobial and antibiofilm activities of common plant polyphenol. *S. Afr. J. Bot.* 146, 521–527. <https://doi.org/10.1016/j.sajb.2021.11.050>
64. Pinheiro PG, Santiago GMP, da Silva FEF, de Araújo ACJ, de Oliveira CRT, Freitas PR, Rocha JE, Neto JBA, da Silva MMC, Tintino SR, Siyadatpanah A, Norouzi R, Dashti S, Wilairatana P, Coutinho HDM, da Costa JGM (2022) Ferulic acid derivatives inhibiting *Staphylococcus aureus* tetK and MsrA efflux pumps. *Biotechnol. Rep. (Amst)* 34, e00717. <https://doi.org/10.1016/j.btre.2022.e00717>
65. Qian W, Liu M, Fu Y, Zhang J, Liu W, Li J, Li X, Li Y, Wang T (2020) Antimicrobial mechanism of luteolin against *Staphylococcus aureus* and *Listeria monocytogenes* and its antibiofilm properties. *Microb Pathog.* 142, 104056. Advance online publication. <https://doi.org/10.1016/j.micpath.2020.104056>
66. Céliz G, Daz M, Audisio MC (2011) Antibacterial activity of naringin derivatives against pathogenic strains. *J. Appl. Microbiol.* 111(3), 731–738. <https://doi.org/10.1111/j.1365-2672.2011.05070.x>
67. Jaradat N, Shawarab N, Hussein F, Al-Masri M, Warad I, Khasati A, Shehadeh M, Qneibi M, Hussein AMA, Makhamreh S (2018) Antibacterial and antioxidant screening of semi-synthetic naringin based hydrazone and oxime derivatives. *Jundishapur J. Microbiol.* 11(6). <https://doi.org/10.5812/jjm.65496>
68. Das MC, Samaddar S, Jawed JJ, Ghosh C, Acharjee S, Sandhu P, Das A, Daware AV, De UC, Majumdar S, Das Gupta SK, Akhter Y, Bhattacharjee S (2022) Vitexin alters *Staphylococcus aureus* surface hydrophobicity to obstruct biofilm formation. *Microbiol. Res.* 263, 127126. <https://doi.org/10.1016/j.micres.2022.127126>
69. Warashina T, Miyase T, Ueno A (1991) Novel acylated saponins from *Tragopogon porrifolius* L. Isolation and the structures of tragopogonsaponins AR. *Chem. Pharm. Bull.* 39(2): 388-396. <https://doi.org/10.1248/cpb.39.388>
70. Hariri EH, El Asmar M, Demirdjian SA, Daher CF, Mroueh MA (2013) Anti-inflammatory effect of the methanol, ethyl acetate and chloroform extracts of *Tragopogon porrifolius* aerial parts. *Planta Med.* 79(13):PN67. <https://doi.org/10.1055/s-0033-1352410>
71. Stabrauskiene J, Kopustinskiene DM, Lazauskas R, Bernatoniene J (2022) Naringin and Naringenin: Their Mechanisms of Action and the Potential Anticancer Activities. *Biomedicines* 10(7), 1686. <https://doi.org/10.3390/biomedicines10071686>
72. Martínez-Rodríguez OP, González-Torres A, Álvarez-Salas LM, Hernández-Sánchez H, García-Pérez BE, Thompson-Bonilla MDR, Jaramillo-Flores ME (2020) Effect of naringenin and its combination with

cisplatin in cell death, proliferation and invasion of cervical cancer spheroids. RSC Adv. 11(1), 129–141. <https://doi.org/10.1039/d0ra07309a>

73. Pereyra-Vergara F, Olivares-Corichi IM, Perez-Ruiz AG, Luna-Arias JP, García-Sánchez JR (2020) Apoptosis Induced by (-)-Epicatechin in Human Breast Cancer Cells is Mediated by Reactive Oxygen Species. Molecules 25(5), 1020. <https://doi.org/10.3390/molecules25051020>

Figures




Figure 1

Petri dish pictures of the antibacterial activity assay of the *T. coelestiacus* methanol extract with various concentrations against 5 bacterial strains: (1) *S. aureus*, (2) *E. aerogenes*, (3) *E. coli*, (4) *P. aeruginosa*, and (5) *K. pneumoniae*. (A) 106.6 (B) 53.3, (C) 26.6, (D) 13.3, (E) 6.6, (F) 3.3, (G) 1.67, (H) 0.83, (I) 0.42, (J) 0.21, (K) 0.1 mg/mL methanol extract of *T. coelestiacus*, (L) Control




Figure 2

Petri dish pictures of the antifungal activity assay of the *T. coelestiacus* methanol extract with different concentrations against 4 yeast strains: (1) *C. albicans*, (2) *C. tropicalis*, (3) *C. krusei*, and (4) *C. parapsilosis*. (A) 106.6 (B) 53.3, (C) 26.6, (D) 13.3, (E) 6.6, (F) 3.3, (G) 1.67, (H) 0.83, (I) 0.42, (J) 0.21, (K) 0.1 mg/mL methanol extract of *T. coelestiacus*, (L) Control




Figure 3

Microplate picture of the methanol extract of *T. coelestis* antibacterial and antifungal activity test against *S. aureus* (A), *E. aerogenes* (B), *E. coli* (C), *P. aeruginosa* (D), *K. pneumonia* (E), *C. albicans* (F), *C. tropicalis* (G), *C. krusei* (H), and *C. parapsilosis* (I). The methanol extract concentration in wells A8 (0.83 mg/mL), B2 (53.3 mg/mL), C2 (53.3 mg/mL), D7 (1.67 mg/mL), E7 (1.67 mg/mL), F2 (53.3 mg/mL), G1 (106.6 mg/mL), H8 (0.83 mg/mL), and I3 (26.6 mg/mL) with no color change were determined as MIC values against *S. aureus*, *E. aerogenes*, *E. coli*, *P. aeruginosa*, *K. pneumonia*, *C. albicans*, *C. tropicalis*, *C. krusei*, and *C. parapsilosis*, respectively. 11. negative control, 12. positive control





Figure 4

Cell viabilities (%) in AR42J (pancreatic cancer) cells (A), HeLa (cervical cancer) cells (B), and MBD-MB-231 (breast cancer) (C) cells after the treatments for 24h, 48h, or 72h with *T. colesyi* extracts obtained either by methanol (up panels) or by water (down panels) at different concentrations. Summary of statistical comparisons is given as tables under each graph of related cell type. $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***) and $p < 0.0001$ (****).



Cell viabilities (%) in ARPE-19 (retinal epithelial) cells (A) and MCF10A (breast epithelial) cells (B) after the treatments for 24h, 48h or 72h with *T. coelestriacus* extracts obtained either by methanol (up panels) or by water (down panels) at different concentrations. Related statistics are given for each graph. $p<0.05$ (*), $p<0.01$ (**), $p<0.001$ (***) and $p<0.0001$ (****).



Page 30/33




Figure 7

Molecular docking pose of naringin-MurB (1HSK) and 2D depiction of their biochemical interaction potential.




Figure 8

Molecular docking pose of naringin-FYN (2DQ7) and 2D depiction of their biochemical interaction potential.




Figure 9

Molecular docking pose of keracyanin chloride-FYN (2DQ7) and 2D depiction of their biochemical interaction potential.




Figure 10

Polar regions of phenolic molecules in water and methanol extracts.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.tif](#)