

Primula meishanensis (Primulaceae), a new species from Sichuan, China

Tingyu Li^{1,2}, Xinyu Chen², Bo Li³, Donglai Hua⁴, Can Luo², Huixian Luo², Yun Liang⁵, Jieli Yue⁵, Xiaodan Xi⁵, Ke Huang², Zhixi Fu^{1,2,6}

1 Ministry of Education, Key Laboratory of Land Resources Evaluation and Monitoring in Southwest (Sichuan Normal University), Chengdu 610101, China

2 College of Life Sciences, Sichuan Normal University, Chengdu 610101, China

3 Sichuan Environmental Monitoring Center, Chengdu 610091, China

4 College of Life Sciences, Mianyang Teachers' College, Mianyang 621000, China

5 Meishan Eco-environmental Monitoring Center Station of Sichuan Province, Meishan, 620010, China

6 Sustainable Development Research Center of Resources and Environment of Western Sichuan, Sichuan Normal University, Chengdu 610101, China

Corresponding authors: Ke Huang (542176689@qq.com); Zhixi Fu (fuzx2017@sicnu.edu.cn)

Abstract

Primula meishanensis K.Huang & Z.X.Fu, **sp. nov.**, a new species of Primulaceae from Meishan City, Sichuan Province, China, is described and illustrated. The morphological data and phylogenetic analysis, based on the complete chloroplast genome, suggest that *Primula meishanensis* is a separate species closely related to *Primula dejuniana*. The complete chloroplast genome of *Primula meishanensis* was 152,175 bp and the complete chloroplast genome of *Primula dejuniana* was 151,988 bp. The new species differs from the latter by the solitary scape, the length of petiole, acute leaf blade apex and pin flower. The distribution map, morphological comparison of related species and conservation status of the new species are also provided.

Key words: Morphological characters, new species, *Primula* sect. *Petiolares*, taxonomy



Academic editor: Bing Liu

Received: 9 May 2024

Accepted: 20 September 2024

Published: 23 October 2024

Citation: Li T, Chen X, Li B, Hua D, Luo C, Luo H, Liang Y, Yue J, Xi X, Huang K, Fu Z (2024) *Primula meishanensis* (Primulaceae), a new species from Sichuan, China. PhytoKeys 248: 73–90. <https://doi.org/10.3897/phytokeys.248.127117>

Copyright: © Tingyu Li et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The genus *Primula* L. is one of the most diverse member within the Primulaceae. It consists of 38 sections and more than 500 species worldwide (Hu and Kelso 1996; Richards 2003). It is widely distributed throughout temperate and cold regions of Asia and Europe, alongside the tropical mountains of the Northern Hemisphere. The genus is composed of herbaceous plants with a basal rosette of leaves, flowers on top of a naked scape, gathered in lateral or perpendicular to the axis umbels. China hosts over 340 species, with particular biodiversity hotspots of *Primula* found in south-western China (Hu 1990; Hu 1998; Richards 2003; Ju et al. 2023; Li et al. 2023). In recent years, several new species of *Primula* from China were reported, for example, *Primula lihengiana* C. M. Hu & R. Li (Li and Hu 2009), *Primula wawushanica* G. Hao, C.M. Hu & Y. Xu, *Primula undulifolia* G. Hao, C. M. Hu & Y. Xu, *Primula pengzhouensis* C.M. Hu, G. Hao & Y. Xu., *Primula surculosa* Y. Xu & G. Hao (Xu et al. 2016a, b, 2017,

2022), *Primula dongchuanensis* Z.K. Wu & Yuan Huang (Wu et al. 2019), *Primula dujiangyanensis* W. B. Ju, Bo Xu & X. F. Gao (Ju et al. 2021), *Primula longipilosa* Ze H. Wang & H. Peng (Wang et al. 2022), *Primula wolongensis* W.B.Ju, Bo Xu & X.F.Gao (Li et al. 2023), *Primula sugongii* J.D.Ya, Bin Yang & Y.H. Tan (Yang et al. 2023) and *Primula xilingensis* K. Huang & Z.X. Fu (Luo et al. 2023).

The *Primula* sect. *Petiolares* Pax comprises more than 60 currently recognised species (Hu 1994; Hu and Kelso 1996; Hu and Geng 2003; Rankin 2010; Wei et al. 2022; Zhang et al. 2023) and are well represented in the region of Himalaya-Hengduan Mountains, with only a few members extending into Kashmir, central China and some other regions (Hu and Kelso 1996). One of the most important diagnostic characters of this section is a globose capsule which does not open by valves, but by crumbling at the membrane apex (Hu 1994).

During the botanical expedition of Zhongyan temple in Qingshen County, Meishan City, Sichuan Province from 2022 to 2024, a population of *Primula* was discovered, photographed and collected. Based on the photographs of the flowering taxon, it appears to be closely related to *Primula dejuniana* G. Hao, C.M. Hu & Y. Xu (Xu et al. 2014) at first sight. After consultation with relevant literature and conducting morphological examination of closely-related taxa, it was determined that it represents an unreported taxon from *P.* sect. *Petiolares*. The new species could be differentiated from other members of the section by the following combination: scape solitary, the length of petiole 2.5–5 cm, a terminal umbel of (1) 2–3 (4) flowers, leaf blade smooth, oblanceolate, 19.5–25 × 2.5–5 cm, apex acute, corolla pastel violet, 5–6 lobes spreading and stamens inserted at middle of corolla tube, corolla tubes 11.0–12.0 mm long, ca. 2 mm in diam., style 4.0–7.0 mm above base of corolla tube, stamens reaching the corolla tube mouth, 7.0–12.0 mm above base of corolla tube.

In addition, in this study, the molecular data of complete chloroplast genomic data were collected and used to identify its relationship with *Primula dejuniana*. In recent years, many new species have been jointly supported by genetic and morphological data, including *Primula sunhangii* T. Deng, D. G. Zhang & Jiao Sun (Sun et al. 2020), *Aster quanzhouensis* M.Tang, G.J.Yan & W.P.Li (Xiao et al. 2022), *Primula undulifolia* G. Hao, C. M. Hu & Y. Xu (Xu et al. 2016), *Aster yaoshanensis* K. Qin, Z.X. Fu and P. Li. (Zheng et al. 2024) and *Saussurea talungensis* S.K.Ghimire & H.K.Rana (Rana et al. 2021).

In this study, we provide a detailed description of this new species, based on observations of living plants in the field and specimens in the herbarium.

Material and methods

Morphological analysis

The observation and collection of both herbarium and living materials of the new species from Qingshen County, Meishan City, Sichuan Province, were conducted in December 2022, January 2023, January 2024 and March 2024. We conducted a morphological comparison using taxonomic literature of closely-related species, i.e. *Primula dejuniana* (description of reference Xu et al. (2014)) and *Primula davidii* Franch. (holotype: *David, A. 1870* (P, image!)), as well as the images of specimens from the Global Plants JSTOR database (<https://plants.jstor.org>), i.e. *Primula epilosa* Craib (holotype: *Sun231* (KUN, image!)) were

consulted. Morphological description and measurements of *P. meishanensis* were based on living plants. The taxonomic description followed the terminology used by Beentje (2016). The holotype voucher specimens were stored at the Herbarium of Sichuan Normal University (SCNU). The conservation status of the new species was assessed following the guidelines of the IUCN Red List Categories and Criteria (IUCN 2024).

DNA extraction and sequencing

Total genomic DNA was obtained using the CTAB method (Doyle and Doyle 1987). We follow the Illumina DNA Library Construction Guide to complete paired-end DNA library construction (Allen et al. 2006). The complete chloroplast genome was sequenced on the Illumina HiSeq XTen platform (San Diego, CA, USA). In this study, SPAdes v.3.10.1 software was used to assemble high-quality data by default parameters (Bankevich et al. 2012). We further used CPGView (<http://47.96.249.172:16085/cpgview/view>) to improve annotation, visualise the structure of the cp genome and identify gene structures including cis-splicing and trans-splicing (Liu et al. 2023). The genome sequence of *Primula dejuniana* and *P. meishanensis* have been deposited in GenBank (accession numbers: [PQ213817](#) and [PQ213816](#)).

Phylogenetic analysis

The phylogenetic analysis of the complete chloroplast genomic dataset of 18 species was performed using the Maximum Likelihood (ML) method implemented in RAxML. The species *Maesa montana* A. DC. ([KU569490](#)) and *Ardisia polysticta* Migo ([KC465962](#)) were selected as outgroups. These chloroplast genome sequences were imported into MAFFT v.7.520 software (Katoh and Standley 2013) for multiple comparisons and the phylogenetic tree was constructed using CIPRES (<https://www.phylo.org/>), with the ML method, based on the GTRGAMMA model and the bootstrap was set to 1,000 (Stamatakis et al. 2008) (Table 2) (Fig. 5).

Result and discussion

Taxonomic treatment

***Primula meishanensis* K.Huang & Z.X.Fu, sp. nov.**

[urn:lsid:ipni.org:names:77350705-1](https://nomenclature.ipni.org/names/77350705-1)

Type. CHINA, Sichuan Province, Meishan City, Qingshen County, Zhongyan temple, grows on moist rock surfaces amidst moss under the forest, at elevations of approximately 417 m; 29°45'47.39"N, 103°50'44.017"E; 26 December 2023 (fl.), *Ke Huang & Zhixi Fu 8200* (holotype SCNU!) (Figs 1–3).

Diagnosis. The new species is morphologically similar to *Primula dejuniana* in having ciliate, sharply and remotely dentate leaf blade margin, while it can be easily recognised by the following combination of characters: scape solitary, the length of petiole 2.5–5 cm, a terminal umbel of (1) 2–3 (4) flowers, leaf blade smooth, oblanceolate, 19.5–25 × 2.5–5 cm, apex acute, corolla pastel




Figure 1. *Primula meishanensis* sp. nov. **A** plant and roots **B** roots **C** petioles **D**, **E** leaves **F** pin flower **G** thrum flower **H** scape and bract (Photos by XC).

violet, 5–6 lobes spreading and stamens inserted at middle of corolla tube, corolla tubes 11.0–12.0 mm long, ca. 2 mm in diam., style 4.0–7.0 mm above base of corolla tube, stamens reaching the corolla tube mouth, 7.0–12.0 mm above base of corolla tube. (Figs 1–3).

Description. Perennial herbs, 12.0–20.0 cm tall. **Roots** numerous, fibrous, without hairs. **Leaves** pilose, forming a spreading rosette, each rosette with only 2–4 leaves of previous year at flowering time; resting bud of rosette clothed by a few small paleaceous scales, basal bud scales ovate to ovate-oblong, ciliate, rose red, apex acute; petiole 2.5–5 cm long, narrowly winged and densely covered with multicellular hairs; leaf blade oblanceolate, smooth, 19.5–25 × 2.5–5 cm, cuneate at base, abaxial surface densely along mid-vein, sparser on lateral veins covered with multicellular hairs; margin ciliate, sharply and remotely dentate; apex acute; mid-vein dull yellow in fresh state, turning brownish when dry; veins impressed adaxially, prominently raised and subalveolate abaxially. **Scape** solitary, 7.2–11.8 cm long, pilose, carrying a terminal umbel of (1) 2–3 (4) flowers, dull yellow in fresh state, turning brownish when dry at base. **Bracts** lanceolate, 4–5.5 mm long, minutely ciliate. **Pedice** 0.4–1.4 cm long, shorter than leaf blade, pilose. **Flowers** distylous. **Calyx** campanulate, 8–9 mm long, parted to 1/2 of its length or slightly below; lobes linear lanceolate to lanceolate, apex acute. **Corolla** pastel violet, annulate, 5–6 lobes spreading, lobes broadly elliptic, 8.0–10.0 mm long, emarginate, densely yellow farinose abaxially, smooth adaxially. **Pin flower:** corolla tubes 9.0–10.0 mm long, stamens inserted at middle of corolla tube, 9.5–10.5 mm long, style slightly exceeding the corolla tube mouth. **Thrum flower:** corolla tubes 11.0–12.0 mm long, ca. 2 mm in diam., 2 times as long as the calyx, style 4.0–7.0 mm above base of corolla tube, stamens reaching the corolla tube mouth, 7.0–12.0 mm above base of corolla tube (Fig. 1G).

Phenology. The flowering period is from December to February and the fruiting period is March to May.

Etymology. The epithet “meishanensis” is derived from Meishan City, located in Sichuan Province, China.

Distribution and habitat. *P. meishanensis* is currently known from its type locality in Zhongyan temple, Ruifeng Town, Qingshen County and roadside of Panjiaozui, forest Hongya Forest Farm, Hongya County, Meishan City, Sichuan Province, China (Fig. 4). This new species probably exists in other localities. It grows on moist rock surfaces amidst moss under the forest, at elevations of approximately ca. 400–1000 m (Fig. 2).

Additional specimens examined. CHINA, Sichuan, Meishan City, Qingshen County, 29°45'47.39"N, 103°50'44.017"E, 26 December 2023 (fl.), Ke Huang & Zhixi Fu 8201 (SCNU!); Hongya County, 29°49'57.23"N, 103°09'50.23"E, 28 May 2015 (fl.), Ya J.D. & Hu X.J. 15CS11038 (KUN!)

Conservation status. Data Deficient (DD). Currently, two populations with more than 100 individuals have been found in the Qingshen and Hongya Counties. The population of *P. meishanensis* inhabits moist rocks. Given its currently limited occurrence near the temple, there is a significant likelihood that the taxon may also be found in other localities. Until we have fully investigated the situation, it would be suggested to assess the conservation status of the species as Data Deficient (DD) following the IUCN Red List Criteria (IUCN 2024).




Figure 2. *Primula meishanensis* sp. nov. **A–C** habitat **D–H** flowers **I, J** fruit (Photos **A, B, D, E** by ZF and **C, F–J** by KH).




Figure 3. Holotype image of *Primula meishanensis* K.Huang & Z.X.Fu, sp. nov.




Figure 4. Location of the population of *Primula meishanensis* in Qingshen County and Hongya County, Meishan City and Sichuan Province (red star).

Relationship with related species. Critical examination of collected specimens, comparison with type material of allied taxa and relevant taxonomic literature revealed that *P. meishanensis* is a new member of the *P.* sect. *Petiolares*. Morphologically, *P. meishanensis* shares certain similarities with *P. dejuniana*. However, *P. meishanensis* differs from *P. dejuniana* in featuring the leaf apex acute (vs. the leaf apex acute, but with a small point at tip), corolla pastel violet (vs. the corolla pale rose-purple), scape solitary, 7.2–11.8 cm long, carrying a terminal umbel of (1) 2–3 (4) flowers (vs. the scape usually one per rosette, 8–12 (18) cm long, carrying a terminal umbel of 2–6 flowers), the length of petiole 2.5–5 cm (vs. 1–3 cm), basal bud scales rose red (vs. flesh pink) and flowering period is from December to February (vs. from early February to early March). The number of chloroplast genes etc. is different in the two species (Table 2, Fig. 6, Suppl. material 1). To some extent, *P. meishanensis* also resembles to *P. davidii* as a perennial herb with calyx and corolla. However, it differs from *P. davidii* in leaf blade margin (sharply vs. erose-dentate) and apex (acute vs. rounded). The species of *P. meishanensis* bears similarities to *P. epilosa*, yet the former is readily distinguished by its leaf apex (acute vs. rounded), corolla (pastel violet vs. rose-purple with a yellow eye) and altitude (400–420 m vs. 2000–2900 m). Further morphological comparisons amongst the species of *P. meishanensis*, *P. davidii*, *P. dejuniana* and *P. epilosa* are shown in Table 1.




Figure 5. Phylogenetic tree reconstruction of 35 taxa using the ML method, based on complete cp genomes (the phylogenetic position of *Primula meishanensis* is marked with red circle.)




Figure 6. The covariance analyses of two species.

Table 1. Morphological characters comparison amongst *P. meishanensis* and closely-related species of *P. dejuniana*, *P. breviscapa* and *P. epilosa*.

Features	<i>P. meishanensis</i>	<i>P. dejuniana</i>	<i>P. epilosa</i>	<i>P. davidii</i>
Roots	numerous, without hairs	numerous, without hairs	few, without hairs	numerous, without hairs
Leaf blade	smooth , oblanceolate, 19.5–25 × 2.5–5 cm	rough, oblanceolate, 8–13(22) × 2–3(5.5) cm	rough, oblong-obovate to oblong-oblanceolate, 5–10 × 2–4 cm	rough, oblong to obovate-oblong, (5–)8–18 × 1.5–4 cm
Petiole	2.5–5 cm long	1–3 cm long	0.5–2.5 cm long	indistinct to nearly obsolete
Leaf apex	acute	Acute, but with a small point at tip	rounded	rounded
Leaf margin	ciliate, sharply and remotely dentate	ciliate, sharply and remotely dentate	hydathode-dentate	erose-dentate
Scape	solitary, 7.2–11.8 cm long, pilose, carrying a terminal umbel of (1)2–3(4) flowers	usually 1 per rosette, 8–12(–18) cm long, pilose, carrying a terminal umbel of 2–6 flowers	3.5–14 cm, sparsely glandular; umbel solitary, 2–5 flowered	8–20 cm, rust-coloured pilose, umbels 2–10-flowered
Corolla	pastel violet	pale rose-purple	rose-purple with a yellow eye	pale rose-purple
Pin flowers	corolla tube 9.0–10.0 mm long, stamens inserted at middle of corolla tube, style slightly exceeding the corolla tube mouth	corolla tube ca. 1.8 cm long, stamens inserted at middle of corolla tube, style reaching annulus	corolla tube ca. 1 cm; stamens ca. 4 mm above base of corolla tube; style slightly exserted	stamens ca. 3.5 mm above base of corolla tube; style ca. as long as tube
Thrum flowers	corolla tubes 11.0–12.0 mm long, ca. 2 mm in diam., style 4.0–7.0 mm above base of corolla tube	corolla tube ca. 2 cm long, style ca. 9 mm long reaching to middle of corolla	corolla tube 1.4–1.7 cm; style 4–5.5 mm	unknown
Altitude	400–1000 m	618–979 m	2000–2900 m	ca. 1000 m
Flowering	December to February	early February to early March	April to May	April

Molecular phylogeny

In this study, 35 chloroplast genome sequences and measured chloroplast genome sequence of *P. meishanensis* were used to construct the ML evolutionary tree (Table 2) (Fig. 5). *Primula meishanensis* is sister to *Primula dejuniana*, forming a highly supported clade. However, the branch lengths of the two species are different, so they should be two species. The number of chloroplast genes etc. is not the same in the two species (Table 3). The complete chloroplast genome of *Primula dejuniana* is 151,988 bp, with the GC content of 37.01% (Fig. 7). The LSC length of *Primula dejuniana* is 83,888 bp, SSC length is 17,730 bp and IR length is 25,185 bp. In the nucleotide sequence of the complete chloroplast gene of the *Primula dejuniana*, the number of A is 47,313, the number of T is 48,431, the number of C is 28,628 and the number of G is 27,616. The complete chloroplast genome of *Primula meishanensis* is 152,175 bp, with the GC content of 36.93% (Fig. 8). The LSC length of *Primula meishanensis* is 84,052 bp, SSC length is 17,773 bp and IR length is 25,175 bp. In the nucleotide sequence of the complete chloroplast gene of the *Primula meishanensis*, the number of A is 47,469, the number of T is 48,508, the number of C is 28,613 and the number of G is 27,585. These complete cp genomes sequences are aligned by MAFFT (Fig. 6). We obtained a fasta file for the comparison of the two species and put it in the additional file (see Suppl. material 1). The cis-splicing genes are the same, but their positions in the sequence are different (Figs 9, 11). The *rps12* is a trans-splicing gene (Figs 10, 12).

Table 2. Molecular analysis for the species.

Species	Family	Genus	GenBank number
<i>Ardisia polysticta</i> Miq.	Myrsinaceae	<i>Ardisia</i> Sw.	KC465962
<i>Maesa montana</i> A.DC.	Myrsinaceae	<i>Maesa</i> Forssk.	KU569490
<i>Primula bracteata</i> Franch.	Primulaceae	<i>Primula</i> L.	NC053592
<i>Primula bulleyana</i> Forrest	Primulaceae	<i>Primula</i> L.	NC046947
<i>Primula calliantha</i> Franch.	Primulaceae	<i>Primula</i> L.	ON804895
<i>Primula chrysoclora</i> Balf.f. & Kingdon-Ward	Primulaceae	<i>Primula</i> L.	KX668178
<i>Primula chungensis</i> Balf.f. & Kingdon-Ward	Primulaceae	<i>Primula</i> L.	NC050245
<i>Primula denticulata</i> Wight	Primulaceae	<i>Primula</i> L.	NC050247
<i>Primula dryadifolia</i> Franch.	Primulaceae	<i>Primula</i> L.	NC053596
<i>Primula filchnerae</i> R.Knuth	Primulaceae	<i>Primula</i> L.	NC051972
<i>Primula forrestii</i> Balf.f.	Primulaceae	<i>Primula</i> L.	NC053602
<i>Primula handeliana</i> W.W.Sm. & Forrest	Primulaceae	<i>Primula</i> L.	MG181221
<i>Primula jiugongshanensis</i> J.W.Shao	Primulaceae	<i>Primula</i> L.	NC056335
<i>Primula knuthiana</i> Pax	Primulaceae	<i>Primula</i> L.	MG181223
<i>Primula kwangtungensis</i> W.W.Sm.	Primulaceae	<i>Primula</i> L.	KX774737
<i>Primula matthioli</i> (L.) V.A.Richt.	Primulaceae	<i>Primula</i> L.	KY235373
<i>Primula moupinensis</i> Franch.	Primulaceae	<i>Primula</i> L.	NC050244
<i>Primula obconica</i> Hance, J. Bot.	Primulaceae	<i>Primula</i> L.	NC046415
<i>Primula odontocalyx</i> Pax	Primulaceae	<i>Primula</i> L.	NC065386
<i>Primula oreodoxa</i> Franch.	Primulaceae	<i>Primula</i> L.	NC050848
<i>Primula pellucida</i> Franch.	Primulaceae	<i>Primula</i> L.	NC050248
<i>Primula persimilis</i> G.Hao, C.M.Hu & Y.Xu	Primulaceae	<i>Primula</i> L.	KX641757
<i>Primula poissonii</i> Franch.	Primulaceae	<i>Primula</i> L.	KF753634
<i>Primula pulchella</i> Franch.	Primulaceae	<i>Primula</i> L.	NC050246
<i>Primula ranunculoides</i> F.H.Chen	Primulaceae	<i>Primula</i> L.	NC056361
<i>Primula sikkimensis</i> Hook.	Primulaceae	<i>Primula</i> L.	NC050243
<i>Primula sinensis</i> Lour.	Primulaceae	<i>Primula</i> L.	KU321892
<i>Primula stenodonta</i> Balf.f. ex W.W.Sm. & H.R.Fletcher	Primulaceae	<i>Primula</i> L.	KX668176
<i>Primula szechuanica</i> Pax	Primulaceae	<i>Primula</i> L.	NC080275
<i>Primula tsiangii</i> W.W.Sm.	Primulaceae	<i>Primula</i> L.	NC046755
<i>Primula veris</i> L.	Primulaceae	<i>Primula</i> L.	KX639823
<i>Primula wilsonii</i> Dunn	Primulaceae	<i>Primula</i> L.	MW442886
<i>Primula woodwardii</i> Balf.f.	Primulaceae	<i>Primula</i> L.	MG181222
<i>Primula dejuniana</i> G.Hao, C.M.Hu & Yuan Xu	Primulaceae	<i>Primula</i> L.	PQ213817
<i>Primula meishanensis</i>	Primulaceae	<i>Primula</i> L.	PQ213816

Table 3. Comparative analyses of cp genomes between the two species.

Species	Genome Size (bp)	LSC (bp)	IR (bp)	SSC (bp)	A	T	C	G	GC Content (%)
<i>Primula dejuniana</i>	151,988	83,888	25,185	17,730	47,313	48,431	28,628	27,616	37.01
<i>Primula meishanensis</i>	152,175	84,052	25,175	17,773	47,469	48,508	28,613	27,585	36.93

Primula dejuniana




Figure 7. Circular map of *Primula dejuniana*. The map of complete chloroplast genome was generated using CPGView (<http://www.1kmpg.cn/cpgview>).

Primula meishanensis




Figure 8. Circular map of *Primula meishanensis*. The map of complete chloroplast genome was generated using CPGView (<http://www.1kmpg.cn/cpgview>).




Figure 9. Schematic map of the cis-splicing genes in the chloroplast genome of *Primula dejuniana*.




Figure 10. Schematic map of the trans-splicing genes in the chloroplast genome of *Primula dejuniana*.




Figure 11. Schematic map of the cis-splicing genes in the chloroplast genome of *Primula meishanensis*.




Figure 12. Schematic map of the trans-splicing genes in the chloroplast genome of *Primula meishanensis*.

Acknowledgements

We are grateful to the staff of KUN and P for providing online access to specimens.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was financially supported by the National Natural Science Foundation of China (No. 32000158), the National Science & Technology Fundamental Resources Investigation Program of China (No. 2021XJKK0702), the Foundation of Sustainable Development Research Center of Resources and Environment of Western Sichuan, Sichuan Normal University (No. 2020CXZYHJZX03), Laboratory equipment research projects, Sichuan Normal University (No. SYJS20220014), Mianyang Normal University Doctoral Start-up Fund Project (QD2023A33) and Meishan Ecological Quality Sample Site Monitoring Project (No. MSJCZ20230007).

Author contributions

ZF and KH collected this species. BL, XC, DH, YL, JY and XX performed the data analysis. TL wrote the manuscript. TL and XC revised the manuscript. All authors have read and approved the final manuscript.

Author ORCIDs

Tingyu Li  <https://orcid.org/0009-0005-6873-2114>

Xinyu Chen  <https://orcid.org/0009-0008-6347-2490>

Bo Li  <https://orcid.org/0000-0002-5131-8639>

Donglai Hua  <https://orcid.org/0000-0002-7778-2608>

Can Luo  <https://orcid.org/0009-0007-1815-6926>

Huixian Luo  <https://orcid.org/0009-0002-8332-5494>

Yun Liang  <https://orcid.org/0009-0009-9005-4662>

Jieli Yue  <https://orcid.org/0009-0002-8786-2473>

Xiaodan Xi  <https://orcid.org/0009-0008-4917-6345>

Ke Huang  <https://orcid.org/0009-0006-2057-5452>

Zhixi Fu  <https://orcid.org/0000-0002-2789-6287>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

Allen GC, Flores-Vergara MA, Krasynanski S, Kumar S, Thompson WF (2006) A modified protocol for rapid DNA isolation from plant tissues using cetyltrimethylammonium bromide. *Nature Protocols* 1(5): 2320–2325. <https://doi.org/10.1038/nprot.2006.384>

- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA (2012) SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* 19(5): 455–477. <https://doi.org/10.1089/cmb.2012.0021>
- Beentje H (2016) *The Kew Plant Glossary, an Illustrated Dictionary of Plant Terms*. Kew Publishing, 184.
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.
- Hu CM (1990) *Primula*. In: Chen FH, CM Hu (Eds) *Fl. Reipubl. Popularis Sin.*, Vol. 59. Science Press, Beijing, 1–277.
- Hu CM (1994) On the geographical distribution of the Primulaceae. *Redai Yaredai Zhiwu Xuebao* 2(4): 1–14.
- Hu CM (1998) Diversity and distribution of *Primula* in China. In: Qiu ST, CI Peng (Eds) *Plant diversity and conservation by the Cross-strait*. National Museum of Natural Science, 24–27.
- Hu CM, Geng YY (2003) Two New Species of *Primula* (Primulaceae) from China. *Novon* 13(2): 196–199. <https://doi.org/10.2307/3393518>
- Hu CM, Kelso S (1996) Primulaceae. In: Wu Z-Y, Raven PH (Eds) *Flora of China*, Vol. 15. Science Press, Beijing and Miss. Bot. Gard. Press, St. Louis, 39–78.
- IUCN (2024) Guidelines for using the IUCN Red List categories and criteria. Version 15.1. Prepared by the Standards and Petitions Committee. <https://www.iucnredlist.org/resources/redlistguidelines> [Accessed March 1, 2024]
- Ju WB, Deng HN, Zhu DH, Gao YD, Gao XF, Xu B (2021) *Primula dujiangyanensis* (Primulaceae) discovered from Sichuan, Southwest China. *Phytotaxa* 510(3): 275–280. <https://doi.org/10.11646/phytotaxa.510.3.7>
- Ju WB, Deng HN, Liu F, He XJ, Gao XF, Xu B (2023) *Primula medogensis*, a new species of Primulaceae from Tibet of China. *PhytoKeys* 230: 107–114. <https://doi.org/10.3897/phytokeys.230.107008>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Li R, Hu CM (2009) *Primula lihengiana* (Primulaceae), a new species from Yunnan, China. *Annales Botanici Fennici* 46(2): 130–132. <https://doi.org/10.5735/085.046.0208>
- Li X, Cheng YH, Lin HQ, Chen C, Gao XF, Deng HN, Yu F, Andelka PM, Ju WB, Xu B (2023) *Primula wolongensis* (Primulaceae), a new species of the primrose from Sichuan, China. *PhytoKeys* 218: 47–57. <https://doi.org/10.3897/phytokeys.218.91161>
- Liu XF, Luo JJ, Zhang MK, Wang Q, Liu J, Wu D, Fu ZX (2023) Phylogenomic analysis of two species of *Parasenecio* and comparative analysis within tribe Senecioneae (Asteraceae). *Diversity* 15(4): 563. <https://doi.org/10.3390/d15040563>
- Luo JJ, Zhang MK, Liu XF, Chen H, Li TY, Ma XD, Huang K, Fu ZX (2023) *Primula xilingensis* (Primulaceae), a new species from Sichuan, China. *PhytoKeys* 234: 135–143. <https://doi.org/10.3897/phytokeys.234.108411>
- Rana HK, Rana SK, Sun H, Fujikawa K, Luo D, Joshi LR, Ghimire SK (2021) *Saussurea talungensis* (Asteraceae), a new species from Humla, Nepal Himalayas. *PhytoKeys* 176: 55–66. <https://doi.org/10.3897/phytokeys.176.61996>
- Rankin DWH (2010) *Primula nghialoensis*. *Curtis's Botanical Magazine* 27(2): 132–139. <https://doi.org/10.1111/j.1467-8748.2010.01689.x>

- Richards J (2003) *Primula*. 2nd ed. Timber Press, London, 1–346.
- Stamatakis A, Hoover P, Rougemont J (2008) A rapid bootstrap algorithm for the RAxML Web servers. *Systematic Biology* 57(5): 758–771. <https://doi.org/10.1080/10635150802429642>
- Sun J, Zhang DG, Huang XH, Tojibaev K, Yang JY, Wang HC, Deng T (2020) *Primula sunhangii* (Primulaceae): A new species from Hubei, Central China. *PhytoKeys* 156: 103–112. <https://doi.org/10.3897/phytokeys.156.49137>
- Wang ZH, Wang Y, Chen L, Peng H, Wu ZK, Guo G (2022) *Primula longipilosa* (Primulaceae), a new species from Yunnan, China. *PhytoKeys* 194: 15–22. <https://doi.org/10.3897/phytokeys.194.81335>
- Wei D, Wang W, Xu Y, Hao G (2022) *Primula tsaiana* (Primulaceae), a New Species from Yunnan, China, and a New Synonym of *P. wenshanensis*. *Annales Botanici Fennici* 59(1): 111–116. <https://doi.org/10.5735/085.059.0117>
- Wu ZK, Zhao FW, Chen JH, Huang Y (2019) *Primula dongchuanensis* (Primulaceae), a new species from northern Yunnan, China. *PhytoKeys* 130: 171–181. <https://doi.org/10.3897/phytokeys.130.35047>
- Xiao JW, Yan GJ, Li WP, Tang M (2022) *Aster quanzhouensis* (Asteraceae), a new riparian species from eastern China. *PhytoKeys* 195: 93–106. <https://doi.org/10.3897/phytokeys.195.82411>
- Xu Y, Yuan S, Hu CM, Hao G (2014) *Primula dejuniana* (Primulaceae), A New Species from Sichuan, China. *Annales Botanici Fennici* 51(6): 372–374. <https://doi.org/10.5735/085.051.0602>
- Xu Y, Yu XL, Hu CM, Hao G (2016) Morphological and molecular phylogenetic data reveal a new species of *Primula* (Primulaceae) from Hunan, China. *PLoS One* 11(8): e0161172. <https://doi.org/10.1371/journal.pone.0161172>
- Xu Y, Li CH, Hu CM, Hao G (2016a) *Primula wawushanica* sp. nov. (Primulaceae) from Sichuan, southwestern China. *Nordic Journal of Botany* 34(2): 156–158. <https://doi.org/10.1111/njb.00894>
- Xu Y, Yu XL, Hu CM, Hao G (2016b) Morphological and molecular phylogenetic data reveal a new species of *Primula* (Primulaceae) from Hunan, China. *PLoS One* 11(10): e0165355. <https://doi.org/10.1371/journal.pone.0165355>
- Xu Y, Huang G, Hu C, Hao G (2017) *Primula pengzhouensis* (Primulaceae), a new species from Sichuan, southwestern China. *Plant Diversity* 39(4): 229–231. <https://doi.org/10.1016/j.pld.2017.08.003>
- Xu Y, He DM, Yang LZ, Hao G (2022) *Primula surculosa* (Primulaceae), a new species from Yunnan, China. *PhytoKeys* 212: 29–35. <https://doi.org/10.3897/phytokeys.212.91133>
- Yang B, Ya JD, Zhang W, Song Y, Wang W, Zhu ZM, He JH, Zuo YJ, Tan YH (2023) Two new species of *Primula* (Primulaceae) from Yunnan, China. *Taiwania* 68(2): 230–240. <https://doi.org/10.6165/tai.2023.68.230>
- Zhang N, Jiang XQ, Wu ZK (2023) *Primula pingbaensis* (Primulaceae), a new species from Guizhou, China. *PhytoKeys* 221: 85–93. <https://doi.org/10.3897/phytokeys.221.97948>
- Zheng X, Qin K, Li T, Qu T, Luo J, Zhang G, Li B, Li P, Fu Z (2024) A new species, *Aster yaoshanensis* (Asteraceae, Astereae), from Guangxi (China), based on morphology and molecular phylogenetic data. *Frontiers in Plant Science* 15: 1367917. <https://doi.org/10.3389/fpls.2024.1367917>

Supplementary material 1

Supplementary data

Authors: Tingyu Li, Xinyu Chen, Bo Li, Donglai Hua, Can Luo, Huixian Luo, Yun Liang, Jieli Yue, Xiaodan Xi, Ke Huang, Zhixi Fu

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.248.127117.suppl1>