

Research Article

Cite this article: Oreshkova NV, Bondar EI, Sharov VV, Dhungana SP, Gailing O, Krutovsky KV (2023). Population genetic variation of microsatellite markers developed for Siberian fir (*Abies sibirica* Ledeb.) and European silver fir (*Abies alba* Mill.) using whole genome sequencing data. *Plant Genetic Resources: Characterization and Utilization* **21**, 149–158. <https://doi.org/10.1017/S1479262123000552>

Received: 3 April 2023

Revised: 25 July 2023

Accepted: 26 July 2023

First published online: 22 August 2023

Keywords:

Abies; conifers; fir; genetic diversity; genome sequencing; microsatellite markers; population structure

Corresponding author:

Konstantin V. Krutovsky;
Email: kkrutov@gwdg.de

Population genetic variation of microsatellite markers developed for Siberian fir (*Abies sibirica* Ledeb.) and European silver fir (*Abies alba* Mill.) using whole genome sequencing data

Natalia V. Oreshkova^{1,2,3,4}, Eugenia I. Bondar^{1,2}, Vadim V. Sharov^{1,5,6},

Sudha P. Dhungana⁷, Oliver Gailing^{7,8} and Konstantin V. Krutovsky^{2,4,7,8,9,10} 

¹Laboratory of Genomic Research and Biotechnology, Federal Research Center “Krasnoyarsk Science Center of the Siberian Branch of the Russian Academy of Sciences”, 660036 Krasnoyarsk, Russia; ²Laboratory of Forest Genomics, Genome Research and Education Center, Institute of Fundamental Biology and Biotechnology, Siberian Federal University, 660041 Krasnoyarsk, Russia; ³Laboratory of Forest Genetics and Selection, V. N. Sukachev Institute of Forest, Siberian Branch of Russian Academy of Sciences, 660036 Krasnoyarsk, Russia; ⁴Department of Genomics and Bioinformatics, Institute of Fundamental Biology and Biotechnology, Siberian Federal University, 660041 Krasnoyarsk, Russia; ⁵Department of High-Performance Computing, Institute of Space and Information Technologies, Siberian Federal University, 660074 Krasnoyarsk, Russia; ⁶Tauber Bioinformatics Research Center, University of Haifa, Haifa 3498838, Israel; ⁷Department of Forest Genetics and Forest Tree Breeding, Georg-August University of Göttingen, 37077 Göttingen, Germany; ⁸Center for Integrated Breeding Research, George-August University of Göttingen, 37075 Göttingen, Germany; ⁹Laboratory of Population Genetics, N. I. Vavilov Institute of General Genetics, Russian Academy of Sciences, 119333 Moscow, Russia and ¹⁰Scientific and Methodological Center, G. F. Morozov Voronezh State University of Forestry and Technologies, 394087 Voronezh, Russia

Abstract

The article presents the results of the search for microsatellite or simple sequence repeat (SSR) loci with tri-, tetra-, penta- and hexanucleotide tandem repeat motifs in the draft *de novo* assembly of the Siberian fir (*Abies sibirica* Ledeb.) genome and the development of convenient relatively highly and moderately polymorphic markers that can be easily genotyped even by simple gel electrophoresis. In total, 64 pairs of oligonucleotide polymerase chain reaction (PCR) primers for 32 detected microsatellite loci were designed and tested. Based on the testing results, 10 most promising polymorphic loci were selected and genotyped in eight natural populations of Siberian fir. Homologous microsatellite loci in the genome of European silver fir (*Abies alba* Mill.) were also identified by mapping Siberian fir contigs, containing SSR loci to the European silver fir genome assembly. A multiplex panel of 14 universal microsatellite loci was developed and genotyped in samples from four natural populations of *A. alba* and a small sample of eight Nordmann fir (*Abies nordmanniana* (Steven) Spach) trees.

Introduction

Over the past 15–20 years, genomic research has been rapidly developing due to the development of high-throughput DNA sequencing methods, called next generation sequencing technologies. This made it possible to sequence giant genomes, including conifer genomes, which are 4–9 times larger [~11–35 Gbp (Leitch *et al.*, 2019)] than the human genome [~3.06 Gbp (Nurk *et al.*, 2022)]. Thus, to date, more than 10 genomes of the conifer family Pinaceae have been fully sequenced, annotated and assembled (see for review Kuzmin *et al.*, 2019; Bondar *et al.*, 2022), including *Abies alba* (Mosca *et al.*, 2019) and partially *Abies sibirica* (Nystedt *et al.*, 2013).

Thanks to whole genome or partial genome sequencing of conifers, it became possible to develop easily genotyped species-specific microsatellite markers for various conifer species, for example, for such main species of the Eurasian boreal forests as *Pinus sibirica* Du Tour (Belokon *et al.*, 2016) and *Larix sibirica* Ledeb. (Oreshkova *et al.*, 2017, 2019). These markers facilitate population genetic studies, identification of the timber and plant material origin in the fight against illegal logging and control over the legal origin of plant material, effective reforestation, identification of clone material, etc. (Krutovsky *et al.*, 2019).

The main object of the study presented here was the Siberian fir (*A. sibirica* Ledeb.), one of the most important forest-forming species of the dark coniferous taiga with a large distribution zone, including the northeast of the European part of Russia, the Urals and most of Siberia. In general, representatives of the genus *Abies* (Mill) are of great interest to foresters, dendrologists, gardeners and landscapers. This genus is considered one of the largest in the Pinaceae



© The Author(s), 2023. Published by Cambridge University Press on behalf of National Institute of Agricultural Botany. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

family, including from 40 to 65 species according to various modern estimates (Semerikova et al., 2018).

However, despite its significance the genetic diversity of Siberian fir has been insufficiently studied. Most of earlier such studies were carried out using allozyme markers, which are less informative than simple sequence repeat (SSR) markers, and did not cover the entire range of the species (Goncharenko and Padutov, 1995; Larionova and Ekart, 2005; Semerikova and Semerikov, 2006; Larionova et al., 2007). Several studies were carried out also in Siberian fir on the dynamics of populations, population size and phylogeography using cytoplasmic DNA and amplified fragment length polymorphism (AFLP) markers (Semerikova and Semerikov, 2014, 2016; Semerikov et al., 2022). However, despite these studies, there is practically no information on the genetic diversity, structure and differentiation of Siberian fir populations in the Krasnoyarsk Territory and the Republic of Khakassia, as well as in many other large areas of its natural distribution. The genetic structure, subdivision and differentiation of populations of this species have not been studied under ecologically heterogeneous environmental conditions. This does not allow assessing the general state of the genetic resources of the species, their spatial distribution, which makes it difficult to develop programmes for the protection, rational use and reproduction of the genetic resources of the species.

Meanwhile, abundant distribution in the genome, high reproducibility, codominance and relatively easy identification have made microsatellite markers popular in modern studies of conifer population genetics. Several microsatellite markers with mostly dinucleotide motifs were previously developed for some fir species (Hansen et al., 2005; Cremer et al., 2006; Cvrčková et al., 2015). However, they are suitable for genotyping only using capillary gel electrophoresis, while microsatellite markers with longer motifs can be easily genotyped in any laboratories with simple slab gel electrophoresis. Therefore, our main objective was to generate universal SSR markers suitable for reliable genotyping in any laboratories with any gel electrophoresis equipment, such as for either simple vertical electrophoresis in polyacrylamide slab gel or much more challenging capillary gel electrophoresis requiring more expensive instruments.

Fir is a good bioindicator of environmental conditions in the areas of their growth (Świercz et al., 2022). Therefore, in addition to the main goal of this study to develop new highly informative microsatellite markers that can be easily genotyped in *A. sibirica* based on the data of whole genome sequencing of this species, we also searched for homologous microsatellite loci in the genome of a closely related species, European silver fir (*A. alba* Mill.), tested them in population samples and developed a multiplex genotyping panel of 14 microsatellite loci, which were successfully tested and genotyped in the samples from eight *A. sibirica* populations, four *A. alba* populations and a small sample of eight trees representing *Abies nordmanniana*.

Both *A. sibirica* and *A. alba* are important species of the boreal and temperate forests in Eurasia. These trees are up to 30 m tall and up to 0.5 m in diameter, living up to 200–250 years in natural conditions. Fir is very shade-tolerant, has low frost resistance and often suffers from late spring frosts. It grows best on well-drained soddy-podzolic loam soils with close underneath limestone occurrence. It avoids strongly podzolized and stagnant-moistened soils and does not grow on poor sandy soils. It can form both pure and mixed (fir-spruce) stands. In the mountains it rises to 2000 m above sea level, where it is present in a bushy form (Mauri et al., 2016; Dobrowolska et al., 2017; Zaitsev et al., 2018).

Materials and methods

Search for microsatellite loci and design of PCR primers

Nucleotide reads obtained as a result of whole genome sequencing of *A. sibirica* Ledeb. with low coverage were used to search for microsatellite loci and design their polymerase chain reaction (PCR) primers. The reads of *A. sibirica* were generated additionally in a project dedicated mainly to the sequencing and assembly of the genome of Norway spruce, *Picea abies* (L.) Karst. (Nystedt et al., 2013; <https://www.ncbi.nlm.nih.gov/bioproject/PRJEB1894>; the NCBI GenBank SRA accession number ERP002568). We assembled reads into contigs using CLC Assembly Cell v.4.4 (QIAGEN, Aarhus, Denmark). The obtained contigs were used to search for microsatellite loci using the GMATo v.1.2 software (Wang et al., 2013). To increase the reliability of genotyping of the prospective microsatellite markers, the search was limited to the search for contigs containing tandem repeats of only three-, four-, five- and six-nucleotide long motifs with thresholds for minimal number of repeats 15, 10, 6 and 7, respectively.

The PCR primers for amplification of the found microsatellite loci within the size range of 140–280 bp were designed using the WebSat online tools (Martins et al., 2009). For successful PCR amplification of microsatellite loci, their primers must be locus-specific, that is, the PCR annealing sites matching primer sequence pairs for a given marker should not occur anywhere else in the genome. Therefore, to design unique, single locus-specific primers, their sequences were mapped to the newly generated assembly of the *A. sibirica* genome using BLAST. Primer pairs that had annealing sites for more than one locus were excluded from further analysis. Single locus-specific PCR primers selected for further testing were synthesized by Evrogen (Moscow, Russia). In addition, Siberian fir contigs containing successful microsatellite markers were used to search for homologous microsatellite loci in the European silver fir (*A. alba* Mill.) genome (Mosca et al., 2019). Then, PCR primers designed for the found homologous loci were tested in four populations of *A. alba* and a small sample of *A. nordmanniana*.

Plant material and DNA isolation

Total DNA was isolated from 100–200 mg of dried needles per individual tree sample collected from 240 Siberian fir trees in eight populations (30 trees per each population) in Central Siberia (Fig. 1) using the cetyltrimethylammonium bromide (CTAB) method (Doyle and Doyle, 1990).

About 50 mg of fresh needles per each sample were used for DNA isolation from 96 individual tree samples of *A. alba* representing four populations of anonymous origin (24 samples per each population, respectively) using the DNeasy[®]96 Plant Mini Kit and following the standard protocol (Qiagen, Hilden, Germany). The same kit was used to isolate DNA from germinated seeds of *A. nordmanniana*. The samples of *A. alba* were provided by Isogen GmbH (Göttingen, Germany; <https://isogen.de>) and the seed samples of *A. nordmanniana* by PlusBaum Samen GmbH (Nagold, Germany; <https://plusbaum-samen.de>).

PCR amplification and genotyping of microsatellite loci

GenePak PCR Core kits (IsoGene Laboratory Ltd, Moscow, Russia) containing hot-start *Taq*-DNA polymerase, deoxynucleoside triphosphates (dNTPs) and magnesium chloride were used for PCR amplification of selected nuclear microsatellite loci in




Figure 1. Area of *Abies sibirica* and geographic location of the eight populations genotyped in the study using microsatellite markers. Krasnoyarsk Region: Boguchanskaya (BB, 58°21' N, 97°30' E, 419), Ordzhonikidzevskaya (MO, 58°19' N, 94°56' E, 293), Kytatskaya (TK, 56°57' N, 91°33' E, 232), Maganskaya (MB, 55°40' N, 93°00' E, 333), Kesovskaya (IK, 55°18' N, 96°43' E, 423), Artemovskaya (KA, 54°23' N, 93°33' E, 542), Shushenskoe (SH, 52°57' N, 91°43' E, 642); Republic of Khakassia: Tabatskaya (BT, 52°48' N, 90° 46' E, 849).

the Axygen MaxyGene II Gradient Thermal Cycler (Axygen Scientific Inc., Union City, California, USA). A touchdown PCR program was used to reduce non-specific amplification, which included first DNA denaturation at 94°C for 1 min, then 9 cycles, including 30 s of denaturation at 94°C, annealing of primers for 30 s at 60°C, with a decrease by 1°C each cycle (up to 50°C), and 1 min of elongation at 72°C. The next 24 cycles included DNA denaturation at 94°C for 30 s, primer annealing at 50°C for 30 s and elongation at 72°C for 30 s. This was followed by an extension step at 72°C for 10 min, and the last stage was cooling at 4°C. Loci that showed stable interpretable PCR amplification spectra were selected for further use.

Amplification products were separated by 6% polyacrylamide gel electrophoresis using Tris-EDTA-borate electrode buffer in vertical electrophoresis chambers. The gel was stained in a solution of ethidium bromide followed by visualization in ultraviolet light. DNA fragments of the pBR322 *E. coli* plasmid digested by the *Hpa*II restriction enzyme were used as standard length markers.

PCR primers that had annealing sites also in homologous microsatellite loci of *A. alba* were also tested and genotyped in samples of 96 trees from four natural populations of *A. alba* and eight seedlings germinated from seeds of eight different *A. nordmanniana* trees using fluorescent labelled primers synthesized by Sigma Aldrich Chemie GmbH (Taufkirchen, Germany) and capillary gel electrophoresis with ABI PRISM 3130 (Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA). Visualization and processing of the results of fragment analysis (electropherograms) were carried out using the GeneMapper 4.0 program (Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA).

Population genetics data analysis

Genetic variation parameters, such as average (N_a) and effective (N_e) number of alleles, observed (H_o) and expected (H_e) heterozygosity, fixation index (F_{IS}) and different parameters of genetic

differentiation (Nei's standardized G_{ST} (G'_{STN}), Hedrick's standardized G_{ST} (G'_{STH} and G''_{ST}), Jost's estimate of differentiation D_{ST}) and assessment of the Hardy–Weinberg equilibrium (HWE) were calculated using GeneAlex v. 6.503 (Peakall and Smouse, 2006, 2012). Polymorphic information content (PIC) was calculated according to Botstein et al. (1980) using the MolMarker program (Jahnke et al., 2022). The frequencies of null alleles were inferred and analysed using MICRO-CHECKER v. 2.2.3 (Van Oosterhout et al., 2004) with data exported from GeneAlex.

Results

Abies sibirica

A total of 264,864 contigs containing tandem microsatellite repeats with three-, four-, five- and six-nucleotide motifs with a specified minimum number of repeats were found in the partial genome assembly of Siberian fir. We specifically focused on selecting loci with longer than dinucleotide motifs in order to improve scorability and comparability. Initially, 475 microsatellite loci with long enough neighbouring sequences on both ends to design PCR primers were selected, 115 of them were used to design PCR primers, and PCR primer pairs were finally synthesized by Evrogen Co. (Moscow, Russia) and further tested for 64 of them (see online Supplementary Table S1 for details and original contig sequences).

An initial test of the selected primers on four DNA samples from each population of Siberian fir showed that out of 64 primer pairs, 11 lacked an amplified product and 7 did not have the expected size of amplified product. The remaining 46 pairs were further tested on an enlarged sample of trees. According to the results of this test, 22 more loci had to be excluded, because 12 of them turned out to be monomorphic and 10 showed unstable amplification. As a result, 24 loci showed stable amplification under the selected conditions and were further used for genotyping entire population samples of Siberian fir trees (Table 1, online Supplementary Table S1).

Based on the results of genotyping of 24 selected loci in eight geographically distant populations of *A. sibirica*, another 14 loci had to be excluded due to poor or nonspecific amplification and a large number of null alleles. Using the MICRO-CHECKER program (Van Oosterhout et al., 2004), hidden null alleles were identified for the following three loci: *As_43741* in the TK and SH samples, *As_563614* in the KA and MB samples and *As_420232* in the BB sample. Hidden null allele frequencies were calculated based on the assumption that populations are in HWE (Chakraborty, 1992). Some of the loci demonstrated significant deviation from HWE, but it was not systematic across all populations. Neither of the loci demonstrated deviation from HWE for all or the majority of populations (online Supplementary Table S2).

Thus, the 10 most reliable loci were selected (Table 1, online Supplementary Table S1), which were easily and unambiguously genotyped using a simple polyacrylamide gel in vertical electrophoresis chambers (online Supplementary Fig. S1).

Abies alba and *Abies nordmanniana*

Based on the results of mapping nucleotide sequences of 64 primer pairs selected for *A. sibirica* to the *A. alba* genome assembly, 36 *A. alba* homologous loci were selected and the corresponding

primers were tested on four *A. alba* samples. Of these, 23 were stably amplified and well genotyped; 14 turned out to be polymorphic and 9 monomorphic. The polymorphic 14 markers were further tested on a larger set of eight samples using capillary electrophoresis on an ABI PRISM 3130 sequencer (Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA). To do this, one of the primers in the pair was labelled with either 6-FAM or HEX fluorescent labels. Then, 14 markers were grouped into four PCR multiplexes and genotyped in 96 trees from four *A. alba* populations and eight seedlings representing eight different *A. nordmanniana* trees. Multiplex PCR primer groups were organized in such a way as to avoid overlapping ranges of allele sizes of different amplified loci labelled with the same fluorescent dye in the same PCR reaction. A maximum of four primer pairs per group were pooled together using two different fluorescent dyes (Table 2). Among 14 selected microsatellite loci tested on samples of Nordmann fir, one was not amplified at all, nine turned out to be polymorphic and four were monomorphic (Table 2).

Population genetic analysis

Polymorphism data for 10 microsatellite loci in eight populations of *A. sibirica* are presented in Table 3 (see also online Supplementary Table S2 for details).

The studied microsatellite markers for Siberian fir turned out to be moderately polymorphic with the number of alleles (N_a) varying from 2 (*As_99437*, *As_2687764*, *As_3458593*) to 4.9 (*As_563614*) per locus on average for eight populations (Table 3). The effective number of alleles (N_e) varied from 1.6 to 3.7 at the *As_3666591* and *As_563614* loci, respectively. Observed (H_o) and expected (H_e) heterozygosity varied from 0.31 to 0.71 and from 0.36 to 0.73 for *As_3666591* and *As_563614*, respectively. The PIC was high and varied from 0.33 to 0.73 for the same pair of markers *As_3666591* and *As_563614*, respectively. The lowest (0.01) and highest (0.12) F_{ST} values were calculated for the *As_99437* and *As_3666591* loci, respectively. G'_{STH} varied from 0.01 to 0.25 for the *As_3458593* and *As_420232* loci, respectively. The highest G'_{STN} , G''_{ST} and D_{ST} values were 0.12, 0.25 and 0.17, respectively (Table 3). F_{IS} values ranged from -0.41 to 0.13, but none of them was statistically significant (Table 3). However, it should be noted that the analysis for hidden null alleles using MICRO-CHECKER revealed null alleles at three loci (*As_43741*, *As_420232*, *As_563614*) in five populations, which were taken into account while calculating the parameters of genetic variation.

Some of the loci demonstrated significant deviation from HWE, but it was not systematic across all populations. Neither of the loci demonstrated deviation from HWE for all or the majority of populations (online Supplementary Table S3).

The results of population clustering are presented in Fig. 2. According to the results of the Mantel test, no significant correlation was found between genetic and geographic distances.

Polymorphism data for 14 microsatellite loci in four populations of *A. alba* and 13 microsatellite loci in a sample of eight trees of *A. nordmanniana* are presented in Table 4 (see also online Supplementary Tables S3 and S4 for details).

Most of the microsatellite markers were highly polymorphic with number of alleles (N_a) ranging from 1.8 (*As_420232*) to 10 (*As_1898225*) alleles per locus in four populations of *A. alba* on average (Table 4). The loci *As_1998225* and *As_2280886* showed the highest genetic variation in all populations (online

Table 1. Characteristics of 24 successfully amplified polymorphic microsatellite loci selected for *Abies sibirica* based on genome sequencing data

Locus*	Motif	Nucleotide sequence of PCR primers (5'-3')		Allele length, bp
		Forward	Reverse	
As_43741 [#]	(TCA) ₁₅	CTCAAAGGTAGTGTTCCTCAGTGGTAT	CAATGATGACAATGATGTTGGC	201, 207, 219, null
As_99437 [#]	(ATG) ₁₇	GGGAGATTGATTATGTGGATCA	CATGCAAGGGGACTAATATCTTC	205, 229
As_279869	(TTA) ₁₅	ACGAAGGTGAAGGGTCTATCA	TTAAGGGGAATTGTTGTGGGTA	253
As_301948	(ATC) ₁₆	CAATGATGATGGACTAGATGGA	AAGTCTGATGGGTTATGCTTGT	214
As_878430	(AAT) ₁₅	TTACCCTCGAATCTAAAGGTGG	CTATCCCAACAAAGGGGT	260
As_1177607	(TAA) ₁₇	GTGATCTATGCCTCTCCAATGA	ACTGGTCACAATGGTTTGAATG	188
As_1563074	(ATT) ₁₈	GTGAGGAGGAGATTGTAAAGTGC	CAGATGAGTACGAACCACTACCC	142
As_420232 [#]	(TAAA) ₁₀	CCTTCCACCTTCCATGATATTC	TCCAAGAAAAGGCACAAGAAGT	168, 176, 182, 188, 192, null
As_563614 [#]	(ATAG) ₁₂	AAGTTGTTTGATCCCTCTCGAC	CATGTAATGCTCCAATCAGGAA	148, 156, 160, 164, 168, 180, 192, 222
As_721750	(TATG) ₁₀	CCCTTACCTCCCTAACACTTGA	GGCAAGATAGATGATTGACACC	147
As_2280886	(TGTA) ₁₀	CACTAAAAGAGGAGCCATGTCC	GACACCTGCAATCACCAAGAC	201
As_2687764 [#]	(TATG) ₁₀	GATGAGAGATGCTTGTGAATCC	GGAGGTTATCATGGAAGTGGAC	218, 222, 226
As_2996373	(ATAC) ₁₁	TCGCTTCCACACTGACCAC	TTACACACAAAAGGGCTCACAC	272
As_3602348	(CATA) ₁₀	GTCGCAGAATGTTGAATTGCT	CATGGATGGAGAGGACAAGTTT	278
As_347605	(GAAAA) ₆	AAGGTTCTTAAATTCACACCCC	GACTTGAGGATGAGGTTCTTGC	253
As_1133359	(GATT) ₆	GGATGCAGGGTTTCATTATTC	CCAGAAGATTGACATGGAAA	238
As_1356647 [#]	(GATAA) ₇	TAGACTTTGAGGTGATGGGCTT	TTGCTCCGTGATCTTAGGTTTT	238, 243, 248, 250
As_2288295 [#]	(AGATA) ₆	TGTTTTCCAGATTGTGAGTGC	AGTCCCGTTGACCTACTTTGTG	167, 172, 177
As_3458593 [#]	(TTATC) ₇	CTCCACATCAATCCATTTC	TGGTTTCTGACGATGACTCTA	225, 230
As_404145 [#]	(CCAGCT) ₈	TAACGACCTGCTGCTATTTTCA	GGGCACCATTCTCAATTTCTAC	173, 179, 185, 187, 191
As_1898225	(TCTTCA) ₉	ATGCTGCCATTGAGAATTAGT	ACCGCTCTTCCCTTACTTTCT	270
As_1970306	(GCAAGA) ₇	TTTGGACTGAGATTTGGGC	CGATTATGAAAGATGGGTAGC	248
As_2631734	(CACTCC) ₇	CCCACCTATCTCTCACTTCA	GGGTTGATATGTAAAAGGGAA	169
As_3666591 [#]	(TATTCC) ₇	CGTTCTTTCATTTTGTGGC	TCCTTGGCATTGTTTCATGT	238, 256, 262

*Number in the name of the loci refers the contig number in the *A. sibirica* genome assembly. [#]Ten most reliable polymorphic microsatellite loci finally selected for genotyping eight *A. sibirica* populations. Allele length in the contig is italicized.

Supplementary Table S3). The effective number of alleles (N_e) varied from 1.3 to 6.6 in the As_420232 and As_1898225 loci, respectively. The observed (H_o) and expected (H_e) heterozygosities varied from 0.21 to 0.84 and from 0.20 to 0.85 for loci As_420232 and As_1898225. The PIC was high and varied from 0.20 to 0.85 for the same pair of markers As_420232 and As_1898225, respectively. The highest (0.09) and lowest (0.01) F_{ST} values were calculated for As_2631734 and As_1970306, respectively. The $G'_{ST}H$ varied from -0.01 to 0.22 for the As_1970306 and As_2280886 loci, respectively. The highest $G'_{ST}N$, G'_{ST} and $Dest$ were 0.10, 0.23 and 0.19, respectively (Table 4). In total, differentiation was not high ranging from 0.03 to 0.08 depending on the parameter, but significant for each parameter (Table 4).

The F_{IS} values varied from -0.37 to 0.69, and significant deficiency of heterozygotes was observed for four loci in single populations, for As_563614 and As_1089795 in more than two populations (online Supplementary Table S5). It is likely caused mainly by a high frequency of hidden null alleles in these a few loci found by MICRO-CHECKER (online Supplementary Table S6).

Discussion

Abies sibirica

In general, the values of the main parameters of genetic polymorphism ($N_a = 2.9$, $N_e = 2.2$, $H_o = 0.55$, $H_e = 0.51$) of *A. sibirica* indicate a relatively lower level of diversity compared to studies of *A. alba* based on other microsatellite markers (Cremer *et al.*, 2006; Cvrčková *et al.*, 2015), but similar to the parameters for *A. alba* obtained in our study and based on a subset of markers used for *A. sibirica* (Table 4). The highest values of almost all parameters were found in the southern populations SH and BT, which are closely located to each other (Fig. 1).

Some of the loci demonstrated significant deviation from HWE, but it is likely due to relatively high self-pollination which is typical for *Abies* species that have a relatively high self-pollination rate among conifers [mean 11% for four populations of *A. balsamea* (Neale and Adams, 1985), 11% for *A. lasiocarpa* (Shea, 1987), 11% (Schroeder, 1989) or 24% (Kormuták and Lindgren, 1996) for *A. alba*, mean 13% for seven populations of *A. amabilis* (Davidson, 1990), 6% for *A. borisii* (Fady and Westfall, 1997), and 10–13% for *A. sibirica* (authors' data, unpublished)], which

Table 2. PCR multiplex panels, number and size range of alleles of the best 14 microsatellite loci used for population genetic genotyping of *Abies alba* and *Abies nordmanniana*

Locus	<i>A. alba</i>		<i>A. nordmanniana</i>		PCR multiplex panel	
	Range, bp	Number	Range, bp	Number	Fluorescent dye	#
As_1898225	241–307	12	247–307	6	HEX	1
As_1356647	263–268	2	268	1	6-FAM	
As_404145	181–211	6	181–217	4	6-FAM	
As_563614	156–162	4	180–240	6	6-FAM	
As_2612077	262–287	5	252–287	6	HEX	2
As_2631734	168–186	3	168	1	HEX	
As_3458593	240–255	4	245	1	HEX	
As_3602348	273–293	5*	No amplification	0	6-FAM	
As_154813	289–316	7*	292–295	2	HEX	3
As_2280886	220–264	10	224–272	9	6-FAM	
As_420232	179–191	2	179–191	2	6-FAM	
As_878430	260–275	4*	272	1	HEX	
As_1089795	273–339	9*	273–324	4	6-FAM	4
As_1970306	254–266	2	254–266	2	HEX	

*Loci with null alleles according to MICRO-CHECKER.

Table 3. Genetic variation parameters of the 10 microsatellite markers averaged for eight populations of *A. sibirica* and mean for all markers

Locus	N_a	N_e	H_o	H_e	PIC	F_{IS}	F_{ST}	$G'_{ST}H$	$G'_{ST}N$	G''_{ST}	Dest
As_43741	3.2	2.6	0.67	0.59	0.56	−0.13	0.04***	0.07***	0.03***	0.07***	0.04***
As_99437	2.0	1.8	0.62	0.44	0.35	−0.41	0.01	0.00	0.00	0.00	0.00
As_420232	3.4	2.6	0.63	0.61	0.61	−0.04	0.10***	0.25***	0.10***	0.25***	0.17***
As_563614	4.9	3.7	0.71	0.73	0.73	0.02	0.05***	0.13***	0.03***	0.14***	0.10***
As_2687764	2.0	1.9	0.46	0.48	0.37	0.04	0.02	0.02	0.01	0.02	0.01
As_1356647	2.5	1.7	0.45	0.42	0.36	−0.09	0.03*	0.02*	0.01*	0.02*	0.01*
As_2288295	2.6	2.0	0.52	0.49	0.42	−0.07	0.03**	0.04**	0.02**	0.05**	0.02**
As_3458593	2.0	1.9	0.59	0.49	0.37	−0.22	0.02	0.01	0.01	0.01	0.01
As_404145	3.6	2.0	0.58	0.50	0.47	−0.16	0.04***	0.06***	0.03***	0.07***	0.04***
As_3666591	2.3	1.6	0.31	0.36	0.33	0.13	0.12***	0.18***	0.12***	0.19***	0.08***
Mean	2.9	2.2	0.55	0.51	0.46	−0.09	0.05***	0.07***	0.04***	0.08***	0.04***
Population	$N_a \pm SE$		$N_e \pm SE$		$H_o \pm SE$		$H_e \pm SE$		$F_{IS} \pm SE$		
BB	2.6 ± 0.3		2.2 ± 0.2		0.55 ± 0.05		0.51 ± 0.04		−0.09 ± 0.07		
MO	2.5 ± 0.2		2.2 ± 0.2		0.62 ± 0.07		0.52 ± 0.04		−0.21 ± 0.09		
TK	2.6 ± 0.3		2.2 ± 0.2		0.48 ± 0.05		0.53 ± 0.03		0.10 ± 0.05		
MB	3.0 ± 0.4		2.3 ± 0.3		0.52 ± 0.05		0.51 ± 0.05		−0.05 ± 0.05		
IK	3.0 ± 0.3		2.1 ± 0.1		0.57 ± 0.05		0.51 ± 0.03		−0.14 ± 0.08		
KA	2.9 ± 0.3		2.0 ± 0.2		0.51 ± 0.05		0.46 ± 0.04		−0.10 ± 0.06		
SH	3.2 ± 0.4		2.3 ± 0.3		0.55 ± 0.04		0.52 ± 0.04		−0.08 ± 0.06		
BT	3.1 ± 0.3		2.4 ± 0.3		0.63 ± 0.07		0.54 ± 0.04		−0.15 ± 0.09		
Mean	2.9 ± 0.1		2.2 ± 0.1		0.55 ± 0.02		0.51 ± 0.01		−0.09 ± 0.03		

N_a , average number of alleles; N_e , effective number of alleles; H_o , observed heterozygosity; H_e , expected heterozygosity; PIC, polymorphic information content; F_{IS} , fixation index; F_{ST} , coefficient of differentiation between populations; $G'_{ST}N$, Nei's standardized G_{ST} ; $G'_{ST}H$, Hedrick's standardized G_{ST} ; G''_{ST} , Hedrick's standardized G_{ST} further corrected for bias when the number of populations is small; Dest, Jost's estimate of differentiation between populations; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Figure 2. Neighbour-joining tree with bootstrap values at the nodes (A) and principal coordinates analysis (PCoA) (B) of eight populations of *A. sibirica* based on the standard Nei's genetic distances (D_N). See Fig. 1 for the full names of the populations and their geographic location.

is comparable to other conifers [2–17% (Adams and Birkes, 1991) or mean ~15% (Table 1 in Muona, 1990; Mitton, 1992)].

Based on clustering presented in Fig. 2 phylogenetic relationships between populations of *A. sibirica* reflect their geographic location in general (Fig. 1). Cluster analysis clearly separated the populations BB, TK and MO located in the northern regions with a high degree of confidence (bootstrap value of 99%) from the distant populations located to the south, among which the southern cluster of populations MB, BT and SH is also distinguished, but with less support (bootstrap value of only 60%).

However, according to the results of the Mantel test, no significant correlation was found between genetic and geographic distances, which may indicate that, in addition to geographic isolation, the genetic differentiation in the studied area of this species could be affected by the ecological heterogeneity of the habitat at different altitudes above sea level related to soil fertility, water availability and mineral nutrition. The obtained results are consistent with similar data obtained for *A. sibirica* using other genetic markers (Larionova *et al.*, 2007; Semerikova and Semerikov, 2016).

Abies alba and *Abies nordmanniana*

In general, the variation of 14 markers developed in this study for *A. alba* was similar to the variation of 11 microsatellite markers developed for this species by Cremer *et al.* (2006): $N_a = 5.2$ vs 4.5 in our study (Table 4), $H_o = 0.31$ vs 0.46, $H_e = 0.53$ vs 0.56, and similar to $N_a = 9.2$, $H_o = 0.43$, and $H_e = 0.49$ obtained in Postolache *et al.* (2014) based on 24 markers including 16 EST-SSRs (calculated from Table 1 in Postolache *et al.*, 2014), but slightly less than $H_e = 0.75$ based on 11 microsatellite markers used by Hrivnák *et al.* (2017), including four markers developed

by Cremer *et al.* (2006), and values obtained for many other populations of *A. alba* (H_o and $H_e = 0.5$ –0.8, see Dalmaris *et al.*, 2022 for references). The lower variation observed in our study can be explained by intentionally not using microsatellite loci with dinucleotide repeats that are mostly used in other studies but notoriously known for typical difficulty regarding consistent and reliable genotyping.

Although very preliminary, variation observed in a small sample of *A. nordmanniana* was less than variation observed in much larger sample in Hansen *et al.* (2005) which was based only on five highly polymorphic markers: $N_a = 30.4$ (calculated from Table 1 in Hansen *et al.*, 2005) vs 3.5 in our study (Table 4), $H_o = 0.70$ vs 0.43 and $H_e = 0.90$ vs 0.40.

The null alleles detected in a few loci of both *A. sibirica* and *A. alba* can potentially lead to an underestimation of allelic richness and heterozygosity for these loci in populations and introduce biases in measures of population differentiation by reducing the observed number of alleles and heterozygosity estimates, but it is unlikely that they would significantly affect these parameters based on all markers, because only a few loci were affected. Moreover, to mitigate the impact of null alleles, we used MICRO-CHECKER program to statistically correct genetic diversity and differentiation parameters for null allele presence.

Conclusions

Based on whole genome sequencing data, 10 polymorphic microsatellite markers were developed for *A. sibirica*, 14 for *A. alba* and 13 for *A. nordmanniana*. Among them, at least five (*As*_1356647, *As*_404145, *As*_563614, *As*_3458593 and *As*_420232) can be used for all three species, but potentially more. The markers were tested

Table 4. Genetic variation and differentiation parameters of the 14 microsatellite markers averaged for four populations of *A. alba* (A, B, C and D) and one population of *A. nordmanniana* and mean for all markers

Locus	N_a	N_e	H_o	H_e	PIC	F_{IS}	F_{ST}	$G'_{ST}H$	$G'_{ST}N$	G''_{ST}	Dest
As_1898225	10.0	6.6	0.84	0.85	0.85	0.00	0.02	0.01	0.00	0.02	0.01
As_1356647	2.0	1.7	0.34	0.40	0.34	0.14	0.08**	0.13**	0.09**	0.14**	0.06**
As_404145	4.8	2.3	0.54	0.56	0.54	0.04	0.05*	0.08*	0.04*	0.09*	0.06*
As_563614	3.3	2.1	0.34	0.51	0.45	0.33	0.06*	0.10*	0.05*	0.11*	0.06*
As_2612077	5.8	4.0	0.31	0.75	0.75	0.58	0.05*	0.14*	0.03*	0.15*	0.12*
As_2631734	2.5	1.3	0.28	0.22	0.22	-0.26	0.09***	0.11***	0.10***	0.13***	0.03***
As_3458593	2.8	1.9	0.50	0.47	0.41	-0.06	0.08**	0.14**	0.08**	0.15**	0.08**
As_3602348	4.0	2.5	0.21	0.59	0.60	0.65	0.04	0.04	0.02	0.04	0.03
As_154813	5.8	3.1	0.66	0.67	0.66	0.01	0.04*	0.10*	0.03*	0.10*	0.07*
As_2280886	8.5	4.7	0.78	0.78	0.80	0.00	0.05***	0.22***	0.05***	0.23***	0.19***
As_420232	1.8	1.3	0.21	0.20	0.20	-0.03	0.07**	0.08**	0.08**	0.10**	0.02**
As_878430	3.8	2.7	0.46	0.62	0.65	0.26	0.03	0.03	0.01	0.03	0.02
As_1089795	5.5	3.0	0.41	0.67	0.68	0.38	0.01	-0.06	-0.02	-0.06	-0.04
As_1970306	2.3	2.0	0.60	0.50	0.38	-0.22	0.01	-0.01	-0.01	-0.01	-0.01
Mean	4.5	2.8	0.46	0.56	0.54	0.17	0.05***	0.07***	0.03***	0.08***	0.04***
Population	$N_a \pm SE$	$N_e \pm SE$	$H_o \pm SE$	$H_e \pm SE$	$F_{IS} \pm SE$						
A	4.3 ± 0.6	2.7 ± 0.3	0.45 ± 0.07	0.56 ± 0.05	0.16 ± 0.10						
B	4.8 ± 0.9	3.0 ± 0.4	0.43 ± 0.06	0.58 ± 0.05	0.25 ± 0.10						
C	4.9 ± 0.7	2.8 ± 0.4	0.44 ± 0.07	0.57 ± 0.06	0.19 ± 0.09						
D	3.9 ± 0.6	2.7 ± 0.4	0.44 ± 0.06	0.53 ± 0.06	0.14 ± 0.10						
Mean	4.5 ± 0.3	2.8 ± 0.2	0.44 ± 0.03	0.56 ± 0.03	0.18 ± 0.05						
<i>Abies nordmanniana</i> *	3.5 ± 0.7	2.4 ± 0.4	0.43 ± 0.11	0.40 ± 0.09	-0.10 ± 0.11						

*Based on 13 loci (As_3602348 with no amplification was excluded).

on samples from eight natural populations of *A. sibirica* and four populations of *A. alba*, and preliminary data on the level of population genetic variation and differentiation were obtained. These markers can potentially be used also in other species in genus *Abies* with the ability to use simple gel electrophoresis, which is very convenient in field research. For genetic laboratories equipped with devices for capillary gel electrophoresis, we have developed multiplex panels of 14 loci. It should be noted that only eight tree samples were studied for *A. nordmanniana*; therefore, the data for this species are preliminary, and it is necessary to significantly increase the number of samples and accessions for more accurate estimates of the genetic variability of this species.

The proposed list of nuclear microsatellite loci will be very useful for studying the variability of natural and artificial populations of different fir species helping to address different problems and questions related to conservation, restoration and reproduction of fir forests.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S1479262123000552>.

Acknowledgements. We thank Alexandra Dolynska, Dr Markus Müller and other colleagues at the Department of Forest Genetics and Forest Tree Breeding (Georg-August University of Göttingen, Germany) for their help with developing and testing microsatellite markers of *A. alba* and *A. nordmanniana*. Funds for the development and genotyping of nSSRs in *A. alba* were provided by the University of Göttingen. We thank also Dr Vladimir

L. Semerikov (Laboratory of Molecular Plant Ecology, Institute of Plant and Animal Ecology, Ural Branch of Russian Academy of Science, Ekaterinburg, Russia) for help with preparing the manuscript.

References

- Adams WT and Birkes DS (1991) Estimating mating patterns in forest tree populations. In Fineschi S, Malvolti ME, Cannata F and Hattermer HH (eds), *Biochemical Markers in the Population Genetics of Forest Trees*. The Hague, The Netherlands: SPB Academic Publishing bv, pp. 157–172.
- Belokon MM, Politov DV, Mudrik EA, Polyakova TA, Shatokhina AV, Belokon YS, Oreshkova NV, Putintseva YA, Sharov VV, Kuzmin DA and Krutovsky KV (2016) Development of microsatellite genetic markers in Siberian stone pine (*Pinus sibirica* Du Tour) based on the *de novo* whole genome sequencing. *Russian Journal of Genetics* 52, 1263–1271.
- Bondar EI, Feranchuk SI, Miroshnikova KA, Sharov VV, Kuzmin DA, Oreshkova NV and Krutovsky KV (2022) Annotation of Siberian larch (*Larix sibirica* Ledeb.) nuclear genome – one of the most cold resistant tree species in the only seasonal senescence GENUS in Pinaceae. *Plants* 11, 2062.
- Botstein D, White RL, Skolnick M and Davis RW (1980) Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American Journal of Human Genetics* 32, 314–331.
- Chakraborty R (1992) Apparent heterozygote deficiencies observed in DNA typing data and their implications in forensic applications. *Annals of Human Genetics* 56, 45–57.
- Cremer E, Liepelt S, Sebastiani F, Buonamici A, Michalczyk IM, Ziegenhagen B and Vendramin GG (2006) Identification and

- characterization of nuclear microsatellite loci in *Abies alba* Mill. *Molecular Ecology Notes* 6, 374–376.
- Cvrčková H, Máchová P and Malá J** (2015) Use of nuclear microsatellite loci for evaluating genetic diversity among selected populations of *Abies alba* Mill. in the Czech Republic. *Journal of Forest Science* 61, 345–351.
- Dalmaris E, Tourvas N and Aravanopoulos FA** (2022) Genetic diversity of silver fir (*Abies alba*) and European beech (*Fagus sylvatica*) populations from the South-Eastern limits of their natural distribution. *Environmental Sciences Proceedings* 22, 37.
- Davidson R** (1990) Patterns of variation in Pacific silver fir on Vancouver Island: a summary. In Joint Meeting of Western Forest Genetics Association and IUFR0 Working Parties S2.02-05, 06, 12 and 14, 20-24 Aug. 1990, Olympia, Washington, pp. 257–270.
- Dobrowolska D, Bončina A and Klumpp R** (2017) Ecology and silviculture of silver fir (*Abies alba* Mill.): a review. *Journal of Forest Research* 22, 326–335.
- Doyle JJ and Doyle JL** (1990) Isolation of plant DNA from fresh tissue. *Focus* 12, 39–40.
- Fady B and Westfall RD** (1997) Mating system parameters in a natural population of *Abies borisii regis* Mattfeld. *Annals of Forest Science* 54, 643–647.
- Goncharenko GG and Padutov VE** (1995) Genetic structure, taxonomic and phylogenetic relationships in firs occurring in the CIS (former Soviet Union). *Doklady Akademii Nauk* 342, 122–126, (in Russian with English Abstract).
- Hansen OK, Vendramin GG, Sebastiani F and Edwards KJ** (2005) Development of microsatellite markers in *Abies nordmanniana* (Stev.) Spach and cross-species amplification in the *Abies* genus. *Molecular Ecology Notes* 5, 784–787.
- Hrivnák M, Paule L, Krajmerová D, Kulač Š, Ševik H, Turna I, Tvaauri I and Gömöry D** (2017) Genetic variation in Tertiary relics: the case of eastern-Mediterranean *Abies* (Pinaceae). *Ecology and Evolution* 7, 10018–10030.
- Jahnke G, Smidla J and Poczar P** (2022) MolMarker: a simple tool for DNA fingerprinting studies and polymorphic information content calculation. *Diversity* 14, 497.
- Kormuťák A and Lindgren D** (1996) Mating system and empty seeds in silver fir (*Abies alba* Mill.). *Forest Genetics* 3, 231–235.
- Krutovsky KV, Putintseva YA, Oreshkova NV, Bondar EI, Sharov VV and Kuzmin DA** (2019) Postgenomic technologies in practical forestry: development of genome-wide markers for timber origin identification and other applications. *Forest Engineering Journal* 9, 9–16.
- Kuzmin DA, Feranchuk SI, Sharov VV, Cybin AN, Makolov SV, Putintseva YA, Oreshkova NV and Krutovsky KV** (2019) Stepwise large genome assembly approach: a case of Siberian larch (*Larix sibirica* Ledeb.). *BMC Bioinformatics* 20(Suppl. 1), 37.
- Larionova AY and Ekart AK** (2005) Genetic structure and differentiation of Siberian fir populations located at varied elevations in Western Sayan. *Ecological Genetics* 3, 22–27, (in Russian with English Abstract).
- Larionova AY, Ekart AK and Kravchenko AN** (2007) Genetic diversity and population structure of Siberian fir (*Abies sibirica* Ledeb.) in Middle Siberia, Russia. *Eurasian Journal of Forest Research* 10, 185–192.
- Leitch IJ, Johnston E, Pellicer J, Hidalgo O and Bennett MD** (2019) Gymnosperm DNA C-values database (release 6.0, Apr 2019) <https://cvalues.science.kew.org>.
- Martins WS, Lucas DCS, Neves KFS and Bertoli DJ** (2009) WebSat – a web software for microsatellite marker development. *Bioinformatics* 3, 282–283.
- Mauri A, de Rigo D and Caudullo G** (2016) *Abies alba* in Europe: distribution, habitat, usage, and threats. In San-Miguel-Ayán J, De Rigo D, Caudullo G, Houston Durrant T and Mauri A (eds), *European Atlas of Forest Tree Species*. Luxembourg, Luxembourg: Publications Office of the European Union, pp. 48–49. Available at <https://doi.org/10.2788/038466>.
- Mittton JB** (1992) The dynamic mating systems of conifers. *New Forest* 6, 197–216.
- Mosca E, Cruz F, Gómez Garrido J, Bianco L, Rellstab C, Brodbeck S, Csilléry K, Fady B, Fladung M, Fussi B, Gömöry D, González-Martínez SC, Grivet D, Gut M, Hansen OK, Heer K, Kaya Z, Krutovsky KV, Kersten B, Liepelt S, Opgenoorth L, Sperisen C, Ullrich KK, Vendramin GG, Westergren M, Ziegenhagen B, Alioto T, Gugerli F, Heinze B, Höhn M, Troggio M and Neale DB** (2019) A reference genome sequence for the European silver fir (*Abies alba* Mill.): a community-generated genomic resource. *G3: Genes Genomes Genetics* 9, 2039–2049.
- Muona O** (1990) Population genetics in forest tree improvement. In Brown AHD, Clegg MT, Kahler AL and Weir BS (eds), *Plant Population Genetics, Breeding, and Genetic Resources*. USA, Sunderland, Massachusetts: Sinauer Associates Inc., pp. 282–298.
- Neale DB and Adams WT** (1985) Allozyme and mating-system variation in balsam fir (*Abies balsamea*) across a continuous elevational transect. *Canadian Journal of Botany* 63, 2448–2453.
- Nurk S, Koren S, Rhie A, Rautiainen M, Bizikadze AV, Mikheenko A, Vollger M.R., Altemose N, Uralsky L, Gershman A, Aganezov S, Hoyt SJ, Diekhans M, Logsdon GA, Alonge M, Antonarakis SE, Borchers M, Bouffard GG, Brooks SY, Caldas GV, Chen N-C, Cheng H, Chin C-S, Chow W, de Lima LG, Dishuck PC, Durbin R, Dvorkina T, Fiddes IT, Formenti G, Fulton RS, Fungtammasan A, Garrison E, Grady PGS, Graves-Lindsay TA, Hall IM, Hansen NF, Hartley GA, Haukness M, Howe K, Hunkapiller MW, Jain C, Jain M, Jarvis ED, Kerpedjiev P, Kirsche M, Kolmogorov M, Korlach J, Kremitzki M, Li H, Maduro VV, Marschall T, McCartney AM, McDaniel J, Miller DE, Mullikin JC, Myers EW, Olson ND, Paten B, Peluso P, Pevzner PA, Porubsky D, Potapova T, Rogaev EI, Rosenfeld JA, Salzberg SL, Schneider VA, Sedlazeck FJ, Shafin K, Shew CJ, Shumate A, Sims Y, Smit AFA, Soto DC, Sovic I, Storer JM, Streets A, Sullivan BA, Thibaud-Nissen F, Torrance J, Wagner J, Walenz BP, Wenger A, Wood JMD, Xiao C, Yan SM, Young AC, Zarate S, Surti U, McCoy RC, Dennis MY, Alexandrov IA, Gerton JL, O'Neill RJ, Timp W, Zook JM, Schatz MC, Eichler EE, Miga KH and Phillipp AM** (2022) The complete sequence of a human genome. *Science (New York, N.Y.)* 376, 44–53.
- Nystedt B., Street N.R., Wetterbom A., Zuccolo A, Lin YC, Scofield DG, Vezzi F, Delhomme N, Giacomello S, Alexeyenko A, et al.** (2013) The Norway spruce genome sequence and conifer genome evolution. *Nature* 497, 579–584.
- Oreshkova NV, Putintseva YA, Sharov VV, Kuzmin DA and Krutovsky KV** (2017) Development of microsatellite genetic markers in Siberian larch (*Larix sibirica* Ledeb.) based on the *de novo* whole genome sequencing. *Russian Journal of Genetics* 53, 1194–1199.
- Oreshkova NV, Bondar EI, Putintseva YA, Sharov VV, Kuzmin DA and Krutovsky KV** (2019) Development of nuclear microsatellite markers with long (tri-, tetra-, penta-, and hexanucleotide) motifs for three larch species based on the *de novo* whole genome sequencing of Siberian larch (*Larix sibirica* Ledeb.). *Russian Journal of Genetics* 55, 444–450.
- Peakall R and Smouse PE** (2006) GENEALX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6, 288–295.
- Peakall R and Smouse PE** (2012) GENEALX 6.5: genetic analysis in Excel. Population genetic software for teaching and research. An update. *Bioinformatics (Oxford, England)* 28, 2537–2539.
- Postolache D, Leonarduzzi C, Piotti A, Spanu I, Roig A, Fady B, Roschanski A, Liepelt S and Vendramin GG** (2014) Transcriptome versus genomic microsatellite markers: highly informative multiplexes for genotyping *Abies alba* Mill. and congeneric species. *Plant Molecular Biology Reporter* 32, 750–760.
- Schroeder S** (1989) Outcrossing rates and seed characteristics in damaged natural populations of *Abies alba* Mill. *Silvae Genetica* 38, 185–189.
- Semerikov VI, Semerikova SA, Khrunyk YY and Putintseva YA** (2022) Sequence capture of mitochondrial genome with PCR-generated baits provides new insights into the biogeography of the Genus *Abies* Mill. *Plants* 11, 762.
- Semerikova SA and Semerikov VI** (2006) Genetic variation and population differentiation in Siberian fir *Abies sibirica* Ledeb. inferred from allozyme markers. *Russian Journal of Genetics* 42, 636–644.
- Semerikova SA and Semerikov VI** (2014) Molecular phylogenetic analysis of the Genus *Abies* (Pinaceae) based on the nucleotide sequence of chloroplast DNA. *Russian Journal of Genetics* 50, 7–19.
- Semerikova SA and Semerikov VI** (2016) Phylogeny of firs (genus *Abies*, Pinaceae) based on multilocus nuclear markers (AFLP). *Russian Journal of Genetics* 52, 1164–1175.

- Semerikova S, Khrunyk Y, Lacoux M and Semerikov V** (2018) From America to Eurasia: a multigenomes history of the genus *Abies*. *Molecular Phylogenetics and Evolution* **125**, 14–28.
- Shea KL** (1987) Effects of population structure and cone production on outcrossing rates in Engelmann spruce and subalpine fir. *Evolution* **41**, 124–136.
- Świercz A, Świątek B and Pietrzykowski M** (2022) Changes in the concentrations of trace elements and supply of nutrients to silver fir (*Abies alba* Mill.) needles as a bioindicator of industrial pressure over the past 30 years in Świętokrzyski National Park (Southern Poland). *Forests* **13**, 718.
- Van Oosterhout C, Hutchinson WF, Wills DPM and Shipley P** (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**, 353–358.
- Wang X, Lu P and Luo Z** (2013) GMATo: a novel tool for the identification and analysis of microsatellites in large genomes. *Bioinformatics* **9**, 541–544.
- Zaitsev GA, Kulagin AY and Davydychev AN** (2018) The particularities of the growth of Siberian fir (*Abies sibirica* Ledeb.) in the first stages of ontogeny in conifer forests (Ufa plateau, Pre-Ural). *Trees* **32**, 511–518.