Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

EHPGS: A Statistical Package for Evaluation of Hybrid Performance in Plant Breeding via Genomic Selection

Szu-Ping Chen
National Taiwan University
Chih-Wei Tung
National Taiwan University
Pei-Hsien Wang
National Taiwan University
Chen-Tuo Liao (ctliao@ntu.edu.tw)
National Taiwan University

Article

Keywords: general combining ability, genomic estimated breeding value, genomic prediction, heterosis, specific combining ability

Posted Date: May 9th, 2023

DOI: https://doi.org/10.21203/rs.3.rs-2890196/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on July 27th, 2023. See the published version at https://doi.org/10.1038/s41598-023-39434-6.

EHPGS: A Statistical Package for Evaluation of Hybrid Performance in Plant Breeding via Genomic Selection

Szu-Ping Chen, Chih-Wei Tung, Pei-Hsien Wang, Chen-Tuo Liao*

Department of Agronomy, National Taiwan University, Taipei, Taiwan.

*Corresponding author: Chen-Tuo Liao, ctliao@ntu.edu.tw

Abstract

Hybrid breeding employs heterosis, which could potentially improve the yield and quality of a crop. Genomic selection (GS) is a promising approach for the selection of quantitative traits in plant breeding. The main objectives of this study are to (i) propose a GS-based approach to identify potential parental lines and superior hybrid combinations from a breeding population, which is composed of hybrids produced by a half diallel mating design; (ii) develop a software package for users to carry out the proposed approach. An R package, designated EHPGS, was generated to facilitate the employment of the genomic best linear unbiased model considering additive plus dominance marker effects for the hybrid performance evaluation. The R package contains a Bayesian statistical algorithm for calculating genomic estimated breeding value (GEBVs), GEBV-based specific combining ability, general combining ability, mid-parent heterosis, and better-parent heterosis. Three datasets that have been published in literature, including pumpkin (*Cucurbita maxima*), maize (*Zea mays*), and wheat (*Triticum aestivum* L.), were analyzed to illustrate the use of EHPGS.

Keywords: general combining ability, genomic estimated breeding value, genomic prediction, heterosis, specific combining ability

Introduction

Hybrid plant breeding can potentially be used as a method that employs heterosis to boost yield stability, allow the combination of dominant major genes, and offer a built-in plant variety protection system (Longin et al., 2012). Several field, vegetable, and flower crops use hybrids, including maize, sorghum, and sunflower. Interestingly,

hybrid rice has been adopted and hybrid wheat research is drawing new attention (Guo et al., 2019). Therefore, it is important and challenging to develop a highly efficient approach for identifying potential parental lines and superior hybrid combinations from many possible candidates. To create such an approach, we constructed a prediction model to screen out the desired individuals based on genomic selection (GS) (Jannink et al. 2010). To facilitate practical application, we also generated a software package to implement our proposed GS-based approach.

Diallel mating designs have been traditionally used to evaluate the combining ability of parental lines in hybrids and to predict hybrid performance on quantitative traits of interest. To analyze diallel crosses, the total genetic variability is often separated into the general combining ability (GCA) for parental lines, and the specific combining ability (SCA) for hybrid combinations. The GCA is a measure of additive gene activity that relates to the average performance of a particular inbred line in hybrid combinations. The SCA is a measure of general combining ability that links to the non-additive effects, including dominance and epistatic effects. In addition, mid-parent heterosis (MPH) is defined as the difference between a hybrid's performance and the average performance of its parental lines, while better-parent heterosis (BPH) is defined as hybrid performance superior to the higher or better parental line (Falconer and Mackay, 1996). However, the number of crossing combinations can be prohibitively high for extensive testing in a field experiment.

Due to the availability of high-density single nucleotide polymorphism (SNP) markers across an entire genome, GS becomes a promising approach to reduce cost and accelerate breeding cycles for plant breeding (Heffner et al. 2010; Nakaya and Isobe, 2012). The conceptual basis of GS is the utilization of a training population with known phenotype and genotype data to build a prediction model that uses individuals with known genotype data only to predict genomic estimated breeding values (GEBVs) (Meuwissen et al. 2001). This GS-based approach has been applied to predict hybrid performance for several crops, such as barley (Schmid and Thorwarth, 2014), maize (Technow et al., 2012, 2014), rice (Xu et al., 2014; Wang et al., 2016), wheat (Zhao et al., 2013; Haile et al., 2018), and pumpkin (Wu et al., 2019). More recently, Sabouri and Sajadi (2022) evaluated hybrid rice performance based on parental characteristics using artificial neural networks, adaptive neuro-fuzzy inference system, and support vector machine.

In this study, we obtained the required estimates for hybrid performance evaluation based on a GBLUP model, which took both additive and dominance marker effects into account. The GBLUP model was built based on a training population with known phenotype and genotype data. Here, we proposed a Bayesian statistical algorithm for the parameter estimation. Three datasets that have been published in literature, including pumpkin (*Cucurbita maxima*), maize (*Zea mays*), and wheat (*Triticum aestivum* L.), were reanalyzed to illustrate the application of our proposed approach.

Materials and Methods

The genomic selection-based approach

The GBLUP model

The GBLUP model considering additive plus dominance effects can be described as follows:

$$\boldsymbol{y} = \mathbf{1}_n\mu + \boldsymbol{g}_A + \boldsymbol{g}_D + \boldsymbol{e}, \quad (1)$$

where $\boldsymbol{y}$ is the vector of the phenotypic values; $\mathbf{1}_n$ is the unit vector of length n (here n is the number of phenotypic values); $\boldsymbol{g}_A$ is the vector of genotypic values for the additive effects; $\boldsymbol{g}_D$ is the vector of genotypic values for the dominance effects; and $\boldsymbol{e}$ is the vector of random errors. It is assumed that $\boldsymbol{g}_A$, $\boldsymbol{g}_D$, and $\boldsymbol{e}$ are mutually independent and follow multivariate normal distributions, denoted by $\boldsymbol{g}_A \sim N(\mathbf{0}, \sigma_A^2\boldsymbol{K}_A)$, $\boldsymbol{g}_D \sim N(\mathbf{0}, \sigma_D^2\boldsymbol{K}_D)$, and $\boldsymbol{e} \sim N(\mathbf{0}, \sigma_e^2\boldsymbol{I}_n)$. Here, $\boldsymbol{K}_A = \frac{1}{p}(\boldsymbol{X}_A\boldsymbol{X}_A^T)$ is the genomic relationship matrix for the additive effects, abbreviated as A-GRM; the variance component σ_A^2 represents the cumulative variability of additive marker effects, abbreviated as A-VC; $\boldsymbol{K}_D = \frac{1}{p}(\boldsymbol{X}_D\boldsymbol{X}_D^T)$ is the genomic relationship matrix for the dominance effects, abbreviated as D-GRM; and the variance component σ_D^2 represents the cumulative variability of the dominance marker effects, abbreviated as D-VC. Moreover, $\boldsymbol{X}_A$ and $\boldsymbol{X}_D$ are the standardized marker score matrices for the additive effects and dominance effects, respectively, and p is the number of the SNP markers.

Estimation for GEBVs and genomic heritability

Let $\hat{\mu}$ be the best linear unbiased estimate (BLUE) for μ, $\hat{\boldsymbol{g}}_A$ be the BLUP for $\boldsymbol{g}_A$,

and $\widehat{\boldsymbol{g}}_D$ be the BLUP for $\boldsymbol{g}_D$. Then, $\hat{\mu}$, $\widehat{\boldsymbol{g}}_A$, and $\widehat{\boldsymbol{g}}_D$ can be obtained from the Henderson's equations (Henderson, 1975):

$$\begin{bmatrix} n & \mathbf{1}_n^T & \mathbf{1}_n^T \\ \mathbf{1}_n & \boldsymbol{I}_n + \boldsymbol{K}_A^{-1}\lambda_A & \boldsymbol{I}_n \\ \mathbf{1}_n & \boldsymbol{I}_n & \boldsymbol{I}_n + \boldsymbol{K}_D^{-1}\lambda_D \end{bmatrix} \begin{bmatrix} \hat{\mu} \\ \widehat{\boldsymbol{g}}_A \\ \widehat{\boldsymbol{g}}_D \end{bmatrix} = \begin{bmatrix} \mathbf{1}_n^T\boldsymbol{y} \\ \boldsymbol{y} \\ \boldsymbol{y} \end{bmatrix}, \quad (2)$$

where $\lambda_A = \sigma_e^2/\sigma_A^2$ and $\lambda_D = \sigma_e^2/\sigma_D^2$. Here, λ_A and λ_D can be replaced with suitable estimates for σ_e^2, σ_A^2, and σ_D^2, respectively denoted by $\hat{\sigma}_e^2$, $\hat{\sigma}_A^2$, and $\hat{\sigma}_D^2$. The estimate for genomic heritability was then obtained as:

$$h^2 = \frac{\hat{\sigma}_A^2 + \hat{\sigma}_D^2}{\hat{\sigma}_A^2 + \hat{\sigma}_D^2 + \hat{\sigma}_e^2}. \quad (3)$$

In this study, the breeding population was composed of all possible hybrid combinations in a half diallel mating design. Let $\boldsymbol{K}_A^{(bp)}$ and $\boldsymbol{K}_D^{(bp)}$ respectively denote the A-GRM and D-GRM between the breeding population and the training population, and let $\widehat{\boldsymbol{g}}_A^{(bp)}$ and $\widehat{\boldsymbol{g}}_D^{(bp)}$ denote the BLUPs for the breeding population of additive and dominance effects, respectively. From Henderson (1977), $\widehat{\boldsymbol{g}}_A^{(bp)}$ and $\widehat{\boldsymbol{g}}_D^{(bp)}$ can be obtained as:

$$\widehat{\boldsymbol{g}}_A^{(bp)} = \boldsymbol{K}_A^{(bp)}\boldsymbol{K}_A^{-1}\widehat{\boldsymbol{g}}_A, \quad (4)$$

and

$$\widehat{\boldsymbol{g}}_D^{(bp)} = \boldsymbol{K}_D^{(bp)}\boldsymbol{K}_D^{-1}\widehat{\boldsymbol{g}}_D. \quad (5)$$

The GEBVs for the individuals in the breeding population were then predicted by:

$$\widehat{\boldsymbol{y}}^{(bp)} = \mathbf{1}_{N_1}\hat{\mu} + \widehat{\boldsymbol{g}}_A^{(bp)} + \widehat{\boldsymbol{g}}_D^{(bp)}, \quad (6)$$

where N_1 is the number of hybrid combinations in the breeding population. Here, $N_1 = C_2^{N_0}$ with N_0 as the number of parental lines.

Estimation for GCA, SCA, MPH, and BPH

Let GCA_i and GCA_j denote the GCAs for the parental lines P_i and P_j, respectively; and SCA_{ij} denote the SCA for their hybrid combination $P_i \otimes P_j$; and $g_A^{(ij)}$ and $g_D^{(ij)}$ are the BLUPs for $P_i \otimes P_j$ of additive and dominance effects, respectively. From

Werner et al. (2018),

$$g_A^{(ij)} = GCA_i + GCA_j, \tag{7}$$

and

$$g_D^{(ij)} = SCA_{ij}. \tag{8}$$

From Eq. (8), the BLUP for SCA_{ij} was obtained as:

$$\widehat{SCA}_{ij} = \hat{g}_D^{(ij)}. \tag{9}$$

Let

$$\bar{G}_A^{(i)} = \frac{\sum_{j\neq i}^{N_0} \hat{g}_A^{(ij)}}{N_0 - 1} \tag{10}$$

and

$$\bar{G}_A = \frac{\sum_{i=1}^{N_0} \sum_{j\neq i}^{N_0} \hat{g}_A^{(ij)}}{N_1} \tag{11}$$

where $\bar{G}_A^{(i)}$ is the average over the additive genotypic values of the parental line *i*, and $\bar{G}_A$ is the average over all of the additive genotypic values. From Eq. (7), the BLUP for GCA_i is given by:

$$\widehat{GCA}_i = \frac{(N_0-1)\bar{G}_A^{(i)}}{N_0 - 2} - \frac{N_0 \bar{G}_A}{2(N_0-2)}. \tag{12}$$

From Wu et al. (2019), the GEBV-based MPH and BPH for $P_i \otimes P_j$ can be estimated by:

$$\widehat{MPH}_{ij} = \widehat{SCA}_{ij} \tag{13}$$

and

$$\widehat{BPH}_{ij} = \widehat{SCA}_{ij} - \left|\widehat{GCA}_i - \widehat{GCA}_j\right| \tag{14}$$

where $\left|\widehat{GCA}_i - \widehat{GCA}_j\right|$ is the absolute value of $(\widehat{GCA}_i - \widehat{GCA}_j)$. Under the positive heterosis assumption, the value of MPH or BPH is larger, and the heterosis of the hybrid combination is stronger.

142

143 **The Bayesian statistical algorithm**

144 For a given training set with known phenotype and genotype data, a Bayesian Gibbs
145 sampling (GBS) algorithm modified from that presented in Xavier et al. (2016) was
146 used to estimate the required parameters. The algorithm can be described as follows.
147 Step 1: Set initial values for the parameters in the model.
148 The default values are given by:
149 $\mu = \bar{y}$ (the sample mean of the phenotypic values), $\boldsymbol{g}_A = \boldsymbol{g}_D = \mathbf{0}$, $\sigma_e^2 = 1$, and $\sigma_A^2 =$
150 $\sigma_D^2 = 0.5$.
151
152 Step 2: Rewrite Eq. (2) as

153 $$\begin{bmatrix} \boldsymbol{C}_{11} & \boldsymbol{C}_{12} & \boldsymbol{C}_{13} \\ \boldsymbol{C}_{21} & \boldsymbol{C}_{22} & \boldsymbol{C}_{23} \\ \boldsymbol{C}_{31} & \boldsymbol{C}_{32} & \boldsymbol{C}_{33} \end{bmatrix} \begin{bmatrix} \boldsymbol{g}_1 \\ \boldsymbol{g}_2 \\ \boldsymbol{g}_3 \end{bmatrix} = \begin{bmatrix} \boldsymbol{\gamma}_1 \\ \boldsymbol{\gamma}_2 \\ \boldsymbol{\gamma}_3 \end{bmatrix}. \quad (15)$$

154 Update $\boldsymbol{g}_i$ by $\boldsymbol{g}_i \sim N(\boldsymbol{g}_i^*, \sigma_e^2 \boldsymbol{C}_{ii}^{-1})$, where $\boldsymbol{g}_i^* = \boldsymbol{C}_{ii}^{-1}(\boldsymbol{\gamma}_i - \boldsymbol{C}_{i,-i}\boldsymbol{g}_{-i})$ for i=1, 2, 3.
155 Here, $\boldsymbol{C}_{i,-i}$ denotes $\boldsymbol{C}_{i,j}$ for all $j \neq i$; and $\boldsymbol{g}_{-i}$ is $\boldsymbol{g}_j$ for all $j \neq i$.
156
157 Step 3: Calculate the vector of residuals as: $\boldsymbol{e} = \boldsymbol{y} - \boldsymbol{g}_1 - \boldsymbol{g}_2 - \boldsymbol{g}_3$.
158
159 Step 4: Update σ_e^2 as $\sigma_e^2 = (\boldsymbol{e}^T\boldsymbol{e} + S^* v^*)/\chi^2_{n+v^*}$, where $\chi^2_{n+v^*}$ is the chi-square
160 random variate with $n + v^*$ degrees of freedom; $S^* = 0.5V$ with V as the sample
161 variance of the values in $\boldsymbol{y}$; and $v^* = 5$.
162
163 Step 5: Update σ_A^2 as $\sigma_A^2 = (\boldsymbol{g}_A^T \boldsymbol{K}_A^{-1} \boldsymbol{g}_A + S^* v^*)/\chi^2_{n+v^*}$; and σ_D^2 as $\sigma_D^2 =$
164 $(\boldsymbol{g}_D^T \boldsymbol{K}_D^{-1} \boldsymbol{g}_D + S^* v^*)/\chi^2_{n+v^*}$.
165
166 Step 6: Update the equations in Eq. (15) with $\lambda_A = \sigma_e^2/\sigma_A^2$, and $\lambda_D = \sigma_e^2/\sigma_D^2$.
167
168 Step 7: Repeat Steps 2–6 K times to generate a series of results over the K iterations,
169 which are denoted by:
170 $\mu^{(k)}$, $\boldsymbol{g}_A^{(k)}$, $\boldsymbol{g}_D^{(k)}$, $\sigma_A^{2(k)}$, $\sigma_D^{2(k)}$, and $\sigma_e^{2(k)}$ for $k = 1, 2, \cdots,$ K.
171

Step 8: Discard the results from the first $0.9K$ iterations, and average the results from the remaining 0.1K iterations. The number of iterations K is defaulted as 10,000.

Step 9: Repeat Steps 1–8 M times to generate M sets of the averages of the parameters generated from Step 8. The number of chains M is defaulted as five.

Step 10: Average the resulting mean values of the parameters over the M chains, and the resulting averages are treated as the estimates for the parameters.

An R package called as EHPGS generated for executing the proposed approach is available from GitHub (https://github.com/spcspin/EHPGS).

The genome datasets

Three datasets that have been published in literature were reanalyzed to illustrate the use of EHPGS.

Pumpkin dataset

Wu et al. (2019) studied a pumpkin dataset which contained 119 intra-crossing hybrid combinations of *C. maxima* with phenotypic values for fruit weight (FWT) (kg). The phenotype data were historical data collected from 1988 to 2016. All the trials were conducted at a single location experiment in southern area of Taiwan. Every hybrid had six to ten observations at each time point, and the average of them was used as the phenotypic observation for the hybrid of the year. Because the phenotypic values of every hybrid were observed for more than one year, the different year effects were therefore removed based on the assumption that they were random effects following a normal distribution.

The germplasm collection of the pumpkin set consisted of 320 parental lines, which were classified into three clusters: *C. maxima* with 142 inbred lines, *C. pepo* with 60 inbred lines and *C. moschata* with 118 inbred lines. After SNP calling, 76,815 SNPs were extracted from the parental lines, and only 4,521 SNPs remaining for *C. maxima* after the filtering by missing rate ≥ 0.05, minor allele frequency (MAF) < 0.05, and a series of operations for determining linkage disequilibrium (LD) blocks. The 142 inbred lines produced $C_2^{142} = 10{,}011$ potential hybrid combinations in a half diallel mating design.

For the additive effects, the SNP at each locus was coded as -1, 0, or 1 for the homozygote of the minor allele, the heterozygote, and the homozygote of the major allele, respectively. For the dominance effects, the marker score was coded as 1 for the heterozygote, and 0 for both homozygotes. The means adjusted from the year effects for the 119 *C. maxima* hybrids were used to build a GBLUP model for evaluating the performance of the 10,011 hybrid combinations.

Maize dataset

Guo et al. (2019) analyzed a maize dataset to study the optimal designs for GS in hybrid crops, which consisted of 276 hybrids derived from 24 parental lines in a half diallel mating design. The 24 diverse parents were classified into two groups according to the germplasm origin and a principal component analysis. The two groups were (i) the temperate and mixed (TM) group, consisting of 11 inbred lines (i.e., B73, B97, Ky21, M162W, Mo17, MS71, Oh43, OH7B, M37W, Mo18W, and Tx303); and (ii) the tropical and sub-tropical (TS) group consisting of the remaining 13 inbred lines (i.e., CML52, CML69, CML103, CML228, CML247, CML277, CML322, CML333, Ki3, Ki11, NC350, NC358, and Tzi8). There were $C_2^{11} = 55$ hybrid combinations in the TM group, $C_2^{13} = 78$ hybrids in the TS group, and 11x13=143 hybrids between the two groups. Three trait values, flowering time, ear height, and grain yield (YLD) (Mg/ha), were evaluated for all of the hybrids at two locations (i.e., Columbia, MO and Clayton, NC) in 2005 and 2006. In this study, the combined BLUP values from the two locations for YLD were evaluated.

Genotype data for the 24 inbred lines were extracted from the Maize HapMap V2 (Chia et al., 2012) at www.panzea.org, which consisted of 10,296,310 SNP markers. The SNP markers were first filtered by missing rate $\geq$ 0.05 and MAF $\leq$ 0.1, resulting in 134,726 SNPs remaining. Missing genotypes were then imputed with the homozygote of the major allele. To screen out reliable SNPs for building a GBLUP model, the retained SNPs were further filtered by LD blocks. The LD parameter r^2 (i.e., the squared Pearson's correlation coefficient) of the SNPs for each chromosome was estimated using TASSEL5.2.41 (Bradbury et al., 2007) with a sliding window = 10. A smooth function between r^2 and the physical distance (bp) was built using an R function *loess.smooth*( ) with a second-degree locally weighted polynomial regression. The LD decay of ten chromosomes is displayed in Figure S1 of the Supplementary

Materials. Filtering the 134,726 SNP markers by the LD block sizes if r^2 approached 0.2, resulting in 46,134 SNPs remaining. The SNP coding was the same as for the pumpkin dataset. A SNP was also deleted if its corresponding column for the dominance effects was a zero vector. Finally, 30,239 SNP markers were retained for further analysis. In this study, all 276 hybrids with known trait values were used as the training population for the prediction model construction.

Wheat dataset

Zhao et al. (2015) investigated the genome-based establishment of a high-yielding heterotic pattern for hybrid wheat breeding, their study was based on 135 advanced elite winter wheat lines. A set of 1604 wheat hybrids produced from crosses among the 15 male lines and 120 female lines were then evaluated for grain yield (YLD) (Mg/ha) in 11 environments. Grain yield data for all $C_2^{135} = 9045$ unique hybrids were predicted based on those of the phenotyped individuals. For the genotype data, the 135 lines were fingerprinted by using a 90,000 SNP array based on an Illumina Infinium array. After quality tests, 17,372 high-quality SNP markers were retained.

Guo et al. (2019) extracted a half diallel of 2556 hybrids, which were crossed from 72 lines selected from the original 135 elite wheat lines, for their study on optimal designs for GS. In this study, an optimal training population with 600 individuals, determined by the r-score criterion presented in Ou and Liao (2019), was used to build the GBLUP model for the performance evaluation on the 2556 hybrid combinations.

Results and Discussion

Pumpkin dataset

By the half diallel mating design, the 119 parental lines produced $C_2^{142} = 10{,}011$ hybrid combinations in the breeding population. For illustration purposes, we only reported the top 25 superior hybrid combinations with the largest GEBVs, together with their SCAs, MPHs, and BPHs in Table 1; and the top 10 potential parental lines with the largest GCAs in Table 2. Table 1 illustrates the important finding that both MPH_{ij} and BPH_{ij} are greater than 0 for all of the selected hybrids, showing that they had better performance in FWT than both of their parents. More interestingly, every superior hybrid presented in Table 1 was derived from one or two of the potential parental lines

presented in Table 2. Particularly, P026, the parental line with the highest GCA, involved the top 11 hybrids with the greatest GEBVs among the 25 selected hybrids.

The estimates for the variance components and genomic heritability are shown in Table 3. From the table, the estimates of the A-VC, D-VC, and genomic heritability are given by $\hat{\sigma}_A^2 = 0.306$, $\hat{\sigma}_D^2 = 0.159$, and $h^2 = 0.807$. The high heritability explains why the values of MPH_{ij} and BPH_{ij} in Table 1 are all positive, and indicates strong heterosis in FWT among the intra-crossing hybrid combinations of *C. maxima*.

Maize dataset

There were $C_2^{24} = 276$ hybrid combinations derived from the 24 parental lines. For illustration purposes, we reported the top 15 superior hybrids with the largest GEBVs, together with their SCAs, MPHs, and BPHs in Table 4; and the top 5 potential parental lines with the largest GCAs in Table 5. From Table 4, both MPH_{ij} and BPH_{ij} are greater than 0 for all of the selected hybrids, showing that they had better performance in YLD than both of their parents. A total of 12 out of the 15 selected hybrids belong to the inter-crossing group between TM and TS. From Table 5, the top five parental lines with the greatest GCAs are CML228, CML103, Mo17, B97, and B73, and involved all of the 15 superior parental lines, with the exception of MS71⊗Tzi8.

The estimates for the variance components and genomic heritability are also displayed in Table 3. From the table, the estimates of the estimates of the A-VC, D-VC, and genomic heritability are given by $\hat{\sigma}_A^2 = 0.434$, $\hat{\sigma}_D^2 = 0.420$, and $h^2 = 0.415$, explaining why the values of MPH_{ij} and BPH_{ij} in Table 4 are all positive, and showing that there is an obvious heterosis in YLD within the breeding population.

Wheat dataset

By the half diallel mating design, the 72 parental lines produced $C_2^{72} = 2556$ hybrid combinations in the breeding population. For illustration purposes, we only reported the top 20 superior hybrids with the largest GEBVs, together with their SCAs, MPHs, and BPHs in Table 6; and the top 10 potential parental lines with the largest GCAs in Table 7. The estimates for the variance components and genomic heritability are also displayed in Table 3. From Table 6, all of the MPH_{ij} are greater than 0, showing that

they had a larger YLD than the mean YLD of their parents. Most of the BPH_{ij} are noticeably smaller than MPH_{ij}, probably because the additive effects ($\hat{\sigma}_A^2$ =0.066, Table 3) were stronger than the dominance effects ($\hat{\sigma}_D^2 = 0.014$, Table 3). Moreover, 11 of the 20 BPHs are negative, showing that the corresponding hybrids were inferior to their better-parents. Every superior hybrid presented in Table 6 was derived from one or two of the potential parental lines presented in Table 7. Particularly, F102, the parental line with the highest GCA (Table 7), involved 17 of the 20 selected superior hybrids (Table 6).

The correlation between phenotypic values and GEBVs

Scatter plots of all available phenotypic values (119, 276, and 2556 individuals in the pumpkin, maize, and wheat datasets, respectively) and their GEBVs in each dataset are displayed in Figures 1–3. The respective Pearson's correlation coefficients are 0.9691, 0.6786, and 0.9445. From the figures, most of the selected superior hybrids appeared in the upper right-hand corners, meaning that the selected hybrids with higher GEBVs also have higher actual phenotypic values. This is a valuable result because phenotypic selection is usually costly and time-consuming for selective breeding. The great consistency exists between the results of genomic selection and phenotypic selection, supporting that the proposed GS-based approach can be recommended for practical applications.

A comparison of results obtained from different approaches

Wu et al. (2019) analyzed the pumpkin dataset using a two-stage approach. The authors first estimated GEBVs, SCAs, GCAs, MPHs, and BPHs based on a whole genome regression model using Bayes C estimation (Perez and de los Campos, 2014). Then, they calculated A-GRM and D-GRM by the formulas presented in Endelman and Jannink (2012) and Su et al. (2012), respectively, and obtained the restricted maximum likelihood estimates for the variance components and genomic heritability based on a GBLUP model using the R package sommer (Covarrubias-Pazaran, 2016). The top 25 superior hybrid combinations identified by the approach of Wu et al. (2019) together with those identified by ours are displayed in Table S1 of the Supplementary Materials, and the corresponding identified 10 potential parental lines are displayed in Table S2. Both sets of the results are highly consistent with each other. From Table S1, 18 hybrids were in common among the 25 hybrids selected by each approach, and the top six

hybrids with the highest GEBVs were the same, even though the order was slightly different. Table S2 indicates seven potential parental lines in common among the 10 selected by each approach. The variance components for additive, dominance, random error effects, and genomic heritability estimated by Wu et al. (2019) are 0.195, 0.119, 0.066, and 0.826, respectively. The corresponding estimates by our approach are 0.306, 0.159, 0.111 and 0.807. Even though the two corresponding estimates are different from each other, the two estimates of the genomic heritability are fairly close.

The Bayesian reproducing kernel Hilbert space (RKHS) method in BGLR (Perez and de los Campos, 2014) has been commonly used to perform GEBV prediction for the GBLUP model in Eq. (1). To compare the use of Bayesian RKHS method with our proposed Bayesian algorithm, we re-run the analysis for the three datasets by BGLR. Note that the priors specified in BGLR were the same as ours, the number of iterations was set to 10,000, the number of burn-in was fixed at 9000, and the number of chains was set to five (the BGLR function was repeatedly run five times). These settings are exactly the same as our algorithm in analyzing the datasets. Overall, the results of the identified top parental lines and hybrid combinations between these two methods were highly consistent with each other. Pearson's correlations between GEBVs and phenotypic values for the datasets are displayed in Table 8. From which, our proposed algorithm led to higher Pearson's correlations in pumpkin and the three maize datasets, but almost equal in wheat dataset. Additionally, the estimates for variance components and genomic heritability by using Bayesian RKHS method are displayed in Table 9. In comparison with those obtained from our algorithm (Table 3), BGLR resulted in relatively low genomic heritability. In our experience, the analysis by using BGLR saved much computing time compared to EHPGS.

Conclusion

In this study, a software package called EHPGS was generated for identifying potential parental lines and superior hybrid combinations from a breeding population, which is composed of all possible hybrids produced by a half diallel mating design. Three real datasets of pumpkin (*Cucurbita maxima*), maize (*Zea mays*), and wheat (*Triticum aestivum* L.) were analyzed to illustrate the use of EHPGS.

Data availability statement

All phenotype and genotype datasets that were analyzed in this study can be downloaded from Figshare (https://doi.org/10.6084/m9.figshare.22359883.v2).

Author contributions

SP Chen: prepared Tables 3-9, and Figures 1-3. CW Tung: wrote the original manuscript. PH Wang: prepared Tables 1-2, S1-S2, and Figure S1. CT Liao: wrote the original manuscript. All authors contributed to the article and approved the submitted version.

Funding

This research was supported by the Ministry of Science and Technology, Taiwan (grand number MOST 110-2118-M-002-002-MY2).

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships.

References

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., and Buckler, E. S. (2007). TASSEL: software for association mapping of complex traits in diverse samples. Genet Pop Analysis 23: 2633-2635.

Chia, J. M., Song, C., Bradbury, P. J., Costich, D., de Leon, N., Doebley, J., Elshire, R. J., Gaut, B., Geller, L., and Glaubitz, J. C. (2012). Maize HapMap2 identifies extant variation from a genome in flux. Nat Genet 44: 803-807.

Covarrubias-Pazaran, G. (2016). Genome-assisted prediction of quantitative traits using the R package sommer. PLOS One 11: e0156744.

Endelman, J. B., and Jannink, J. L. (2012). Shrinkage Estimation of the Realized Relationship Matrix. G3: Genes Genomes Genet 2: 1405–1413.

Falconer, D. S., and Mackay, T. F. C. (1996). Introduction to quantitative genetics. 4th ed. San Francisco: Benjamin-Cummings Pub Co.

Guo, T., Yu, X., Li, X., Zhang, H., Zhu, C., Flint-Garcia, S., Michael, D., McMullen, J. B., Holland, S. J., Szalma, R. J., and Yu, W. J. (2019). Optimal Designs for Genomic Selection in Hybrid Crops. Mol Plant 12: 390-401.

Haile, J. K., N'Diaye, A., Clarke, F., Clarke, J., Knox, R., Rutkoski, J., Bassi, F. M., Pozniak, C. J. (2018). Genomic selection for grain yield and quality traits in durum wheat. Mol Breed 38: 75.

Heffner, E. L., Lorenz, A. J., Jannink, J. L., and Sorrells, M. E. (2010). Plant breeding with genomic selection: Gain per unit time and cost. Crop Sci. 50: 1681-1690.

Henderson, C. R. (1975). Best linear unbiased estimation and prediction under a selection model. Biometrics 32: 69-84.

Henderson, C. R. (1977). Best linear unbiased prediction of breeding values not in the model for records. J Diary Sci 60: 783-787.

Jannink, J. L., Lorenz, A. J., and Iwata, H. (2010). Genomic selection in plant breeding: from theory to practice. Brief Funct Genomics 9: 166 -177.

Longin, C. F. H., Muhleisen, J., Maurer, H. P., Zhang, H., Gowda, M., Reif, J. C. (2012). Hybrid breeding in autogamous cereals. Theor Appl Genet 125: 1087-1096.

Meuwissen, T. H. E., Hayes, B. J., and Goddard, M. E. (2001). Prediction of total genetic value using genome-wide dense marker maps. Genetics 157: 1819–1829.

Nakaya, A., and Isobe, S. N. (2012). Will genomic selection be a practical method for plant breeding? Ann Bot 110: 1303-1316.

Ou, J. H., and Liao, C. T. (2019). Training set determination for genomic selection. Theor. Appl. Genet 132: 2781–2792.

Perez, P., and de los Campos, G. (2014). Genome-wide regression and prediction with the BGLR statistical package. Genetics 198: 483–495.

Sabouri, H. and Sajadi, S. J. (2022). Predicting hybrid rice performance using AIHIB model based on artificial intelligence. Sci Rep 12: 9709.

Schmid, K. J., and Thorwarth, P. (2014). Genomic selection in barley breeding. Biotechnological Approaches to Barley Improvement 69: 367-378.

Su, G., Christensen, O. F., Ostersen, T., Henryon, M., and Lund, M. S. (2012). Estimating additive and non-additive genetic variances and predicting genetic merits

using genome-wide dense single nucleotide polymorphism markers. PLOS One 7: e45293.

Technow, F., Riedelsheimer C., Schrag, T. A., Melchinger, A. E. (2012). Genomic prediction of hybrid performance in maize with models incorporating dominance and population specific marker effects. Theor Appl Genet 125: 1181–1194.

Technow, F., Schrag, T. A., Schipprack, W., Bauer, E., Simianer, H., Melchinger, A. E. (2014). Genome properties and prospects of genomic prediction of hybrid performance in a breeding program of maize. Genetics 197: 1343–1355.

Wang, X., Li, L., Yang, Z., Zheng, X., Yu, S., Xu, C., Hu, Z. (2016). Predicting rice hybrid performance using univariate and multivariate GBLUP models based on North Carolina mating design II. Heredity 118: 302–310.

Werner, C. R., Qian, L., Voss-Fels, K. P., Abbadi, A., Leckband, G., Frisch, M., and Snowdon, R. J. (2018). Genome-wide regression models considering general and specific combining ability predict hybrid performance in oilseed rape with similar accuracy regardless of trait architecture. Theor Appl Genet 131: 299–317.

Wu, P. Y., Tung, C. W., Lee, C. Y., and Liao, C. T. (2019). Genomic prediction of pumpkin hybrid performance. Plant Genome 12: 180082.

Xavier, A., Muir, W. M., Craig, B., Rainey, M. (2016). Walking through the statistical black boxes of plant breeding. Theor Appl Genet 129: 1933-1949.

Xu, S., Zhu, D., and Zhang, Q. (2014). Predicting hybrid performance in rice using genomic best linear unbiased prediction. Proc Natl Acad Sci 111: 12456–12461.

Zhao, Y., Zeng, J., Fernando, R., Reif, J. C. (2013). Genomic prediction of hybrid wheat performance. Crop Sci 53: 802-810.

Zhao, Y., Li, Z., Liu, G., Jiang, Y., Maurer, H. P., Wurschum, T., Mock, H. P. Matros, A., Ebmeyer, E., Schachschneider, R. (2015). Genome-based establishment of a high-yielding heterotic pattern for hybrid wheat breeding. Proc Natl Acad Sci 112: 15624-15629.

Table 1. The top 25 superior hybrid combinations with the largest GEBVs for fruit weight (FWT) within a pumpkin population. Note that $GEBV_{ij}$ is the genomic estimated breeding value; SCA_{ij} is the specific combining ability; MPH_{ij} is the mid-parent heterosis; BPH_{ij} is the better-parent heterosis for hybrid $P_i \otimes P_j$.

$P_i \otimes P_j$	$GEBV_{ij}$	SCA_{ij}	MPH_{ij}	BPH_{ij}
P026⊗P236	3.432	0.686	0.686	0.648
P026⊗P234	3.396	0.676	0.676	0.612
P026⊗P235	3.385	0.660	0.660	0.601
P026⊗P027	3.362	0.656	0.656	0.578
P026⊗P028	3.321	0.626	0.626	0.537
P026⊗P237	3.315	0.646	0.646	0.531
P026⊗P302	3.107	0.527	0.527	0.322
P007⊗P026	3.105	0.526	0.526	0.321
P026⊗P254	3.062	0.440	0.440	0.277
P026⊗P253	3.060	0.433	0.433	0.276
P026⊗P255	3.034	0.425	0.425	0.250
P227⊗P236	3.012	0.569	0.569	0.302
P227⊗P235	3.005	0.584	0.584	0.337
P227⊗P234	2.999	0.582	0.582	0.341
P100⊗P234	2.998	0.532	0.532	0.340
P100⊗P235	2.990	0.520	0.520	0.323
P100⊗P236	2.982	0.490	0.490	0.272
P026⊗P252	2.974	0.356	0.356	0.189
P234⊗P313	2.959	0.619	0.619	0.300
P235⊗P313	2.950	0.605	0.605	0.281
P027⊗P100	2.947	0.495	0.495	0.318
P028⊗P227	2.936	0.544	0.544	0.328
P236⊗P313	2.930	0.564	0.564	0.219
P028⊗P100	2.926	0.485	0.485	0.318
P027⊗P227	2.923	0.521	0.521	0.293

Table 2. The top 10 potential parental lines with the largest GCAs for fruit weight (FWT) within a pumpkin population. GCA_i is the general combining ability for parental line P_i.

P_i	GCA_i
P026	0.6143
P236	0.5766
P235	0.5552
P234	0.5506
P027	0.5360
P028	0.5251
P237	0.4993
P253	0.4565
P254	0.4514
P252	0.4473

Table 3. The estimates for the variance components and genomic heritability in fruit weight (FWT) for a pumpkin dataset and in yield (YLD) for maize, and wheat datasets.

Dataset	$\hat{\sigma}_A^2$	$\hat{\sigma}_D^2$	$\hat{\sigma}_e^2$	h^2
Pumpkin	0.306	0.159	0.111	0.807
Maize	0.434	0.420	1.202	0.415
Wheat	0.066	0.014	0.002	0.976

Table 4. The top 15 superior hybrid combinations with the largest GEBVs for grain yield (GYD) within a maize population. Note that $GEBV_{ij}$ is the genomic estimated breeding value; SCA_{ij} is the specific combining ability; MPH_{ij} is the mid-parent heterosis; BPH_{ij} is the better-parent heterosis for hybrid $P_i \otimes P_j$.

$P_i \otimes P_j$	$GEBV_{ij}$	SCA_{ij}	MPH_{ij}	BPH_{ij}
OH7B⊗CML228*	12.975	0.587	0.587	0.385
B73⊗CML228*	12.697	0.635	0.635	0.452
Mo17⊗Tzi8*	12.589	0.582	0.582	0.454
Mo18W⊗CML103*	12.512	0.447	0.447	0.228
M162W⊗CML228*	12.504	0.460	0.460	0.239
Tx303⊗CML228*	12.499	0.358	0.358	0.132
B73⊗CML69*	12.476	0.347	0.347	0.308
Oh43⊗CML228*	12.463	0.389	0.389	0.182
Mo17⊗OH7B	12.460	0.466	0.466	0.400
Mo17⊗CML228*	12.450	0.401	0.401	0.265
B73⊗M162W	12.432	0.336	0.336	0.298
CML52⊗CML103	12.361	0.294	0.294	0.076
MS71⊗Tzi8*	12.360	0.474	0.474	0.462
MS71⊗CML228*	12.331	0.342	0.342	0.065
B97⊗CML103*	12.311	0.317	0.317	0.188

* represents a hybrid belonging to the inter-group between TM and TS.

Table 5. The top five potential parental lines with the largest GCAs for grain yield (GYD) within a maize population. GCA_i is the general combining ability for parental line P_i.

P_i	GCA_i
$CML228^{TS}$	0.2361
$CML103^{TS}$	0.1827
$Mo17^{TM}$	0.0999

B97TM	0.0538
B73TM	0.0527

TS: the tropical and subtropical group; TM: the temperate and mixed group.

Table 6. The top 20 superior hybrid combinations with the largest GEBVs for grain yield (GYD) within a wheat population. Note that $GEBV_{ij}$ is the genomic estimated breeding value; SCA_{ij} is the specific combining ability; MPH_{ij} is the mid-parent heterosis; BPH_{ij} is the better-parent heterosis for hybrid $P_i \otimes P_j$.

$P_i \otimes P_j$	$GEBV_{ij}$	SCA_{ij}	MPH_{ij}	BPH_{ij}
F6⊗F102	11.491	0.354	0.354	0.083
F102⊗M6	11.436	0.070	0.070	0.032
F102⊗M9	11.435	0.151	0.151	0.029
F97⊗F102	11.414	0.191	0.191	0.007
F1⊗F102	11.409	0.250	0.250	0.002
F46⊗F102	11.401	0.344	0.344	-0.008
F39⊗F102	11.378	0.316	0.316	-0.031
F20⊗F102	11.370	0.141	0.141	-0.037
F102⊗XM14	11.361	0.090	0.090	-0.045
F100⊗F102	11.359	0.051	0.051	-0.047
F97⊗M6	11.355	0.169	0.169	0.024
F101⊗F102	11.350	0.172	0.172	-0.057
F44⊗F102	11.342	0.200	0.200	-0.066
F98⊗F102	11.338	0.143	0.143	-0.070
F99⊗F102	11.337	0.221	0.221	-0.072
F115⊗M6	11.321	0.137	0.137	-0.010
F61⊗F97	11.320	0.315	0.315	0.277
F102⊗F115	11.311	0.090	0.090	-0.096
F102⊗M1	11.295	0.150	0.150	-0.113
F102⊗M8	11.290	0.162	0.162	-0.119

Table 7. The top 10 potential parental lines with the largest GCAs for grain yield (GYD) within a wheat population. GCA_i is the general combining ability for the parental line P_i.

P_i	GCA_i
F102	0.311
M6	0.272
F100	0.213
M9	0.188
M14	0.176
F20	0.133
F97	0.127
F3	0.126
F115	0.125
F84	0.122

Table 8. Pearson's correlations between GEBVs and phenotypic values for the datasets obtained from the Bayesian RKHS method in BGLR and our proposed Bayesian algorithm.

	Datasets				
Methods	Pumpkin	Maize-A[1]	Maize-B[1]	Maize-C[1]	Wheat
BGLR	0.9461	0.6167	0.6580	0.6039	0.9439
Our algorithm	0.9691	0.6786	0.7560	0.6546	0.9445

[1]Maize-A: the combined data from the two locations; Maize-B: the data from Columbia, MO; Maize-C: the data from Clayton, NC.

514

515 Table 9. The estimates for the variance components and genomic heritability in fruit
516 weight (FWT) for a pumpkin dataset and in yield (YLD) for maize, and wheat datasets
517 by using the Bayesian RKSH method in BGLR.


Dataset	$\hat{\sigma}_A^2$	$\hat{\sigma}_D^2$	$\hat{\sigma}_e^2$	h^2
Pumpkin	0.156	0.144	0.132	0.694
Maize	0.042	0.284	1.253	0.207
Wheat	0.029	0.014	0.003	0.935

518

519

520

521



522

523 Figure 1. The scatter plot for all available phenotypic values (i.e., 119 individuals) and
524 their GEBVs in the pumpkin dataset. The colored points represent the two hybrids out
525 of the 25 selected superior hybrids. Note that the remaining 23 selected hybrids didn't
526 appear in the plot, because their phenotypic values were not available. Pearson's
527 correlation for these 119 points was calculated as $r = 0.9691$.




Figure 2. The scatter plot for all available phenotypic values (i.e., 276 individuals) and their GEBVs in the maize dataset. The triangle points represent the top 15 superior hybrid combinations with the highest GEBVs. Pearson's correlation for these 276 points was calculated as $r = 0.6786$.




Figure 3. The scatter plot for all available phenotypic values (i.e., 2556 individuals) and their GEBVs in the wheat dataset. The colored points represent the top 20 superior hybrids with the highest GEBVs. Pearson's correlation for these 2556 points was calculated as $r = 0.9445$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- SuppMaterialsSciRep.docx