

Ethanomedicinal Properties of *Butea Monosperma* (Palas)

Rohini Sharma¹, B. K. Mehta²

^{1,2}Natural Products Research Division, School of Studies in Chemistry and Biochemistry, Vikram University, Ujjain 456010, M.P., India

Abstract

Butea monosperma (Lam.) Taub., commonly known as Palas or “Flame of the Forest,” is a widely distributed medicinal plant in India and other parts of South Asia, deeply rooted in traditional healthcare systems such as Ayurveda, Siddha, Unani, and various folk practices. Ethanomedicinally, different parts of the plant—including flowers, leaves, bark, seeds, and gum—are extensively used by indigenous communities for the treatment of numerous ailments. The flowers are traditionally employed as anti-inflammatory, antipyretic, and hepatoprotective agents, while the bark is used for its astringent and antimicrobial properties. Seeds of *B. monosperma* are well known for their anthelmintic and anti-parasitic activities, particularly in treating intestinal worm infections. Leaf extracts are used in wound healing, ulcers, and skin disorders, whereas the gum (Bengal kino) is utilized in diarrhea and dysentery. Phytochemical investigations reveal the presence of bioactive compounds such as flavonoids, chalcones, steroids, tannins, and glycosides, which may be responsible for its therapeutic efficacy. The rich ethnomedicinal knowledge associated with *Butea monosperma* highlights its significant potential as a source of natural therapeutic agents and supports the need for further pharmacological and clinical studies to validate its traditional uses.

Keywords: *Butea monosperm*, Ethanomedicinal properties, medicinal compounds.

Introduction:

Butea monosperma (Lam) Kuntze commonly known as Flame of the Forest belongs to family leguminoaceae. It is locally called as Palas, Dhak and Khakhra. It is medium size tree commonly found throughout India¹, except in the arid region. Bark, flowers, seeds, gum and leaves of *B. monosperma* are used in indigenous system of medicine for the treatment of various diseases. *B. monosperma* is one of the most showy flowering tree. The flowers in the dried condition are known as Tesu, Kesu and Palas ke phool, have a bright orange colour but *B. monosperma* tree with pale yellow colour flowers has been reported for the first time from Bhalua forest in Gaya forest division², although they are much larger, closely resemble in appearance to the common gorse (*Wex european*) with, they are botanically allied. *B. monosperma* is also called as medicinal dye bearing tree³.
Ethanomedicinal properties

Leaves

Leaves of *B. monosperma* are astringent, anti-inflammatory, diuretic, tonic and aphrodisiac. The leaves induced fall of blood sugar level and given in the treatment of glycosuria. An infusion or decoction of

the leaves is given in diarrhoea, dysentery, as a vaginal douche in leucorrhoea and as a gargle for mouthwash and in septic and congested throat⁴⁻⁵. The fresh juice of leaves is taken internally in phthisis, haemorrhagic affection, diarrhea, dysentery, fever and leaf juice is dropped into eyes to treat conjunctivitis⁶. Extract of fresh leaves is employed to kill intestinal worms and petiole is chewed and the juice is sucked to cure cough, cold and stomach disorder⁷.

Green leaves of this plant are good fodder for domestic animals and are used for making dining plates and bowls. The complete mature leaves are also used for making Ghongda to protect from rains. Fresh twigs are tied on horns of bullocks on occasion of 'Pola' festival⁴.

Leaf powder, about two spoonfuls mixed with a cup of water per day for a month is drunk mixed with a cup of water to cure diabetes⁶. A hot poultice of the leaves is effectively used as antiphlogistic or dispersing boils, pimples, tumours haemorrhoids, ulcer, swelling etc. in case of retention of urine the region of the bladder is fermented with hot leaves. The leaves are also used in inflammation in eye diseases⁸⁻⁹. A euphane triterpenoid ester and a pterocapan were reported from the leaves of this plant¹⁰.

Flower

Its dried flowers of this plant are called as tisso, tesoo, keeso, kesaree flower and Tesu ke Phool in Hindi and are used in making colour. In India, where they are used in the production of beautiful yellow and orange, red dyes, but the tints are not permanent¹¹. Flowers and young fruits are used as vegetables by tribals and flower are boiled in water and cooked to obtain a dye⁴.

Flowers are astringent, sweet, laxative, anthelmintic and tonic, aphrodisiac and diuretic¹². The flowers are used for the treatment of liver disorder and to cure Kapha and Vata, skin diseases, swelling, bone fracture, as antifertility and abortifacient¹³. The flowers boiled in water are applied as a poultice over the abdomen pubes to promote diuresis in different painful discharges of urine and to promote menstrual flow as an antiphlogistic.

The antihyperglycemic activity of the ethanolic extract of *B. monosperma* was studied in glucose-loaded and alloxan-induced diabetic rats. Single dose treatment of the extract (200 mg/kg, p.o.) significantly improved glucose tolerance and caused reduction in blood glucose level in alloxan-induced rats. Repeated oral treatment with this extract (200 mg/kg day) for 2 weeks significantly reduced blood glucose, serum cholesterol and improved HDL-cholesterol and albumin as compared to diabetic control group¹⁴.

The flowers yielded several new anthocyanin, flavones and flavonoids glycoside¹⁵ viz cyanin chloride, cyanidin, butein, butin, palasonin, isobutein¹⁶, coreopsin, isocoreopsin¹⁷, allophanic acid (Methyl ester), gallic acid,¹⁸ sulphurien, monospermoside and isomonospermoside¹⁹⁻²¹. The chalcone and aurones are responsible for the bright colour of the flowers. From the flowers terpenes, waxes, gums containing leucoanthocyanidin were also reported²². Four compounds isolated from the flowers of *B. monosperma* have been characterized as butrin, butein, 3', 4', 7'-trihydroxyflavone and stigmasterol-3- β -D-glucopyranoside²³.

Bark

The bark of this plant is astringent, pungent, alternative, aphrodisiac, anthelmintic and used in ulcer and tumours. It is also used against house flies²⁵. Decoction of bark is given in cold, cough, fever, haemorrhage, menstrual disorder, piles to cure "Vata and Kapha". Alcoholic extract of bark is given in bloody diarrhoea. Bark fibers are obtained from stem for making cordage and stem powder is used to st-

upely fishes.

The antidiarrhoeal potential of the ethanolic extract of stem bark of *B. monosperma* has been evaluated using several experimental models in white albino rats. The extract inhibited castor oil induced diarrhea and PGE₂ induced enteropooling in rats, it also reduces gastrointestinal motility after charcoal meal administration. The result obtained established the efficacy and substantiate the use of this herbal remedy as a non-specific treatment for folk medicine²⁵.

The antifungal compound isolated from the petroleum and ethylacetate extracts of the stem bark from was identified as (-)-3-hydroxy-9-methoxypterocarpan, (-)-medicarpin. Both the compounds and its acetate were active against *Cladosporioides*. The petroleum ether and ethylacetate extract of bark of this plant yielded lupeol, lupenone, two isoflavones (5-methoxy genistein and prunelin) and an agent having antifungal activity²⁶.

Seeds:

Seed contain oil called as moodooga oil. The fresh seeds are mainly used as a powerful anthelmintic for roundworms²⁷, 10 to 20 grains of the seeds made into a paste with honey are given thrice a day for three days, followed by a dose of castor oil on the fourth day. However, a considerable difference of opinion present regarding their efficacy as an anthelmintic. A decoction of the seeds is given with nitre in gravel complaints. Externally the seeds are used as a paste for ringworm. Seeds are crushed in milk and this mixture about two spoons is taken orally to treat urinal complaints and also against urinary stone²⁸. A paste made with lime juice is useful for the cure of dhobic itch. Two to three seeds are powdered and consumed by children as a remedy against intestinal worms³⁰. The powdered seeds are used for the killing maggots from infected sores and wounds. The pulverized seeds also act as a rubefacient.

The methanolic extract of seed coat and seed preparation prevented pregnancy in adult female albino rats at daily level of 120, 440 and 500 mg/kg body weight, when administered for 10 days dose post coitum²⁴. The seed preparation of *B. monosperma* as such also showed estrogenic activity at dose level of 500 mg/kg body weight on female white albino rats³⁰. Significant antifertility activity i.e. 75% and 67.25% have been observed from the alcoholic extract of outer seed coat and seed powder of *B. monosperma* respectively at the dose level of 200 mg/kg body weight in female rats³¹.









The invitro antimicrobial efficacy of *B. monosperma* outer seed coat extracts was studied by filter paper disk method against several human pathogenic gram positive and gram negative bacteria and phytopathogenic fungi. The hexane extract of seed showed excellent antibacterial effect, while the ethanol extract of seeds showed significant fungicidal effects³². Siddiqui and coworkers screened the seed and its petroleum ether extract for their biological activity and found low hypertensive activity³³⁻³⁶.

A new imide palasoimide has been isolated from the pods of *B. monosperma* and identified as palasonin-N-phenylimide³⁷. Kurup and co-workers have isolated three alkaloids from the seeds of *B. monosperma*³⁸⁻³⁹. Savita Sharma, and coworkers from this laboratory have been investigated seed and seed coat of *B. monosperma* and have isolated following compounds-

1. 1-Carbomethoxy-2-carbonyl hydrazine⁴⁰
2. 15-hydroxy pentacosanoic acid⁴⁰
3. 1-acetyl-4-methoxy-4-imidazolin-2-one⁴¹
4. 1-N-acetyl-2-oxo-4-methoxy-3H,5H-imidazole⁴²
5. 4-carbomethoxy-3,6-dioxo-5-hydro-1,2,4-triazine⁴³

6. 2-hydroxy- ω -methyl allophanic acid⁴¹
7. Phosphonic acid bis (methyl)⁴⁴
8. Patchulia alcohol⁴⁵⁻⁴⁶
9. Methyl aristollene⁴⁷⁻⁴⁸

Phenolic compounds are the main class of natural antioxidants. One of these compounds is a chalcone compound known as butein (20,40,3,4-tetrahydroxy chalcone) (1). Butein is one of the major active components of *B. monosperma*⁴⁹ (Wang, Weng, & Cheng, 2000). It is also found in *Acacia* heartwood, and has been reported to be an inhibitor of xanthine oxidase, and has inhibitory effects on lipid peroxidation in rat liver microsomes⁵⁰. In addition, it has been shown to be a specific protein tyrosine kinase inhibitor and also to induce apoptosis in human leukaemic HL-60 cells. Butein also shows antioxidant activity⁵¹⁻⁵².

 Butein	 Monospermoside
 Isomonospermoside	 Palasonin
 L actone-n-heneicosanoic acid-δ-lactone	 Allophanic acid  Gallic acid
 Ref No. (11) Jalaric ester I R = OH, Z = CH = CH Jalaric ester II R = OH, Z = (CHOH)₂ Laccijalaric ester III R = H, Z = CH = CH Laccijalaric ester IV R = H, Z = (CHOH)₂	

Flower

Triterpene, butein, butin, isobutrin, coreopsin, isocoreopsin (butin glucoside), sulphurein, monospermoside (butein 3- α -D-glucoside) and somonospermoside, chalcones, aurones, flavonoids (palasitrin, prunetin) and steroids were reported from the flower.

Gum: The gum contains tannins, mucilaginous material, pyrocatechin.

Seed Oil

(Yellow, tasteless), proteolytic and lypolytic enzymes, plant proteinase and polypeptidase. (Similar to yeast tripsin). A nitrogenous acidic compound, along with palasonin is present in seeds. It also contains monospermoside (butein 3- α -D-glucoside). The seeds were rich in fatty oil and several fatty acids were reported⁵².

Resin

Beside it also contains jalaric esters I, II and laccijalaric esters III, IV; Z- amyrin, β -sitosterol its glucoside and sucrose; lactone-n-heneicosanoic acid-{-lactone}.

Sap

Chalcones, butein, butin, colourless isomeric flavanone and its glucosides, butrin.

Leaves

Glucoside, Kino-oil containing oleic and linoleic acid, palmitic and lignoceric acid.

Bark

Kino-tannic acid, gallic acid, pyrocatechin, palasitrin, and major glycosides as butrin, alanind, allophanic acid, butolic acid, cyanidin, histidine, lupenone, lupeol, (-)-medicarpin, miroestrol, palasimide and shellolic acid. Stem: 3-Z-hydroxyeuph-25-ene and 2, 14-dihydroxy-11, 12- dimethyl-8-oxo-octadec-11-enylcyclohexane. Stigmasterol- α -D-glucopyranoside and nonacosanoic acid⁵⁰⁻⁵¹.

Botanical description

An erect tree 12-15 m high with crooked trunk and irregular branches, bark rough, ash coloured, young parts downy. Leaves 3-foliate, petioles 10-15 cm long, stipules linearlanceolate. Leaflets coriaceous (the terminal 10-20 cm long, broadly ovate from a cuneate base, the lateral smaller, 10-15 by 7.5 – 10 cm, obliquely rounded at the base, equilateral, the lower side is larger), all obtuse, glabrous above when old, finely silky and conspicuously reticulately veined beneath; petioles 6 mm long, stout-stipels subulate, deciduous. Flowers large, in a rigid racemes 15 cm long, 3 flowers together form the tumid nodes of the dark olive-green velvety rhachis: pedicels about twice as long as the calyx, densely brown-velvety: bracts and bracteoles small, deciduous. Calyx 13 mm long, dark olive-green, densely velvety outside, clothed with silky hairs within: teeth short, the 2 upper connate, the 3 lower equal, deltoid. Corolla 3.8-5 cm long, clothed outside with silky, silvery hairs, orange or salmon coloured, standard 2.5 cm broad, veined. Pods stalked 12.5-20 by 2.5-5 cm, thickened at the sutures, reticulately veined argente cane ascent stalked 2 cm long⁵³⁻⁵⁴.

References

1. Mehta B.K. and Batra A., Nagarjun, **28(4-6)**, 13-14 (1984).
2. Prasad U.N., Indian for., **108(3)**, 239 (1982).
3. Zaidi S.T.H., Ahmad S., Ali Z. and Khan J.U., National Symposium on Emerging Trends in Indian Medicinal Plants, p.1(1-0/1), 10-12 (2003).
4. Patil M.V., Patil D.A. and Pawar S., Natural Product Radiance, **5(4)**, 323-325 (2006).

5. Iqbal Zafar, Lateef Muhammad, Jabbar Abdul, Ghayur M.N. and Gilani A.H., *Fitoterapia*, **77**, 137-140 (2006).
6. Wagner H., Geyer B., Fiebig M., Kiso M., Hikino H., *Planta Med.*, **52**, 77-79 (1986).
7. Kirtikar K.R. and Basu B.D., *Indian Med. Plants* (Lalit Mohan Basu, Allahabad), **1-4**, 896 (1936).
8. Chopra R.N., Nayer S.C. and Chopra I.C., *Glossary of Indian Medicinal Plants* (CSIR, New Delhi), (1956).
9. Shukla Y.N., Mishra M. and Kumar S., *Ind. J. Chem.*, **41B**, 881-883 (2002).
10. Manjul V., Shrilakshmi D., Mustaszi J. and Mehta B.K., *Ind. J. Chem.*, **43B**, 442-426, (2004).
11. Arya V.R. and Kotalkkal V., *Indian Medicinal Plants* (Lalit Mohan Basu, Allahabad), **1**, 914 (1994).
12. Kirtikar K.R. and Basu B.D., *Indian Med. Plants* (Lalit Mohan Basu, Allahabad), **1-4**, 896 (1936).
13. Gupta S.R., Ravindranath B. and Seshadri T.R., *Phytochem.*, **9**, 2231-2235 (1970).
14. Somani R., Kasture S. and Singhai A.K., *Fitoterapia*, **77**, 86-90 (2006).
15. Puri B. and Seshadri T.R., *J. Chem. Soc.*, 1589 (1955).
16. Puri B. and Seshadri T.R., *J. Sci. Ind. Res.*, India, **13B**, 321(1954).
17. K.M. Nadkarni's, *Indian Materia Medica.*, **Vol. I**, pp. 223-25.
18. Seshadri T.R. and Trikha R.N., *Ind. J. chem.*, **9(11)**, 1201-1203 (1971).
19. Madhav R., Seshadri T.R. and Subramanim, G.D.V., *Curr. Sci.*, **331(9)**, 264-266 (1966).
20. Muyri R., Bhaskar K. and Krishna Swami N., *Proc. Ind. Acad. Sci.*, **12(B)**, 472-476 (1946).
21. Hummel James and Perkin G.A., *Chem. Abstr.*, **29**, 133 (1984).
22. Lal J.B. and Dutta S., *J. Ind. Chem. Soc.*, **12**, 262-67 (1935).
23. Robinson G.M., *J. Chem. Soc.*, **12**, 262-267 (1935).
24. Mishra Mamta, Shukla Y.N. and Kumar Sushil, *J. Med. Arom. Plant Sci.*, **24(1)**, 19-22 (2002).
25. Asolkar L.V., Kakkar K.K. and Chakre O.J., *Glossary of Indian Medicinal Plants with Active Principles* (CSIR, New Delhi), Part I, 148-149.
26. Gunakkunru A., Padmanaban K., Thirumal P., Parimala G., Vengatesan N., Gnanasekar N., Perianayagam J. B., Sharma S.K. and Pillai K.K., *J. Ethnopharmacol.*, **98**, 241-244 (2005).
27. Bondra B.M.R., Savita Kumar N. and Willmalasi K.M.S., *J. National Science Council of Srilanka*, **81(2)**, 97-103 (1990).
28. Nankarni K.N., *Indian Meteria Medica* (Dhoot Papeswar Prakashan Ltd., Bombay), III edition, 224, 135 (1954).
29. Watt George, *A Dictionary of Economic Product of India*, Vol; I-VI (1893).
30. Jain S.K., *Dictionary of Indian Folk Medicine and Ethanobotany* (Deep Publication, New Delhi, India), (1991).
31. Greter Buchi and Tokoroyama, *Tetrahedron*, 822 (1962).
32. Mehta B.K., Dubey A., Bokadia M.M., *Ind. J. Chem.*, **22(B)**, 282 (1983).
33. Mehta B.K., Dubey A., Bokadia M.M. and Mehta S.C., *Acta Microbial, Hungarica*, **30(1)**, 75 (1983).
34. Siddique H.H., Kalaysaraj R. and Kurup P.A., *Ind. J. Pharma.*, **25**, 270-271 (1963).
35. Siddque H.H., *Ind. J. Expt. Biol.*, **1**, 11 (1973).
36. Mehta B.K., Jaiswal S. and Batra A., *Phytopathol.*, **109**, 90 (1984).
37. Mehta B.K., Dubey A., Bokadia M.M., Mehta S.C., *Acta Microbiol. Hungarica*, **30(1)**, 75 (1983).
38. Guha D.K., Rai R. and Bhattacharya A., *Phytochemistry*, **126(6)**, 2017 (1990).
39. Kurup P.A. and Kaleysaraj R.J., *J. Pharm.*, **24(63)**, 1962 (1980).
40. Mehta B.K. and Bokadia M.M., *J. Chem. Ind. (Lodon)*, 98 (1981).

41. Sharma S., Batra A. and Mehta B.K., Indian J. Chem., **30(B)**, 715-716 (1991).
42. Porwal M., Sharma S. and Mehta B.K., Fitoterapia, **LIX(2)**, 134-135 (1988).
43. Mehta B.K., Bokadia M.M. and Dhar K.L., Chem. Ind. (Lodon), 862 (1988).
44. Porwal M., Mehta B.K. and Gupta D.N., Nat. Acad. Sci. Letters, **2(3)**, (1988).
45. Gotedell I.J., Can. J. Chem., **47**, 2461 (1962).
46. Mirrgtan R.N., J. Org. Chem., **37**, 2871 (1972).
47. Yamada K., Kyotani Yamada K., Kyotani Y., Manabe S. and Suzuki M., Tetrahedron, **35**, 293 (1979).
48. Wang W., Weng X. and Cheng D., Food Chemistry, **71**, 45–49, (2000).
49. Sogawa S., Nihro Y., Ueda H., Miki T., Matsumoto H. and Satoh T., Pharmacology Bulletin, **17**, 251–256, (1994).
50. Yang E. B., Zhang K., Cheng L. Y., and Mack P., Biochemical Research Communication, **245**, 435–438, (1998).
51. Kim N. Y., Pae H. O., Oh G. S., Kang T. H., Kim Y. C., Rhew H. Y. and Chung H. T., Pharmacology and Toxicology, **88**, 261–266, (2001).
52. Kapoor L.D., Handbook of Ayurvedic Medicinal Plants, (Replica Press Pvt. Ltd., India, pp.86, (2005).
53. The Wealth of India, A dictionary of India raw material and Industrial products, (Publication and Information Directorate, CSIR, New Delhi, 1988) Vol. II, pp. 1-344.
54. Naf F., Helv Chem. Acta., **64**, 1387 (1981).