



In Vitro Evaluation of Salt Tolerance and Melatonin-Mediated Stress Amelioration in Wild Potato (*Solanum okadae*)

Amar Hundare^{1,2} · Aneesh Lale² · Neelu Joshi³ · Timothy P. Robbins²



Received: 22 June 2025 / Accepted: 16 October 2025 / Published online: 6 February 2026
© The Author(s) 2026

Abstract

Salinity is a critical abiotic stress influencing agricultural crop productivity in arid and semiarid environments. This study reports an in vitro evaluation of NaCl tolerance of wild diploid potato species *Solanum okadae* and the effect of exogenous MT application on reducing the salt stress effects. In vitro grown plantlets stressed with 50, 100, 150, and 200 mM NaCl had stress tolerance indices of 84.67%, 76.88%, 81.84%, and 67.94%, respectively. Compared to the control, 200 mM NaCl had severe adverse effects on growth parameters and decreased total chlorophyll and K⁺ content by 88.12% and 25.91%, respectively, and increased Na⁺ content by 50.17-fold. The rise in catalase activity, Ca²⁺ and Mg²⁺ contents, was 1.08-fold, 1.59-fold, and 1.25-fold, respectively. Melatonin (100 µM) treatment mitigated the growth inhibition caused by 200 mM NaCl, as demonstrated by an increase of 17.28%, 150%, 25.39%, 36.36%, and 6.72% in fresh weight, shoot length, total chlorophyll, chlorophyll b, and K⁺ content, along with decreased Na⁺ content and catalase activity by 1.79-fold and 1.08-fold, respectively. This preliminary in vitro study shows that *Solanum okadae* has natural salt tolerance, which is enhanced by exogenous melatonin improving physiological and biochemical responses under salinity stress. These findings highlight *S. okadae*'s potential for breeding salt-tolerant potato cultivars and for understanding the genetic and molecular mechanisms of salinity tolerance. However, further validation in greenhouse and field conditions is needed to confirm the stability and practical relevance of these responses in real agricultural settings.

Keywords Catalase · Chlorophyll · Melatonin · Potato · Salinity · *Solanum okadae*

✉ Amar Hundare
A.hundare@soton.ac.uk

¹ University of Southampton, Southampton, Hampshire SO17 1BJ, UK

² Division of Plant and Crop Science, School of Biosciences, University of Nottingham, Sutton Bonington, Loughborough LE12 5RD, UK

³ School of Biotechnology & Bioinformatics, D.Y. Patil Deemed to be University, CBD Belapur, Sector-15, Navi Mumbai 400614, India

Introduction

Salinity is a significant environmental stress factor that frequently poses a serious obstacle to the production of crops (Peethambaran et al. 2018). The most common salt that causes toxicity and is responsible for saline soils is NaCl. It affects the growth and productivity of crops by inducing ion imbalance and osmotic and oxidative stress (Jiang et al. 2021). In response, plants have developed several adaptive mechanisms including ion compartmentalization, osmolyte synthesis, and antioxidative metabolism, which depend upon activation of molecular processes involved in stress recognition, signal transduction, and expression of stress-related genes and metabolites (Gupta and Huang 2014). Growth, physiological, and biochemical traits are generally measured to screen for salt tolerance. Besides, the stress tolerance index (STI) is another indicator of salt tolerance (Fernandez, 1992).

Melatonin (MT, N-acetyl-5-methoxytryptophan), a lipophilic and hydrophilic physiological regulator, plays a significant role in plants' response to abiotic stress by inducing stress-responsive pathways (Arnao et al. 2022). Exogenous application of MT has emerged as an effective strategy to mitigate salt stress in various plants, including tomato, rice, and cotton (Zhou et al. 2016; Liang et al. 2017; Zhang et al. 2021). It works by regulating processes such as antioxidant metabolism, photosynthesis, polyamine metabolism, ionic balance, and NO signalling.

Under in vivo and in vitro conditions, *S. tuberosum* is categorized as a moderately salt sensitive crop, with a threshold electrical conductivity (EC) level of 1.7 dS/m, as its tuber yield may decline at field salinity EC levels of 2.0–3.0 dS/m (approximately 22–33 mM NaCl) (Shrestha et al. 2023). In the reproductive phase, salinity stress can alter the normal K^+/Na^+ ratio in the leaves, adversely affect the ultrastructure of the leaves, weaken photosynthesis in the leaves, and hamper the transfer of carbon from the vegetative tissues to the tubers, as is the quantity and size of the tubers, consequently (Levy and Veilleux, 2007). To control this loss, application of stress ameliorating agents to the potato crop could be an efficient, cost-effective and eco-friendly way. In vitro system is generally preferred for salt tolerance studies as it can provide quicker, reproducible, and more accurate assessments of plant growth considering the environmental variations in conventional field trials (Banu et al. 2015). However, there are limited reports on the use of such system in potato. In one instance, Zaki and Radwan (2022) performed in vitro evaluation of diploid potato *Solanum chacoense* Bitt and identified its clones 'A-6', 'C-8', and 'D-2' as highest salt tolerant. They also determined the effect of exogenous applications of salicylic acid, proline, and glycine betaine to reduce salt stress injury in the potatoes. Other instances include in vitro screening for salt tolerance (Ahmed et al. 2020) in commercial potato varieties like Innovator, Lady Claire, Kennebec, Burren, Marabel, and Hermes and evaluating the effect of MT in relieving salt stress symptoms (Efimova et al. 2023) in *Solanum tuberosum* cv. Lugovskoi. Wild potato species serve as crucial genetic resources for enhancing abiotic stress tolerance in cultivated varieties. *Solanum okadae*, a wild diploid species native to Bolivia, is known for its resilience to environmental stresses like drought (Watanabe et al. 2011) and its compatibility with *Solanum tuberosum* (Camadro et al. 2008), making it an ideal candidate for pre-breeding efforts. These wild relatives often harbour

unique physiological and biochemical mechanisms such as improved ion homeostasis, antioxidant regulation, and osmotic adjustment that are not present or are diminished in modern cultivars. Despite this potential, studies evaluating salt tolerance in wild diploid potato species remain limited. Such investigations are essential, as they could contribute significantly to the development of salt-tolerant potato lines through conventional or molecular breeding strategies. Therefore, our study aimed to evaluate the salt tolerance of *S. okadae* under in vitro conditions and to investigate the potential of exogenously applied melatonin (MT) in ameliorating salt-induced stress symptoms.

Materials and Methods

Plant Material

True seeds of *Solanum okadae* (accession: OKA7129) acquired from the Commonwealth Potato Collection curated by the James Hutton Institute (Invergowrie, Scotland) were germinated in the greenhouse at the Plant and Crop Science department, University of Nottingham, UK. From a 4-week-old seed-derived plant, nodal explants were collected to establish shoot cultures (Hundare 2023). Explants were washed under running tap water, surface sterilised using 70% ethanol for 1 min, followed by a 10-min treatment with 5% NaOCl, and were rinsed thrice with sterile distilled water. Subsequently, explants (2 cm) were cultured on solidified Murashige and Skoog (MS) medium containing 8 g/L agar, 30 g/L sucrose along with 3 mg/L BA (6-benzylaminopurine), 2 mg/L NAA (1-naphthaleneacetic acid), and 1 mg/L KIN (kinetin). The pH was adjusted to 5.7 ± 0.1 prior to the addition of agar and subsequent autoclaving at 121 °C and 15 psi for 20 min. Tissue excision, implantation, and transition were conducted under sterile conditions. Cultures were incubated in culture room with a temperature of 22 ± 2 °C, 16-h photoperiod and $\sim 30 \mu\text{mol m}^{-2} \text{s}^{-1}$ light intensity and sub-cultured every week on fresh media to obtain shoot cultures. To minimize genetic variability, all in vitro cultures were established from nodal explants derived from a single seedling originating from one true seed of *S. okadae*. Once established, these clonal shoot cultures were used for all experimental treatments, ensuring genetic uniformity across replicates.

Melatonin (MT) (Sigma-Aldrich, St. Louis, USA) was dissolved in 0.1% (v/v) ethanol to a stock concentration of 10 mM and diluted with sterile water to obtain a final working concentration of 100 μM . The final ethanol concentration in the culture medium did not exceed 0.1%.

Experimentation

Five individual stem nodes derived from *Solanum okadae* shoot cultures were placed in Magenta boxes containing MS medium supplemented with seven different treatments:

- (i) NaCl at concentrations of 0 (control), 50, 100, 150, and 200 mM and

- (ii) 200 mM NaCl combined with 100 μM MT. Cultures were maintained under standard growth room conditions (22 ± 2 °C, 16-h photoperiod, $\sim 30 \mu\text{mol m}^{-2} \text{s}^{-1}$ light intensity).

All measurements were taken 30 days after inoculation. No pre-treatment or acclimation phase was applied before the start of the experiment. For each treatment, five explants were placed in each of six Magenta boxes, with each box considered an independent biological replicate. The experiment was repeated three times, giving a total of 90 explants per treatment for statistical analysis.

The selected NaCl concentrations (50–200 mM) were based on previous *in vitro* studies in potato, which reported growth inhibition at ≥ 100 mM. The 200 mM level was used to represent severe salinity stress, as it reliably induces physiological and biochemical responses without causing complete lethality (Zaki and Radwan 2022). The melatonin (MT) concentration of 100 μM was selected based on multiple studies demonstrating its effectiveness in improving plant survival and reducing oxidative damage under salt and heat stress conditions in tomato and cotton (Martínez et al. 2018). In this study, melatonin treatment was applied only at 200 mM NaCl to evaluate its effect under high salinity stress. Due to resource and experimental limitations, melatonin was not tested across the full range of salinity treatments (0, 50, 100, 150 mM NaCl), which restricts the interpretation of its effectiveness at varying salt levels. Future studies should apply melatonin treatments across all salinity concentrations to provide a more comprehensive assessment of its role in salt tolerance.

Measurement of Growth Traits

Shoot height (SH, cm), root length (RL, cm), number of shoots (SN), roots (RN), and leaves (LN), fresh weight (FW, g), and dry weight (DW, g) were measured for plantlets from control and each treatment. The percent plant water content (PWC) was calculated using the equation (Rashid et al. 2019):

$$\text{PWC} = ((\text{FW} - \text{DW}) / \text{FW}) \times 100$$

Photosynthetic Pigments

Determination of carotenoids (Car), chlorophyll a (Chl a), and b (Chl b) was conducted following the methodology established by Arnon (1949) and Lone et al. (2018). One hundred milligrams of leaf sample was crushed in 10 ml of 80% acetone using mortar and pestle, followed by centrifugation at 5000 rpm for 5 min. The supernatant was transferred to a fresh tube following the absorbance (A) of the solution at 480 nm, 645 nm, and 663 nm against the blank (acetone) using a spectrometer (CE 2041 UV/VIS, Buck Scientific, Inc.). Chl a, Chl b, total chlorophyll (total Chl), and Car contents were expressed as micrograms per gram of fresh weight ($\mu\text{g/g}$ FW). The concentrations were calculated using the following equations:

$$\begin{aligned}
 \text{Total chlorophyll} &= 20.2(A_{645}) + 8.02(A_{663}) \\
 \text{Chlorophyll } a &= 12.7(A_{663}) - 2.69(A_{645}) \\
 \text{Chlorophyll } b &= 22.9(A_{645}) - 4.68(A_{663}) \\
 \text{Carotenoid} &= [A_{480} + \{(0.114 \times A_{663}) - (0.638 \times A_{645})\}] \times V / 1000 \times W
 \end{aligned}$$

where V is the volume of acetone and W is the weight of sample in mg.

Proline Determination

The proline concentration in the extracts was determined following the protocol of Stoleru et al. (2019). Fresh leaves (500 mg) were homogenised in 3% (w/v) 5-sulfosalicylic acid and centrifuged at 4000 g for 10 min at 4 °C. The supernatant was mixed with acid ninhydrin and glacial acetic acid, heated for 60 min at 100 °C, and subsequently cooled, and then 4 mL of toluene was added before measuring the absorbance at 520 nm, using a spectrophotometer and toluene as the blank.

CAT Activity Assay

Leaf tissue (200 mg) was homogenised in 3 ml of 100 mM PBS buffer (pH 7.8) and centrifuged at 10,000 × g for 20 min at 4 °C, and the supernatant collected and kept on ice. For determining catalase activity (Chen and Zhang 2016), the supernatant was mixed with 100 mM PBS buffer (pH 7.8) containing 30% H₂O₂, and the change in absorbance at 240 nm was recorded for 1 min using a spectrophotometer. One unit of CAT activity was quantified as the amount of enzyme required to catalyse the decomposition of 1 μmole of H₂O₂ per minute at pH 7.0, and it was expressed as U/mg protein.

Histochemical Detection of Peroxide (H₂O₂)

H₂O₂ in leaves was detected histochemically by staining the leaf samples with 1 mg/mL DAB (3,3-diaminobenzidine) as described by Daudi and O'Brien (2012). Leaves were then mounted on clean glass slides and covered with a coverslip and visualized under a microscope (LAS AF, Leica Microsystems) on a plain white background under uniform lighting.

Elemental Profiling Using ICP-MS

The ionomics analysis of samples was carried out as described (Danku et al. 2013). Whole plantlets were harvested and rinsed twice in deionised water and quickly sealed. Samples were dried at 80 °C for 24 h and then ground. Ions were extracted from samples in 0.5% HNO₃. Na⁺, K⁺, Ca²⁺, and Mg²⁺ were assayed using inductively coupled plasma mass spectrometry (ICP-MS) (PerkinElmer NexION 2000 equipped with an Elemental Scientific Inc. auto-sampler and the Syngistix software in collision mode (He)). The concentrations of minerals were expressed in mmol/g on a dry weight basis.

Salt Tolerance Trait Index (STTI)

To evaluate the limit of salt tolerance (i.e. maximum salt concentration in which in vitro plantlets could survive) and check the effect of MT on stressed plantlets, the average of each trait (morphological and biochemical) as STI was estimated using the following equation:

$$\text{STTI} = \frac{\text{Value of trait under stressed condition}}{\text{Value of trait under no stress}} \times 100$$

For each salt concentration, average STTIs were computed separately for (i) morphological parameters, (ii) pigment traits, (iii) ionic concentrations, and (iv) biochemical traits. No weighting was applied to individual traits. Trait values were not normalized across datasets. The rationale for selecting these parameters was based on their known involvement in salt stress responses and their ease of quantification under in vitro conditions.

Statistical Analysis

For each treatment, 30 explants were used, distributed across 6 magenta boxes (5 explants per box), with each box considered 1 experimental unit. Experiments were conducted in a randomized design with three independent runs. All measurements were taken from individual explants, and data were presented as mean $\pm$ standard deviation (SD). Statistical analysis was performed using SPSS software (version 21.0, SPSS Inc., Chicago, USA). Differences between treatments were assessed using one-way ANOVA followed by Duncan's Multiple Range Test (DMRT) at a significance level of $P < 0.05$.

Results

Growth Parameters

All NaCl concentrations reduced growth parameters to varying degrees, with ≥ 100 mM causing broad and severe effects. Even the lowest concentration (50 mM) significantly reduced fresh weight, leaf number, and shoot number compared to the control, although shoot height, root number, root length, and dry weight were not significantly affected (Table 1). Leaf necrosis was observed at higher NaCl levels. Severe salt stress (200 mM) resulted in significant reductions in fresh and dry weight, leaf number, and shoot length by 64%, 83.33%, 64.94%, and 82.67%, respectively, alongside a 100% decrease in the number of shoots, roots, and root length compared to control. Although root and shoot numbers were reduced to near zero in some replicates at 200 mM NaCl, plantlets retained some physiological activity as evidenced by measurable proline content, CAT activity, and ionic changes, suggesting survival under stress rather than complete lethality. PWC values

Table 1 Effect of various NaCl concentrations (mM) and Melatonin (MT) treatment on different morphological parameters and photosynthetic pigments of *Solanum okadae* in vitro

Treatment	0	50	100	150	200	NaCl + MT
FW (g)	2.25 ± 0.6 ^a	1.50 ± 0.1 ^b	1.32 ± 1.2 ^{bc}	1.10 ± 0.7 ^{bc}	0.81 ± 0.3 ^c	0.95 ± 0.4 ^c
DW (g)	0.18 ± 0.0 ^a	0.14 ± 0.0 ^a	0.08 ± 0.0 ^b	0.06 ± 0.0 ^b	0.03 ± 0.0 ^b	0.10 ± 0.0 ^{ab}
PWC (%)	85.38 ± 0.9 ^a	85.83 ± 1.5 ^a	85.92 ± 2.9 ^a	86.26 ± 2.3 ^a	86.46 ± 1.2 ^a	85.00 ± 1.2 ^a
LN	9.50 ± 4.35 ^a	6.67 ± 2.1 ^b	5.83 ± 0.9 ^b	4.33 ± 2.0 ^b	3.33 ± 1.7 ^c	6.33 ± 2.3 ^b
SN	11.17 ± 4.2 ^a	4.67 ± 4.0 ^b	3.33 ± 3.8 ^b	0.33 ± 0.5 ^c	0.00 ± 0 ^c	1.23 ± 1.1 ^{bc}
SH (cm)	7.3 ± 0.3 ^a	7.2 ± 0.4 ^a	2.30 ± 0.2 ^c	1.9 ± 0.7 ^c	1.2 ± 0.3 ^c	3.0 ± 0.2 ^b
RN	3.50 ± 0.8 ^a	3.40 ± 1.5 ^a	2.00 ± 1.0 ^b	0.0 ^c	0.0 ^c	1.19 ± 0.2 ^b
RL (cm)	1.30 ± 0.0 ^b	1.4 ± 0.1 ^b	1.7 ± 0.2 ^a	0.0 ± 0.0 ^c	0.00 ± 0 ^c	1.1 ± 0.1 ^b
Total Chl	37.74 ± 5.4 ^a	6.42 ± 2.6 ^c	4.60 ± 1.2 ^c	11.81 ± 3.6 ^b	4.48 ± 1.6 ^c	5.58 ± 1.9 ^c
Chl a (µg/g FW)	14.44 ± 3.2 ^a	4.59 ± 1.1 ^b	3.02 ± 1.6 ^c	4.86 ± 1.5 ^b	2.17 ± 1.3 ^d	2.08 ± 1.1 ^d
Chl b (µg/g FW)	23.31 ± 4.4 ^a	1.56 ± 0.1 ^c	1.58 ± 0.8 ^c	7.22 ± 2.4 ^b	2.31 ± 1.1 ^d	3.51 ± 0.4 ^c
Car (µg/g FW)	1.53 ± 1.1 ^c	1.38 ± 0.4 ^c	2.60 ± 1.4 ^b	3.75 ± 0.7 ^a	4.20 ± 1.1 ^a	2.69 ± 0.2 ^b

All values represent the means ± S.E. Different letters within a column indicate significant differences at $P < 0.05$ by Duncan's multiple range test. *FW* fresh weight, *DW* dry weight, *SH* shoot height, *RL* root length, *SN* shoot number, *RN* root number, *LN* leaf number, *Chl* chlorophyll, *Car* carotenoid

did not differ significantly among treatments (Table 1); however, a numerical maximum of 86.46% was observed at 200 mM NaCl, suggesting that osmotic adjustment may help *S. okadae* maintain water content even under severe salt stress. Under 200 mM NaCl stress, MT application significantly increased shoot length (150%) and leaf number (90.09%) compared to untreated stressed plantlets, while fresh weight showed a non-significant increase of 17.28%. MT also enhanced shoot and root formation significantly.

Photosynthetic Pigments

Chlorophyll a, b, and total chlorophyll contents generally declined with increasing NaCl concentration, but the trend was not strictly dose-dependent (Table 1). For example, Chl b levels at 50 mM and 100 mM were lower than at 200 mM, and the highest mean values for Chl a, Chl b, and total chlorophyll (4.86 ± 1.5 , 7.22 , and 11.81 ± 3.6 µg/g FW, respectively) were recorded at 150 mM NaCl. Several of these differences were not statistically significant. Carotenoid content showed a less consistent pattern, with the highest mean value (4.20 ± 1.12 µg/g FW) at 200 mM NaCl; however, statistical groupings did not always correspond to the absolute values, indicating variability among replicates. Under 200 mM NaCl stress, MT application significantly increased total chlorophyll (by 24.55%) and chlorophyll b (by 51.94%) compared with stressed plantlets without MT. Carotenoid content decreased by 35.95% under MT treatment, which was statistically significant. Chlorophyll a did not differ significantly between MT-treated and untreated stressed plantlets.




Fig. 1 Effect of various NaCl concentrations (mM) and melatonin (MT) treatment on Na⁺, K⁺, Ca²⁺, and Mg²⁺ content of *Solanum okadae*. Means with different letters are significantly different ($P < 0.05$)

Ionic Content

The analysis revealed a rise in Na⁺, Ca²⁺, and Mg²⁺ and a decline in K⁺ levels with salt stress compared to the control (Fig. 1). The rise was highest for Na⁺ (50.18-fold) and Ca²⁺ (1.59-fold) at 200 mM NaCl and for Mg²⁺ (2.12-fold) at 150 mM, over control. A slight decline in K⁺ level (0.74-fold) was recorded at 200 mM NaCl. MT treatment at 200 mM NaCl notably enhanced the concentrations of K⁺ (6%), Ca²⁺ (13%), and Mg²⁺ (17%), while also leading to a significant reduction in Na⁺ (44%) concentrations compared to untreated stressed plantlets.




Fig. 2 Effect of various NaCl concentrations and melatonin (MT) treatment on proline content and CAT activity of *Solanum okadae*. Means with different letters are significantly different ($P < 0.05$)

CAT Activity, H_2O_2 and Proline Content

NaCl stress significantly elevated CAT activity and proline content, with the highest rise of 7.77% in CAT activity at 200 mM NaCl and 27% proline content at 150 mM NaCl; over control was recorded (Fig. 2). DAB staining of the leaf tissues of stressed plantlets showed small dark spots under the microscope, indicating increased contents of H_2O_2 (Fig. 3). Supplying exogenous MT helped in lowering the H_2O_2 content and ultimately lowered the CAT activity by 7.09% and proline content by 34% compared to untreated stressed plantlets.

Salt Tolerance Trait Index (STTI)

The highest STI for morphological traits (89.76%) and photosynthetic pigments (31.29%) was recorded at 50 mM NaCl, whereas it was highest at 150 mM NaCl




Fig. 3 Visualization of H_2O_2 microbursts in response to salt stress using DAB staining. **a** Control, **b** 50 mM NaCl, **c** 200 mM NaCl, and **d** 200 mM NaCl + 100 μM melatonin

for ionic content (145.78%) and for CAT activity, proline, and protein content (108.01%) (Table 2). Plantlets growing in 150 mM NaCl had the highest overall stress tolerance index (81.84%). At 200 mM NaCl stress, exogenous MT treatment significantly enhanced the STI for morphological parameters and ionic content from 38.58% to 63.31% and 103.42% to 118.70%, respectively.

Discussion

The present study investigated the effects of NaCl stress and stress reliever MT in *S. okadae* under in vitro conditions. In vitro screening of potato genotypes for water stress tolerance has been proposed as an efficient alternative to field-based testing (Gelmesa et al. 2017), and results were found to be like those observed in field conditions (Rashid et al. 2019).

Morphological and Growth Responses

Salinity stress reduced the morphology and biomass accumulation of *S. okadae* plantlets, with concentrations ≥ 100 mM NaCl significantly inhibiting shoot length, root length, number of shoots, leaves, and overall biomass. At 50 mM NaCl, only fresh weight and shoot number were significantly reduced, while other traits remained statistically similar to the control. This aligns with previous studies

Table 2 Salt tolerant trait indices (STTI, %) for various traits of in vitro raised *Solanum okadae* under NaCl concentrations (mM) and Melatonin (MT) treatment

Treatment	Morphological	Photosynthetic pigments	Ionic content	CAT activity and proline content	Mean STTI
50 mM NaCl	89.76 ^a	31.29 ^a	101.17 ^c	98.44 ^{ab}	80.16 ^a
100 mM NaCl	78.44 ^b	21.72 ^{bc}	107.48 ^c	99.87 ^{ab}	76.88 ^b
150 mM NaCl	43.81 ^d	29.75 ^b	145.78 ^a	108.01 ^a	81.84 ^a
200 mM NaCl	38.58 ^d	28.19 ^b	103.42 ^c	101.55 ^{ab}	67.94 ^c
200 mM NaCl+ 100 µM MT	63.31 ^c	23.22 ^{bc}	118.70 ^b	90.55 ^c	73.94 ^b

All values represent the means $\pm$ S.E. Different letters within a column indicate significant differences at $P < 0.05$ by Duncan's multiple range test

indicating that salinity adversely affects cell elongation and division due to osmotic and ionic stress (Sudharsan et al. 2012; Zaman et al. 2015). The inhibition of leaf formation and root development under high salt conditions could be attributed to energy diversion towards osmotic regulation and ion homeostasis, as maintaining cellular equilibrium becomes metabolically expensive under saline stress (Biswas et al. 2017; Evers et al. 1999).

Plant water content (PWC) did not differ significantly among treatments. The numerically highest value at 200 mM NaCl may suggest that *S. okadae* can maintain water balance under stress, but given the lack of statistical significance, this should be interpreted cautiously and confirmed in further studies. This contrasts with reports in *S. tuberosum* and *S. chacoense*, where PWC decreased under salt stress (Zaki and Radwan 2022). The maintenance of PWC in *S. okadae* suggests that this species possesses an intrinsic ability for osmotic adjustment, potentially involving effective solute accumulation, xylem water transport, and stomatal regulation (Sack et al. 2018). Such mechanisms allow the retention of cellular turgor and continued metabolic function, preventing premature senescence and supporting survival at 150 mM NaCl, a level at which many cultivars fail.

At a mechanistic level, salinity-induced reductions in turgor pressure result from progressive dehydration, leading to membrane destabilization and loss of cell viability (Tyree et al. 2003). Plants that prevent this through osmotic regulation, such as by maintaining water potential equilibrium between cells and xylem, can continue physiological processes under stress. Our data suggest that *S. okadae* may achieve this via solute accumulation and cellular compartmentalization, maintaining turgidity and delaying growth cessation.

Exogenous application of MT (100 µM) further promoted shoot and root development under 200 mM NaCl stress, with significant increases in shoot length and leaf number, and numerical (non-significant) increases in fresh weight, root length, and shoot number compared to salt-only treatment. These selective improvements suggest MT may preferentially enhance above-ground growth processes under severe stress. This supports melatonin's known role in promoting cell division and

elongation under abiotic stress, potentially through the modulation of auxin pathways, ROS homeostasis, and stress hormone signalling (Tan and Reiter 2020).

While MT did not significantly affect fresh weight, dry weight, PWC, or root number compared to salt-only treatments, the improvements in above-ground morphological traits suggest selective reinforcement of growth-related processes. This indicates that MT may preferentially target shoot meristem activity and leaf expansion, possibly by stabilizing cytoskeletal elements or enhancing carbon assimilation efficiency. These findings align with those of Altaf et al. (2022), who demonstrated that MT-treated tomato seedlings under drought stress exhibited superior morphological traits due to improved antioxidant status and hormone balance.

Taken together, these results indicate that MT confers partial protection to growth-related traits under salt stress in *S. okadae* by enhancing cellular integrity, maintaining hormonal signalling, and promoting shoot meristem activity, mechanisms that collectively improve growth performance despite osmotic and ionic constraints.

Photosynthetic Apparatus and Pigment Stability

Salinity imposes multifaceted stress on the photosynthetic machinery by disrupting pigment synthesis, degrading chlorophyll, altering stomatal behaviour, and reducing gas exchange. It inhibits nitrogen and magnesium uptake, both crucial for chlorophyll biosynthesis, thereby compromising photosynthetic efficiency (Kaya et al. 2020; Sarker et al. 2018). Moreover, salinity-induced stomatal closure reduces internal CO₂ concentration, and lower stomatal density further impairs photosynthesis (Charfeddine et al. 2019).

In the present study, increasing NaCl concentrations led to a significant reduction in chlorophyll a, b, and total chlorophyll content in *S. okadae* plantlets, indicating stress-induced pigment degradation. These results are consistent with previous reports in both cultivated and wild potato species (Daneshmand et al. 2010) and tomato (Juan et al. 2005). The reduction in chlorophyll under salt stress is linked to elevated chlorophyllase activity and ion toxicity-induced oxidative stress, which accelerates pigment breakdown (Sultana et al. 1999).

The decline in chlorophyll content under salinity was not strictly dose-dependent in *S. okadae*. Chl b values at 50 mM and 100 mM were lower than at 200 mM, and the relatively higher chlorophyll content observed at 150 mM may reflect variability among replicates rather than a consistent adaptive response. As several differences were statistically non-significant, this pattern should be interpreted with caution. Nonetheless, similar transient pigment retention under moderate salinity has been noted in some tolerant genotypes of rice and pea (Misra et al. 1997; Hernandez et al. 2000), suggesting that further replicated studies are needed to confirm whether this occurs consistently in *S. okadae*. The capacity to retain or even upregulate chlorophyll under moderate stress suggests that *S. okadae* activates compensatory mechanisms to maintain photosynthetic capacity, likely involving antioxidant protection of chloroplast structures and regulatory control over pigment biosynthetic genes.

Chlorophyll b, primarily located in the light-harvesting complexes of photosystem II (PSII), increased under stress in this study. This may indicate a structural adaptation to improve light absorption and energy transfer under suboptimal conditions (Kitajima and Hogan 2003). Given that PSII is highly sensitive to salinity-induced photoinhibition (Kolomeichuk et al. 2020), the maintenance of Chl b content in *S. okadae* may contribute to preserving photosynthetic efficiency under salt stress.

Carotenoids, essential for photoprotection and ROS scavenging, increased with rising NaCl concentrations. Their role in quenching singlet oxygen, stabilizing thylakoid membranes, and dissipating excess light energy is vital under abiotic stress. The increase observed here aligns with findings in *Amaranthus tricolor* and *Solanum acaule* (Wang and Nii 2000; Daneshmand et al. 2010) and underscores the contribution of carotenoids to stress resilience by preventing lipid peroxidation and preserving photosystem integrity.

Regarding melatonin application, MT-treated *S. okadae* plantlets at 200 mM NaCl exhibited a marked increase in total chlorophyll and Chl b content. This suggests that MT stabilizes the photosynthetic apparatus, likely through upregulation of chlorophyll biosynthesis genes (*CHLH*, *CHLM*, *PORA*) and protection of PSII (Arnao and Hernández-Ruiz 2019). In this study, MT treatment under 200 mM NaCl stress significantly increased total chlorophyll content, primarily due to a marked rise in chlorophyll b, while chlorophyll a showed no significant difference compared to untreated stressed plantlets. Carotenoid content, however, decreased significantly with MT application. These findings suggest that melatonin's effect on pigment stabilization is selective, enhancing components associated with light-harvesting complexes (where chlorophyll b predominates) rather than uniformly increasing all chlorophyll types. These results differ from tomato (Altaf et al. 2021) and cucumber (Wang et al. 2015), where MT enhanced both Chl a and carotenoids, possibly reflecting species-specific responses or differential sensitivity of pigment types to MT regulation.

In terms of underlying process, the observed effects may be linked to chloroplast-localized melatonin biosynthesis. Chloroplasts, originating from endosymbiotic cyanobacteria, retain the machinery for melatonin production, including the key enzyme N-serotonin acetyltransferase (SNAT) (Tan et al. 2012). Under salinity, stress-induced ROS accumulation triggers melatonin synthesis within the chloroplast, providing local antioxidant defence and stabilizing pigment-protein complexes (Murch and Erland 2021). Therefore, the MT-induced increase in Chl b and total Chl under salt stress in *S. okadae* likely reflects a coordinated response involving both enhanced pigment biosynthesis and ROS mitigation within the chloroplast.

In summary, *S. okadae* exhibits a unique capacity to retain or recover chlorophyll under moderate salt stress, likely through regulatory and structural adjustments in pigment biosynthesis and light-harvesting complexes. Melatonin enhances this adaptive capacity by protecting chloroplast function, mitigating ROS damage, and preserving PSII integrity, contributing significantly to photosynthetic resilience under salinity.

Mineral Composition

Salt stress, often described as hyper-ionic stress, disrupts plant cellular ion homeostasis primarily through excessive Na^+ accumulation, which inhibits K^+ uptake and alters the K^+/Na^+ balance—crucial for enzymatic activity and protein synthesis (Ishikawa and Shabala 2019; Maathuis and Amtmann 1999). In this study, increasing NaCl concentrations elevated Na^+ levels while reducing K^+ levels in *S. okadae*, although K^+ remained relatively stable at lower stress levels (50 mM), indicative of intrinsic regulatory mechanisms.

Notably, MT application at 200 mM NaCl significantly reduced Na^+ accumulation and enhanced K^+ , Ca^{2+} , and Mg^{2+} concentrations, thereby improving the cytosolic K^+/Na^+ ratio. This suggests that MT confers salt tolerance by modulating ion transport and compartmentalization pathways. On a cellular level, melatonin may upregulate tonoplast-localized Na^+/H^+ antiporters (*NHX1*) and K^+ channels (*AKT1*), as demonstrated in other species like crab apple, maize, and rapeseed (Li et al. 2012; Jiang et al. 2016; Zhao et al. 2018). These proteins mediate Na^+ efflux and K^+ influx to restore ionic equilibrium under salt stress. Additionally, MT likely influences ion flux through ABA and Ca^{2+} -mediated signalling, as seen in the upregulation of Ca^{2+} transporters and SOS pathway components (e.g., *SOS1*, *SOS2*) in other model systems (Vafadar et al. 2020).

The observed increase in Ca^{2+} content under salt stress in *S. okadae* may reflect the activation of Ca^{2+} -dependent signalling pathways. Ca^{2+} sensors such as calmodulin (CaM), CBLs, and CDPKs activate downstream targets like non-selective cation channels (NSCCs), cyclic nucleotide-gated channels (CNGCs), and Ca^{2+} -ATPases, contributing to ionic balance and stress signal transduction (Conde et al. 2011; Adams and Shin 2014).

Likewise, Mg^{2+} levels increased under salt stress, particularly at 150 mM NaCl, aligning with chlorophyll enhancement observed under the same treatment. As Mg^{2+} is integral to chlorophyll structure and enzyme function, its accumulation may support both photosynthesis and protein synthesis under stress (Gerendás and Führes 2013).

Together, these results suggest that *S. okadae* employs a coordinated ionic stress mitigation strategy involving selective ion uptake, intracellular sequestration, and signalling crosstalk. Melatonin strengthens this mechanism by enhancing ion homeostasis, reducing ionic toxicity, and modulating transporter gene expression. Future studies should validate the expression of key transporters (*NHX1*, *HKT1*, *AKT1*, *SOS1/2*) in *S. okadae* to substantiate the proposed pathways and further elucidate the melatonin-mediated ionic regulatory network under salt stress.

Ionic Composition

Salt stress imposes hyper-ionic conditions in plant tissues, primarily due to elevated sodium accumulation and disrupted potassium uptake. In potato, the accumulation of Na^+ in shoot tissues and the antagonistic suppression of K^+ uptake under high salinity disrupt cellular ion balance, affecting membrane integrity, enzymatic activity, and inducing oxidative stress through reactive oxygen species (ROS) such as superoxide radicals and hydrogen peroxide (Ishikawa and Shabala 2019). The

maintenance of cytosolic K^+/Na^+ homeostasis is thus a critical determinant of salinity tolerance.

In the present study, salt stress led to a clear increase in Na^+ concentrations and a progressive decline in K^+ levels in *S. okadae* plantlets, particularly beyond 50 mM NaCl, consistent with observations in *S. tuberosum* and *S. chacoense* (Zaki and Radwan 2022; Altaf et al. 2021). Interestingly, despite significant Na^+ accumulation, the K^+ concentrations in *S. okadae* remained relatively stable across treatments, suggesting an intrinsic capacity to buffer Na^+ toxicity and preserve K^+ -dependent metabolic processes. This trait is associated with higher salt tolerance and has been attributed to active ion transport and compartmentalization mechanisms (Maathuis and Amtmann 1999).

MT application under 200 mM NaCl significantly reduced Na^+ levels and improved K^+ retention, increasing the $[K^+]/[Na^+]$ ratio, a recognized marker of salinity tolerance (Flowers and Läuchli, 1983). On a genetic level, melatonin is known to regulate key transporter genes, including *NHX1* and *AKT1*, facilitating Na^+ sequestration into vacuoles and enhanced K^+ uptake (Jiang et al. 2016). In *S. okadae*, the improvement in ionic balance by melatonin likely reflects the upregulation of these ion transport systems, possibly through ABA-dependent or Ca^{2+} -mediated signalling cascades (Kaya et al. 2019). This is further supported by increased Ca^{2+} concentrations in MT-treated plants, which are known to activate SOS pathway components (e.g., *SOS3*, *SOS2*) that regulate *SOS1*-mediated Na^+ efflux (Qiu et al. 2002).

The Ca^{2+} influx observed in MT-treated plants may also trigger downstream signalling through calmodulin, CDPKs, and CBL-CIPK modules, enhancing the expression or activity of plasma membrane and tonoplast transporters (Conde et al. 2011). These interactions likely contribute to reduced cytosolic Na^+ concentrations and stable cellular function under high salt.

Magnesium, a key structural and enzymatic cofactor, also increased significantly under salt stress, particularly at 150 mM NaCl, correlating with elevated chlorophyll content and improved growth. This may reflect effective uptake and utilization of Mg^{2+} in *S. okadae*, possibly supported by reduced antagonism from Ca^{2+} at the root interface (Grattan and Grieve 1999). The positive association between Mg^{2+} and photosynthetic capacity highlights its contribution to salinity tolerance by supporting chlorophyll biosynthesis, protein stability, and enzymatic reactions (Chen et al. 2018; Ishfaq et al. 2021).

MT enhanced Mg^{2+} and Ca^{2+} levels in stressed plants, which may further stabilize membranes, regulate ROS levels, and promote photosynthesis. These results align with earlier findings in wheat and cotton where melatonin increased K^+ and Ca^{2+} and suppressed Na^+ accumulation under salt and metal stress (Jiang et al. 2021; Kaya et al. 2019). Melatonin's role in ion regulation appears to extend beyond direct transporter activation to include crosstalk with hormonal and redox signalling, influencing both transporter expression and ion partitioning.

Additionally, MT-mediated protection of ion homeostasis may involve vacuolar compartmentalization of Na^+ via V-ATPase-driven NHX-type antiporters, preventing cytosolic ion toxicity while maintaining osmotic balance (Yang et al. 2010; Yuan et al. 2016). The coordination between Ca^{2+} signalling and MT may facilitate long-distance stress signalling, as reported in *Dracocephalum kotschy* (Vafadar

et al. 2020), further underscoring the integrative role of melatonin in systemic salt tolerance.

In summary, *S. okadae* exhibits a robust capacity for ion homeostasis under salt stress, which is significantly enhanced by melatonin through modulation of key transporter systems (*NHX1*, *AKT1*, *SOS1*), signalling elements (Ca^{2+} /CaM, ABA), and cytosolic $[\text{K}^+]/[\text{Na}^+]$ maintenance. These results move beyond phenotypic observations to provide mechanistic insight into the molecular and physiological pathways through which melatonin confers salt tolerance in *S. okadae*. Future research should target the expression analysis of ion transporter and signalling genes to validate the regulatory network underlying MT-mediated salt stress resilience.

Proline Accumulation

Proline is one of the most critical osmolytes accumulated in plants in response to salinity and other abiotic stresses. Beyond its traditional role in osmotic adjustment, proline functions as a molecular chaperone, ROS scavenger, redox buffer, and signalling molecule (Sharma et al. 2011). The accumulation of proline is typically regulated by increased expression of biosynthetic genes such as *P5CS* and *P5CR* and/or repression of degradation pathway genes like *PDH* and *P5CDH* (Marco et al. 2015).

In the present study, proline concentration increased significantly in *Solanum okadae* plantlets at 150 mM NaCl, indicating its functional role in osmotic protection and stress adaptation. The elevated proline levels suggest either enhanced biosynthesis through the glutamate pathway or suppressed degradation, consistent with metabolic shifts under moderate salt stress. This accumulation likely aids in ROS detoxification and cellular homeostasis, particularly in leaf tissues where photosynthetic activity must be preserved (Silva-Ortega et al. 2008).

Interestingly, under severe salt stress (200 mM NaCl), proline levels declined markedly, which contrasts with the common expectation of continued accumulation. This reduction may signify a threshold beyond which the metabolic cost of osmolyte synthesis becomes unsustainable, or it could indicate activation of catabolic pathways due to cellular damage or altered hormonal signalling. These findings challenge the notion that proline accumulation is linearly correlated with tolerance and support the view that proline dynamics are tightly regulated in a context-dependent manner.

Melatonin application further modulated proline responses. At 200 mM NaCl, exogenous MT significantly decreased proline content relative to NaCl-only treatment. This decline suggests a shift in stress mitigation strategy from osmolyte-based protection to melatonin-mediated ROS detoxification. MT's antioxidant properties likely reduce the cellular need for proline accumulation as a secondary defence. This is consistent with observations in *Brassica juncea* and fenugreek, where MT decreased proline under drought or salt stress while still improving plant tolerance (Park et al. 2021; Zamani et al. 2020).

Functionally, MT may regulate proline metabolism by downregulating *P5CS* or upregulating *PDH*, possibly through redox or hormone-mediated signalling pathways, including interactions with ABA and nitric oxide (NO). The reduced proline

content in MT-treated *S. okadae* under high salt stress thus reflects an integrated regulatory network where MT substitutes for or suppresses traditional osmotic stress responses by enhancing antioxidant capacity and maintaining ion homeostasis. This underscores the potential of melatonin as a key regulator of cellular reprogramming under abiotic stress in wild potato genotypes.

Catalase Activity and Hydrogen Peroxide (H_2O_2) Accumulation

Under salt stress, the overproduction of reactive oxygen species (ROS), particularly hydrogen peroxide (H_2O_2), can overwhelm the plant's antioxidant defences, leading to lipid peroxidation, protein oxidation, membrane instability, and impaired cellular metabolism (Yazici et al. 2007). In this study, *Solanum okadae* exhibited a salt concentration-dependent increase in catalase (CAT) activity, indicating the activation of its enzymatic ROS-scavenging system in response to oxidative stress. This adaptive response mirrors that observed in other salt-tolerant species such as *Cassia angustifolia*, *Sesamum indicum*, and *Zea mays* (Agarwal and Pandey 2004; Koca et al. 2008; de Azevedo Neto et al. 2006) and aligns with reports in salt-tolerant potato genotypes where antioxidant enzyme activities increased proportionally with stress severity (Sanwal et al. 2022).

Catalase, one of the primary ROS-detoxifying enzymes, decomposes H_2O_2 into water and oxygen, thereby preventing oxidative damage to organelles and maintaining redox balance. The observed increase in CAT activity in *S. okadae* under salinity highlights its potential for maintaining cellular homeostasis during stress, supporting its classification as a salt-tolerant species.

Exogenous melatonin further modulated the oxidative response. Melatonin treatment led to a significant reduction in H_2O_2 accumulation, as evidenced by histochemical DAB staining, accompanied by a moderate decrease in CAT activity. This inverse relationship between ROS content and CAT activity in MT-treated plantlets suggests that melatonin not only enhances antioxidant enzyme expression but also acts directly as a potent ROS scavenger. As a multifunctional antioxidant, melatonin's amphiphilic structure enables it to neutralize ROS such as H_2O_2 and superoxide radicals across various subcellular compartments (Szafrńska et al. 2016).

At the level of physiological mechanism, MT application may prime antioxidant defences through both transcriptional regulation and feedback inhibition. MT has been shown to upregulate genes encoding *SOD*, *CAT*, and *APX* under salinity in model systems like *Arabidopsis* and wheat (Chen et al. 2018; Ke et al. 2018), while also downregulating ROS production by stabilizing plasma membranes and suppressing Na^+ uptake, factors that otherwise exacerbate oxidative stress. The reduced H_2O_2 content observed in MT-treated *S. okadae* may thus be attributed not only to enhanced enzymatic detoxification but also to the upstream regulation of ion homeostasis and membrane integrity.

Furthermore, MT has been reported to increase the expression of *TaSNAT*, a key gene in its biosynthesis pathway, promoting endogenous MT accumulation under salt stress (Ke et al. 2018). This endogenous elevation could create a feedback loop

that enhances systemic ROS tolerance without triggering excessive antioxidant enzyme activity, thereby conserving energy for growth and development.

Taken together, the findings from this study demonstrate that melatonin confers oxidative stress protection in *S. okadae* through a dual mechanism: (1) direct scavenging of ROS and (2) transcriptional modulation of enzymatic antioxidant systems. These results extend beyond confirmatory observations by illustrating how melatonin orchestrates redox homeostasis and contributes to the broader physiological adaptation of *S. okadae* under high salinity stress.

Salt Tolerance Trait Index (STTI)

Salt stress disrupts multiple physiological processes in plants, including water balance, ion homeostasis, redox stability, and hormonal regulation. Plants adapt by activating a suite of interconnected mechanisms such as hormonal signalling, ion exclusion and compartmentalization, osmotic adjustment, and detoxification of reactive oxygen species (ROS) (Rana et al. 2019). In wild relatives and landraces, these responses tend to be more robust and integrated than in commercial cultivars, which have undergone intensive selection for yield and marketability at the expense of abiotic stress resilience (Morton et al. 2019). For instance, landraces of tomato have shown superior salinity tolerance compared to commercial hybrids, linked to higher K^+/Na^+ ratios, and improved ion regulation (Massaretto et al. 2018).

The Salt Tolerance Trait Index (STTI) was employed in this study as a trait-specific metric to evaluate the relative tolerance of *Solanum okadae* to different salt concentrations and to assess the effect of exogenous MT. The STTI provided insights into how different functional traits respond to salt stress, revealing that *S. okadae* tolerates salinity through distinct physiological mechanisms at different salt levels.

At 50 mM NaCl, *S. okadae* exhibited the highest STTI for morphological traits (89.76%) and photosynthetic pigments (31.29%), indicating that early stress conditions allow the plant to maintain growth and pigment stability. Several morphological traits at 50 mM NaCl were statistically like the control. This suggests efficient osmotic adjustment and possibly minimal ion toxicity at lower salt levels. According to the overall STTI analysis, growth and pigment traits were also most stable at this concentration. These findings support earlier studies by Sanwal et al. (2022), who associated salt tolerance in potato genotypes with sustained photosynthetic activity, osmolyte accumulation, and antioxidant enzyme activity.

At 150 mM NaCl, the highest STTI values were observed for ionic content (145.78%) and biochemical traits such as CAT activity, proline, and protein content (108.01%). This implies that under moderate-to-high stress, the plant shifts to a biochemical and ionic adaptation mode, engaging antioxidant defences, osmoprotectant accumulation (e.g., proline), and enhanced ion regulation (e.g., K^+ retention and Na^+ sequestration). The strong ionic STTI values suggest the activation of efficient transport or compartmentalization mechanisms, potentially involving vacuolar Na^+ sequestration or membrane-bound ion channels. Such responses may be mediated by

upregulation of genes like *SOS1*, *NHX1*, and *CAT*, which are known to contribute to ion compartmentalization and ROS detoxification in salt-tolerant species.

The highest overall STIT value (81.84%) at 150 mM NaCl further supports that this concentration represents an upper physiological threshold where *S. okadae* can still maintain integrated functional performance across diverse traits. This aligns with the notion that wild potato species possess adaptive traits that are often lost during domestication (Morton et al. 2019), yet in rare cases, such traits have been retained or reintroduced through breeding (Chourasia et al. 2021). Thus, *S. okadae* demonstrates a high potential as a donor for salt tolerance traits in potato improvement programs.

At 200 mM NaCl, STTI values dropped for morphological and pigment traits, indicating substantial growth suppression and pigment degradation. However, relatively high STTI values for ionic content (103.42%) and antioxidant-related traits (101.55%) were still observed, suggesting that the plants, although morphologically impaired, retained physiological activity. This was further supported by measurable levels of proline and CAT activity, indicating sustained biochemical adaptation in response to severe stress.

By quantifying STTI across functional trait categories and under melatonin treatment, this study provides a mechanistic understanding of how *S. okadae* manages salinity stress and highlights trait-specific physiological strategies that may be exploited for crop improvement. Rather than merely confirming prior knowledge, this approach contributes new insights into the specific adaptive pathways activated in a wild diploid potato species under salt stress, laying the foundation for targeted breeding and molecular validation efforts.

Conclusion and Future Prospects

This study highlights the inherent salt tolerance of *Solanum okadae*, a wild diploid potato species, and demonstrates how this resilience can be further enhanced through the exogenous application of melatonin. Even at high salinity levels (up to 200 mM NaCl), *S. okadae* maintained physiological activity, suggesting robust intrinsic mechanisms for osmotic adjustment, ion homeostasis, and oxidative stress mitigation. The application of melatonin significantly improved shoot and root development, chlorophyll retention, and ionic balance while reducing oxidative damage markers such as hydrogen peroxide and catalase activity. These results underscore the dual advantage of utilizing *S. okadae* as a pre-breeding resource for salt tolerance traits and melatonin as an effective biostimulant to bolster stress responses under saline conditions.

A limitation of this study is that melatonin was applied only at the highest salinity level (200 mM NaCl), which restricts conclusions about its effects across different degrees of salt stress. Applying melatonin across a broader range of salinity treatments would provide more robust insights into its dose-dependent efficacy and potential threshold effects. Future work should expand melatonin application to all salinity levels tested to fully characterize its protective role and interaction with

stress severity. Such studies will better inform the use of melatonin in crop improvement under diverse saline conditions.

Looking ahead, further studies are needed to unravel the molecular and genetic bases of both *S. okadae*'s natural salt tolerance and melatonin-mediated stress alleviation. Transcriptomic and metabolomic profiling could identify key regulatory genes and pathways involved, while field-based evaluations are essential to validate these findings under real agronomic conditions. Since the current work was conducted exclusively under controlled in vitro conditions, the results should be considered preliminary, and further testing in greenhouse and field environments is necessary to confirm their applicability in practical agriculture. Moreover, exploring the genetic engineering of melatonin biosynthesis genes into cultivated varieties, along with investigating melatonin's interactions with other plant hormones, could broaden its potential for crop improvement. Combining the resilience of wild species with plant-derived signalling molecules like melatonin offers a promising approach to enhance stress tolerance and sustain crop productivity in increasingly saline environments.

Acknowledgements Amar Hundare was supported by a Vice Chancellor's scholarship from the University of Nottingham to conduct the research. The manuscript was prepared and finalized at the University of Southampton, where the corresponding author is currently affiliated. The authors thank both institutions for their support. We thank the Morley Agricultural Foundation (Norfolk, UK) for financial assistance. We thank the Commonwealth Potato Collection (CPC) maintained by the James Hutton Institute (JHI, Invergowrie, Scotland) for all potato germplasm used in this study.

Author Contribution Amar H: Conceptualization, experimental execution, data collection and analysis, and writing original draft of the manuscript. Aneesh L: Histochemical analysis, manuscript reviewing, and editing. Neelu J: Data analysis, manuscript reviewing, and editing. Tim R: Manuscript reviewing and editing. The final manuscript was read and approved by all authors.

Data Availability The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adams E, Shin R (2014) Transport, signalling, and homeostasis of potassium and sodium in plants. *J Integr Plant Biol* 56:231–249. <https://doi.org/10.1111/JIPB.12159>

- Agarwal S, Pandey V (2004) Antioxidant enzyme responses to NaCl stress in *Cassia angustifolia*. *Biol Plant* 48:555–560. <https://doi.org/10.1023/B:BIOP.0000047152.07878.E7/METRICS>
- Ahmed HAA, Şahin NK, Akdoğan G et al (2020) Variability in salinity stress tolerance of potato (*Solanum tuberosum* L.) varieties using in vitro screening. *Cienc Agrotecnol* 44:1–14. <https://doi.org/10.1590/1413-7054202044004220>
- Altaf MA, Shahid R, Ren MX et al (2021) Melatonin mitigates nickel toxicity by improving nutrient uptake fluxes, root architecture system, photosynthesis, and antioxidant potential in tomato seedling. *J Soil Sci Plant Nutr* 21:1842–1855. <https://doi.org/10.1007/S42729-021-00484-2/FIGURES/13>
- Altaf MA, Shahid R, Altaf MM et al (2022) Melatonin: first-line soldier in tomato under abiotic stress current and future perspective. *Plant Physiol Biochem* 185:188–197. <https://doi.org/10.1016/J.PLAPHY.2022.06.004>
- Arnao MB, Hernández-Ruiz J (2019) Melatonin and reactive oxygen and nitrogen species: a model for the plant redox network. *Melatonin Res* 2:152–168. <https://doi.org/10.32794/11250036>
- Arnao MB, Hernández-Ruiz J, Cano A (2022) Role of melatonin and nitrogen metabolism in plants: implications under nitrogen-excess or nitrogen-low. *Int J Mol Sci* 23:15217. <https://doi.org/10.3390/IJMS232315217>
- Arnon DI (1949) Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiol* 24:1. <https://doi.org/10.1104/PP.24.1.1>
- Banu MSA, Huda KMK, Sahoo RK et al (2015) Pea p68 imparts salinity stress tolerance in rice by scavenging of ROS-mediated H₂O₂ and interacts with Argonaute. *Plant Mol Biol Report* 33:221–238. <https://doi.org/10.1007/S11105-014-0748-7/FIGURES/8>
- Biswas MS, Islam MR, Zakaria M (2017) Evaluation of indigenous potato Challisha (*Solanum tuberosum* L. cv. Challisha) somaclonals tolerance to salinity in vitro. *J Trop Life Sci* 7:77–82. <https://doi.org/10.11594/JTLS.07.01.13>
- Camadro EL, Saffarano SK, Espinillo JC et al (2008) Cytological mechanisms of 2n pollen formation in the wild potato *Solanum okadae* and pollen-pistil relations with the cultivated potato, *Solanum tuberosum*. *Genet Resour Crop Evol*. <https://doi.org/10.1007/S10722-007-9254-1>
- Charfeddine M, Charfeddine S, Ghazala I et al (2019) Investigation of the response to salinity of transgenic potato plants overexpressing the transcription factor StERF94. *J Biosci*. <https://doi.org/10.1007/S12038-019-9959-2>
- Chen T, Zhang B (2016) Measurements of proline and malondialdehyde content and antioxidant enzyme activities in leaves of drought stressed cotton. *Bio-protocol* 6(17):e1913. <https://doi.org/10.21769/BioProtoc.1913>
- Chen ZC, Peng WT, Li J, Liao H (2018) Functional dissection and transport mechanism of magnesium in plants. *Semin Cell Dev Biol* 74:142–152. <https://doi.org/10.1016/J.SEMCDB.2017.08.005>
- Chourasia KN, Lal MK, Tiwari RK et al (2021) Salinity stress in potato: understanding physiological, biochemical and molecular responses. *Life*. <https://doi.org/10.3390/LIFE11060545>
- Conde A, Chaves MM, Gerós H (2011) Membrane transport, sensing and signalling in plant adaptation to environmental stress. *Plant Cell Physiol* 52:1583–1602. <https://doi.org/10.1093/PCP/PCR107>
- Daneshmand F, Arvin MJ, Kalantari KM (2010) Physiological responses to NaCl stress in three wild species of potato *in vitro*. *Acta Physiol Plant* 32:91–101. <https://doi.org/10.1007/S11738-009-0384-2>
- Danku JMC, Lahner B, Yakubova E, Salt DE (2013) Large-scale plant ionomics. In: Maathuis F (ed) *Plant mineral nutrients. Methods in molecular biology*, vol 953. Humana Press, Totowa, NJ. https://doi.org/10.1007/978-1-62703-152-3_17
- Daudi A, O'Brien, JA (2012) Detection of hydrogen peroxide by DAB staining in *Arabidopsis* leaves. *Bio-protocol* 2(18):e263. <https://doi.org/10.21769/BioProtoc.263>
- de Azevedo Neto AD et al (2006) Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salt-tolerant and salt-sensitive maize genotypes. *Environ Exp Bot* 56(1):87–94
- Efimova MV, Danilova ED, Zlobin IE et al (2023) Priming potato plants with melatonin protects stolon formation under delayed salt stress by maintaining the photochemical function of photosystem II, ionic homeostasis and activating the antioxidant system. *Int J Mol Sci*. <https://doi.org/10.3390/IJMS24076134>
- Evers D, Overney S, Simon P et al (1999) Salt tolerance of *Solanum tuberosum* L. overexpressing a heterologous osmotin-like protein. *Biol Plant* 42:105–112. <https://doi.org/10.1023/A:1002131812340/METRICS>
- Fernandez GC (1992) Effective selection criteria for assessing stress tolerance. In: Kuo CG (ed) *Proceedings of the international symposium on adaptation of vegetables and other food crops in temperature and water stress*, AVRDC Publication, Tainan, p 257–270

- Flowers TJ, Läuchli A (1983) Sodium versus potassium: substitution and compartmentation. In: Läuchli A, Bielecki RL (eds) Inorganic plant nutrition, vol. 15B. Springer-Verlag, Berlin, pp 651–681
- Gelmesa D, Dechassa N, Mohammed W et al (2017) In vitro screening of potato genotypes for osmotic stress tolerance. *Open Agric* 2:308–316. <https://doi.org/10.1515/OPAG-2017-0035>
- Gerendás J, Führs H (2013) The significance of magnesium for crop quality. *Plant Soil* 368:101–128. <https://doi.org/10.1007/S11104-012-1555-2>
- Grattan S, Grieve C (1999) Mineral nutrient acquisition and response by plants grown in saline environments. In: Pessarakli M (ed) Handbook of plant and crop stress, 2nd ed. Marcel Dekker, New York . pp 203–229. <https://doi.org/10.1201/9780824746728.ch9>
- Gupta B, Huang B (2014) Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. *Int J Genomics* 2014:701596. <https://doi.org/10.1155/2014/701596>
- Hernandez JA et al (2000) Antioxidant systems and $O_2^{\cdot-}/H_2O_2$ production in chloroplasts of pea plants exposed to salt stress. *Plant Cell Environ* 23(5):515–524
- Hundare A (2023) *Solanum okadae*: a wild diploid potato species with the potential to improve cultivated potato. Ph.D. Thesis, University of Nottingham. Available at: <https://eprints.nottingham.ac.uk/73456/>
- Ishfaq M, Zhong Y, Wang Y, Li X (2021) Magnesium limitation leads to transcriptional down-tuning of auxin synthesis, transport, and signaling in the tomato root. *Front Plant Sci* 12:802399. <https://doi.org/10.3389/FPLS.2021.802399>
- Ishikawa T, Shabala S (2019) Control of xylem Na^+ loading and transport to the shoot in rice and barley as a determinant of differential salinity stress tolerance. *Physiol Plant* 165:619–631. <https://doi.org/10.1111/PPL.12758>
- Jiang C, Cui Q, Feng K et al (2016) Melatonin improves antioxidant capacity and ion homeostasis and enhances salt tolerance in maize seedlings. *Acta Physiol Plant* 38:1–9. <https://doi.org/10.1007/S11738-016-2101-2/FIGURES/6>
- Jiang D, Lu B, Liu L et al (2021) Exogenous melatonin improves the salt tolerance of cotton by removing active oxygen and protecting photosynthetic organs. *BMC Plant Biol* 21:1–19. <https://doi.org/10.1186/S12870-021-03082-7/FIGURES/11>
- Juan M, Rivero RM, Romero L, Ruiz JM (2005) Evaluation of some nutritional and biochemical indicators in selecting salt-resistant tomato cultivars. *Environ Exp Bot* 54:193–201. <https://doi.org/10.1016/J.ENVEXPBOT.2004.07.004>
- Kaya C et al (2019) Melatonin-mediated nitric oxide improves tolerance to cadmium toxicity by reducing oxidative stress in wheat plants. *Chemosphere* 225:627–638. <https://doi.org/10.1016/j.chemosphere.2019.03.026>
- Kaya C, Higgs D, Ashraf M et al (2020) Integrative roles of nitric oxide and hydrogen sulfide in melatonin-induced tolerance of pepper (*Capsicum annuum* L.) plants to iron deficiency and salt stress alone or in combination. *Physiol Plant* 168:256–277. <https://doi.org/10.1111/PPL.12976>
- Ke Q, Ye J, Wang B et al (2018) Melatonin mitigates salt stress in wheat seedlings by modulating polyamine metabolism. *Front Plant Sci* 9:391686. <https://doi.org/10.3389/FPLS.2018.00914/BIBTEX>
- Kitajima K, Hogan KP (2003) Increases of chlorophyll a/b ratios during acclimation of tropical woody seedlings to nitrogen limitation and high light. *Plant Cell Environ* 26:857–865. <https://doi.org/10.1046/J.1365-3040.2003.01017.X>
- Koca H et al (2008) Responses of pepper to salinity and the role of exogenous glycine betaine. *Biol Plant* 52(4):626–630
- Kolomeichuk L, Efimova MV, Zlobin IE et al (2020) 24-epibrassinolide alleviates the toxic effects of NaCl on photosynthetic processes in potato plants. *Photosynth Res* 146:151–163. <https://doi.org/10.1007/s11120-020-00708-z>
- Levy D, Veilleux RE (2007) Adaptation of potato to high temperatures and salinity—a review. *Am J Potato Res* 84(6):84–506. <https://doi.org/10.1007/BF02987885>
- Li C, Wang P, Wei Z, Liang D, Liu C, Yin L, Jia D, Fu M, Ma F (2012) The mitigation effects of exogenous melatonin on salinity-induced stress in *Malus hupehensis*. *J Pineal Res*. 53(3):298–306. <https://doi.org/10.1111/j.1600-079X.2012.00999.x>
- Liang C, Li A, Yu H et al (2017) Melatonin regulates root architecture by modulating auxin response in rice. *Front Plant Sci* 8:134. <https://doi.org/10.3389/fpls.2017.00134>
- Lone JA, Lone FA, Suseela MR (2018) Photosynthetic pigment quantification from two green microalgal species of *Scenedesmus*. *IJARSE* 7:2220–2230
- Maathuis FJM, Amtmann A (1999) K^+ nutrition and Na^+ toxicity: the basis of cellular K^+/Na^+ ratios. *Ann Bot* 84:123–133. <https://doi.org/10.1006/ANBO.1999.0912>

- Marco F, Bitrián M, Carrasco P et al (2015) Genetic engineering strategies for abiotic stress tolerance in plants. *Plant Biol Biotechnol Vol II Plant Genomics Biotechnol*, pp 579–609. https://doi.org/10.1007/978-81-322-2283-5_29/COVER
- Martinez V, Nieves-Cordones M, Lopez-Delacalle M et al (2018) Tolerance to stress combination in tomato plants: new insights in the protective role of melatonin. *Molecules* 23(3):535. <https://doi.org/10.3390/molecules23030535>
- Massaretto IL, Albaladejo I, Purgatto E et al (2018) Recovering tomato landraces to simultaneously improve fruit yield and nutritional quality against salt stress. *Front Plant Sci* 871:412396. <https://doi.org/10.3389/FPLS.2018.01778/BIBTEX>
- Misra AN, Sahu SM, Misra M et al (1997) Genotypic differences in salinity tolerance of rice seedlings. *Indian J Plant Physiol* 2(2):126–130
- Morton MJL, Awlia M, Al-Tamimi N et al (2019) Salt stress under the scalpel – dissecting the genetics of salt tolerance. *Plant J* 97:148–163. <https://doi.org/10.1111/TPJ.14189>
- Murch SJ, Erland LE (2021) Melatonin in plants: biosynthesis and physiological roles. *Plant Signal Behav* 16(8):1901963
- Park HS, Kazerooni E, Kang SM, Al-Sadi AM, Lee IJ (2021) Melatonin enhances the tolerance and recovery mechanisms in *Brassica juncea* (L.) Czern. under saline conditions. *Front Plant Sci* 12:593717. <https://doi.org/10.3389/fpls.2021.59371777>
- Peethambaran PK, Glenz R, Höninger S et al (2018) Salt-inducible expression of OsJAZ8 improves resilience against salt-stress. *BMC Plant Biol*. <https://doi.org/10.1186/S12870-018-1521-0>
- Qiu QS et al (2002) SOS1 functions in Na⁺ efflux in *Arabidopsis*. *Proc Natl Acad Sci* 99(12):8436–8441.81
- Rana MM, Takamatsu T, Baslam M et al (2019) Salt tolerance improvement in rice through efficient SNP marker-assisted selection coupled with speed-breeding. *Int J Mol Sci* 20:2585. <https://doi.org/10.3390/IJMS20102585>
- Rashid MHO, Islam S, Bari M (2019) In vitro screening for salt stress tolerance of native and exotic potato genotypes by morphological and physiological parameters. *J Bio-Science* 28:21–32. <https://doi.org/10.3329/JBS.V28I0.44707>
- Sack L, John GP, Buckley TN (2018) ABA accumulation in dehydrating leaves is associated with decline in cell volume, not turgor pressure. *Plant Physiol* 176:489. <https://doi.org/10.1104/PP.17.01097>
- Sanwal SK, Kumar P, Kesh H et al (2022) Salinity stress tolerance in potato cultivars: evidence from physiological and biochemical traits. *Plants* 11:1842. <https://doi.org/10.3390/PLANTS11141842>
- Sarker U, Islam MT, Oba S (2018) Salinity stress accelerates nutrients, dietary fiber, minerals, phytochemicals and antioxidant activity in *Amaranthus tricolor* leaves. *PLoS One* 13:e0206388. <https://doi.org/10.1371/JOURNAL.PONE.0206388>
- Sharma S, Villamor JG, Verslues PE (2011) Essential role of tissue-specific proline synthesis and catabolism in growth and redox balance at low water potential. *Plant Physiol* 157:292–304. <https://doi.org/10.1104/PP.111.183210>
- Shrestha B, Darapuneni M, Stringam BL et al (2023) Irrigation water and nitrogen fertilizer management in potato (*Solanum tuberosum* L.): a review. *Agronomy* 13(13):2566. <https://doi.org/10.3390/AGRONOMY13102566>
- Silva-Ortega CO, Ochoa-Alfaro AE, Reyes-Agüero JA et al (2008) Salt stress increases the expression of p5cs gene and induces proline accumulation in cactus pear. *Plant Physiol Biochem* 46:82–92. <https://doi.org/10.1016/j.plaphy.2007.10.011>
- Stoleru V, Slabu C, Vitanescu M et al (2019) Tolerance of three Quinoa Cultivars (*Chenopodium quinoa* Willd.) to salinity and alkalinity stress during germination stage. *Agron* 9:9:287. <https://doi.org/10.3390/AGRONOMY9060287>
- Sudharsan C, Manuel SJ, Ashkanani J, Al-Ajeel A (2012) In vitro screening of potato cultivars for salinity tolerance. *J Sustain Agric* 6:344–348
- Sultana N, Ikeda T, Itoh R (1999) Effect of NaCl salinity on photosynthesis and dry matter accumulation in developing rice grains. *Environ Exp Bot* 42:211–220. [https://doi.org/10.1016/S0098-8472\(99\)00035-0](https://doi.org/10.1016/S0098-8472(99)00035-0)
- Szafránska K et al (2016) Melatonin as a plant antioxidant: implications for crop improvement. *Front Plant Sci* 7:1989
- Tan D et al (2012) Functional roles of melatonin in plants and perspectives in nutritional and agricultural science. *J Exp Bot* 63(2):577–597. <https://doi.org/10.1093/jxb/err256>

- Tan DX, Reiter RJ (2020) An evolutionary view of melatonin synthesis and metabolism related to its biological functions in plants. *J Exp Bot* 71:4677–4689. <https://doi.org/10.1093/JXB/ERAA235>
- Tyree MT, Engelbrecht BMJ, Vargas G, Kursar TA (2003) Desiccation tolerance of five tropical seedlings in Panama. Relationship to a field assessment of drought performance. *Plant Physiol* 132:1439. <https://doi.org/10.1104/PP.102.018937>
- Vafadar F et al (2020) Melatonin and Ca^{2+} signaling in salt stress adaptation. *Environ Exp Bot* 176:104106
- Wang Y, Nii N (2000) Changes in chlorophyll, ribulose biphosphate carboxylase-oxygenase, glycine betaine content, photosynthesis and transpiration in *Amaranthus tricolor* leaves during salt stress. *J Hortic Sci Biotechnol* 75:623–627
- Wang P et al (2015) Melatonin alleviates oxidative stress under salinity in cucumber. *Plant Physiol Biochem* 94:100–107
- Watanabe KN, Kikuchi A, Shimazaki T, Asahina M (2011) Salt and drought stress tolerances in transgenic potatoes and wild species. *Potato Res* 54:319–324
- Yang DH et al (2010) NHX1 involvement in salt stress responses. *Plant Mol Biol Rep* 28(2):195–202
- Yazici I, Türkan I, Sekmen AH, Demiral T (2007) Salinity tolerance of purslane (*Portulaca oleracea* L.) is achieved by enhanced antioxidative system, lower level of lipid peroxidation and proline accumulation. *Environ Exp Bot* 61:49–57. <https://doi.org/10.1016/J.ENVEXPBOT.2007.02.010>
- Yuan F, Leng B, Wang B (2016) Progress in studying salt secretion from the salt glands in recretohalophytes: how do plants secrete salt? *Front Plant Sci* 7:173278. <https://doi.org/10.3389/FPLS.2016.00977>
- Zaki HEM, Radwan KSA (2022) The use of osmoregulators and antioxidants to mitigate the adverse impacts of salinity stress in diploid and tetraploid potato genotypes (*Solanum spp.*). *Chem Biol Technol Agric* 9:1–21. <https://doi.org/10.1186/S40538-022-00286-3>
- Zaman MS, Ali GM, Muhammad A et al (2015) In vitro screening of salt tolerance in potato (*Solanum tuberosum* L.) varieties. *Sarhad J Agric* 31:106–113. <https://doi.org/10.17582/JOURNAL.SJA/2015/31.2.106.113>
- Zamani Z, Amiri H, Ismaili A (2020) Improving drought stress tolerance in fenugreek (*Trigonella foenum-graecum*) by exogenous melatonin. *Plant Biosyst* 154:643–655. <https://doi.org/10.1080/11263504.2019.1674398>
- Zhang Y, Fan Y, Rui C et al (2021) Melatonin improves cotton salt tolerance by regulating ROS scavenging system and Ca^{2+} signal transduction. *Front Plant Sci* 12:693690. <https://doi.org/10.3389/fpls.2021.693690>
- Zhao G et al (2018) Nitric oxide is required for melatonin-enhanced tolerance against salinity stress in rapeseed (*Brassica napus* L.) seedlings. *Int J Mol Sci* 19(7):1912
- Zhou X, Zhao H, Cao K, Hu L, Du T, Baluška F and Zou Z (2016) Beneficial roles of melatonin on redox regulation of photosynthetic electron transport and synthesis of D1 protein in tomato seedlings under salt stress. *Front Plant Sci* 7:1823. <https://doi.org/10.3389/fpls.2016.01823>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.