

Portulaca oleracea exhibited anti-coccidian activity, fortified the gut microbiota of Hu lambs

shiheng li

1575615649@qq.com

Henan Agricultural University

Senyang li

Henan Agricultural University

Shuaiqi liu

Henan Agricultural University

shunli lu

Henan Agricultural University

jing li

Henan Agricultural University

Shuqi cheng

Henan Agricultural University

sumei zhang

Laboratory INNOTECH International: Laboratoire INNOTECH International

Shucheng huang

Henan Agricultural University

junqiang li

Henan Agricultural University

Fuchun jian

Henan Agricultural University

Research Article

Keywords: Anticoccidial Activity, Gut Microbiota, Intestinal, Health Portulaca oleracea

Posted Date: January 24th, 2024

DOI: <https://doi.org/10.21203/rs.3.rs-3763788/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at AMB Express on May 3rd, 2024. See the published version at <https://doi.org/10.1186/s13568-024-01705-4>.

Abstract

Coccidia of the genus *Eimeria* are important pathogens that cause coccidiosis in livestock and poultry. Due to the expansion of intensive farming, coccidiosis has become more difficult to control. In addition, the continued use of anti-coccidiosis drugs has led to drug resistance and residue. Some herbs used in traditional Chinese medicine (TCM) have been shown to alleviate the clinical symptoms of coccidiosis, while enhancing immunity and growth performance (GP) of livestock and poultry. Previous *in vitro* and *in vivo* studies have reported that the TCM herb *Portulaca oleracea* exhibited anti-parasitic activities. In total, 36 female Hu lambs were equally divided into six treatment groups: PL (low-dose *P. oleracea*), PH (high-dose *P. oleracea*), PW (*P. oleracea* water extract), PE (*P. oleracea* ethanol extract), DIC (diclazuril), and CON (control). The treatment period was 14 days. The McMaster counting method was used to evaluate the anti-coccidiosis effects of the different treatments. Untargeted metabolomics and 16S rRNA gene sequencing were used to investigate the effects of treatment on the gut microbiota (GM) and GP. The results showed that *P. oleracea* ameliorated coccidiosis, improved GP, increased the abundances of beneficial bacteria, and maintained the composition of the GM, but failed to completely clear coccidian oocysts. The *Firmicutes* to *Bacteroides* ratio was significantly increased in the PH group. *P. oleracea* increased metabolism of tryptophan as well as some vitamins and cofactors in the GM and decreased the relative content of arginine, tryptophan, niacin, and other nutrients, thereby promoting intestinal health and enhancing GP. As an alternative to the anti-coccidiosis drug DIC, *P. oleracea* effectively inhibited growth of coccidia, maintained the composition of the GM, promoted intestinal health, and increased nutrient digestibility.

Introduction

Coccidiosis is generally a self-limiting infection caused by coccidia of the genus *Eimeria* (Kemp LE et al. 2013). Coccidia are widely distributed globally, highly host-specific, and can infect a wide range of vertebrates, including cattle, sheep, pigs, rabbits, chickens, and ducks (Yun CH et al. 2000; Keeton STN et al. 2018). Massive parasitic reproduction of coccidia causes severe damage to the mucosa of the host digestive tract, which reduces digestion and absorption of nutrients, resulting in diarrhea, stunted growth, and even death (Barba E et al. 2022). The global economic cost of coccidiosis control is estimated at around 13 billion USD annually (Blake DP et al. 2020). With the change to animal husbandry from free-range to large-scale intensive farming, the stocking density continues to increase, which leads to rapid spread of coccidia. Besides, it is difficult to prevent and control coccidiosis because of the high reproductive potential of ubiquitous coccidian oocysts (Allen PC et al. 2002). At present, antibiotics and triazines are commonly used for treatment of coccidiosis. However, the use of chemical-based drugs in animal husbandry is no longer sufficient due to widespread drug resistance, drug residues in food, and threats to the environment (Dauguschies A et al. 1998; Waller PJ 2006; Paul SM et al. 2010; Bacanlı M. Başaran N. 2019). And the cost of drug development continues to rise, finding alternative anti-coccidian drugs has become an urgent priority.

Various natural plant products used in traditional Chinese medicine (TCM) are effective for treatment of parasitic diseases. Herbs used in TCM for control of coccidiosis have the advantages of wide availability, broad-spectrum applications, safety, environmental compliancy, and economic efficiency. The antimalarial drug artemisinin is reported to significantly alter wall formation of coccidian oocysts, resulting in impaired oocyst development and reduced sporulation rates in chickens (del Cacho E et al. 2010). Moreover, the combination of the plant extract berberine and amprolium was shown to significantly decrease the oocysts per gram (OPG) values of coccidia as compared to amprolium alone. In addition, the combination of berberine and amprolium was found to improve growth performance (GP) (Malik TA et al. 2016). The active ingredients of herbs used in TCM primarily include flavonoids, biophenols, alkaloids, terpenoids, and polyphenols. However, various extraction processes are required to fully activate and enhance the efficacy of these active compounds. And the roots, stems, and leaves of some plants used in TCM are toxic and must be processed before use. Some medicinal plants used in TCM are toxic in relatively large dosages, but have few side effects that after appropriate processing. Hence, the uncertainty in the dosage and form of TCM is challenging for treatment of coccidiosis.

Portulaca oleracea is an annual succulent in the family Portulacaceae used in various medicines and cosmetics (Juraimi AS et al. 2009; Uddin K et al. 2017). The stems and leaves of *P. oleracea* have anti-inflammatory and analgesic effects, and are used in TCM for treatment of gastrointestinal and liver diseases (Liu L et al. 2000). *P. oleracea* has also been used to prevent and treat gastrointestinal parasitic diseases in sheep (Cai Y et al. 2004). However, most previous studies have been conducted using avian or non-ruminant species or only in comparison to chemical deworming agents. Various *in vivo* and *in vitro* studies have shown that the gut microbiota (GM) and related metabolites play important roles in resistance to pathogens. However, relatively few studies have investigated the use of *P. oleracea* to modulate the composition of the GM and metabolite levels to influence the GP and intestinal health of ruminants.

Coccidia infection alters the composition of the GM both directly and indirectly by inducing changes to physiological characteristics, permeability, and antimicrobial peptide production (Zaiss MM, Harris NL. 2016). The pathophysiology of coccidia mainly includes damage to the intestinal mucosa, immunological inflammation, and changes to the GM (Lu C et al. 2021). The GM is a dynamic population of microorganisms that maintains a symbiotic relationship with the gut mucosa (Sebastián Domingo JJ. 2018). Changes to the structural composition of the GM, particularly the ratio of *Firmicutes* to *Bacteroidetes* (F/B ratio), can impact colonization of parasites in the host, infection status, and management of parasitic diseases (White EC et al. 2018; Wen JJ et al. 2021). Besides, some metabolites produced by bacteria in the host gut are relevant to the intestinal barrier. Alterations to the microbial metabolites tryptophan, indoleacetic acid, and indole acetaldehyde regulate intestinal barrier function via the aryl hydrocarbon receptor (Lamas B et al. 2016). However, further investigations are needed to elucidate the mechanism of action employed by *P. oleracea* to alter the composition of the GM and related metabolites during coccidia infection.

In the study, the impact of low and high doses of *P. oleracea*, as well as water and ethanol extracts, on the number of coccidia in lambs were investigated. Additionally, the impacts of *P. oleracea* on the composition of the GM and related metabolites during coccidia infection were analyzed. The results showed that *P. oleracea* decreased the OPG values of coccidia among the different treatment groups, but not as significantly the anti-coccidian drug diclazuril (DIC) in the early stages of treatment. However, the duration of efficacy was relatively longer for *P. oleracea*, which resulted in lower OPG values. The use of *P. oleracea* improved the mean daily weight gain (DWG), diversity of the GM, and F/B ratio of sheep. Overall, the findings of this study support dietary supplementation with *P. oleracea* as a substitute for anti-coccidian drugs and antibiotics.

Methods

Study approval

The study protocol was approved by the Institutional Animal Care and Use Committee of the College of Animal Medicine, Henan Agricultural University (approval no. 17–0126), and conducted in accordance with the Guide for the Care and Use of Agricultural Animals in Research and Teaching. A flowchart of the experimental design is provided in Fig. 1.

Materials and reagents

P. oleracea was purchased from Yuzhou Qianhe Pharmaceutical Co., Ltd. (Yuzhou, China). DIC premix (0.5%) was obtained from Hefei Zhonglong Shenli Animal Pharmaceutical Co., Ltd. (Hefei, China). Food-grade ethanol (75%) was acquired from Shandong Xintai Huayuan Medical and Sanitary Products Co., Ltd. (Xintai, China). A McMaster counting chamber to quantify coccidian oocysts was procured from Henan Agricultural University (Zhengzhou, China).

Preparation of herbal extracts

The herbal extract was prepared by solvent extraction. In brief, the herbs were crushed, macerated for 6 days in a mixture of water and 70% ethanol (1:10), and filtered several times through four layers of gauze and a 60 mesh until there was no residue. Then, the filtrate was evaporated in a boiling water bath and the resulting paste was fixed in saline. The final concentration of the extract was 1 g/1 mL of solvent.

Animals and experimental design

In total, 36 female Hu lambs (age, 1.5 months; mean body weight [BW], 13.08 ± 0.31 kg) were obtained from a large-scale sheep farm and divided into six groups of six animals each. Fecal samples were collected two times per day for 3 days before the experiment and coccidia OPG values were calculated. Lambs in the blank control (CON) group were fed a basal diet. Lambs in the DIC group were fed the basal diet and orally administered DIC at 1 mg/kg BW for 2 consecutive days. Lambs in the low-dose *P. oleracea* (PL) group and high-dose *P. oleracea* (PH) group were fed the basal diet supplemented with *P. oleracea* at 30 g/day and 120 g/day respectively for 14 days. Lambs in the *P. oleracea* water extraction

(PW) group and ethanol extraction (PE) group were fed the basal diet and orally administered a water and ethanol extract of *P. oleracea* respectively at 30 mL/day for 14 days. Feed was offered in equal amounts at 07:00 and 18:00 h daily. All animals had *ad libitum* access to drinking water. The study period was 42 days.

Sample collection

Fecal samples were collected and OPG values were calculated on days 0, 7, 14, 21, 28, 35, and 42. Fecal samples collected on day 14 were stored at -80°C for genomic sequencing of the GM. The BW of each lamb was measured in the morning on days 0, 14, and 42.

Calculation of coccidia OPG

The OPG value was calculated using the McMaster method [12] and reported as the mean number of oocysts in both chambers multiplied by 200.

DNA extraction and 16S rRNA sequencing

DNA was extracted from the fecal samples using the FastDNA™ SPIN Kit for Soil (MP Biomedicals, Heidelberg, Germany) as described in the manufacturer's instructions but with slight modifications. Briefly, 250 mg of each fecal sample were homogenized in the buffer provided with the kit using a FastPrep-24™ Instrument (MP Biomedicals). Cell lysates were separated by centrifugation ($14,000 \times g$, 15 min) and proteins were precipitated from the supernatant. DNA was diluted with pre-warmed (55°C) purified water and quantified using a NanoDrop™ spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA).

The V1-V2 region of the 16S rRNA gene of each sample was amplified and sequenced as previously described (Camarinha-Silva A et al. 2014). The primer pair 27F and 338R was used to amplify the target region, with a slightly modified sequence of the primer 27F (AGRGTTTGATYMTGGCTCAG). The quality of the obtained amplicons was assessed by agarose gel electrophoresis. The qualified amplicons were sequenced with an Illumina MiSeq™ instrument (Illumina, Inc., San Diego, CA, USA) in paired-end mode (2×250 base pairs). Sequence reads were quality filtered and assembled using mothur software (Schloss PD et al. 2009). Reads with an mean quality score < 20 , total length > 355 base pairs (bp), any primer or barcode mismatch, and homopolymer stretches with > 8 Oran characters were excluded. The reads were checked for the presence of chimeras and clustered into operational taxonomical units (OTUs) at 97% identity (Wang Q et al. 2007). OTUs appearing only once across all samples or with fewer than 10 reads were manually deleted. The remaining OTUs were finally assigned to the closest taxonomical representative using the Sequence Match tool (Lighten J et al. 2014).

The abundance of each OTU was assessed using Primer 6 v.6 software (PRIMER-E Ltd., Plymouth, UK). Statistical differences across time points among the treatment groups were identified using analysis of variance (ANOVA) with permutations.

Metabolomic profiling of fecal samples

Fecal samples (50 mg) were homogenized in extraction buffer (1000 µL; methanol, acetonitrile, and water at 2:2:1 chilled to – 20°C; and 20 µL of an internal standard) using a ball mill for 4 min at 45 Hz, ultrasonicated at 5 min, and incubated for 1 h at – 20°C to precipitate the proteins. Following centrifugation for 15 min at 12000 rpm and 4°C, 20 µL aliquots of the supernatants were used for liquid chromatography–mass spectrometry.

Follow-up analysis was performed after normalizing the original peak area to the total peak area. Principal component analysis and Spearman correlation analysis were performed to assess the repeatability of the samples within groups and the quality of the samples. Classification and associated pathways of the identified compounds were determined in reference to the Kyoto Encyclopedia of Genes and Genomes (KEGG; <https://www.genome.jp/kegg/>), Human Metabolome Database (<https://hmdb.ca/>), and Lipid Maps database (<https://www.lipidmaps.org/>). The variable importance in projection (VIP) score was calculated by Orthogonal Projections to Latent Structures Discriminant Analysis (OPLS-DA) using the multiple cross-validation method. The screening criteria for differential metabolites were fold change (FC) > 1, probability (*p*) value < 0.05, and VIP score > 1. The significance of differential metabolites associated with KEGG pathways was calculated using the hypergeometric distribution test.

Analytical procedures and statistical analysis

The efficacy of *P. oleracea* and extracts was assessed based on survival rates, changes to OPG values, DWG, and other indicators. Pooled oocyst counts for each group were performed in accordance with the McMaster counting method. Changes to OPG values were assessed using one-way ANOVA. The average DWG was calculated as (mean final weight of live lambs in the pen) – (mean initial weight of all lambs in the pen) / number of days. The data are presented as the mean ± standard deviation. Statistical analyses were conducted with IBM SPSS Statistics for Windows, version 19.0. (IBM Corporation, Armonk, NY, USA). Figures were generated using Prism 7 software (GraphPad Software, Inc., San Diego, CA, USA). The significance of differences among the six groups was determined by one-way ANOVA. The least significant difference test was used for data exhibiting homogeneity of variance; otherwise, Tamhane's T2 analysis was selected. The Pearson's correlation coefficient was determined to identify linear correlations. A *p* value ≤ 0.05 was considered statistically significant.

Results

Effect of *P. oleracea* on excretion of coccidian oocysts

As shown in Table 1, the OPG values of the PL, PH, PW, PE, and DIC groups on day 7 were 11200, 7883, 5367, 5783, and 733, respectively. The OPG values were significantly lower in the PW and DIC groups than the CON group (*p* ≤ 0.05 and 0.01, respectively). On day 14, the OPG values were significantly lower in the PW and DIC groups than the CON group (*p* ≤ 0.01). On the final day of treatment (day 28), the OPG values were significantly lower in the PH, PW, and PE groups than the CON group (*p* ≤ 0.05), while there was no significant difference between the DIC and CON groups (*p* > 0.05). The OPG values of each *P. oleracea* treatment group at specific time points are shown in the line graph presented in Fig. 2a. The

OPG values of the different *P. oleracea* treatment groups had decreased as compared to the initial values, with the most obvious decrease in the DIC group. After 14 days, the OPG value of the DIC group suddenly increased, while the OPG values of the other drug groups were relatively stable. DIC had the strongest anti-coccidian effect, but efficacy was only maintained for 14 days, followed by the PW group. This finding suggests that DIC has a relatively short-term anti-coccidian effect. A heat map of the OPG values over time is presented in Fig. 2b. After medication, the color of the lambs gradually turned white, with the highest proportion and degree observed in the DIC group. Some of the sheep appeared blue on day 14. However, the heat map of the different treatment groups indicates a trend of turning white. Changes to the OPG values of individual sheep collectively reflect the overall change to the group.

Treatment with *P. oleracea* and the extracts tended to decrease the OPG values. However, anti-coccidian effect was weaker in the PL group than the PH, PW, and PE groups. There were significant differences in OPG values between the PW and CON group on days 7, 14, and 21, with the lowest OPG value on day 42 ($p \leq 0.05$). There were also significant differences in OPG values between the PE and CON groups on days 7 and 14, and between the PH and CON groups on day 28 ($p \leq 0.05$). These results show the *P. oleracea* extracts had better anti-coccidian effects, which may be due to greater release of effective components of *P. oleracea* after extraction. Although DIC initially exhibited strong anti-coccidian effects on days 7 and 14 ($p \leq 0.01$), the OPG value immediately increased with no difference as compared to the CON group. Notably, the coccidia were not completely eradicated in the PH, PW, and PE groups, but infection was ameliorated for prolonged periods.

Table 1

Coccidia oocyst excretion of different treatment groups of *P. oleracea* (a, b, c values with different superscripts differ significantly ($P \leq 0.05$))

Group	Mean OPG before medication	Mean OPG after medication					
		7 d	14 d	21 d	28 d	35d	42d
PL	13800 ± 7200 ^a	11200 ± 8679 ^{ab}	6200 ± 3704 ^{ab}	8150 ± 9328 ^{ab}	9000 ± 7934 ^{ab}	10267 ± 5493 ^a	9800 ± 4726 ^a
PH	13883 ± 9757 ^a	7883 ± 6940 ^{ab}	6967 ± 6869 ^{ab}	6450 ± 7378 ^{ab}	4317 ± 3719 ^b	6700 ± 3899 ^a	6067 ± 4046 ^a
PW	12350 ± 11068 ^a	5367 ± 4100 ^b	2950 ± 2944 ^B	4567 ± 3135 ^b	5167 ± 3925 ^b	7050 ± 1938 ^a	6633 ± 2220 ^a
PE	12650 ± 5747 ^a	5783 ± 3624 ^{ab}	3883 ± 4447 ^b	6783 ± 6532 ^{ab}	6667 ± 7872 ^b	7633 ± 6746 ^a	7550 ± 5014 ^a
DIC	13483 ± 8720 ^a	733 ± 1700 ^B	683 ± 1140 ^B	9067 ± 6323 ^{ab}	13050 ± 8206 ^{ab}	12350 ± 5532 ^a	10583 ± 5795 ^a
CON	12233 ± 10504 ^a	13150 ± 5616 ^a	10883 ± 5648 ^a	13750 ± 6843 ^a	15800 ± 10226 ^a	13033 ± 7816 ^a	11616 ± 6728 ^a

Effect of *P. oleracea* on GP

As shown in Fig. 3a, the mean BW of lambs on days 14 and 42 was higher in each *P. oleracea* treatment group as compared to the CON group. On day 14, the mean BW was significantly higher in the PE group than the DIC group ($p \leq 0.05$). As shown in Fig. 3b, the mean DWG during the 14-day treatment period was significantly higher in the PH and PW groups than the PL group ($p \leq 0.05$), and significantly higher in the PE group than the PL and DIC groups ($p \leq 0.01$). As shown in Fig. 3c, the mean DWG was higher in each treatment group than the CON group throughout the entire 42-day experimental period. The mean DWG was highest in the PH, PW, and PE groups, but there was no significant difference as compared to the CON group ($p > 0.05$). Collectively, these results demonstrate that supplementation with *P. oleracea* can improve the GP of sheep and a single dose of DIC has relatively limited effects.

P. oleracea is beneficial to maintain the composition of the GM

The roles of the GM in health and disease are well established. In the present study, 16S rRNA analysis was conducted to clarify the effects of dietary additives on the composition of the GM. The term “clean reads” refers to high-quality reads obtained after quality control of the original sequence (Fig. 4a). As shown in Fig. 4b, the richness of the GM (ACE index) was significantly higher in the PH, PW, and PE groups than the CON group ($p \leq 0.05$), while there was no difference in the PL and DIC groups as compared to the CON group. The diversity of the GM (Shannon index) was significantly higher in the PH and PW groups than the CON group ($p \leq 0.05$; Fig. 4b; Fig. 4c), and even more pronounced in the PE group than the CON group ($p \leq 0.01$), while there was no significant difference between the PL and CON groups. These results indicate that the high-dose of *P. oleracea* as well as the water and ethanol extracts increased the diversity of the GM, while the low-dose of *P. oleracea* also improved diversity, but to a lesser extent. A dilution curve is generally used to reflect the reliability of the diversity results of sequencing data (Amato, K.R et al., 2013). As shown in Fig. 4d, the dilution curve gradually flattened, indicating that the diversity results are reliable.

Clustering OTUs at 97% similarity reflects differences in sequencing data. As shown in Fig. 4e, the number of OTUs was lower in the CON group than the PH and PW groups, demonstrating that *P. oleracea* influenced the composition of the GM. In order to further study the similarities and differences in the composition of the GM among the groups, non-metric multidimensional scaling (NMDS) and principal co-ordinate analysis (PCoA) were performed. Cluster analysis uses a tree structure to describe and compare similarities among multiple samples. As demonstrated by the results of NMDS (Fig. 4f) and PCoA (Fig. 4g), the CON group is similar in distance to the PH and PE groups demonstrate that the colony structure of the lambs, when treated with *P. oleracea*, is similar to the colony composition of normal lambs. The large distance between the DIC group and other groups indicates that the use of DIC has changed the colony composition of GM. The effects of different properties of chemical drugs and traditional Chinese medicine on GM may vary.

Differences in dominant microbial communities among the treatment groups

At the phylum level (Fig. 5a), 90% of the GM of the six groups were mainly composed of *Firmicutes* and *Bacteroides*. The abundances of *Firmicutes* in the PL, PH, PW, PE, DIC, and CON groups were 65.07%, 81.13%, 72.99%, 76.32%, 68.30%, and 64.00%, respectively, while the relative abundances of *Bacteroides* were 15.13%, 10.94%, 15.91%, 14.64%, 21.05%, and 18.07%. As shown in Fig. 5b, the relative abundance of *Firmicutes* was significantly higher in the PH group than the CON and PL groups ($p \leq 0.01$).

Meanwhile, the abundance of *Firmicutes* was significantly lower in the DIC group than the PH group ($p \leq 0.05$). The relative abundance of *Bacteroides* was lower in the TCM groups than the CON group, but this difference was not significant ($p > 0.05$). *Firmicutes* is believed to help the absorb food calories and leads to weight gain, suggesting that weight gain associated with *P. oleracea* might be related to an increase in the abundance of *Firmicutes*. Also, the relative abundance of *Bacteroides* was higher in the DIC group than the CON group and gradually decreased with an increase in the dose of *P. oleracea*.

At the genus level, the GM of the six groups were mainly composed of *Lachnospiraceae*, *Rikenellaceae*, and *Oscillospiraceae*. As shown in Fig. 5c, the abundances of *Lachnospiraceae* in the PL, PH, PW, PE, DIC, and CON groups were 18.31%, 15.49%, 16.90%, 17.50%, 12.62%, and 15.99%, respectively, while the abundances of *Rikenellaceae* were 7.26%, 6.00%, 8.62%, 7.11%, 13.17%, and 11.47%. Although there was no significant difference among the six groups, DIC was found to reduce the relative abundance of *Lachnospiraceae* ($p \leq 0.01$; Fig. 5d). However, the abundance of *Rikenellaceae* was increased in the DIC group and decreased in the other TCM treatment groups (Fig. 5d).

The F/B ratio is as an index to evaluate imbalances in the GM caused by various illnesses. As compared to the CON group, the F/B ratio was significantly increased in all of the TCM groups ($p \leq 0.05$), but decreased in the DIC group (Fig. 5e). These results indicate that *P. oleracea* can improve the structure and maintain the stability of the GM.

LEfSe (linear discriminant analysis effect size) analysis identified significant differences in GM structure at the phylum level among the treatment groups (Fig. 5f). The results of linear discriminant analysis showed that *o__Monglobales*, *c__clostridia*, and *p__Fimicutes* were the dominant bacteria in the PH group, whereas *o__Oscillospiraceae*, *g__Monglobales*, and *g__unclassified__Veillonellales__Selenomonadales* were the dominant bacteria in the PW group. The main differentiated microbial species between the DIC and other groups were *o__Bacteroidales*, *f__Planococcaceae*, and *g__unclassified__Peptostreptococcaceae* (Fig. 5g). These results suggest that *P. oleracea* enhanced the proportions of beneficial bacteria, increased the F/B ratio, and restored the natural balance of the GM.

P. oleracea increased the gut metabolite content

The main purpose of metabolomics analysis is to identify metabolites with significant biological activities in order to elucidate the related metabolic processes and underlying mechanisms. The results

of OPLS-DA identified various differential metabolites among the treatment groups. As shown in Fig. 6a, there were significant differences in the differential metabolites between the PH and CON groups. The permutation test ($R^2Y = 0.996$ and $Q^2 = 0.866$) confirmed that the OPLS-DA results were reliable. There were also significant differences in metabolites between the PW and CON groups ($R^2Y = 0.993$ and $Q^2 = 0.528$) (Fig. 6b).

In addition, the PLS-DA results showed significant separation among the PH, PW, DIC, and CON groups (Fig. 6c), indicating the reliability of the alternative experimental model. In total, 1675 metabolites were identified and quantified in this study, which included lipids, amino acids, carbohydrates, organic acids, and esters. Cluster heat map analysis identified differences in metabolites and relative changes to metabolite concentrations among the groups. The metabolite concentrations were similar between the PW and PH groups, while there were significant differences between the DIC and CON groups (Fig. 6d).

After conducting qualitative and quantitative analyses of the detected metabolites, Fold Change values and abundance scores of the groups were compared (Fig. 6e and 6f). As shown in Fig. 6e, the metabolites kedaricidin chromophora, levanbiose, prunasin, and 3-hydroxycosanoyl-CoA were 2.17, 1.98, 1.72, and 1.55 times higher respectively in the PH group than the CON group. Levanbiose is reported to prevent dental caries and weight gain, and correct GM imbalance. Prunasin also has the ability to nourish the stomach and diuresis. High doses of *P. oleracea* may increase the abundance of metabolites that promote digestion, growth, and improve GM. Meanwhile, the metabolites biotin sodium, prunasin, and acadesine (drug) were 1.03, 0.67, and 0.66 times higher, respectively, in the PW group than the CON group, dihydroxyl aminic acid, gamma aminobutyric acid were 2.15, 1.02 times lower. The PW group is similar to the PH group, the content of bioactive components levambios, prunasin, and biotin sulfone are relatively increased compared to the CON group. And central nervous system inhibitory neurotransmitters that play an important role in metabolites γ - The decrease in the abundance of aminobutyric acid and acute toxic dihydroxy amino acid suggests that *purslane* may upregulate the abundance of substances that promote cell growth, fatty acid production, and fat and amino acid metabolism, thereby promoting body growth.

As shown by the map of metabolite abundance scores presented in Fig. 6f, the dots are distributed on the right side of the axis and the longer line segments indicate greater overall expression of the pathway. The differences in metabolite abundance scores between the PH and CON groups showed that the Arginine synthesis pathway, Biosynthesis of type II polyketide products, Tetracycline biosynthesis, and Biosynthesis of vancomycin group antibiotics pathways were upregulated, while the thiamine metabolism pathway was downregulated (Fig. 6f). The metabolic pathways of Pantothenate and CoA biosynthesis and Naphthalene degradation were upregulated in the PW group, while Atrazine degradation, Clavulanic acid biosynthesis, and Sulfur relay system were down-regulated. Similar to the results of Fold Change values, the use of *P. oleracea* can up-regulate metabolic pathways such as arginine, Pantothenate and CoA that provide energy and active substances. The DIC group showed down-regulation of phenylalanine, tyrosine, and tryptophan biosynthesis; biosynthesis of enediyne antibiotics, *Pseudomonas aeruginosa*, Fluor benzoate degradation. In summary, *P. oleracea* has

increased the abundance of certain beneficial bacteria, which promote the up-regulation of metabolic pathways of nutrients and active substances, maintain intestinal health, promote growth and development, and resist pathogens.

Discussion

In this study, the effects of different doses of *P. oleracea*, as well as water and ethanol extracts, on fecal coccidian concentrations, BW, GM, and intestinal metabolites were investigated. The results demonstrate that *P. oleracea* reduced fecal coccidian concentrations, improved GP, maintained the composition of the GM, and reduced the F/B ratio. These findings provide a more comprehensive understanding of the therapeutic effects of *P. oleracea* against coccidiosis.

TCM is used for treatment of various diseases, including parasitic infections (Yu L et al. 2019; Liang Y et al. 2019). In the present study, low and high doses of *P. oleracea*, as well as water and ethanol extracts, ameliorated coccidiosis in sheep within 28 days in a dose-dependent manner. In the first two weeks, the efficacy of DIC was much higher than that of *P. oleracea*, while after two weeks, the effect of *P. oleracea* was better than that of DIC. This result is consistent with the findings of a previous study of root extract of the deciduous undershrub traditional Chinese herbal *Dichroa febrifuga* Lour (Zhang DF et al. 2012), which did not achieve complete eradication of coccidia in the early disease stage, similar to *P. oleracea*. Also, the effect of DIC had basically disappeared by short-term use and discontinuation after one developmental cycle of coccidia.

DIC is an anti-coccidian drug widely used in poultry production (El-Ashram S et al. 2019; Wang Z et al. 2020). However, due to the long-term and large-scale use of DIC, drug resistance has become problematic (Ahmadi P et al. 2022). *P. oleracea* poses a low risk of drug resistance and has shown long-lasting anti-coccidian effects, thereby demonstrating advantages as a promising treatment option. However, as compared to some herbs used in TCM with antiparasitic effects, such as Changshan, *Brucea javanica*, and *Artemisia annua*, *P. oleracea* is less effective against coccidiosis. A previous study (Lan L et al. 2016) found that as compared to hydrobromide, *B. javanica* significantly ameliorated coccidiosis, the diarrhea index, and bloody stool score of chickens. Young et al (2001) found that after 10 days of treatment with 15 herbs used in TCM for treatment of coccidiosis in chickens, *Artemisia annua* significantly reduced the lesion score and excretion of coccidian oocysts. Although the anti-coccidiocidal effect was inferior to that of chemical drugs, *P. oleracea* has the advantages of widely available sources and convenience (Kumar A et al. 2022). As compared to other non-medicinal or edible herbs that require processing into essential oils and cumbersome extraction procedures, *P. oleracea* can be directly added to feed or simply extracted with water.

In this study, dietary supplementation with *P. oleracea* effectively improved the GP of sheep. In the first two weeks, the mean DWG was significantly higher in the PE group than the DIC group, and significantly higher in PW group than the CON group. These results are consistent with the findings of a previous study that dietary supplementation with 2% and 3% *P. oleracea* significantly improved GP and increased

feed conversion (Wang C et al. 2021). A prior study found that *P. oleracea* increased the GP of broilers via modulation of the GM, significantly increased the BW, improved the concentrations of immunoglobulin M and superoxide dismutase, and decreased the number of harmful bacteria (Abd El-Hack et al. 2022). The GP of sheep was found to be positively impacted by consumption of *P. oleracea*, which may be attributed to the presence of biological phenols and other active substances, reduction in parasites, and changes to the GM, or the combined actions of many factors. *P. oleracea* is widespread throughout temperate regions of China and welcome "health" edible herb in Chinese and cheap and easy to gain, it has great potential as feed additives in animal health breeding.

The roles of the GM in health and disease have been established (Kenny DJ et al. 2020). In the present study, 16S rRNA analysis was conducted to better clarify the effect of dietary *P. oleracea* on the GM of sheep. The results showed that the ACE and Shannon indices were higher in the PE, PW, and PH groups than the CON group, demonstrating that dietary *P. oleracea* improved the diversity and richness of the GM, consistent with the findings of a previous study (Yi S et al. 2022). Improving the GM can also increase absorption of dietary fibers and GP (Pascale A et al. 2019).

Meanwhile, the difference in species diversity between samples can be better illustrated through principal coordinate analysis. The closer the distance between samples on the coordinate diagram, the more similar they are. The PCoA results, based on Beta diversity analysis (Fig. 6c), indicate that the data distance matrix of the PH, PW, PE group and the CON group overlapped, while the DIC group did not overlap. The result demonstrates that the colony structure of the lambs, when treated with *P. oleracea*, is similar to the colony composition of normal lambs. This discrepancy may be attributed to the variation in the composition of TCM and Western medicine. *P. oleracea* had a positive regulatory effect on the relative abundance and diversity of the GM. Stability of the GM is necessary for the host to resist invasion of pathogenic bacteria (Wang Y et al. 2019; Liu Z et al. 2020). Hence, it is particularly important that drugs are not only effective, but also do not interfere with the stability of the GM.

Bacteroidetes and *Firmicutes* account for 90% of the human GM (Jandhyala SM et al. 2015). The proportions of *Firmicutes*, including *Listeria* and *Streptococcus*, are reported to decrease in response to diseases (Lanza VF et al. 2015; Wronowski MF et al. 2021). Gram-positive bacteria, such as *Lactococcus*, *Lactobacillus*, and *Listeria*, help to maintain the composition of the GM and prevent pathogen invasion (Garneau JE et al. 2008). However, *Bacteroidetes* include bacteria negatively correlated with inflammatory cytokines and are known to cause intestinal diseases (Yang X et al. 2021). In this study, the abundance of *Firmicutes* at the phylum level was significantly increased in the PH and PE groups ($p \leq 0.5$), while the abundance of *Bacteroidetes* was decreased in the *P. oleracea* treatment groups and increased in the DIC group. These results indicate that *P. oleracea* may promote the proliferation of beneficial bacteria, but not harmful bacteria. The F/B ratio serves as an index to evaluate imbalances in the GM in response to various diseases (Liu X et al. 2020). Both *Firmicutes* and *Bacteroidota* play important roles in the digestion of carbohydrates and proteins, as well as maturation of the intestinal immune system. In addition, relatively high abundances of *Firmicutes* and *Bacteroidota* in the GM have been correlated to the energy and nutritional requirements of the host (Spence C et al.

2006). *P. oleracea* can increase the F/B ratio and improve the balance of the GM. Moreover, high levels of *Firmicutes* have been correlated to weight gain (Koliada A et al. 2017), consistent with the *P. oleracea* groups in the present study. At the genus level, the abundance of members of the *Lachnospiraceae* family was increased in the *P. oleracea* treatment groups and decreased in the DIC group. All members of the *Lachnospiraceae* family are anaerobic, fermentative, and chemoorganotrophic, and some exhibit strong hydrolyzing activities (Vacca M et al. 2020). *Lachnospiraceae* can produce short chain fatty acids with potential anti-inflammatory effects (Facchin S et al. 2020). The effect on GP may be related to the ability of *P. oleracea* to improve the structure of the GM and increase the abundance of beneficial bacteria, such as *Lachnospiraceae*. In this study, *P. oleracea* improved the structure of the GM, the F/B ratio, and the abundance of some probiotics, which promoted GP and digestion.

Various dietary components can induce significant changes to the composition and metabolism of the GM. In the present study, arginine content was 1.63 times higher in the PH group than the CON group, while the contents of valine and tyrosine transporters were 4.05 times lower in the CON group. Arginine is used for urea synthesis and beneficial to maintain kidney function (Morris SM. 2006). L-type amino acid transporter 1 (LAT1) is a sodium-independent transporter and mainly responsible for the transport of branched-chain amino acids, such as valine and tyrosine. Up-regulation of LAT1 has been linked to cancer progression and metastases by providing essential amino acids to tumor cells (Lu X. 2019). The contents of biotin sulfone and (-)-sprucenol were higher in the PW group than the CON group. Notably, (-)-sprucenol is a diterpenoid organic compound associated with cytotoxicity and anti-tumor activity (Gunasekera SP et al. 1979). Beauvericin is a naturally occurring mycotoxin that can severely damage the cell membrane, thereby allowing the release of intracellular substances (Mallebrera B et al. 2018). In the present study, the content of beauvericin was 1.35 times higher in the DIC group than the CON group, suggesting that DIC increased the content of beauvericin and is likely toxic to host cells.

The addition of *P. oleracea* can improve the structure of the GM and promote the synthesis of some bioactive metabolites. The addition of *P. oleracea* increased metabolism of tryptophan and nicotinic acid in the PW group as compared to the PH and DIC groups, reduced the contents of arginine and other substances, and up-regulated some vitamins and vitamin co-factors, demonstrating that TCM can impact the production of certain metabolites linked to overall health. For instance, nicotinic acid has the ability to soften blood vessels and can be used in conjunction with lipid-lowering drugs for treatment of atherosclerosis (Saggini A et al. 2011; Li T. 2015). In addition, *P. oleracea* can up-regulate production of metabolically active vitamins and certain amino acids to change the composition of the GM and improve weight gain of sheep. However, the underlying mechanisms remain unclear. The specific pathways or gut barrier effects are also unknown. Therefore, further studies are warranted to elucidate the roles of the composition of the GM in the GP and defense against parasitic diseases of sheep.

Abbreviations

TCM: traditional Chinese medicine

GP: growth performance

GM: the gut microbiota

GP: growth performance

OPG: the oocysts per gram

PL: low-dose *P. oleracea*)

PH: high-dose *P. oleracea*

PW: *P. oleracea* water ex-tract

PE: *P. oleracea* ethanol extract

DIC: diclazuril

Declarations

Ethics approval and consent to participate

The Research Ethics Committee of Henan Agricultural University reviewed and approved the experimental protocol. The collection of samples was performed following the Guide for the Care and Use of Laboratory Animals (Ministry of Health, China). All samples tested in this experiment were collected with permission from herders, while no special permission was required for sampling locations.

Consent for publication

Not applicable.

Availability of data and materials

All raw sequences were submitted to the NCBI Sequence Read Archive (<https://www.ncbi.nlm.nih.gov/sra/>) under BioProject PRJNA1061842. The data supporting the conclusion of this article are included in this article.

Declaration of Competing Interest

The authors declare that there is no conflict of interest in this study.

Funding

This work was partly supported by Key research and development projects of Henan , China(231111111600), the China Agriculture Research System of MOF and MARA (CARS-38) and

Author statement

FCJ, JQL and SMZ contributed to the conception and design of the experiments; SHL, SQL and SLL performed the experiments; SHL, and JL were involved in data collection and analysis. SQC assist in sample collection. SYL and SCH helped interpreting the results and drafting the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgements

We thank Mr. weiqiang Chang, Mr. yanfeng He and Ms. pan Chen, Ms. kaili Liu for technical assistance and sample collection in the Sheep breeding process.

Author detail

^aCollege of Animal Veterinary Medicine, Henan Agricultural University, Zhengzhou 450046, China.

^bKey Laboratory of Quality and Safety Control of Poultry Products, Ministry of Agriculture and Rural Affairs, P.R. China.

References

1. Abd El-Hack ME, Alabdali AYM, Aldhalmi AK, Reda FM, Bassiony SS, Selim S, El-Saadony MT, Alagawany M (2022) Impacts of *Purslane* (*Portulaca oleracea*) extract supplementation on growing Japanese quails' growth, carcass traits, blood indices, nutrients digestibility and gut microbiota. *Poult Sci* 101, 102166. doi:10.1016/j.psj.2022.102166
2. Ahmadi P, Baakhtari M, Yasuda M, Nonaka N, Yoshida A (2022) Toltrazuril and diclazuril comparative evaluation of anti-coccidial drugs using a murine model. *J Vet Med Sci* 84, 1345-1351. doi:10.1292/jvms.22-0136
3. Allen PC, Fetterer RH (2002). Recent advances in biology and immunobiology of *Eimeria* species and in diagnosis and control of infection with these coccidian parasites of poultry. *C Clin Microbiol Rev* 15, 58-65. doi:10.1128/CMR.15.1.58-65.2002
4. Amato KR, Yeoman CJ, Kent A, Righini N, Carbonero F, Estrada A, Gaskins HR, Stumpf RM, Yildirim S, Torralba M, Gillis M, Wilson BA, Nelson KE, White BA, Leigh SR (2013) Habitat degradation impacts black howler monkey (*Alouatta pigra*) gastrointestinal microbiomes. *ISME J* 7, 1344-1353. doi:10.1038/ismej.2013.16
5. Bacanlı M, Başaran N (2019) Importance of antibiotic residues in animal food. *Food Chem Toxicol* 125, 462-466. doi:10.1016/j.fct.2019.01.033
6. Barba E, Guedes AC, Molina JM, Martín S, Muoz MC, Ferrer O, Lara PC, Hermosilla C, Taubert A, Ruiz A (2022) Immunoprotection against mixed *Eimeria* spp. infections in goat kids induced by X-

- irradiated oocysts. *Parasitol Res* 121(5):1517-1525
7. Blake DP, Knox J, Dehaeck B, Huntington B, Rathinam T, Ravipati V, Ayoade S, Gilbert W, Adebambo AO, Jatau ID, Raman M, Parker D, Rushton J, Tomley FM (2020) Re-calculating the cost of coccidiosis in chickens. *Vet Res* 51, 115. doi:10.1186/s13567-020-00837-2
 8. Cacho E, Gallego M, Francesch M, Quílez J, Sánchez-Acedo C (2010) Effect of artemisinin on oocyst wall formation and sporulation during *Eimeria tenella* infection. *Parasitol Int* 59, 506-511. doi: 10.1016/j.parint.2010.04.001
 9. Cai Y, Luo Q, Sun M, Corke H (2004) Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci* 74, 2157-2184. doi:10.1016/j.lfs.2003.09.047
 10. Camarinha-Silva A, Jáuregui R, Chaves-Moreno D, Oxley AP, Schaumburg F, Becker K, Wos-Oxley ML, Pieper DH (2014) Comparing the anterior nares bacterial community of two discrete human populations using Illumina amplicon sequencing. *Environ Microbiol* 16, 2939-2952. doi:10.1111/1462-2920.12362
 11. Dauschies A, Gässlein U, Rommel M (1998) Comparative efficacy of anticoccidials under the conditions of commercial broiler production and in battery trials. *Vet. Parasitol* 76, 163-171.
 12. El-Ashram S, Aboelhadid SM, Arafa WM, Gadelhaq SM (2019) Protective potential of diclazuril-treated oocysts against coccidiosis in layer chicks. *Vet Parasitol* 273, 105-111. doi:10.1016/j.vetpar.2019.08.010
 13. Facchin S, Vitulo N, Calgaro M, Buda A, Romualdi C, Pohl D, Perini B, Lorenzon G, Marinelli C, D'Inca R, Sturniolo GC, Savarino EV (2020) Microbiota changes induced by microencapsulated sodium butyrate in patients with inflammatory bowel disease. *Neurogastroenterol Motil* 32, e13914. doi:10.1111/nmo.13914
 14. Garneau JE, Tremblay DM, Moineau S (2008) Characterization of 1706, a virulent phage from *Lactococcus lactis* with similarities to prophages from other *Firmicutes*. *Virology* 373, 298-309. doi:10.1016/j.virol.2007.12.002
 15. Gunasekera SP, Cordell GA, Farnsworth NR (1979) Potential anticancer agents. XIV. Isolation of spruceanol and montanin from *Cunuria spruceana* (Euphorbiaceae). *J Nat Prod* 42, 658-662. doi:10.1021/np50006a012
 16. Jandhyala SM, Talukdar R, Subramanyam C, Vuyyuru H, Sasikala M, Nageshwar D (2015) Role of the normal gut microbiota. *World J Gastroenterol* 21, 8787-8803. doi:10.3748/wjg.v21.i29.8787
 17. Juraimi AS, Begum M, Ismail MR, Rahim AA, Othman R (2009) Floristic composition of weed community in turf grass area of west peninsular Malaysia. *INT J AGRIC BIOL* 11, 13-20.
 18. Keeton STN, Navarre CB (2018) Coccidiosis in Large and Small Ruminants. *Vet Clin North Am Food Anim Pract* 34(1):201-208. doi:10.1016/j.cvfa.2017.10.009
 19. Kemp LE, Yamamoto M, Soldati-Favre D (2013) Subversion of host cellular functions by the apicomplexan parasites. *FEMS Microbiol Rev* 37(4):607-31. doi:10.1111/1574-6976.12013

20. Kenny DJ, Plichta DR, Shungin D, Koppel N, Hall AB, Fu B, Vasan RS, Shaw SY, Vlamakis H, Balskus EP, Xavier RJ (2020) Cholesterol Metabolism by Uncultured Human Gut Bacteria Influences Host Cholesterol Level. *Cell Host Microbe* 28, 245-257.e246. doi:10.1016/j.chom.2020.05.013
21. Koliada A, Syzenko G, Moseiko V, Budovska L, Puchkov K, Perederiy V, Gavalko Y, Dorofeyev A, Romanenko M, Tkach S, Sineok L, Lushchak O, Vaiserman A (2017) Association between body mass index and *Firmicutes/Bacteroidetes* ratio in an adult Ukrainian population. *BMC Microbiol* 17, 120. doi:10.1186/s12866-017-1027-1
22. Kumar A, Sreedharan S, Kashyap AK, Singh P, Ramchiary N (2022) A review on bioactive phytochemicals and ethnopharmacological potential of *purslane* (*Portulaca oleracea* L). *Heliyon*. 8, e08669. doi:10.1016/j.heliyon.2021.e08669
23. Lamas B, Richard ML, Leducq V, Pham HP (2016) *CARD9* impacts colitis by altering gut microbiota metabolism of tryptophan into aryl hydrocarbon receptor ligands. *Nat Med* 22, 598-605. doi:10.1038/nm.4102
24. Lan L, Zuo B, Ding H, Huang Y, Chen X, Du A (2016) Anticoccidial evaluation of a traditional Chinese medicine--*Brucea javanica*--in broilers. *Poult Sci* 95, 811-818. doi:10.3382/ps/pev441
25. Lanza VF, Tedim AP, Martínez JL, Baquero F, Coque TM (2015) The Plasmidome of *Firmicutes*: Impact on the Emergence and the Spread of Resistance to Antimicrobials. *Microbiol Spectr* 3, PLAS-0039-2014. doi:10.1128/microbiolspec.PLAS-0039-2014
26. Li T, Chiang JY (2015) Bile acids as metabolic regulators. *Curr Opin Gastroenterol*. 31, 159-165. doi:10.1097/MOG.0000000000000156
27. Liang Y, Lin C, Huang S, Xu Y (2019) Traditional Chinese Medicine and Intestinal Microbiota: A Complementary and Integrative Health Approach to Ameliorate Obesity-Related Diseases. *Holistic nursing practice* 33, 259-265. doi:10.1097/HNP.0000000000000311
28. Lighten J, van Oosterhout C, Bentzen P (2014) Critical review of NGS analyses for de novo genotyping multigene families. *Mol Ecol* 23, 3957-3972. doi:10.1111/mec.12843
29. Liu L, Howe P, Zhou YF, Xu ZQ, Hocart C, Zhan R (2000) Fatty acids and beta-carotene in australian *purslane* (*Portulaca oleracea*) varieties. *J Chromatogr A* 893, 207-213. doi:10.1016/S0021-9673(00)00747-0
30. Liu X, He S, Li Q, Mu X, Hu G, Dong H (2020) Comparison of the Gut Microbiota Between Pulsatilla Decoction and Levofloxacin Hydrochloride Therapy on *Escherichia coli* Infection. *Front Cell Infect Microbiol* 10, 319. doi:10.3389/fcimb.2020.00319
31. Liu Z, Li A, Wang Y, Iqbal M, Zheng A, Zhao M, Li Z, Wang N, Wu C, Yu D (2020) Comparative analysis of microbial community structure between healthy and *Aeromonas veronii*-infected *Yangtze finless porpoise*. *Microb Cell Fact* 19, 123. doi:10.1186/s12934-020-01383-4
32. Lu C, Yan Y, Jian F, Ning C (2021) Coccidia-Microbiota Interactions and Their Effects on the Host. *Front Cell Infect Microbiol* 11, 751481. doi:10.3389/fcimb.2021.751481
33. Lu X (2019) The Role of Large Neutral Amino Acid Transporter (LAT1) in Cancer. *Curr Cancer Drug Targets* 19, 863-876. doi:10.2174/1568009619666190802135714

34. Malik TA, Kamili AN, Chishti MZ, Tanveer S, Ahad S (2016) Synergistic approach for treatment of chicken coccidiosis using berberine—A plant natural product. *Microb Pathog* 93, 56-62. doi:10.1016/j.micpath.2016.01.012
35. Mallebrera B, Prosperini A, Font G, Ruiz MJ (2018) In vitro mechanisms of Beauvericin toxicity: A review. *Food Chem Toxicol* 111, 537-545. doi:10.1016/j.fct.2017.11.019
36. Morris SM (2006) Arginine: beyond protein. *The American journal of clinical nutrition*. 83, 508s-512s. doi:10.1093/ajcn/83.2.508S
37. Pascale A, Marchesi N, Govoni S, Coppola A, Gazzaruso C (2019) The role of gut microbiota in obesity, diabetes mellitus, and effect of metformin: new insights into old diseases. *Curr Opin Pharmacol* 49, 1-5. doi:10.1016/j.coph.2019.03.011
38. Paul SM., Mytelka DS, Dunwiddie CT, Persinger CC, Munos BH, Lindborg SR, Schacht AL (2010) How to improve R&D productivity: the pharmaceutical industry's grand challenge. *Nat Rev Drug Discov* 9, 203-214. doi:10.1038/nrd3078
39. Saggini A, Anogeianaki A, Angelucci D, Cianchetti E, D'Alessandro M, Maccauro G, Salini V, Caraffa A, Teté S, Conti F, Tripodi D, Fulcheri M, Frydas S, Shaik-Dasthagirisahab YB (2011) Cholesterol and vitamins: revisited study. *J Biol Regul Homeost Agents* 25, 505-515.
40. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75, 7537-7541. doi:10.1128/AEM.01541-09
41. Sebastián Domingo JJ, Sánchez Sánchez C (2018) From the intestinal flora to the microbiome. *Rev Esp Enferm Dig* 110 51-56. doi:10.17235/reed.2017.4947/2017
42. Shabat SK, Sasson G, Doron-Faigenboim A, Durman T, Yaacoby S, Berg Miller ME, White BA, Shterzer N, Mizrahi I (2016) Specific microbiome-dependent mechanisms underlie the energy harvest efficiency of ruminants. *Isme j* 10, 2958-2972. doi:10.1038/ismej.2016.62
43. Spence C, Wells WG, Smith CJ (2006) Characterization of the primary starch utilization operon in the obligate anaerobe *Bacteroides fragilis*: Regulation by carbon source and oxygen. *J Bacteriol* 188, 4663-4672. doi:10.1128/JB.00125-06
44. Uddin K, Juraimi AS, Ismail MR, Brosnan JT (2017) Characterizing Weed Populations in Different Turfgrass Sites throughout the Klang Valley of Western Peninsular Malaysia. *WEED TECHNOL* 24, 173-181.
45. Vacca M, Celano G, Calabrese FM, Portincasa P, Gobbetti M, De Angelis M (2020) The Controversial Role of Human Gut *Lachnospiraceae*. *Microorganisms* 8(4), 573. doi:10.3390/microorganisms8040573
46. Waller PJ (2006) From discovery to development: current industry perspectives for the development of novel methods of helminth control in livestock. *Vet Parasitol* 139, 1-14. doi:10.1016/j.vetpar.2006.02.036

47. Wang C, Liu Q, Ye F, Tang H, Xiong Y, Wu Y, Wang L, Feng X, Zhang S, Wan Y, Huang J (2021) Dietary *purslane* (*Portulaca oleracea* L.) promotes the growth performance of broilers by modulation of gut microbiota. *AMB Express*. 11, 31. doi:10.1186/s13568-021-01190-z
48. Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 73, 5261-5267. doi:10.1128/AEM.00062-07
49. Wang Y, Li A, Zhang L, Waqas M, Mehmood K, Iqbal M, Muyou C, Li Z, Lian Y, Sizhu S, Li J (2019) Probiotic potential of *Lactobacillus* on the intestinal microflora against *Escherichia coli* induced mice model through high-throughput sequencing. *Microb Pathog*. 137, 103760. doi:10.1016/j.micpath.2019.103760
50. Wang Z, Wu X, Liu L, Xu L, Kuang H (2020) Rapid and sensitive detection of diclazuril in chicken samples using a gold nanoparticle-based lateral-flow strip. *Food Chem* 312, 126116. doi:10.1016/j.foodchem.2019.126116
51. Wen JJ, Li MZ, Gao H, Hu JL, Nie QX, Chen HH, Zhang YL, Xie MY, Nie SP (2021) Polysaccharides from fermented *Momordica charantia* L. with *Lactobacillus plantarum* NCU116 ameliorate metabolic disorders and gut microbiota change in obese rats. *Food Funct* 12, 2617-2630. doi:10.1039/d0fo02600j
52. White EC, Houlden A, Bancroft AJ, Hayes KS (2018) Manipulation of host and parasite microbiotas: Survival strategies during chronic nematode infection. *Sci Adv* 4, eaap7399. doi:10.1126/sciadv.aap7399
53. Wronowski MF, Kotowska M, Banasiuk M, Kotowski A, Kuzmicka W, Albrecht P (2021) Bovine Lactoferrin in the Prevention of Antibiotic-Associated Diarrhea in Children: A Randomized Clinical Trial. *Front Pediatr* 9, 675606. doi:10.3389/fped.2021.675606
54. Yang X, He Z, Hu R, Yan J, Zhang Q, Li B, Yuan X, Zhang H, He J (2021) Dietary β -Carotene on Postpartum Uterine Recovery in Mice: Crosstalk Between Gut Microbiota and Inflammation. *Front Immunol* 12, 744425. doi:10.3389/fimmu.2021.744425
55. Yi S, Jin X, Liu B, Wu P, Xiao W, Chen W (2022) *Portulaca oleracea* extract reduces gut microbiota imbalance and inhibits colorectal cancer progression via inactivation of the Wnt/ β -catenin signaling pathway. *Phytomedicine*. 105, 154279. doi:10.1016/j.phymed.2022.154279
56. Youn HJ, Noh JW (2001) Screening of the anticoccidial effects of herb extracts against *Eimeria tenella*. *Vet Parasitol* 96, 257-263. doi:10.1016/s0304-4017(01)00385-5
57. Yu L, Xing ZK, Mi S, Wu X (2019) Traditional Chinese Medicine and Intestinal Microbiota: A Complementary and Integrative Health Approach to Ameliorate Obesity-Related Diseases. *Holistic nursing practice* 33, 259-265. doi:10.1097/HNP.0000000000000311
58. Yun CH, Lillehoj HS, Lillehoj EP (2000) Intestinal immune responses to coccidiosis. *Dev Comp Immunol* 24(2-3):303-24. doi:10.1016/s0145-305x(99)00080-4
59. Zaiss MM, Harris NL (2016) Interactions between the intestinal microbiome and helminth parasites. *Parasite Immunol* 38, 5-11. doi:10.1111/pim.12274

60. Zhang DF, Sun B, Yue YY, Zhou QJ, Du A (2012) Anticoccidial activity of traditional Chinese herbal *Dichroa febrifuga* Lour. extract against *Eimeria tenella* infection in chickens. Parasitol Res 111, 2229-2233. doi:10.1007/s00436-012-3071-y

Figures




Figure 1

A flowchart of the experiment.

In total, 36 female Hu lambs (age, 1.5 months; mean body weight [BW], 13.08 ± 0.31 kg) were divided into 6 groups of 6 animals each. Lambs in the CON group were fed a basal diet. Lambs in the DIC group were fed the basal diet and orally administered DIC at 1 mg/kg BW for 2 consecutive days. Lambs in the PL group and PH group were fed the basal diet supplemented with *P. oleracea* at 30 g/day and 120 g/day for 14 days. Lambs in the PW group and PE group were fed the basal diet and orally administered a water extract and ethanol extract of *P. oleracea* respectively at 30 mL/day for 14 days. Lambs in the were fed the basal diet and orally administered ant of *P. oleracea* at 30 mL/day for 14 days.




Figure 2

Effects of different treatments on coccidia in lambs.

(a) line chart of mean OPG values of the different treatment groups. (b) Heat map of changes to OPG values, where higher OPG values are indicated by brighter colors. (c) Line graph of changes to OPG

values. Analysis of variance was used to compare the OPG values. Asterisks indicate significant differences (* $0.01 < p \leq 0.05$; ** $0.001 < p \leq 0.01$; *** $p \leq 0.001$)




Figure 3

Effect of different treatments on GP of sheep.

(a) Mean weight of sheep in each group at different times. (b) Mean DWG in each group during the dosing period (14 days). (c) Mean DWG of each group throughout the trial period (42 days). (* $0.01 < p \leq 0.05$; ** $0.001 < p \leq 0.01$; *** $p \leq 0.001$)




Figure 4

Effect of *P. oleracea* on the overall structure of the GM.

(a) Sample coverage of the six treatment groups. (b) Chao1 indices of the six treatment groups. (c) Shannon indices of the six treatment groups. (d) Dilution curves to compare the richness of species among samples and to assess the size of the cohort. (e) Venn diagrams of the OTUs of each group. (f) PCoA plots based on Spearman distances are colored by time point. The significance of differences among the groups was determined by analysis of similarity. (g) NMDS among groups was conducted by one-way ANOVA (* $p < 0.05$; ** $p < 0.01$).




Figure 5

The dominant members of the GM at the phylum and genus levels.

(a) Composition of the GM (top 8 phyla) of each group. (b) Relative abundance of *Firmicutes* and *Bacteroides* of the six groups. (c) Relative abundances of the top 30 genera of each group. (d) Relative abundances of *Lachnospiraceae* and *Rikenellaceae* of the six groups/ (e) F/B ratio. (f and g) Taxonomic cladogram obtained by LEfSe of the six groups. Biomarker taxa are highlighted with colored circles and shaded areas. The diameter of each circle reflects the abundance of the taxa. (b–d), Relative abundance data were analyzed by one-way ANOVA (* $p < 0.05$; ** $p < 0.01$).




Figure 6

Effects on different fecal metabolites in sheep

(a) OPLS-DA score chart and permutation test (PH vs. CON). (b) OPLS-DA score chart and permutation test (PW vs. CON). The OPLS-DA score chart is often used to directly show the classification effect of the model. (c) PLS-DA of the microbial metabolites. (d) Heat map of tissue metabolites. (e) Log converted values of the differential metabolites among the groups. (f) Map of differential metabolite abundance, where the name of the differential pathway is shown on the vertical axis and the DA score is shown on the horizontal axis. The DA score reflects the overall change to all metabolites of the metabolic pathway, where a score of 1 indicates an upward trend and a score of -1 indicates a downward trend in the expression of all annotated metabolites in the pathway.