

Biological nitrogen fixation by legume cover plants in oil palm plantations: calibration of the ureide technique and effects of plantation age and soil nitrate

Ann McNeill (✉ ann.mcneill@adelaide.edu.au)

University of Adelaide <https://orcid.org/0000-0002-6060-4661>

Rachel Pipai

Murray Unkovich

Murom Banabas

Paul N. Nelson

Research Article

Keywords: Calapogonium, Pueraria, Mucuna, standing biomass, litter, nodulation

Posted Date: April 28th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2839206/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Plant and Soil on July 17th, 2023. See the published version at <https://doi.org/10.1007/s11104-023-06147-8>.

Abstract

1. Background and Aims

To sustainably manage N in oil palm systems quantities of N fixed by cover legumes need to be understood. Current values are scarce, based on shoot N measures and do not include litter which releases nitrate as it decomposes. We aimed to quantify N_2 fixed by legumes under oil palm systems in PNG and to determine if soil nitrate influenced dependence on N_2 fixation (Ndfa).

2. Methods

The ureide technique for estimating tropical legume Ndfa was calibrated for *Calapogonium mucunoides*, *Pueraria phaseoloides* and *Mucuna pruriens* using ^{15}N isotope dilution, and then used to assess Ndfa for legume cover under oil palms (2-25 years old) in Papua New Guinea. Amounts of fixed N in above-ground legume biomass (shoot plus litter) were calculated incorporating % groundcover. Soil nitrate under the legume litter was also measured.

3. Results

Legume Ndfa was highly negatively correlated with soil nitrate concentration but independent of palm age. Legume groundcover, shoot and litter dry matter, and quantity of fixed N were greater under oil palms less than 5 years old, decreasing under older plantations where solely *C. caeruleum* was present. DM and N content of litter were similar to shoots for legumes in plantations less than 6 years old.

4. Conclusion

The calibrated ureide technique can be used, together with estimates of annual legume N accumulation, to quantify N input from legume groundcover during the life cycle of oil palm plantations and other tropical ecosystems, in order to support more sustainable management of N.

Introduction

Oil palm (*Elaeis guineensis*) has become one of the world's major agricultural crops, driven by unprecedented increase in consumer demand (Corley 2009) coupled with rapid commercial development, with production occurring in lowland moist humid tropical zones in Central Africa, South America and Southeast Asia (Byerlee et al. 2016). Although 85% of global production currently comes from Indonesia and Malaysia, the oil palm industry is a highly important component of the export economy for many other countries, including Papua New Guinea (PNG), where it is the largest non-government employer and underpins rural livelihoods (Allen et al. 2009; Cramb and Curry 2012). There is increasing pressure from consumers and governments for agriculture and food production systems to be more sustainable, with oil palm under particular public scrutiny to improve productivity per current area rather than encroach further on important natural environments (Teoh 2010; Sayer et al. 2012). Many palm oil producing countries, including PNG, are committed to increase the sustainability of the oil palm industry via activities undertaken as members of the Roundtable on Sustainable Palm Oil (RSPO, <https://rspo.org/as-an-organisation/>).

A key factor for sustainability of any agricultural system is the management of nitrogen (N) in order to maximise productivity and minimise off-site effects, particularly eutrophication of waterways caused by run off and nitrate (NO_3^-) leaching, and greenhouse gas (GHG) emissions from denitrification and volatilisation (Galloway et al. 2013). Nitrogen fertilizers, a significant component of the cultivation costs in oil palm systems (Corley and Tinker 2016), are considered a major potential source for such N losses (Choo et al. 2011; Comte et al. 2012). Amounts of mineral fertiliser N applied to oil palm plantations vary widely, ranging from 48–90 kg N ha/year for immature palms to 56–206 kg N/ha/yr for mature palms (see references cited in Pardon et al 2016). Whilst there is a dearth of comprehensive data to describe the complete N cycle of an oil palm system from planting to harvest, modelling using the limited data available suggests potential for N losses is greatest during the immature phase (first 2–3 years) when N uptake by palms is low, and then later during the mature phase when groundcover under palms tends to be sparse and N fertiliser rates higher (Pardon et al. 2016). The most influential parameters for N loss from an oil palm system were identified as mineral N fertiliser rate, drainage and fraction of legume in groundcover (Pardon et al. 2017).

Perennial legume species such as *Pueraria phaseoloides*, *Calapogonium mucunoides* *Calopogonium caeruleum*, and *Mucuna pruriens* have long been utilised as groundcover for control of soil erosion and suppression of weeds under oil palm plantations (Fairhurst and Hardter 2003; Samedani et al. 2015), as well as for N input via biological N_2 fixation or BNF (Giller and Fairhurst 2003). Amounts of N fixed by cover legumes, including *P. phaseoloides* under 2–3 year old oil palm in Malaysia, were estimated at 150 kg N/ha/yr (Agamuthu and Broughton 1985) based on the mean difference in N content between the vegetation of the N_2 fixing legume and non-fixing “natural” covers (N difference method). Another Malaysian oil palm study using ^{15}N isotope-dilution methodology reported similar amounts of N_2 fixation by *P.*

phaseoloides (Zaharah et al. 1986). However, published measures of BNF for cover legumes in oil palm systems *in situ* remain scant (Giller and Fairhurst 2003; Pardon et al. 2016), despite a number of methodologies available for field use (Unkovich et al. 2008).

One approach for assessing the dependence of a legume cover on BNF that does not require the presence of a non-fixing cover is the ureide technique (Herridge 1982). This method is based on the premise that many tropical legume plants, from the tribes *Phaseoleae* and *Desmoideae* within the subfamily *Papilionoideae*, transport identifiable N_2 fixation products in the xylem sap as ureides in contrast to N from mineral N assimilated into amides (Peoples et al. 2009). While the method itself is inexpensive and relatively simple, it must be calibrated with another technique such as the natural abundance $\delta^{15}N$ (‰) or ^{15}N isotope dilution technique (Unkovich et al. 2008). Ureide calibrations have been carried out for a number of legume species, including *Desmodium ovalifolium* and *Centrosema spp.* that are commonly used as oil palm covers. Stem relative ureide N (RUN) for these species has been shown to be highly correlated with the proportion of N derived from N_2 fixation (pNdfa) assessed using ^{15}N isotope dilution, proving the reliability of the calibrated ureide technique (Unkovich et al. 2008). Although there is mention of an unpublished ureide calibration for *C. caeruleum* (Wilson et al. 1995), there appear to be no published calibrations for this species, or for *C. mucunoides*, *P. phaseoloides* and *M. pruriens* (Unkovich et al. 2008) which are widely used as cover legumes in oil palm systems.

As plantations age and the palm canopy closes, the legume cover often declines due to shading, although some species thrive under older plantations (Giller and Wilson 1991). Reduced legume cover growth means lower amounts of N_2 fixed (Zaharah et al. 1986), but it is unclear how plantation age and accompanying increase in shade, affects the proportional dependence on N_2 fixation of the legumes. Also, since pNdfa tends to be reduced if nitrate is present in the legume root zone (Streeter and Wong 1988), it might be expected that accumulation of nitrate in soil, resulting from increased accumulation and decomposition of cover legume litter as plantations age, would reduce legume dependence on N_2 fixation. However, a glasshouse study to determine the effect of litter quantity on pNdfa for *P. phaseoloides* suggested that effects were small as it was only reduced from 87–84% when litter input was increased from zero to the equivalent of 8 Mg DM/ha (Vesterager et al. 1995). Legume covers in tropical environments accrue litter rapidly within 6–9 months of establishment (Broughton 1977; Chiu 2004; Samedani et al. 2015), but reports of the rate at which this litter decomposes vary widely from a few months up to a year or more (Vesterager et al. 1995; Turner and Gillbanks 2003; Chiu 2004; Philip and Abraham 2009). Furthermore, although incubation experiments with ^{15}N labelled residues of the cover legumes *P. phaseoloides* and *M. bracteata* demonstrate rapid net N mineralisation (Mendham et al. 2004), it has been suggested the flux may be reduced in the plantation situation by N immobilisation due to the presence of other residues from living and previously felled trees in the plantation (Pardon et al. 2016). It has been reported that the presence of cover legumes actually reduced leaching of N in plantations (Agamuthu and Broughton 1985; Lehmann et al. 1999) but there appears to be no direct investigation of soil nitrate under cover legumes in the field specifically related to BNF inputs in oil palm systems.

Therefore, this study aimed to quantify N_2 fixation by legume cover crops under oil palm systems in PNG and to determine possible relationships between pNdfa and amount of nitrate in the soil under the cover legumes. The ureide technique was calibrated for legume cover species in the glasshouse, subsequently used to estimate pNdfa of legume covers in the field under different ages of palm, and then amounts of fixed N in total above-ground biomass (shoot plus litter) of legume covers were calculated.

Materials and Methods

1. Glasshouse experiment to calibrate the xylem ureide technique using ^{15}N isotope dilution

Plant culture and growing conditions

A glasshouse experiment was conducted to measure changes in the relative ureide abundance in stems of nodulated legumes grown at varying nitrate supply, with N_2 fixation measured using ^{15}N isotope dilution. The experiment took place under natural daylight over 9 months (May-January) at the Waite campus of the University of Adelaide, South Australia (34.9688° S, 138.6339° E). Average day/night temperature in the glasshouse for the growth period was 25/17°C. During winter (June - August) supplementary glasshouse lighting was used to ensure a daylight period of 12h. Seeds of *C. mucunoides*, *P. phaseoloides* and *M. pruriens* were sourced from the Australian Tropical Grains Germplasm Centre (Queensland). Prior to sowing *C. mucunoides* seeds were soaked in hot (75°C) water for three minutes to break dormancy (Giller and Fairhurst 2003) *P. phaseoloides* seeds were soaked for one hour in glycerol at 50°C. *M. pruriens* seeds required no pre-treatment. Single seeds of each species were sown in free-draining 4-L pots containing fine acid-washed river sand in May (*C. mucunoides* and *M. pruriens*) and July 2011 (*P. phaseoloides*). All pots were inoculated with Group M *Bradyrhizobium* strain (CB756, Beckerwood, Australia), by mixing peat inoculum in water and pouring a small amount of slurry into each seeding hole before the seeds were sown.

Nutrient solution and nitrate treatments

The glasshouse experiment had five nitrate (NO_3^-) concentration treatments including zero, four replicates and two harvests for each species, giving a total of 120 pots, arranged in a randomized complete block design. Pots were watered as needed with tap water until the first true leaves were observed on the plants, and until 10 weeks of age fed with a basal minus N nutrient solution for tropical legume culture (Herridge and Peoples 2002b) containing (as g/L) 0.0625 MgSO_4 , 0.0368 CaCl_2 , 0.02188 KH_2PO_4 , 0.0188 KCl, and 0.0259 FeEDTA and as (mg/L) 0.715 H_3BO_3 , 0.4525 MnCl_2 , 0.0275 ZnCl_2 , 0.0125 CuCl_2 and 0.0063 NaMoO_4 , initially diluted 1/8 or 1/4 according to the growth rate of the seedlings. Nitrate feeding of plants commenced ten weeks after sowing using ^{15}N labelled potassium nitrate (5.0 atom% ^{15}N) mixed with unlabelled $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ and unlabelled KNO_3 to make 1, 3, 5 and 10 mM NO_3^- solutions. Resulting enrichments of these nitrate solutions were 1 (1mM) 0.33 (3 mM), 0.2 (5mM) and 0.1 (10mM) atom % ^{15}N excess in the above nutrient solutions. Pots were fed 1L of nutrient solution twice weekly until harvest.

Sampling

Plants were harvested on different dates due to different growth rates, with *M. pruriens* harvested 16 and 21, *C. caeruleum* 29 and 34 and *P. phaseoloides* 23 and 28 weeks after sowing. At harvest, plants were cut just above the sand surface and 7 stem pieces (12–15cm long) excised and oven dried at 85°C for 48 hours and weighed, as was the remainder of the shoot. Each pot was tipped over carefully to remove loose sand and ensure the root system plus nodules remained as intact as possible. Roots were washed clean using tap water and a 2-mm sieve used to capture the remaining roots and nodules from the wash bucket. Each intact root system was divided into 0–5 cm, and > 5 cm sections in relation to depth in the sand. Nodules were separated from the root and counted, before roots and nodules were oven-dried at 85°C for 48 hours and weighed. All dried plant samples were ground in a ball mill.

Analyses of plant material

Dried samples of legume shoots, roots and nodules were analysed for total N and ^{15}N using an automated combustion analyser linked to a mass spectrometer (Dawson and Brooks 2001). Ground stem samples were subject to solute extraction and the extracts stored in a freezer prior to analysis for ureide, amino and nitrate-N as described in Unkovich et al. (2008). Briefly, ureide-N was analysed using the Rimini-Schryver reaction (Young and Conway 1942), amino compounds were analysed using the ninhydrin method (Yemm and Cocking 1955) and nitrate was analysed using the salicylic acid method (Cataldo et al. 1975).

Calculations

The concentration of ureides in the stem solutes relative to the amino and nitrate N concentrations, which derive principally from soil nitrate uptake (Herridge and Peoples 2002a), can be expressed as relative ureide N (RUN). Given that there are four N atoms in the ureide molecule, the RUN (%) can then be represented as:

$$\text{RUN (\%)} = (4a / (4a + b + c)) \times 100$$

where *a* is the concentration of ureides; *b* nitrate-N and *c* amino-N (Herridge 1982).

The relationship between RUN (%) and the proportion of plant N derived from the atmosphere (pNd_{fa}, %) has been shown to be close to unity for a range of legumes (Unkovich et al. 2008).

In the present study the pNd_{fa} (%) was calculated using the ^{15}N isotope dilution technique with the ^{15}N analysis data and the following equation (Unkovich et al. 2008)

$$\text{pNd}_{\text{fa}} (\%) = [1 - (\text{atom\%}^{15}\text{N excess in plant} / \text{atom\%}^{15}\text{N excess in nutrient solution})] \times 100$$

2. Field survey of N_2 fixation by legumes under oil palm

Site information

Field studies were carried out in thirteen plantations across three commercial oil palm estates (Kapiura, Mosa and Numundo) near Kimbe (5.5512° S, 150.1387° E), West New Britain Province, PNG. The climate in the region is tropical with rain every month and an average annual rainfall of 3600 mm. Oil palm ages in the 13 sampled plantations ranged from 2 to 25 years with each plantation approximately 27 ha. Fertiliser application in the year prior to sampling (2011) varied with age of palm and management approach (Table 1). Although a mix of legume cover species had generally been sown immediately after the palms were planted as per best practice recommendation (Fairhurst and Hardter 2003) there were only three plantations, two on the Kapiura estate and one on the Mosa estate, where more than one species (i.e. *C. caeruleum* with *P. phaseoloides*) contributed to the ground cover (Table 1), whereas at all other plantations only one species was

present. (Table 1). The soils are volcanic ash in origin (Banabas et al. 2008). Sampled soils pH_{CaCl2} ranged from 4.3 to 5.6 (Table 1) and were sandy loam to clay texture in the top 30 cm.

Soil and plant sampling and analyses

Soil and plant sampling was conducted in January and February 2012. In Kapiura and Mosa estates, for each legume cover species present, ten replicate samples of above-ground biomass (green shoots and litter separately) within 1m² quadrats were taken at each plantation. Quadrats were placed randomly in areas containing the specific legume cover. Stem segments (10 cm length) were also obtained from each quadrat. Sampling at the Numundo estate plantations was restricted to stem segments and assessment of legume ground cover. Shoot, stem and litter samples were oven dried (85°C), weighed, ground, subsampled and transported to Australia for analysis of total N and N solutes extracted from stem segments as described for the glasshouse study. Soil was sampled to a depth of 20 cm using an auger (15 cm diameter) in four of the quadrats at each plantation. The four samples were thoroughly mixed, subsampled (~ 500g) and oven-dried at 60°C. Sub-samples of the dried soils (~ 100g) were sent to Australia. To comply with Australian quarantine regulations all dried plant and soil samples were gamma irradiated at a dosage of 25kGray (Steritech Pty Ltd, Queensland), prior to release for analysis. Sub-samples of each soil were then extracted with 2M KCl (40ml/10g soil) for one hour and the extracts analysed on an Alpchem Flow Solution II Analyser for nitrate-N and ammonium-N using colorimetry (Rayment and Lyons 2011). Further sub-samples were extracted in 0.01M CaCl₂ (1:5) for one hour and after allowing sediment to settle for 30 min, pH in the supernatant was measured with a Thermo ORION 960 pH/conductivity electrode (Rayment and Lyons 2011).

Table 1 Palm age (years), legume cover species present, soil pH at sampling and fertiliser applied (kg/palm) in the year 2011 prior to the sampling in Jan-Feb 2012, for plantations sampled on three commercial oil palm estates near Kimbe, West New Britain Province PNG

Estate	Palm Age (yrs)	Legume	Soil pH _{CaCl2}	Fertilizers applied in 2011 (kg/palm)				
		Cover Species		N	P ₂ O ₅	K ₂ O	MgO	B
Kapiura								
	3	<i>C. caeruleum</i>	4.3	0.73	0.25	1.27	0.18	0.009
		<i>P. phaseoloides</i>	4.4					
	5	<i>C. caeruleum</i>	4.9	0.87	0.28	1.74	0	0
		<i>P. phaseoloides</i>	5.2					
	7	<i>C. caeruleum</i>	4.6	0.48	0	0	0	0
	25	<i>C. caeruleum</i>	4.7	0	0	0	0	0
Mosa								
	2	<i>P. phaseoloides</i>	4.4	0.89	0.44	1.22	0.16	0.019
	4	<i>C. caeruleum</i>	5.1	1.05	0.24	1.71	0.23	0
		<i>P. phaseoloides</i>	5					
	6	<i>M. pruriens</i>	5.6	1.04	0.40	1.51	0.24	0.011
	13	<i>C. caeruleum</i>	5	1.04	0.40	1.75	0.24	0.011
	18	<i>C. caeruleum</i>	4.8	1.37	0.57	1.75	0.16	0
Numundo*								
	4	<i>P. phaseoloides</i>	nd	1.01	0	1.75	0	0.011
	8	<i>C. caeruleum</i>	nd	0.82	0	1.00	0.15	0.010
	9	<i>C. caeruleum</i>	nd	0.65	0	0	0	0.010
	16	<i>C. caeruleum</i>	nd	0.24	0	0.40	0	0
nd = not determined								

* Sampling in Numundo estate was restricted to legume cover and stem segments for solute analysis whereas locations in the other two estates were also sampled for shoot and litter dry matter.

Estimation of legume ground cover (%) and amount of fixed N in standing shoot biomass

Ground cover (%) for each legume species present in a plantation was assessed using a representative hexagonal 'palm unit' (Fig. 1) that accounted for the triangular layout of palms, a weeded circle, the cleared harvest path and the frond piles. A quadrat (1m²) was moved from place to place within the hexagonal area to assess the total area (m²) covered by the legume species. Cover was observed to grow across harvest paths and on top of frond piles but the weeded circle usually remained as bare soil. The amount of N fixed by the legumes in standing shoot biomass in the plantation was calculated as the product of estimated % legume cover, pNdfa (%) as determined from stem ureide analysis and application of the calibration equation generated in the glasshouse study, and measured N contents of the shoots. To calculate the total amount of above-ground fixed N (in standing biomass plus litter) a simple assumption was made that the pNdfa (%) measured for the shoots was also applicable to the N in the litter under those shoots.

Legume nodulation assessment

Legume nodulation was visually assessed on randomly selected plants at all plantations on the Mosa and Kapiura estates using a scoring system based on number and position of nodules on the excavated root system, modified from Corbin et al. (1977). Categories were poor (score 0–2); fair (2–3), good (3–4) and excellent (5) nodulation, with the lowest considered indicative of little or no N₂ fixation and the highest implying that N₂ fixation potential is near optimal.

Statistical analysis of data

Analysis of variance was used to explore differences in species responses to NO₃⁻ supply in the glasshouse experiment. Linear regressions were established between RUN (%) and pNdfa (%) from the glasshouse experiment. Firstly a general linear homogeneity of slopes analysis was used to determine if there were differences between species in the slopes of the regressions (MANCOVA, P < 0.05). If there were no differences between species then a single linear regression was derived using the mean species data. All analyses were conducted with the Statistica™ software package (v13.5.0.17, TIBCO Software Inc., CA USA). For the field data, general relationships between variables were explored using visual lines of best fit (power curves) using Microsoft Excel. The field experiments were based on a need to gather quantitative data on N₂ fixation and as such were descriptive rather than hypothesis driven.

Results

1. Glasshouse experiment to calibrate the relative ureide N technique for cover legumes

Proportional dependence on nitrogen fixation (pNdfa, %) measured by ¹⁵N isotope dilution

Since the only source of N in the zero NO₃⁻ treatments was N₂ fixation, these plants are assumed to have fixed 100% of their N. The NO₃⁻ supply regime was effective in providing a wide range of dependencies on N₂ fixation (Fig. 2Figure 1A). Across the two harvests pNdfa (%) decreased with increasing N supply, especially for *C. mucunoides* and *P. phaseoloides* (Fig. 2Figure 2A) which appeared very similar in their response, with 5mM NO₃⁻ being sufficient to suppress N₂ fixation by approximately 90%. In contrast, *M. pruriens* maintained a significantly (P < 0.01) higher dependence on N₂ fixation compared with the other two species under all NO₃⁻ additions.

Nodule mass per plant (Fig. 2B) was much greater for *M. pruriens* than the other two species, and as for N₂ fixation, was suppressed less than for the others under increasing NO₃⁻ supply. Nodule number (data not shown) also significantly decreased with increasing nitrate supply in *P. phaseoloides*, but not in the other two species.

Stem ureide N calibration

Stem RUN (%) indices were very similar in the two harvests (data not shown) and thus mean values across harvests within a species and NO₃⁻ supply are presented (Fig. 3). Mean RUN was the same (P < 0.01) for *C. mucunoides*, and *P. phaseoloides* at each rate of NO₃⁻ supply (Fig. 3A), except for the 1 mM treatment, in which *C. mucunoides* had a higher RUN (40%, P < 0.01) than the other two species. The RUN for

M. pruriens (Fig. 3B) was much less sensitive to low NO_3^- supply than the other species, which displayed more rapid increases in RUN as NO_3^- supply fell below 1mM.

Plotting the ^{15}N -estimated Ndfa (%) against the measured RUN (%) for each NO_3^- concentration again showed the similarities between *C. mucunoides* and *P. phaseoloides* (Fig. 4A) and the disparate response of *M. pruriens* (Fig. 4B). Homogeneity of slopes analysis showed that the regressions for *C. mucunoides* and *P. phaseoloides* were not significantly different from each other ($P > 0.05$ by analysis of variance), and thus they were combined to produce a single calibration curve relating Ndfa (%) to stem RUN (%) for use in the field study (viz. $\text{Ndfa (\%)} = 1.514 \times \text{RUN (\%)} + 4.9848$). While the correlation between Ndfa and stem RUN for *M. pruriens* was no weaker than for the other two species (Fig. 3), the slope was very much steeper, meaning that an absolute change of 1% in stem RUN for *M. pruriens* results in a shift in Ndfa of 7.5%, with the comparable shift for the other species of 1.5% ($\text{Ndfa (\%)} = 7.4857 \times \text{RUN (\%)} - 2.5601$).

2. Field Survey of legume cover species in oil palm plantations

Legume cover shoot and litter dry matter and N content

Shoot dry matter for total legume cover sampled in each plantation ranged from 144 g/m² to 792 g/m² (mean 347 g/m²) across all plantations (Fig. 5A). Shoot biomass was greater in plantations with young oil palm, especially where cover was a mixture of two species (*P. phaseoloides* and *C. caeruleum*), and less in plantations with older (7–25 year) palms where cover was mainly *C. caeruleum*. As a result, there was a strong correlation ($R^2 = 0.67$) between legume cover shoot dry matter and age of palm across plantations (Fig. 5A), with shoot biomass decreasing rapidly in the first 5–10 years, after which it remained at around 150 g/m². Litter collected directly under the legume shoots averaged 762 g/m² in plantations 6 years old or less and 139 g/m² for those with palms 7–25 years old. The litter mass was almost always greater than the standing shoot biomass, regardless of palm age or legume species present, and was also strongly correlated with oil palm plantation age (Fig. 5B).

Shoot N concentration for legumes ranged from 2.2% (*C. caeruleum*, Kapiura 25 year old plantation) to 2.9% (*M. pruriens*, Mosa 6 year old). There were no significant differences ($P = 0.44$) between shoot N concentrations of *C. caeruleum* and *P. phaseoloides*, however species differences remain unclear because *P. phaseoloides* was only found in young palm plantations and there was a strong negative correlation ($\text{Adj } R^2 = 0.64, P < 0.01$) between the age of the stand and the shoot N concentration. The litter N concentrations for individual species (1.6 to 3.2%) were marginally lower than those of the associated legume shoots, and more markedly so in older plantations. Legume shoot N content was 3.6 (*C. caeruleum*, Kapiura, 25 year old) to 18.6 (*P. phaseoloides*, Mosa 2 year old) g/m² and legume litter N content was 2.0 (*C. caeruleum*, Kapiura, 25 year old) to 21.8 (*M. pruriens*, Mosa 6 year old) g/m². There were good correlations between legume shoot N content and palm age (Fig. 5C, $R^2 = 0.71$) and litter N content and palm age (Fig. 5D, $R^2 = 0.65$), as reported for dry matter.

Stem RUN

C. caeruleum generally had higher stem RUN than *P. phaseoloides*, ranging from 17 to 43% (average 28%), while the range for *P. phaseoloides* was 13 to 38% (average 18%). The RUN for *M. pruriens* sampled in the field survey was 28%, exceeding the maximum (12%) recorded for that species in the glasshouse experiment. Hence, the calibration for RUN (%) against pNdfa (%) derived in the glasshouse study for *C. mucunoides* and *P. phaseoloides* (Fig. 4A) was also used to derive pNdfa (%) for the single *M. pruriens* cover in the field survey. Furthermore, in the glasshouse study RUN in *C. mucunoides* obtained at zero N and 10mM N were very similar to those reported for fully nodulated (80% RUN) and fully nitrate-dependent (10% RUN) *C. caeruleum* (Gault and Peoples 1993), suggesting the two species of that genera are very similar in ureide activity. This gave confidence for applying the calibration derived for RUN in *C. mucunoides* to the *C. caeruleum* cover present in the field.

Legume proportional dependence on N₂ fixation (pNdfa, %) for cover species

Mean pNdfa was significantly higher ($P < 0.05$) for *C. caeruleum* (49%) than for *P. phaseoloides* (32%), although for both species a wide range of values were observed (Table 2). The single measure of pNdfa for *M. pruriens* was 49% based on the calibration from the other species but, given the glasshouse observations, it could be higher. There was no correlation between plantation age and pNdfa for *P. phaseoloides* or *C. caeruleum* (Fig. 6A) although *P. phaseoloides*, which was only present in younger plantations, generally had low pNdfa (14–27%) and consistently appeared to be deriving a larger proportion of its N from soil (72–82%) than *C. caeruleum* (25–72%) or *M. pruriens* (51%, Table 2).

Soil extractable N

Concentrations of nitrate-N in the soil extracts ranged from 2–25 mg NO₃⁻-N/kg. The highest NO₃⁻-N occurred in soil from under *P. phaseoloides* in a 3 year old plantation on the Kapiura estate and the lowest NO₃⁻-N in soil under *C. caeruleum* from a 25 year old plantation on the same estate. There was a strong negative relationship ($R^2 = 0.55$) between soil nitrate concentration and pNdfa (%) of individual cover legumes (Fig. 6B), with a marked decline in dependence on N₂ fixation as nitrate increased above 5 mg NO₃⁻-N/kg.

There were exceptionally high concentrations of ammonium-N (95–292 mg NH₄⁺-N/kg) in all the soil extracts and a poor correlation between pNdfa (%) and ammonium-N concentration in soil ($R^2 = 0.02$). It was suspected that ammonium concentrations in the soils may have been influenced by the irradiation treatment of the samples on import to Australia.

Nodulation score

A high nodulation score (4 or 5) indicating an excellent potential for N₂ fixation was recorded for both *P. phaseoloides* and *C. caeruleum* in plantations that were 5 years old or less (Table 2). A nodulation score of 4 was also recorded for *C. caeruleum* in a 13 year old Mosa plantation, whereas in the majority of plantations with older palms the nodulation scores for all three legumes were 2–3 indicating fair nodulation but reduced N₂ fixation potential. A nodule score of 1 was assigned to *C. caeruleum* in the 18 year old Kapiura plantation suggesting little or no N₂ fixation activity by this cover under mature trees. However, there were no obvious correlations between nodule score and either soil nitrate-N or pNdfa (%) of individual legume species (Table 2).

Plantation legume ground cover (%) and total standing biomass (DM), N and fixed N (kg/ha)

The proportion of ground covered by legumes in different plantations was highly variable and ranged from 0.6% in the oldest plantation sampled (*C. caeruleum*, Kapiura 25 year old) to 44% (*P. phaseoloides*, Mosa 2 year old). There were insufficient samples in the survey (Table 2) to be able to ascertain if there were trends in species presence with age of oil palm plantation, particularly as *M. pruriens* was only encountered in one plantation. However, it was evident from the restricted dataset (Table 2) that greater legume ground cover was recorded in plantations that were 5 years old or less, and that only *C. caeruleum* was present under the older plantations. Generally the percentage legume ground cover decreased as oil palm age increased ($R^2 = 0.67$).

Assuming legume ground cover (%) across the entire plantation was similar to that in the sampled regions then the total biomass of above-ground legume cover (standing shoot plus litter) could be 1.2–5.6 Mg DM/ha containing 30–135 kg N/ha in plantations five years old or less (Table 2). This includes three plantations where mixtures of cover species occurred. Whereas in plantations older than five years standing shoot plus litter biomass (16–670 kg DM/ha) and N (< 1-21kg N/ha) would be markedly lower, corresponding with observations of less cover and lower shoot/litter N concentrations in these plantations.

The amount of fixed N in shoots plus litter ranged from 2–34 kg N/ha (Table 2) but was not correlated with plantation age. Nevertheless, amounts of fixed N in total legume above-ground biomass (sum of standing shoot plus litter of all species present) were generally higher in the younger plantations where dry matter was greater due to more legume cover, and in some cases two cover species present (Table 2). Fixed N in total legume above-ground biomass was partitioned between standing shoots and litter in a ratio of 1:1 or 1:>1 in plantations less than 6 years old and in older plantations (with one exception) at a ratio of 1:1 or 1:<1. The proportion of N in total legume above-ground biomass derived from N₂ fixation was greater than 30% for 6 of 9 plantations across two estates (Table 2), partly due to relatively high pNdfa for *C. caeruleum* (Table 2).

Table 2

Groundcover (%), proportional dependence on N₂ fixation (pNdfa, %)¹, nodulation (Nod) score² and DM, total N and N derived from N₂ fixation (kg/ha) for above ground biomass (standing shoot plus litter) of the legume cover species present under oil palm plantations of different ages in West New Britain Province, PNG.

Estate		Above-ground legume biomass (kg/ha)												
Plantation ID	Palm Age (yrs)	Legume Species	% ground cover	s.d. ³	pNdfa (%)	s.d.	Nod score	DM	s.d.	Total N	s.d.	Fixed N	s.d.	Fixed N as % Tot N
Kapiura														
K3	3	<i>C. caeruleum</i>	9	2	39	13	5	779	416	18	10	7	5	
		<i>P. phaseoloides</i>	8	3	18	7	5	787	423	18	9	4	2	
		Total	17					1566		36		11		31
K5	5	<i>C. caeruleum</i>	19	3	75	60	2	625	206	15	10	12	10	
		<i>P. phaseoloides</i>	10	5	28	16	4	610	201	15	5	4	2	
		Total	29					1235		30		16		53
K7	7	<i>C. caeruleum</i>	7	6	37	15	2	148	108	3	<1	1	<1	33
K25	25	<i>C. caeruleum</i>	<1	1	53	28	3	16	26	<1		<1	<1	nd
Mosa														
M2	2	<i>P. phaseoloides</i>	44	8	22	7	3	5593	2020	135	52	34	8	25
M4	4	<i>C. caeruleum</i>	20	13	51	16	4	1986	1761	43	10	25	6	
		<i>P. phaseoloides</i>	2	3	26	19	3	233	301	5	7	2	4	
		Total	22					2219		48		27		56
M6	6	<i>M. pruriens</i>	5	5	49	17	2	670		21	32	3	4	14
M13	13	<i>C. caeruleum</i>	5	6	28	19	4	172	193	4	2	2	3	50
M18	18	<i>C. caeruleum</i>	10	16	55	28	1	475	840	9	10	6	10	67
¹ Ndfa (%) derived from measured ureides in solute extracts from field stems using a ¹⁵ N isotope dilution calibration (Unkovich et al 2008)														
² Nodulation score based on a modified Corbin et al 1977; ³ s.d. Standard deviation nd not determined														

Discussion

A calibrated ureide technique was used, for the first time globally as far as we are aware, to estimate proportional dependence on N₂ fixation (pNdfa) by cover legumes in oil palm plantations. Stem RUN for *P. phaseoloides* and *C. caeruleum* in a field survey in PNG were within the range (11–76%) reported for these species from a Malaysian rubber plantation study (Norhayati et al. 1988). Dependence on N₂ fixation by the legume covers in this study ranged from 18% (*P. phaseoloides*) to 75% (*C. caeruleum*) and was highly variable between and within species, although overall *C. caeruleum* exhibited higher pNdfa than the other species. The pNdfa obtained for *M. pruriens* (one site only) and *P. phaseoloides* (4 sites) in this field survey in PNG were generally also lower than reported for these species in other tropical field studies using ¹⁵N-based methods (Cadisch et al. 1989; Sanginga et al. 1996a; Sanginga et al. 1996b; Giller 2001b; Fosu 2003; Sanginga 2003). The glasshouse study provided a robust calibration of the ureide technique (using ¹⁵N isotope dilution) for *C. mucunoides* and *P. phaseoloides*, but further investigation is required to confirm a specific calibration for *M. pruriens* which appeared to export less ureide than the other two legumes, and was only sampled at one location in the field survey. The individual calibrations for *C. mucunoides* and *P. phaseoloides* were similar to those of pigeon pea (Peoples et al. 1989) and common bean (Unkovich et al. 2008), whereas that for *M. pruriens* was closer to the calibration obtained for the tree legume species *Codariocalyx gyroides* (Herridge et al. 1996). The ureide calibrations from this study provide

an avenue to more widely use this reliable and fairly simple technique in future N₂ fixation studies of legume covers in oil palm plantations or other agroecosystems.

It was apparent from the field survey that pNdfa for legume covers did not show any relationship with age of oil palm plantation, and hence age of these perennial legume covers. However, as one might expect given evidence in the literature (cited in the introduction) and data from the glasshouse study, pNdfa was negatively correlated with NO₃⁻ concentration in the soil under the cover legumes. Across all species and ages of plantation pNdfa was significantly lower where soil NO₃⁻-N concentrations were above 5mg/kg. Data from the glasshouse study also indicates that N₂ fixation in *C. mucunoides* and *P. phaseoloides* was very sensitive to low levels of available NO₃⁻. *M. pruriens* on the other hand appears less sensitive and may continue to fix N₂ under conditions of higher available N in the root zone. Thus, where mineral N in the soil is high, *C. mucunoides* and *P. phaseoloides* are likely to be more dependent on uptake of NO₃⁻ from soil than *M. pruriens* and this may decrease the potential for loss of N by leaching or through denitrification under these two species. Studies of *P. phaseoloides* as a cover crop with fruit trees in an Amazonian rubber plantation, using ¹⁵N-labelled fertiliser, similarly concluded that it was a good scavenger of applied N (Lehmann et al. 2000) and reduced N leaching losses compared to under the trees (Lehmann et al. 1999).

A decrease in nodule mass for all three legume cover species in response to increasing NO₃⁻-N supplied was observed in the glasshouse study. This is consistent with many observations that legume nodulation, as well as legume-rhizobium symbiotic function, is inhibited by NO₃⁻ (Murray et al. 2016). The results for *C. mucunoides* contrasted with the observation Camargos and Sodek (2010) that nodule mass and N₂ fixation in this species were unaffected by NO₃⁻-N, although that was after only a 5 day exposure to NO₃⁻, and the same study did report a reduction in nodule number for *C. mucunoides* after 35 days exposure. A simple scoring method to assess nodulation of cover legumes in the field survey indicated poor nodulation for legume cover plants in a few of the older plantations, but generally fair to excellent nodulation across most plantations, irrespective of age and soil nitrate concentration. This was not necessarily expected given the highly acidic nature of the soils (Table 1), and that negative effects of low pH in the root zone on both growth of the legume and activity of rhizobia are well documented (Hungria and Vargas 2000). It is likely that inoculation practices using appropriately adapted rhizobium strains may have been implemented, or that at least the indigenous rhizobia are effective at nodulation in these soils as has been observed for *C. mucunoides* in low pH soils of the Cerrado in Brazil (Ferreira et al. 2016). Nodule scoring is a useful rapid assessment of the potential N₂ fixing ability of a legume but it is subjective and does not provide any indication of whether nodules are active or moribund. Perhaps this explains why there was no relationship between Ndfa (%) of legume species and nodule score for this field survey. Indeed, it can be concluded from the results of the glasshouse study that nodule mass is probably a better indicator than nodule score or nodule number in relation to pNdfa (%) and the response of legume N₂ fixation to mineral N.

The measures of ammonium in these soils were extremely high and are considered to be an artefact from the mandatory gamma irradiation rather than representative of field conditions. While irradiation is thought to have little effect on NO₃⁻ concentrations in soils (McNamara et al. 2003), an increase in ammonium might result from amino N released from soil organisms which are all killed during irradiation, with no subsequent conversion to nitrate due to the absence of any active nitrifying bacteria (Lensi et al. 1991). A substantial increase (5 fold) in ammonium concentration in soils has been reported over a 55 day period following similar irradiation (Lensi et al 1991), particularly where soils were moist. Although the soils for this study had been oven dried at 60°C in PNG it is possible that the samples retained some moisture during that process, or under the humid atmospheric conditions prevalent in PNG the stored samples re-absorbed moisture prior to transport and subsequent treatment in Australia. Delayed release of the imported samples from the Australia irradiation facility meant it was 3 months post-irradiation before the soils were analysed. The effects of ammonium on BNF are not as well defined as those for nitrate (Murray et al. 2016) and may merit consideration in future studies, although clearly valid measures need to be obtained using non-irradiated soils.

The standing shoot biomass (443–792 g/m²) obtained for legume cover from the field survey for the four plantations 5 years or less in age is similar to that recorded for 3–4 months growth of the same legume cover species in other tropical field environments (Broughton 1977; Sylvester-Bradley 1984; Zaharah et al. 1986; Sanginga et al. 1996a; Sanginga et al. 1996b; Tian et al. 1999; Giller 2001a; Fosu 2003; Sanginga 2003; Franke et al. 2008; Samedani et al. 2015). Therefore, it could be considered reasonable to assume that amounts of fixed N in standing shoot biomass for each of these four plantations represents 4 months production, and could thus be multiplied by a factor of 3 to obtain annual N₂ fixation estimates of 48–102 kg N /ha/year for legume cover in young PNG plantations. These quantities of fixed N in standing shoot biomass are less than the 150–200 kg N /ha/year reported for the same legume covers under oil palm and rubber plantations in Malaysia (Broughton 1977; Agamuthu and Broughton 1985; Zaharah et al. 1986), but are similar to that for *C. mucunoides* and *M. pruriens* cover crops in Ghana (Fosu 2003). Values reported in the literature appear to be from studies with close to 100% legume cover compared with the lower cover (17–44%) recorded in the present field survey in PNG. A reduced proportion of legume groundcover in plantations is not only due to shading as palm canopy closes, but also the result of other management factors, including poor cover crop establishment, slashing of cover crop vegetation to create harvest paths and removal of plants in the weeded circle. In the PNG survey total

standing above-ground biomass (DM) and N content of cover legumes, and by default calculated amounts of N fixed, decreased in plantations more than five years old because legume groundcover markedly decreased in the older plantations. *C. caeruleum* was the sole species surviving in plantations > 5 years old, perhaps indicating a greater tolerance to shade and other management influences.

It has been pointed out that the few reported amounts of N fixed by cover legumes in oil palm plantations do not include N in litter (Giller and Fairhurst 2003). Data from the field survey in PNG demonstrate that in plantations less than six years old there can be as much or more DM and N in the litter as there is in the corresponding standing shoot biomass of cover legumes. Indeed, assuming the same pNd_fa for litter as shoots, the estimated amount of fixed N in legume litter for young plantations in this study either equalled (5.1–7 kg N/ha, Kapiura estate plantations) or was greater than that in shoots (15.5–19.3 kg N/ha, Mosa estate plantations). Accounting for this fixed N in litter would substantially increase annual estimates of BNF by cover legumes. Furthermore, this litter represents a significant pool of potentially mineralisable N in the warm moist climate of PNG, and is likely to be a factor in the observed accumulation of soil NO₃⁻ in the field survey that was related to reduction of pNd_fa. Fixed N in legume litter sampled under older plantations in the survey was substantially lower at only 0.1–3.4 kg N/ha, often due to poor groundcover. It should be remembered that these amounts of N are for litter sampled at a single point in time and not annual values which are likely to be greater. Nitrogen transfer to litter from living biomass of cover legumes in plantations (oil palm and rubber) varies with age of cover and has been estimated at 120–160 kg N/ha/year for the first 3 years to less than 40 kg N/ha/year over years 4–7 (Agamuthu and Broughton 1985; Pushparajah and Chew 1998). However, annual litter accumulation and turnover cannot be quantified from the limited measures in this study and clearly requires further research to improve estimates of BNF inputs by cover legumes. Furthermore, a commonality between this study and all others that quantify BNF by tropical perennial cover legumes is that N accumulated below-ground in roots has not been measured. Studies with annual crop legumes have indicated that on average 30% of total plant N may be located below-ground (Herridge et al. 2008), and where legumes are dependent on BNF this represents input of fixed N to the system that needs to be included when considering the effect of cover legumes on the N cycle of oil palm plantations.

Conclusions

This study used a calibrated ureide technique for the first time in oil palm plantations to estimate pNd_fa by cover legumes. Proportional dependence on N₂ fixation by legume cover in oil palm plantations in PNG varied widely and was not related to age of the plantation but was strongly negatively correlated with increasing soil nitrate. Total amount of N derived from N₂ fixation in above-ground biomass of legume cover was positively correlated with legume dry matter production which was greatest under young plantations. Legume cover declined with age, due to shading and other management influences, and *C. caeruleum* was the only cover species present in plantations that were seven years or older. Further research is recommended, using the calibrated RUN technique to obtain annual estimates of BNF inputs over those obtained simply from standing shoot biomass by quantifying total legume biomass production over time, including litter and root accumulation. With this knowledge, more informed decisions can be made regarding the effective management of N inputs from fertilisers and legumes, to better synchronise fluxes of N in the system, and move towards more sustainable oil palm cultivation.

Declarations

Acknowledgements

Staff and students at The University of Adelaide, South Australia and the Oil Palm Research Association in Papua New Guinea (PNG) who generously assisted with sampling and other support. Mark Peoples and David Herridge for advice. The Biogeochemistry Centre at The University of Western Australia for isotope analyses. The Australian Centre for International Agricultural Research provided a John Allwright Fellowship for Rachel Pipai and technical support in PNG via Project SMCN-2009-013 'Sustainable management of soil and water resources for oil palm production systems in PNG'.

The Australian Centre for International Agricultural Research funded a John Allwright Fellowship for Rachel Pipai and technical support in PNG via Project SMCN-2009-013 'Sustainable management of soil and water resources for oil palm production systems in PNG'. Research infrastructure support was provided by The University of Adelaide and the Oil Palm Research Association in PNG.

The authors have no relevant financial or non-financial interests to disclose.

All authors contributed to the study conception and design. Set up of experiments, data collection and analysis were performed by Rachel Pipai with supervision and input by Murray Unkovich and Ann McNeill. The first draft of the manuscript was written by Ann McNeill and all authors commented on the initial and subsequent versions. All authors read and approved the final manuscript.

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

References

1. Agamuthu, P, Broughton, WJ (1985) Nutrient cycling within the developing oil palm-legume ecosystem. *Agriculture, Ecosystems & Environment* **13**, 111-123.
2. Allen, M, Bourke, RM, McGregor, A (2009) Cash Income from Agriculture In 'Food and Agriculture in Papua New Guinea.' (Eds RM Bourke, T Harwood.) pp. 283-424. (The Australian National University Canberra)
3. Banabas, M, Turner, MA, Scotter, DR, Nelson, PN (2008) Losses of nitrogen fertiliser under oil palm in Papua New Guinea: 1. Water balance, and nitrogen in soil solution and runoff. *Australian Journal of Soil Research* **46**, 332-339.
4. Broughton, WJ (1977) Effect of various covers on soil fertility under *Hevea brasiliensis* muell. arg. and on growth of the tree. *Agro-Ecosystems* **3**, 147-170.
5. Byerlee, D, Falcon, WP, Naylor, RL (2016) The Tropical Oil Crop Revolution: Food, Feed, Fuel, and Forests. (Oxford University Press Available at <https://doi.org/10.1093/acprof:oso/9780190222987.001.0001> [Accessed
6. Cadisch, G, Sylvester-Bradley, R, Nösberger, J (1989) ¹⁵N-Based estimation of nitrogen fixation by eight tropical forage-legumes at two levels of P:K supply. *Field Crops Research* **22**, 181-194.
7. Camargos, L, Sodek, L (2010) Nodule growth and nitrogen fixation of *Calopogonium mucunoides* L. show low sensitivity to nitrate. *Symbiosis* **51**, 167-174.
8. Cataldo, D, Maroon, M, Schrader, L, Youngs, V (1975) Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Communications in Soil Science and Plant Analysis* **6**, 71-80.
9. Chiu, SB (2004) *Mucuna bracteata* - dry matter conversion and decay rate of litter. *The Planter* **80**, 461-464.
10. Choo, YM, Muhamad, H, Hashim, Z, Subramaniam, V, Puah, CW, Tan, Y (2011) Determination of GHG contributions by subsystems in the oil palm supply chain using the LCA approach. *The International Journal of Life Cycle Assessment* **16**, 669-681.
11. Comte, I, Colin, F, Whalen, JK, Grünberger, O, Caliman, J-P (2012) Chapter three - Agricultural Practices in Oil Palm Plantations and Their Impact on Hydrological Changes, Nutrient Fluxes and Water Quality in Indonesia: A Review. In 'Advances in Agronomy.' (Ed. LS Donald.) Vol. Volume 116 pp. 71-124. (Academic Press:
12. Corbin, E, Brockwell, J, Gault, R (1977) Nodulation studies on chickpea (*Cicer arietinum*). *Australian Journal of Experimental Agriculture* **17**, 126-134.
13. Corley, RHV (2009) How much palm oil do we need? *Environmental Science & Policy* **12**, 134-139.
14. Corley, RHV, Tinker, PBH (2016) Chapter 12 Mineral Nutrition of Oil Palms. In 'The Oil Palm.' pp. 329-398. (John Wiley and Sons Ltd: Chichester UK)
15. Cramb, R, Curry, GN (2012) Oil palm and rural livelihoods in the Asia–Pacific region: An overview. *Asia Pacific Viewpoint* **53**, 223-239.
16. Dawson, T, Brooks, PD (2001) Fundamentals of stable isotope chemistry and measurement In 'Application of stable isotope techniques to study biological processes and functioning of ecosystems.' (Eds MJ Unkovich, JS Pate, A McNeill, DJ Gibbs.) pp. 1-18. (Kluwer Academic: Dordrecht)
17. Fairhurst, T, Hardter, R (Eds) (2003) 'Oil palm: Management for large and sustainable yields.' (IPNI and IPI: Singapore)
18. Ferreira, TC, Aguilar, JV, Souza, LA, Justino, GC, Aguiar, LF, Camargos, LS (2016) pH effects on nodulation and biological nitrogen fixation in *Calopogonium mucunoides*. *Brazilian Journal of Botany* **39**, 1015-1020.
19. Fosu, M (2003) Nitrogen accumulation and release by Sunn Hemp, Calopo, *Mucuna* and devil bean in semi-arid Ghana. *Agricultural and Food Science Journal of Ghana* **2**, 141-153.
20. Franke, A, Laberge, G, Oyewole, BD, Schulz, S (2008) A comparison between legume technologies and fallow, and their effects on maize and soil traits, in two distinct environments of the West African savannah. *Nutrient Cycling in Agroecosystems* **82**, 117-135.
21. Galloway, JN, Leach, AM, Bleeker, A, Erisman, JW (2013) A chronology of human understanding of the nitrogen cycle. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368**, 20130120.
22. Gault, R, Peoples, MB (1993) 'Development of on-farm methods for measuring N₂ fixation. , Proceedings of XVIII International Grassland Congress.' New Zealand.
23. Giller, KE (2001a) Legumes as green manures and cover crops. In 'Nitrogen Fixation in Tropical Cropping Systems.' (Ed. KE Giller.) (CAB International Publishing:
24. Giller, KE (2001b) 'Nitrogen fixation in tropical cropping systems ' (CABI Publishing: London)
25. Giller, KE, Fairhurst, T (2003) Legume cover plants. In 'Oil palm: management for large and sustainable yields.' (Eds T Fairhurst, R Hardter.) pp. 384. (Potash & Phosphate Institute (PPI)/Potash: Singapore)

26. Giller, KE, Wilson, KJ (1991) 'Nitrogen fixation in tropical cropping systems.' (C.A.B International: London)
27. Herridge, DF (1982) Relative abundance of ureides and nitrate in plant tissues of soybean as a quantitative assay of nitrogen fixation. *Plant Physiology* **70**, 1-6.
28. Herridge, DF, Palmer, B, Nurhayati, DP, Peoples, MB (1996) Evaluation of the xylem ureide method for measuring N₂ fixation in six tree legume species. *Soil Biology and Biochemistry* **28**, 281-289.
29. Herridge, DF, Peoples, MB (2002a) Calibrating the xylem-solute method for nitrogen fixation measurement of ureide-producing legumes: cowpea, mungbean, and black gram. *Communications in Soil Science and Plant Analysis* **33**, 425-437.
30. Herridge, DF, Peoples, MB (2002b) Timing of xylem sampling for ureide analysis of nitrogen fixation. *Plant and Soil* **238**, 57-67.
31. Herridge, DF, Peoples, MB, Boddey, R (2008) Global inputs of biological nitrogen fixation in agricultural systems. *Plant and Soil* **311**, 1-18.
32. Lehmann, J, Da Silva, J, Trujillo, L, Uguen, K (1999) 'Legume cover crops and nutrient cycling in tropical fruit tree production, II ISHS Conference on Fruit Production in the Tropics and Subtropics 531.'
33. Lehmann, J, Pereira da Silva, J, Schroth, G, Gebauer, G, Da Silva, LF (2000) Nitrogen use in mixed tree crop plantations with a legume cover crop. *Plant and Soil* **225**, 63-72.
34. Mendham, DS, Kumaraswamy, S, Balasundaran, M, Sankaran, KV, Corbeels, M, Grove, TS, O'Connell, AM, Rance, SJ (2004) Legume cover cropping effects on early growth and soil nitrogen supply in eucalypt plantations in south-western India. *Biology and Fertility of Soils* **39**, 375-382.
35. Murray, JD, Liu, C-W, Chen, Y, Miller, AJ (2016) Nitrogen sensing in legumes. *Journal of Experimental Botany* **68**, 1919-1926.
36. Norhayati, M, Noor, S, Chong, K, Faizah, A, Herridge, DF, Peoples, MB, Bergersen, F (1988) Adaptation of methods for evaluating N₂ fixation in food legumes and legume cover crops. *Plant and Soil* **108**, 143-150.
37. Pardon, L, Bessou, C, Nelson, PN, Dubos, B, Ollivier, J, Marichal, R, Caliman, J-P, Gabrielle, B (2016) Key unknowns in nitrogen budget for oil palm plantations. A review. *Agronomy for Sustainable Development* **36**, 20.
38. Pardon, L, Huth, NI, Nelson, PN, Banabas, M, Gabrielle, B, Bessou, C (2017) Yield and nitrogen losses in oil palm plantations: Main drivers and management trade-offs determined using simulation. *Field Crops Research* **210**, 20-32.
39. Peoples, MB, Hebb, DM, Gibson, AH, Herridge, DF (1989) Development of the xylem ureide assay for the measurement of nitrogen fixation by pigeonpea (*Cajanus cajan* (L.) Millsp.). *Journal of Experimental Botany* **40**, 535-542.
40. Peoples, MB, Unkovich, MJ, Herridge, DF (2009) Measuring symbiotic nitrogen fixation by legumes. In 'Nitrogen fixation in crop production.' (Ed. DW Emerich, Krishnan, H. B.) (American Society of Agronomy: United States of America)
41. Philip, A, Abraham, J (2009) Litter chemistry and decomposition in rubber plantations. *Natural Rubber Research* **22**, 10-16.
42. Pushparajah, E, Chew, PS (1998) 'Integrated nutrient management for sustaining high yield of plantation tree crops in tropical Asia, CD ROM Proceedings of 16th World Congress of Soil Science' Montpellier. (ISSS and FSSS:
43. Rayment, GE, Lyons, DJ (2011) 'Soil chemical methods Australasia.' (CSIRO: Melbourne)
44. Samedani, B, Juraimi, A, Rafii, M, Sheikh Awadz, S, Anwar, M, Anuar, A (2015) Effect of cover crops on weed suppression in oil palm plantation. *Int J Agric Biol* **17**, 251-260.
45. Sanginga, N (2003) Role of biological nitrogen fixation in legume based cropping systems; a case study of West Africa farming systems. *Plant and Soil* **252**, 25-39.
46. Sanginga, N, Ibewiro, B, Houngnandan, P, Vanlauwe, B, Okogun, JA, Akobundu, IO, Versteeg, M (1996a) Evaluation of symbiotic properties and nitrogen contribution of mucuna to maize grown in the derived savanna of West Africa. *Plant and Soil* **179**, 119-129.
47. Sanginga, N, Okogun, A, Akobundu, IO, Carsky, RJ, Tian, G, Wirkom, LE (1996b) Nodulation and estimation of symbiotic nitrogen fixation by herbaceous and shrub legumes in Guinea savanna in Nigeria. *Biology and Fertility of Soils* **23**, 441-448.
48. Sayer, J, Ghazoul, J, Nelson, P, Boedhihartono, AK (2012) Oil palm expansion transforms tropical landscapes and livelihoods. *Global Food Security* **1**, 114-119.
49. Streeter, JG, Wong, PP (1988) Inhibition of legume nodule formation and N₂ fixation by nitrate. *Critical Reviews in Plant Sciences* **7**, 1-23.
50. Sylvester-Bradley, R (1984) Rhizobium inoculation trials designed to support a tropical forage legume selection programme. *Plant and Soil* **82**, 377-386.
51. Teoh, CH (2010) Key Sustainability Issues in the Palm Oil Sector. World Bank Group Washington USA.
52. Tian, G, Kolawole, GO, Salako, FK, Kang, BT (1999) An improved cover crop-fallow system for sustainable management of low activity clay soils of the tropics. *Soil Science* **164**, 671-682.
53. Turner, P, Gillbanks, R (2003) 'Oil Palm Management and Cultivation.' (The Incorporated Society of Planters: Kuala Lumpur Malaysia)

54. Unkovich, MJ, Herridge, DF, Peoples, MB, Cadisch, G, Boddey, B, Giller, KE, Alves, BJR, Chalk, P (2008) 'Measuring plant-associated nitrogen fixation in agricultural systems.' (Australian Centre for International Agricultural Research (ACIAR): Canberra)
55. Vesterager, JM, Osterby, S, Jensen, ES, Schjoerring, JK (1995) Symbiotic N₂-fixation by the cover crop *Pueraria phaseoloides* as influenced by litter mineralization. *Plant and Soil* **177**, 1-10.
56. Wilson, KJ, Peoples, MB, Jefferson, RA (1995) New techniques for studying competition by *Rhizobia* and for assessing nitrogen fixation in the field. *Plant and Soil* **174**, 241-253.
57. Yemm, EW, Cocking, EC (1955) The determination of amino-acids with ninhydrin *Analyst* **80** 209-214.
58. Young, EG, Conway, CF (1942) On the estimation of allantoin by the Rimini-Schryver reaction. *Journal of Biological Chemistry* **142**, 839-853.
59. Zaharah, AR, Sharifuddin, HAH, Razley, MN, Mohd Saidi, AK (1986) Measurement of nitrogen fixed by *Pueraria phaseoloides* by N-15 dilution technique. *Pertanika* **9**, 45-49.

Figures




Figure 1

Hexagonal 'palm unit' assessed for % legume species cover at each location. Palm spacing varied with planting density but at all locations was close to the standard triangular distance between adjacent palms of 9 x 9x 9m (Corley and Tinker 2008). The weeded circle diameter is 1m radius at time of palm planting and is gradually expanded to 2m radius by palm age 2 years.




Figure 2

(A) Legume dependence on N_2 fixation (Ndfa, %) as a function of NO_3^- supplied in the glasshouse, and (B) nodule mass (g/plant) as a function of NO_3^- supply for *C. mucunoides*, *P. phaseoloides* and *M. pruriens*. Values are means across two harvests. Error bars are standard error of the mean for $n=8$.




Figure 3

Relative stem ureide N (RUN, %) for (A) *C. mucunoides* and *P. phaseoloides* as a function of NO_3^- supplied in the glasshouse, and (B) the same for *M. pruriens*. Values are means across two harvests. Error bars are standard error of the mean for $n=8$. A single (power) curve has been fitted across species in (A) and a simple linear regression provided the best fit for *Mucuna* (B).




Figure 4

Correlation of Ndfa (%), derived using ^{15}N isotope dilution technique, and stem RUN (%) for (A) *C. mucunoides* and *P. phaseoloides* and (B) *M. pruriens*. Data are means from two sampling times in the glasshouse experiment.




Figure 5

Correlations between age of oil palm plantation and (A) shoot or (B) litter dry matter (DM), and (C) shoot or (D) litter N content for total legume cover sampled from different aged palm plantations in West New Britain Province, PNG. Power curves have been fitted between age of oil palm and total legume cover DM or N per plantation.




Figure 6

Scatterplots showing legume dependence on N_2 fixation (Ndfa,%) in relation to (A) oil palm age or (B) nitrate-N in 0-20cm soil depth under cover legumes (*C. caeruleum*, *P. phaseoloides* and *M. pruriens*) in different aged oil palm plantations in West New Britain Province, PNG.