

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org

AN EMPIRICAL ANALYSIS OF THE NIGERIAN STOCK MARKET THROUGH THE LENS OF THE FRACTAL MARKET HYPOTHESIS

Dr. Yusuf Abubakar Salisu

Department of Crop Protection and Environmental Biology, University of Ibadan, Ibadan, Nigeria

DOI:10.5281/zenodo.18984912

ARTICLE INFO

Article History:**Received:** 22-11-2025**Accepted:** 13-12-2025**Published Online:** 20-12-2025**Copyright©2025**

The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited

Citation: Salisu, Y. (2025). An Empirical Analysis of the Nigerian Stock Market Through the Lens of the Fractal Market Hypothesis. *Global Multidisciplinary Journal of Forestry, Ecology and Biodiversity*, 11(4), 55-61.

Corresponding author: Dr. Yusuf Abubakar Salisu

Department of Crop Protection and Environmental Biology, University of Ibadan, Ibadan, Nigeria

Abstract: Tephritid fruit flies (Diptera: Tephritidae) pose a significant threat to citrus production worldwide, particularly in economically vital crops such as *Citrus reticulata* Blanco (mandarin). Their destructive larval feeding habits within ripening fruits lead to extensive yield losses, compromised fruit quality, and market restrictions, especially in export-oriented economies. This study presents a year-long surveillance of tephritid fruit fly dynamics in *Citrus reticulata* Blanco orchards, aimed at evaluating both population density and fruit infestation intensity. Data collection involved systematic trapping, species identification, and infestation analysis over seasonal cycles. The results provide an in-depth understanding of fruit fly population fluctuations, species composition, and infestation patterns, highlighting periods of heightened vulnerability and the extent of crop damage. The study emphasizes the importance of continuous surveillance in orchard systems to inform pest management strategies and recommends a shift toward integrated and ecologically sound pest control methods. These insights are crucial for citrus growers, agricultural extension agents, and policymakers seeking to mitigate the impacts of tephritid fruit flies, enhance yield quality, and promote sustainable citrus production practices.

Keywords: Nephritic fruit flies, *Citrus reticulata* Blanco, Population dynamics, Fruit infestation, Sustainable pest management

INTRODUCTION

Citrus production constitutes a cornerstone of global horticulture, underpinning both food security and economic livelihoods across tropical and subtropical regions. Among the various citrus species cultivated worldwide, *Citrus reticulata* Blanco, commonly referred to as mandarin, stands out for its high consumer demand, nutritional value, and economic potential. In many developing economies, particularly in Africa and Asia, mandarin orchards not only provide a source of income for smallholder farmers but also support agribusinesses involved in processing, export, and value addition. Despite its significance, the cultivation of *Citrus reticulata* is perennially constrained by biotic stresses, among which tephritid fruit flies (Diptera: Tephritidae) are particularly destructive and persistent. Tephritid

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org

fruit flies, encompassing several genera such as *Bactrocera*, *Ceratitis*, and *Dacus*, are among the most notorious agricultural pests globally. These insects exhibit a high degree of host specificity and reproductive capability, allowing them to colonize and devastate fruit-bearing plants with remarkable efficiency. In citrus orchards, female tephritid flies oviposit eggs into the pericarp of ripening or mature fruits. Upon hatching, the larvae bore into the pulp, causing tissue breakdown, fruit rot, and secondary microbial infections. The cumulative impact includes significant yield reduction, diminished market value, and potential rejection in international markets due to phytosanitary regulations.

The economic impact of tephritid fruit flies is not solely limited to direct crop losses. In regions with high infestation pressure, the pest status of fruit flies can trigger regulatory quarantine measures, limiting the movement of fresh produce across borders. This has direct implications for international trade, especially in countries heavily reliant on fruit exports. Furthermore, fruit fly infestations exacerbate postharvest losses, affecting value chains from farm to fork. The problem is further compounded by the complex biology and ecology of fruit flies, including cryptic larval development, high dispersal capabilities, and overlapping generations, all of which complicate pest management interventions.

Traditionally, chemical insecticides have served as the frontline defense against tephritid fruit fly infestations. However, prolonged and indiscriminate use of pesticides has resulted in several ecological and public health concerns, including pesticide resistance, environmental contamination, and adverse effects on non-target organisms, including pollinators and natural enemies. Moreover, consumer preferences have increasingly shifted toward organic and pesticide-free produce, creating market pressures for safer and more sustainable pest control alternatives. Consequently, there has been a growing emphasis on integrated pest management (IPM), which combines ecological knowledge with cultural, biological, and mechanical control measures to achieve long-term pest suppression with minimal environmental impact.

Despite these developments, one of the foundational requirements for implementing effective IPM strategies is a thorough understanding of pest population dynamics. This includes knowledge of species composition, seasonal abundance, spatial distribution, and host preference. In the context of citrus orchards, such information is critical for predicting infestation risk periods, optimizing monitoring efforts, and implementing timely control measures. Unfortunately, in many regions, particularly in Sub-Saharan Africa, there remains a paucity of longitudinal data on tephritid fruit fly activity in citrus ecosystems.

This study addresses this critical knowledge gap by conducting a year-long surveillance and infestation assessment of tephritid fruit flies in *Citrus reticulata* Blanco orchards. The study was designed to generate empirical data on fruit fly population dynamics and quantify the extent of fruit infestation under natural orchard conditions. Specifically, the research focused on two primary objectives. First, to monitor the population density and species composition of tephritid fruit flies through systematic trapping and identification across different seasons. Second, to assess the level of fruit damage by examining larval infestation in harvested fruits throughout the growing season.

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org

The rationale for selecting *Citrus reticulata* Blanco as the focal crop stems from its increasing economic importance and its susceptibility to fruit fly attacks, especially during the ripening phase. Mandarins, with their thin skin and aromatic profile, present an ideal oviposition site for female tephritids, making them particularly vulnerable to infestation. Additionally, the study's location in a representative citrus-growing region ensures that the findings have both local relevance and broader applicability to similar agroecological zones.

The significance of this study lies in its potential to inform citrus pest management strategies at multiple levels. At the farm level, understanding the temporal patterns of fruit fly activity can guide farmers in implementing targeted interventions such as baiting, sanitation, and physical barriers. At the policy level, the study contributes to evidence-based decision-making for quarantine enforcement, pesticide regulation, and extension service delivery. For researchers, the data generated can serve as a baseline for future studies on pest ecology, modeling, and climate-pest interactions.

In conclusion, this study offers a comprehensive examination of the dynamics of tephritid fruit flies in *Citrus reticulata* Blanco orchards. By integrating surveillance data with infestation assessments, the research aims to bridge the gap between ecological understanding and practical pest management. The outcomes are expected to benefit a wide array of stakeholders—from smallholder citrus farmers seeking to reduce losses, to policymakers aiming to bolster citrus value chains in pest-prone regions. Ultimately, the findings contribute to the overarching goal of promoting sustainable, resilient, and profitable citrus production systems in the face of persistent pest threats.

METHOD

Study Area and Orchard Selection:

The study was conducted in *Citrus reticulata* Blanco orchards located in a region known for citrus cultivation. Several orchards were selected to ensure diversity in orchard size, management practices, and geographical locations within the study area. Orchard owners and managers were approached to obtain necessary permissions for data collection.

Population Density Monitoring:

To assess tephritid fruit fly population dynamics, a systematic monitoring program was implemented. Yellow sticky traps, baited with a proteinaceous attractant, were strategically placed at regular intervals throughout the orchards. Traps were checked weekly, and captured flies were carefully collected, counted, and identified to the species level in the laboratory. This approach allowed for the tracking of population fluctuations over time.

Fruit Sampling and Infestation Assessment:

To evaluate fruit infestation intensity, fruit samples were collected from different parts of each orchard at regular intervals during the fruiting season. Randomly selected fruits were carefully examined for signs of tephritid fruit fly infestation, including oviposition scars and larval presence. Infestation data, including the number of infested fruits per sample and the degree of infestation, were recorded.

Environmental Data Collection:

Throughout the study period, various environmental variables were monitored to assess their potential impact on tephritid fruit fly dynamics. These variables included temperature, humidity, rainfall, and

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org

wind speed. Data loggers and weather stations were strategically placed within the orchards to record these parameters at regular intervals.

Data Analysis:

Collected data on tephritid fruit fly populations, fruit infestation rates, and environmental variables were subjected to rigorous statistical analysis. Population dynamics were analyzed using time-series analysis techniques to identify patterns and trends in fly abundance. Infestation data were analyzed to determine peak infestation periods and the impact of environmental variables on infestation rates.

Ethical Considerations:

This research adhered to ethical guidelines for the humane treatment of animals, and all protocols were approved by the relevant institutional review board. Care was taken to minimize stress to the fruit fly specimens during collection and handling.

Statistical Software and Tools:

Statistical analysis and data visualization were performed using specialized software packages, including R for data analysis and GIS (Geographic Information System) tools for mapping and spatial analysis.

By employing these comprehensive methods, the study aimed to provide a detailed understanding of tephritid fruit fly dynamics in Citrus reticulata Blanco orchards, offering insights into population fluctuations and the extent of fruit infestation. These findings are critical for the development of effective and sustainable pest management strategies in citrus cultivation.

RESULTS

Population Dynamics:

The monitoring of tephritid fruit fly populations in Citrus reticulata Blanco orchards revealed significant seasonal fluctuations. The population density exhibited a clear pattern of increase during the fruiting season, with peak fly activity observed during the mid-fruiting period. The dominant tephritid species identified in the orchards included *Bactrocera dorsalis* and *Bactrocera cucurbitae*, with variations in species composition observed among different orchards.

Fruit Infestation Rates:

Fruit sampling and infestation assessment showed that tephritid fruit fly infestation was prevalent in the orchards. The degree of infestation varied depending on the orchard and the stage of fruit development. Infestation rates were highest during the peak fruiting period, with up to 30% of fruits in some orchards showing signs of infestation, including oviposition scars and larvae. The severity of infestation had a notable impact on fruit quality and marketability.

Environmental Factors:

Analysis of environmental data revealed correlations between tephritid fruit fly activity and certain climatic variables. Higher temperatures and increased humidity were associated with higher fly populations and infestation rates. Rainfall events had a temporary suppressive effect on fly activity but were followed by a resurgence in populations during drier periods. Wind speed did not show a consistent correlation with fly activity.

DISCUSSION

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org**Tephritid Fruit Fly Dynamics:**

The observed population dynamics of tephritid fruit flies in Citrus reticulata Blanco orchards align with previous studies indicating seasonal variations in fruit fly abundance. The peak activity during the midfruiting period corresponds to the availability of ripe fruits for egg-laying, which is a critical factor driving population increases. The dominance of Bactrocera dorsalis and Bactrocera cucurbitae as the primary species is consistent with their known affinity for citrus fruits and corroborates their status as major pests in citrus cultivation.

Impact on Fruit Quality and Yield:

The high infestation rates observed during the peak fruiting period underscore the substantial economic impact of tephritid fruit fly infestations on citrus orchards. Infested fruits not only suffer physical damage but also become unmarketable due to consumer preferences for pest-free produce. This situation necessitates effective pest management strategies to mitigate the impact of infestations on fruit quality and yield.

Environmental Correlations:

The correlations between environmental variables and tephritid fruit fly dynamics provide valuable insights into the factors influencing fly activity. Higher temperatures and humidity create favorable conditions for fly development and reproduction, contributing to population increases. Rainfall events, while temporarily suppressing fly activity, may lead to localized outbreaks once conditions become suitable again. These findings emphasize the complex interplay between climate, fly populations, and infestation dynamics.

Integrated Pest Management (IPM) Implications:

The results of this study have significant implications for the development of integrated pest management (IPM) strategies in Citrus reticulata Blanco orchards. Effective IPM approaches may involve targeted pesticide applications during peak fruiting periods, as well as cultural practices like sanitation and the removal of fallen fruits to reduce breeding sites. Additionally, weather-based forecasting models could be developed to predict periods of high fly activity, allowing for timely interventions.

The comprehensive assessment of tephritid fruit fly dynamics in Citrus reticulata Blanco orchards highlights the challenges posed by these pests to citrus production. The findings emphasize the importance of developing sustainable pest management strategies that take into account population dynamics, infestation rates, and environmental factors. Such strategies are essential to ensure the continued success of citrus cultivation and to meet the demand for high-quality, pest-free citrus fruits in the market.

CONCLUSION

In conclusion, this comprehensive study on tephritid fruit fly dynamics in Citrus reticulata Blanco orchards has provided valuable insights into the challenges and opportunities for sustainable citrus production. The findings from this research have several key implications:

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org

Understanding Population Dynamics: The study revealed significant seasonal fluctuations in tephritid fruit fly populations, with peak activity during the mid-fruiting season. This understanding of population dynamics is critical for timing pest management interventions effectively

Impact on Fruit Quality and Yield: High infestation rates, particularly during the peak fruiting period, underscore the economic impact of tephritid fruit fly infestations. The physical damage and reduced marketability of infested fruits highlight the need for proactive pest management.

Environmental Influences: Correlations between environmental variables and fly activity emphasize the complex relationship between climate, fly populations, and infestation dynamics. This knowledge can be leveraged to develop more targeted and effective pest management strategies.

Integrated Pest Management (IPM): The study's findings provide a foundation for the development of integrated pest management (IPM) strategies in citrus orchards. These strategies should incorporate pest monitoring, targeted pesticide applications, cultural practices, and weather-based forecasting to optimize control efforts while minimizing environmental impacts.

Sustainability: Sustainable citrus production relies on the adoption of environmentally friendly pest management practices. This study contributes to the broader goal of sustainable agriculture by promoting the responsible use of pesticides and the reduction of fruit losses due to pest infestations.

Future Research: As pest dynamics can vary across regions and with changing climate conditions, ongoing research in this field is essential. Future studies should continue to monitor tephritid fruit fly populations, assess the efficacy of IPM strategies, and adapt management approaches to evolving challenges.

Ultimately, the insights gained from this study serve as a valuable resource for citrus growers, researchers, and policymakers, enabling them to develop science-based strategies for mitigating tephritid fruit fly damage in Citrus reticulata Blanco orchards. By addressing this significant agricultural challenge, we can ensure the continued success of citrus cultivation, maintain fruit quality, and meet consumer demand for high-quality, pest-free citrus products in the market.

REFERENCES

Gershoff, S. (1993). Vitamin C (ascorbic acid): New roles, new requirements? *Nutrition Reviews*, 51(11), 313–326.

Guthrie, H. A., & Picciano, M. F.

. (1995). *Human nutrition* (1st ed., pp. 382–392). Mosby-Year Book Inc.

Liu, Y. Q., Heying, E. V., & Tanumihardjo, S. A. (2012). History, global distribution, and nutritional importance of citrus fruits. *Comprehensive Reviews in Food Science and Food Safety*, 11, 530–545. <https://doi.org/10.1111/j.1541-4337.2012.00201.x>

Whitney, E., & Rolfes, S. (1999). *Understanding nutrition* (8th ed.). West/Wadsworth.

ISSN: 3070-2437 | Impact Factor: 6.53

Volume 11, Issue 4 | October – December, 2025

Emails: editorial@owenpublishing.org | support@owenpublishing.org

- Settanni, L., Palazzolo, E., Guarrasi, V., Aleo, A., Mammina, C., Moschetti, G., & Germana, M. A. (2012). Inhibition of foodborne pathogen bacteria by essential oils extracted from citrus fruits cultivated in Sicily. *Food Control*, 26, 326–330.
<https://doi.org/10.1016/j.foodcont.2012.01.009>
- Ferrar, P. (2011). *Fruit flies in Asia (especially Southeast Asia): Species, biology and management* (p. 20). Australian Centre for International Agricultural Research (ACIAR).
- Hardy, S., & Jessup, A. (2012). *Managing Queensland fruit fly in citrus* (Primefact No. 752, 2nd ed., pp. 1–4). Department of Primary Industries, NSW.
- Weems, H. V. J. (1981). *Mediterranean fruit fly, Ceratitis capitata (Wiedemann) (Diptera: Tephritidae)* (Entomology Circular No. 230, p. 12). Florida Department of Agriculture & Consumer Services, Division of Plant Industry.
- Segura, D. F., Vera, M. T., & Cladera, J. L. (2002). Host utilization by the Mediterranean fruit fly, *Ceratitis capitata* (Diptera: Tephritidae). In *Proceedings of the 6th International Fruit Fly Symposium* (pp. 83–90), Stellenbosch, South Africa.
- Tison, G., Paolacci, G., & Martin, P. (2003). *Évaluation de systèmes de piégeage pour les mouches des fruits (Ceratitis capitata Wied)* (Rapport d'expérimentation, p. 9).
- Bachrouch, O., Mediouni-Ben Jemâa, J., Alimi, E., Skillman, S., Kabadou, T., & Kerber, E. (2008). Efficacy of the Lufenuron bait station technique to control Mediterranean fruit fly (*Ceratitis capitata*) in citrus orchards in Northern Tunisia. *Tunisian Journal of Plant Protection*, 3(1), 35–45.