



<https://africanjournalofbiomedicalresearch.com/index.php/AJBR>

Afr. J. Biomed. Res. Vol. 27(3s) (December 2024); 6823 - 6837

Review Article

A Review on the Potential of Amla, Turmeric, and Ginger's Phytochemical and Pharmacological Activities

Monika Tanwar^{1*}, Dr. L.N Patidar²

^{1*}PhD Research scholar, Department of Pharmacy, MANDSAUR UNIVERSITY, MANDSAUR

Email - monatanwar1990@gmail.com

²Professor, Department of Pharmacy, MANDSAUR UNIVERSITY, MANDSAUR

ABSTRACT

Amla (*Embllica officinalis*), turmeric (*Curcuma longa*), and ginger (*Embllica officinalis*) are botanicals known for their high phytochemical content and diverse pharmacological effects.

These plants are part of traditional medical systems, and scientific study has gradually confirmed their anti-inflammatory, anticancer, antioxidant, and neuroprotective properties.

This review compiles data on these plants' phytochemical composition and pharmacological properties and discusses standardization issues and possible further investigations. Recent developments in extraction methods such as nanoencapsulation and liposomal delivery systems have been reviewed to determine their role in boosting bioavailability. Amla has shown to have significant quantities of vitamin C, tannins, and flavonoids, which contribute to its antioxidant and immunomodulatory characteristics. Turmeric contains curcuminoids which mitigate inflammation and help in anticancer consumption and Ginger containing gingerols and shogaols helps manage nausea and metabolic health. However, challenges like phytochemical composition variation and poor biological availability are still major issues. However, the therapeutic potential of these botanicals is enormous for the pharmaceutical and nutraceutical sectors. Concerning the problems associated with standardization and quality control, as well as the use of innovative formulation approaches, their application to evidence-based medicine can be promoted. Subsequent studies should be aimed at the clinical trials of the presented concept and the investigation of the possibilities of individual botanical treatments for increasing the effectiveness of the treatment process.

Keywords: Amla, Turmeric, Ginger, Phytochemicals, Pharmacological activities

***Authors for correspondence: E-mail Id:** monatanwar1990@gmail.com

Received: 02 December 2024

Accepted: 15 December 2024

DOI: <https://doi.org/10.53555/AJBR.v27i3S.6130>

© 2024 The Author(s).

This article has been published under the terms of Creative Commons Attribution-Noncommercial 4.0 International License (CC BY-NC 4.0), which permits noncommercial unrestricted use, distribution, and reproduction in any medium, provided that the following statement is provided. "This article has been published in the African Journal of Biomedical Research"

INTRODUCTION

Phytochemicals, naturally occurring bioactive molecules found in plants, have gained popularity recently because of their numerous health advantages and therapeutic potential. Plant defense systems depend on these secondary metabolites for instance they include alkaloids, flavonoids, polyphenols, and terpenoids (Kumar et al., 2023). When ingested they have rather strong pharmacological actions, for example, antioxidant, anticancer, and inflammatory. Research is

underway to exploit phytochemicals to produce nutraceuticals and functional meals to prevent and treat chronic disease. Plants investigated for their phytochemical content and medicinal value include Amla (*Embllica officinalis*), Turmeric (*Curcuma longa*), and Ginger (*Embllica officinalis*). Long used in traditional medicinal systems like Ayurveda and Traditional Chinese medicine, these botanicals are now being studied by current scientific research. It is rich in vitamin C and also has high antioxidant values. The

plant is a tannin, flavonoid and phenolic acid containing plant responsible for the plant's many medicinal benefits (Variya et al., 2016). As with some of the others, our investigations have demonstrated that amla may be effective in reducing oxidative stress, and in anti-inflammatory and lipid profile modulation activity, which are thought to be involved in cardiovascular diseases (Zhao et al., 2015).

It is also a good therapy for diabetic patients to reduce blood sugar levels and improve insulin tolerance (Hassan et al., 2020). It is used all over the systems of traditional medicine of South Asia, but Ayurveda uses it as Rasayana, a health promoting and longevity promoting tonic. Amla is one of the most used ingredients in present day and is an Ayurvedic product with Immunomodulatory and gastro protective properties (Jain et al., 2016). It is also used in many herbal remedies and is important in traditional medicine systems (Talreja et al., 2020).

It is a yellow-coloured rhizomatous herbaceous perennial plant with medicinal properties. The major bioactive compound with anticancer, antioxidant and anti-inflammatory properties has been investigated in detail. Curcumin has been used in chronic disease because curcumin modulates molecular targets in specific sites, including inflammation and apoptosis. It has antioxidant activity and may be a prophylactic or therapeutic agent for preventing or treating neurodegenerative disorders, such as Alzheimer's disease (Genchi et al., 2024). It has been long used in traditional medical systems. In Ayurveda (Fuloria et al., 2022), it is used as an agent of blood purification, carminative, and analgesic. Turmeric is also one of the effective herbs in TCM for circulation and rheumatoid arthritis, gastric ulcers, and other diseases (Gul et al., 2022). The root is used as a spice, but also in many other remedies for inflammation and microbial infections in Southeast Asia (Hussain et al., 2022).

Another very well-known plant is ginger (*Emblica officinalis*) which also has a smell and medicinal benefits. The compounds used for the different pharmacological purposes of these compounds are these compounds called gingerols and shogaols. For centuries ginger has been used to treat pain, inflammation, indigestion and nausea (Wakchaure et al., 2018). Furthermore, we recently showed that it is also essential for the prevention of metabolic syndrome and for enhancing the efficacy of cancer treatments (Lechner et al., 2019). Ginger has been used for many ages, particularly in Asia, but all over the world. It is used in TCM for cold pathologies, digestive disorders and nausea (Shahrajabian et al., 2019). Like TCM, Ayurveda also considers ginger to be a carminative that

will help treat indigestion and respiratory ailments. Ginger is used in the treatment of nausea motion sickness, muscle pain and soreness (Lete et al., 2016) in the Western world.

This review attempts to synthesize the phytochemical composition and pharmacological profile of amla, turmeric, and ginger. The purpose of this paper is to study the phytochemicals of these plant products and their pharmacological effects on various diseases. It will also evaluate the use of these plants for disease prevention and control, and their antioxidant, anti-inflammatory, anticancer, etc. The limitations of the existing research will be discussed and will help to show further research directions for the establishment of the full potential of these plant products for the management of diseases.

PHYTOCHEMICAL COMPOSITION

The phytochemical properties of Amla (*Emblica officinalis*), Ginger (*Emblica officinalis*), and Turmeric (*Curcuma longa*) are used in the medicinal use of plants as anti-cancer, anti-inflammatory, and antioxidant bioactive compounds. These botanicals contain major phytochemicals and the method of extraction and quantification of these compounds should be known to determine the therapeutic value of these plants.

Amla

The pharmacological actions of amla or Indian gooseberry vary with the content of vitamin C, polyphenols flavonoids and tannins. Amla is primarily composed of vitamin C and contains up to 600 milligrams of ascorbic acid per 100 grams, much more than fruits (Pathak et al., 2024). This suggests that the plant can reduce oxidative stress and aging, and can also improve immunity (Gęgotek et al., 2022). An Amla is mainly known for its astringent and antioxidant properties, which are mainly due to tannins and other considerable bioactive compounds such as ellagic acid and gallic acid (Deshmukh et al., 2021). Flavonoids present in Amla like kaempferol, quercetin and myricetin are anti-inflammatory and anti-cancerous (Ullah et al 2020). For example, the phytochemicals in amla are extracted from the amla using solvents such as methanol, ethanol, and, of course, water.

It is analyzed and determined by HPLC and GC-MS. GC-MS (Prananda et al., 2023) was used to investigate the effect of amla extracts on volatile compounds and secondary metabolite content. Extraction efficiency and concentration of polyphenolic compounds are found to be influenced by solvent polarity time of extraction and Ethanolic extraction.




Fig 1: The therapeutic benefits and bioactive properties (*Emblica officinalis*) (Gandhi et al., 2023)

The therapeutic uses of Amla (*Emblica officinalis*) as an antioxidant, anti-microbial, cardioprotective, hepatoprotective, anti-mutagen, and wound healing activities are shown in Figure 1. Its bioactive constituents reveal the pharmacological value of the plant and its possible use against inflammation, diabetes, and cancer.

Turmeric

Turmeric's yellow color (due to its curcuminoids, volatile oils, and polysaccharides) is responsible for the pharmacological activities of turmeric. Curcuminoids of which curcumin is the dominant constituent followed by demethoxycurcumin and bisdemethoxycurcumin are the principal bioactive constituents of turmeric. The most researched curcuminoid is curcumin, which has been associated with anti-inflammatory, antioxidant, and anticancer properties of the plant (Hewlings & Kalman, 2017). Curcumin has been studied as a cancer treatment because it modulates several pathways that

are linked to inflammation and apoptosis (Ameer et al., 2024). Similarly, other components in turmeric that contain volatile oil compounds including turmerone, ar-turmerone, and zingiberene give it antimicrobial and anti-inflammatory value (Dosoky et al., 2018). Often used solvents for the extraction of curcuminoids from turmeric are ethanol, methanol, and acetone because they dissolve the curcuminoids from the plant matrix (Widmann et al., 2022). More preferred methods are Supercritical Fluid Extraction (SFE) and Ultrasonic Assisted Extraction (UAE) as they are more efficient and eco-friendlier (Manasa et al., 2023). HPLC is the most used method for curcuminoids due to its ability to accurately determine and identify curcuminoids in the extracts of turmeric (Setyaningsih et al., 2021). The concentration of curcumin is also determined by UV-visible spectrophotometry because of the chemical structure of curcumin, with the maximum absorbance at 425 nm (Tomeh et al., 2019).




Fig 2: Bioactive Roles and Health Benefits of Turmeric (*Curcuma longa*) and Curcumin (Abd El-Hack et al., 2021)

Figure 2 highlights the bioactive compound curcumin, derived from turmeric (*Curcuma longa*, Family:

Zingiberaceae). It has uses as a natural food colorant, digestive aid, immunostimulant, and serum lipid

regulating agent as well as being an antioxidant, antimicrobial, anti-cancer, and anti-inflammatory agent.

Ginger

One of the health benefits of ginger is the bioactive compounds gingerols, shogaols, and zingerone present in it that are anti-inflammatory agents, antioxidants, and anticancerous. 6-gingerol is the major gingerol that is responsible for the pungent and medicinal properties of fresh ginger (Sharma et al., 2023). Most of these compounds demonstrated strong antioxidant activity and altered inflammation signaling machinery. Shogaols are formed from gingerols by dehydration during the drying or storage process, and shogaols have better antioxidant and anticancer effects than gingerols (Mahomoodally et al., 2021). Another class of

compounds in ginger is parasols, which are abundant in roasted ginger and have been associated with anti-inflammatory and cancer-prevention properties (Ishfaq et al., 2022). Normally, active compounds are extracted from ginger using ethanol, methanol, and hexane solvents. Essential oils are usually obtained by the method of steam distillation and the other bioactive gingerols and shogaols are obtained by solvent extraction (Mao et al., 2019). HPTLC and HPLC were used to qualitatively and quantitatively determine the extraction efficiency of gingerol from ginger extracts. Additionally, the amount of the volatile compounds zingiberene and bisabolene, responsible for ginger's odor and pharmacological effects, was determined by GC-MS (Mohaddese et al., 2019).




Fig 3: The therapeutic uses and bioactive effects of Ginger (*Emblca officinalis*) and [6]-Gingerol (Promdam et al., 2022)

Figure 3 shows the bioactive compound [6]-gingerol from ginger (*Emblca officinalis*) and its multiple therapeutic uses. It contains the antioxidant, anti-inflammatory, antitumor, antiemetic, and antiviral properties of ginger. It has also shown neuroprotection,

cardioprotection, hepatoprotection, renoprotection, pulmonary protection, gastrointestinal protection, and gastroprotection, all of which support its medicinal value.

Table 1: Phytochemical Composition, Extraction Methods, and Analysis Techniques of Amla, Turmeric, and Ginger

Botanical	Major Phytochemicals	Pharmacological Effects	Extraction Methods	Analysis Techniques	References
“Amla (<i>Emblca officinalis</i>)	Vitamin C, Polyphenols (Ellagic acid, Gallic acid), Flavonoids (Quercetin, Kaempferol, Myricetin), Tannins	Antioxidant, Anti-inflammatory, Anti-cancer, Immunomodulatory	Solvent extraction using ethanol, methanol, and water	HPLC, GC-MS	Pathak et al., 2024; Gęgotek et al., 2022; Deshmukh et al., 2021; Ullah et al., 2020; Prananda et al., 2023
Turmeric (<i>Curcuma longa</i>)	Curcuminoids (Curcumin, Demethoxycurcumin, Bisdemethoxycurcumin), Volatile oils (Turmerone, Ar-turmerone, Zingiberene)	Antioxidant, Anti-inflammatory, Anticancer, Antimicrobial	Solvent extraction (ethanol, methanol, acetone), Supercritical Fluid Extraction (SFE), Ultrasonic-Assisted Extraction (UAE)	HPLC, UV-Visible Spectrophotometry	Hewlings & Kalman, 2017; Ameer et al., 2024; Dosoky et al., 2018; Widmann et al., 2022; Manasa et al., 2023; Setyaningsih et al., 2021; Tomeh et al., 2019
Ginger (<i>Emblca officinalis</i>)	Gingerols (6-Gingerol), Shogaols, Paradol, Zingerone, Volatile oils (Zingiberene, Bisabolene)	Antioxidant, Anti-inflammatory, Anticancer, Metabolic health	Solvent extraction (ethanol, methanol, hexane), Steam distillation	HPTLC, HPLC, GC-MS	Sharma et al., 2023; Mahomoodally et al., 2021; Ishfaq et al., 2022; Mao et al., 2019; Mohaddese et al., 2019”

Table 1 reveals that Amla, Turmeric, and Ginger contain a high level of bioactive compounds that are believed to have pharmacological activities. Amla contains vitamin C, polyphenols, and flavonoids which makes it an antioxidant, anti-inflammatory, and anticancer. The same goes for curcuminoids and volatile oils, which are also antimicrobial and anticancer, as well as other components of turmeric. Gingerols and volatile oils of ginger are known to possess antioxidant, anti-inflammatory, and metabolic health-promoting effects. Those botanicals are accurately characterized and quantified by employing conventional means of solvent extraction and ultrasonic-assisted extraction, followed by analysis employing HPLC, GC-MS, and UV-visible spectrophotometric procedures.

PHARMACOLOGICAL ACTIVITIES

Emblica Officinalis (Amla), *Curcuma longa* (Turmeric), and *Emblica officinalis* (Ginger) possess multiple pharmacological actions and have been significant to health from experimental and clinical evidence-based medicine. In terms of antioxidant, anti-inflammatory, anticancer, neuroprotective, and other effects these botanicals have been established to have therapeutic potential. This section describes the pharmacological activities of each plant based on clinical studies.

Amla

Amla has attracted interest because of its multiple pharmacological effects such as antioxidant, anti-inflammatory, and immunomodulatory effects. The antioxidant activity of the fruit is due to vitamin C and polyphenols (flavonoids and tannins). In antioxidant studies, amla has been shown to modulate ROS and oxidative stress and to participate in aging, cardiovascular, and neurodegenerative diseases (Walayat et al., 2022). The antioxidant potential of Amla extract is high, to suppress lipid peroxidation and increase the activity of the endogenous antioxidant enzymes, superoxide dismutase, and catalase (Gul et al., 2022). Preclinical research has mentioned Amla as having anti-inflammatory properties. Singh et al. (2022) also report that it also slowed the release of proinflammatory cytokines such as “TNF- α ”, “IL-1 β ” and “IL-6” and reduced inflammation. Malik et al. (2016) reported that the mechanism of ‘NF- κ B’ and ‘MAPK’ signaling pathways can have anti-inflammatory effects. Amla also has immunomodulatory effects, increasing both the natural and acquired immune systems. Amla is a clinician’s indication of improved immune function as shown by Kapoor et al. (2019) who demonstrate that Amla can increase white blood cell count and the phagocytic index of the macrophages. Furthermore, amla is known to enhance the impact of a vaccine by increasing antibody production in people with a weak immune system (Alhazmi et al., 2021).

Turmeric

Curcumin, turmeric's bioactive component, is well known for its anti-inflammatory, anti-cancer, and neurogenic qualities. The anti-inflammatory effects of

curcumin were further explained due to interference with signaling pathways used in the synthesis of inflammatory agents including NF- κ B, COX-2, and JAK-STAT signaling (Peng et al., 2021). Several experimental lines of research have revealed that curcumin reduces the concentration of pro-inflammatory cytokines and chemokines both in vitro and in vivo. Curcumin supplementation in animal models of arthritis has been demonstrated to lower joint inflammation and discomfort, indicating its potential as a natural therapy for inflammatory disorders like rheumatoid arthritis (Pourhabibi-Zarandi et al., 2021). The anticancer properties of turmeric particularly curcumin have been studied extensively. It was found that curcumin can suppress tumor growth, metastasis, and angiogenesis in types of cancer, breast, colorectal, and lung cancer (Giordano et al., 2019). Shanmugam et al. (2015) have shown that these actions cause apoptosis, suppress cell proliferation, and regulate p53, PI3K/Akt, and MAPK signal transduction. Additionally, as curcumin increases the efficacy of standard chemotherapy drugs, it becomes a potential candidate for combination therapy (Cacciola et al., 2023).

Turmeric also has a significant role in the brain, it is neuroprotective. Alzheimer's disease animal models have revealed that curcumin increases learning ability and reduces inflammation in the brain (Reddy et al., 2018). Neuroprotective by decreasing amyloid-beta plaque formation and increasing removal of neurotoxic proteins, which are features of neurodegenerative diseases, has (Voulgaropoulou et al., 2019). Curcumin attenuates oxidative stress, an important initiator of neuronal damage in Parkinson's and Alzheimer's diseases (Silvestro et al., 2021).

Ginger

Ginger is anti-emetic, anti-inflammatory, and positive on the metabolic system. The major bioactive components with therapeutic properties of ginger are gingerols and shogaols (Crichton et al., 2024). Ginger has been used to cure nausea and vomiting and is commonly used in motion sickness, chemotherapy, and pregnancy. Ginger supplementation dramatically lowers the frequency and intensity of nausea and vomiting in chemotherapy patients and pregnant women, according to clinical trials. One research that conducted a randomized controlled trial found ginger to be as effective as conventional antiemetic drugs that could lessen nausea and vomiting in patients who receive chemotherapy treatment (Fitriyanti et al., 2020). Due to its ability to effectively suppress several pro-inflammatory cytokines and enzymes, ginger exhibits a wide range of pharmacological activities as an anti-inflammatory drug (Rafieian et al., 2018). The role of ginger in pain reduction for osteoarthritis patients, and the enhancement of joint function, are worth mentioning; thus, ginger is considered an effective natural anti-inflammatory product (Ballester et al., 2022). Secondly, preclinical and clinical evidence also supports the use of ginger in the downregulation of the “NF- κ B/MAPK” signaling pathway (Shoaib et al., 2020).

Besides its anti-inflammatory uses, ginger is quite relevant to metabolic health. Some research has shown that patients with type 2 diabetes can benefit from ginger by having higher insulin levels and lower blood glucose and cholesterol levels (Zhao et al., 2017) as shown in Figure 4. Conducting a clinical study on diabetic patients, daily ginger supplementation showed

a decrease in fasting blood glucose and HbA1c levels and thus may be useful in the parallel management of diabetes (Salaramoli et al., 2022). Further, the weight loss effect and the reduction of visceral fat by ginger have been studied and it was found that ginger could improve fat oxidation and thermogenesis (Liao et al., 2019).




Fig 4: Pharmacological Activities of Amla, Turmeric, and Ginger: Effects, Mechanisms, and Applications

SYNERGISTIC EFFECTS AND FORMULATIONS

Amla (*Emblica officinalis*), Turmeric (*Curcuma longa*), and Ginger (*Emblica officinalis*) are well-known botanicals in Ayurveda and Chinese treatment, where these botanicals are utilized together to improve health and fight diseases. These plants have been analyzed separately and as a whole in modern research, and the results have shown that the synergy of the two plants' pharmacological activities is increased.

Combination of Amla, Turmeric, and Ginger in Traditional Medicine

Amla, Turmeric, and Ginger are used in combination by Ayurveda to treat many diseases including digestive disorders and inflammation. Amla is one of the fruits that is rich in vitamin C and antioxidants and an anti-aging fruit. It is also a carminative, anti-inflammatory, and antiemetic herb. In addition, it works with other plant agents, for instance, turmeric, which has its own anti-inflammatory and anti-cancer properties (Kunnumakkara et al., 2023). Treated diseases (Zhou et al., 2022) such as general health and arthritic disorders, gastrointestinal disorders, and respiratory disorders. Ginger and turmeric are combined in traditional Chinese medicine for cold and dampness and Amla (Shahrajabian et al., 2019) is used for energy and immunity. They are used together more therapeutically than when used alone and are more effective in the management of health disorders.

Modern Formulations and Their Enhanced Pharmacological Effects

In contemporary scientific society, the combined effects of Amla (*Emblica officinalis*), Turmeric (*Curcuma longa*), and Ginger (*Emblica officinalis*) have been focused on better pharmacological activity. Poor bioavailability is the major challenge in creating these botanicals. Products like curcumin (turmeric), gingerols (ginger) and vitamin C (Amla) are rich in active compounds that have low bioavailability. The current formulations use nanoencapsulation and delivery systems of liposomes and adjuvants such as piperine, a black pepper alkaloid (Bertoncini et al., 2024). Taken together, these three plants have more therapeutic value than the other two plants combined. Kulsum et al. (2018) showed that Amla, Turmeric and Ginger formulations have higher anti-inflammatory and antioxidant activity than the single formulation of each. Patients with rheumatoid arthritis had decreased oxidative stress and inflammation, increased joint mobility, and decreased pain after treatment with this formulation. It probably is due to the modulation of different inflammation processes such as "NF-κB" and "MAPK" signaling (Kahkhaie et al., 2019). A preclinical study of Amla, Turmeric and Ginger formula by Fernando et al. (2023) showed that the formula reduced the size of the tumor and inhibited the growth of mice breast cancer cells. The effects of the botanicals on apoptosis, angiogenesis and metastatic characteristics are thought to be responsible for this enhanced anticancer activity. This combination has been used as a neuroprotectant in models of Alzheimer's disease. Amla, Turmeric, and Ginger helps improve brain performance and decrease the inflammation of the

mind. It has also been known to reduce amyloid beta plaque formation in cells and increase neurotrophic factors, including the BDNF 'protein' that is essential for neuron survival (Halder et al., 2021). It has also been established in clinical trials that it has a metabolic effect. Amla, Turmeric, and Ginger have been reported to reduce fasting blood glucose levels and first-phase insulin response in Type 2 diabetic patients. These effects may be attributed to botanicals that may possess insulin-regulating properties and anti-oxidant properties that decrease inflammation, and oxidative

embarrassment (Chavda et al., 2017). Furthermore, the formulation of Amla, Turmeric, and Ginger has been discovered to benefit the digestive system of the body Jafarzadeh et al. (2022) in a controlled randomized study showed that the combination enhances gastrointestinal motility and IBS symptoms. The formulation technologies like nanotechnology and liposomal encapsulation have enhanced this formulation and as such Amla, Turmeric, and Ginger are more effective in the treatment of various diseases due to their improved efficacy and bioavailability.

Table 2: Synergistic Effects and Modern Formulations of Amla, Turmeric, and Ginger

Combination	Traditional Use	Modern Formulation Benefits	References
Amla (<i>Emblica officinalis</i>)	Rich in vitamin C and antioxidants, anti-aging, digestive health, energy & immunity.	Enhances anti-inflammatory and antioxidant activity, improves joint flexibility, and decreases pain in rheumatoid arthritis.	Kunnumakkara et al., 2023; Zhou et al., 2022
Turmeric (<i>Curcuma longa</i>)	Anti-inflammatory, anticancer, used for cold and damp conditions.	Boosts anti-inflammatory, and anticancer properties, helps in cancer prevention, and reduces oxidative stress.	Kunnumakkara et al., 2023; Kahkhaie et al., 2019
Ginger (<i>Emblica officinalis</i>)	Anti-nausea, anti-inflammatory, and digestive properties.	Enhances neuroprotection, improves cognitive ability, and reduces neuroinflammation in Alzheimer's models.	Shahrajabian et al., 2019; Halder et al., 2021
Combination (Amla + Turmeric + Ginger)	Used for inflammation, digestive disorders, respiratory diseases, and energy.	Synergistic effect in anti-inflammatory, antioxidant, and anticancer activities. Improves metabolic health and digestive motility.	Bertoncini et al., 2024; Kulsum et al., 2018; Fernando et al., 2023; Chavda et al., 2017; Jafarzadeh et al., 2022

Table 2 shows that Amla, Turmeric, and Ginger when used individually or in combination, offer various therapeutic benefits. Amla, rich in vitamin C and antioxidants, enhances anti-inflammatory and antioxidant activity, benefiting joint flexibility. Turmeric's anti-inflammatory and anticancer properties help in cancer and reduce oxidative stress. Ginger, with its anti-nausea and digestive properties, also provides neuroprotection. The combination of these botanicals synergistically boosts anti-inflammatory, antioxidant, and anticancer activities, improving metabolic health and digestive motility, making it a potent therapeutic approach for multiple health conditions.

IN-SITU GELS AND THEIR APPLICATIONS IN ALLERGIC RHINITIS

Innovative drug delivery formulations, in situ gel systems, undergo a sol-to-gel transition when exposed to physiological conditions such as temperature, pH, or ionic concentration. They are particularly useful for the treatment of nasal diseases like allergic rhinitis since they provide sustained release of bioactive compounds at the site of action prolonged local effects and higher bioavailability. In-situ gels allow therapeutic compounds to be delivered directly to the nasal mucosa, where they can exert antioxidant, anti-inflammatory, and immune-modulatory effects.

Amla (*Emblica officinalis*):

Amla's antioxidant and anti-inflammatory properties were used to develop sodium alginate-based in situ gels incorporating Amla extract. The Amla's active compounds have increased bioavailability with these gels that adhere to the nasal mucosa for extended

periods. Amla has large quantities of vitamin C and polyphenols, that help to decrease oxidative stress very significantly in allergic rhinitis, as reaction to allergies is connected with higher oxidative stress (Saini et al., 2022). Amla helps stabilize mast cells, reduces histamine release, and prevents the generation of pro-inflammatory cytokines (TNF- α and IL-6). They help reduce the symptoms of sneezing, nasal congestion, and itching (Jantrapirom et al., 2021).

Turmeric (*Curcuma longa*):

Turmeric and its main ingredient curcumin are well known for their anti-inflammatory, antioxidant, and immune-modulatory qualities. Curcumin-loaded in situ gels, often with excipients such as chitosan and β cyclodextrin, enable the local delivery of curcumin in the nasal cavity (Garg et al., 2024). These gels lower inflammation by inhibiting important inflammatory pathways like NF- κ B and COX-2. Curcumin has antioxidant effects that protect nasal tissues from the damage done by allergens. The antimicrobial activity of turmeric is also beneficial in preventing secondary infections of the nasal mucosa beyond its anti-inflammatory effects (Wu et al., 2016). The in-situ gels loaded with turmeric can provide symptomatic relief in allergic rhinitis through both inflammatory and oxidative stress.

Ginger (*Emblica officinalis*):

Allergic rhinitis has been successfully treated with ginger — a known anti-inflammatory agent. Bioactive components of ginger namely gingerols and shogaols show effective action at reducing inflammation by blocking prostaglandin and leukotriene synthesis. In-

situ gels containing ginger extract or 6-gingerol have been formulated in hydroxypropyl methylcellulose (HPMC) based gels to extend the residence time of ginger in the nasal cavity. The gels reduce the symptoms of nasal discharge, sneezing, and itching by stabilizing mast cells and preventing histamine release. Furthermore, ginger's strong antioxidant properties also help heal inflamed mucosal tissues and help neutralize the oxidative damage done by allergens (Kawamoto et al., 2016).

MECHANISMS OF ACTION IN ALLERGIC RHINITIS

Amla, Turmeric, and Ginger have significant therapeutic benefits for allergic rhinitis due to their anti-inflammatory, antioxidant, and immunomodulatory properties. Amla inhibits oxidative stress and inflammation pathways and stabilizes mast cells preventing histamine release. Turmeric's antioxidant activity protects nasal tissues and inhibits COX2 and NF- κ B pathways to target inflammation. Instead, Ginger restricts proinflammatory mediators such as prostaglandins and leukotrienes and promotes mucosal healing.

Amla (*Emblica officinalis*)

Amla is a strong antioxidant fruit rich in vitamin C and polyphenolic compounds, and has great therapeutic potential. The mechanisms by which Amla provides relief in allergic rhinitis are multifactorial:

Antioxidant Activity: Amla is well known for its high vitamin C, polyphenols and flavonoids and its

significant antioxidant properties. These compounds may serve to neutralize reactive oxygen species (ROS) generated during allergic inflammation. Oxidative stress will increase inflammatory responses in nasal tissues. Amla scavenges ROS and reduces the damage of cellular in the nasal mucosa, thereby reducing the severity of allergic symptoms (Li et al., 2023).

Anti-Inflammatory Effects: Blocking main inflammatory signaling pathways has shown amla to have a significant anti-inflammatory effect. One of its main actions is to downregulate (NF- κ B), a key switch of the inflammatory response in allergic diseases. NF- κ B activation has been identified as linking to allergic rhinitis, and subsequent production of pro inflammatory cytokines that include TNF- α , IL-1 β , and IL-6 (Wang et al., 2020). In addition, Amla suppresses the mitogen-activated protein kinase (MAPK) signaling pathway and down-regulates inflammatory mediators (Nagaraju et al., 2023).

Immune Modulation: Immune modulation by Amla consists of suppression of Th2 type cytokines, which are essential for allergic immune response. Amla inhibits IL-4, IL-5, and IL-13 from releasing chemicals resulting in the activation of eosinophils and mast cells, the central managers of allergy in the nasopharyngeal passage. Amla also stabilizes mast cells and prevents histamine release, all of which help relieve symptoms as they occur such as nasal congestion, itching, and sneezing as shown in Figure 5 (Maheshwari et al., 2022).




Fig 5: Mechanism of action of Amla in allergic Rhinitis

Turmeric (*Curcuma longa*)

Turmeric contains curcumin, a bioactive molecule that is responsible for its therapeutic effects in inflammatory diseases including allergic rhinitis. Curcumin exerts its effects through the mechanisms listed below:

Anti-Inflammatory Pathways: Several critical pathways involved in inflammation are modulated by

curcumin. It (COX-2) and (LOX), are two enzymes that synthesize pro-inflammatory prostaglandins and leukotrienes. The swelling, congestion, and irritation of allergic rhinitis are due to these mediators. Furthermore, curcumin down-regulates NF- κ B, a central regulator of the inflammatory cascade induction (Chainoglou et al., 2019).

Antioxidant Properties: One of the most prominent activities of curcumin is its antioxidant activity. ROS scavenging activity of curcumin and its reduction of oxidative stress associated with the inflammatory process in allergic rhinitis (Boskabady et al., 2020). Curcumin may prevent oxidative damage to nasal epithelial cells and hence protect against additional tissue destruction, alleviating symptoms such as nasal blockage and hyperresponsiveness.

Immune Regulation: Curcumin balances the Th1/Th2 response of the immune system. It down-regulates the Th2 immune response, which is responsible for allergic inflammation. Curcumin inhibits the Th2 cytokines (IL-4 and IL-13) and helps in the reduction of eosinophils and mast cell activation as shown in Figure 6. It reduces the release of histamines and other mediators of allergic rhinitis (Haftcheshmeh et al., 2022).




Fig 6: Mechanism of action of Turmeric in allergic Rhinitis

Ginger (*Emblca officinalis*)

Ginger is well-known for its antioxidant and anti-inflammatory effects. Its active compounds, gingerols, and shogaols, exert powerful effects on the molecular mechanisms implicated in allergic rhinitis:

Inhibition of Pro-Inflammatory Mediators: Gingerols and shogaols inhibit the activities of COX and LOX enzymes that lead to the formation of inflammatory mediators, e.g. prostaglandins and leukotrienes. Allergic rhinitis is characterized by vasodilation, edema, and mucus production, and these mediators contribute to these events. Ginger acts by blocking these pathways reduces inflammation and clears nasal congestion and discharge (Rudrapal et al., 2023).

Mast Cell Stabilization: One of the important components of allergic rhinitis is mast cell activation, which causes the release of histamine and other inflammatory mediators. Ginger has been proven to stabilize mast cells, preventing the release of histamine and thus relieving symptoms such as itching, sneezing, and congestion (Gupta et al., 2023). In particular, this action is important in reducing hypersensitivity of the nasal mucosa to allergens.

Antioxidant Support: The bioactive compounds of ginger, and gingerols have antioxidant properties scavenge ROS, and protect tissues from oxidative damage in nasal tissues. This antioxidant action is critical for lowering inflammation and stimulating the repair of injured epithelial cells in the nasal mucosa,

resulting in a reduction in symptoms such as nasal irritation and post-nasal drip as shown in Figure 7 (Alolga et al., 2022).




Fig 7: Mechanism of action of Ginger in allergic Rhinitis

CHALLENGES AND FUTURE DIRECTIONS

Many therapeutic uses of Amla (*Emblica officinalis*), Turmeric (*Curcuma longa*), and Ginger (*Emblica officinalis*) are associated with some difficulties, especially regarding standardization, quality control, and research. Therefore, while these botanicals demonstrate great pharmacological promise, various challenges limit their use in clinical medicine and inclusion in contemporary systems of healthcare delivery. A major limitation in the use of these botanicals is the variation in phytochemical composition that is not well defined. The levels of active constituents, like curcumin in turmeric, gingerols in ginger, and vitamin C in Amla, may also differ due to the geographical location of the plant, and the way it was grown and processed. Such fluctuation complicates the reliability and stability of quality in subsequent batches of goods and services. Moreover, due to the lack of internationally acceptable standards for quality control of herbal products, there are variations in the strength and purity of the products. Adulteration and contamination with heavy metals, pesticides, and microbial agents add another dimension to the quality control issue. These issues are dangerous to consumers and lower the reliability of botanical products. HPLC, mass spectral analysis, and GC/MS are all other analytical methods employed for accurate measurements and identification of phytochemicals. However, the costs and technical skills needed for these approaches make them impractical in most developing countries and other low-resource environments. Although the curative properties of Amla, Turmeric, and Ginger have been established, much is still

unknown about how they work and what the long-term consequences of their use are. For example, the bioavailability of active phytochemicals including curcumin and gingerols is still a problem. Thus, although there are improvements in delivery systems, more studies are required to develop formulations that improve solubility and stability in the gastrointestinal tract. Moreover, most of the previous researches are based on single compounds instead of the interaction of whole extracts or the mix of these plants. The ability to understand how the mixture of phytochemicals in the plant interacts with biological systems may uncover new uses and potential treatments. In addition, most related studies were conducted on animal models and explants, and few randomized clinical trials to prove effectiveness and safety in human beings. Few longitudinal research in understanding the impacts of these plant products on the diseases that develop over an extended period; hence diabetes, cancer, and neurodegenerative diseases.

Nevertheless, the specific pharmacological characteristics of Amla, Turmeric, and Ginger present a vast opportunity for the synthesis of new therapeutic compounds. The development of nanotechnology, bioinformatics, and system biology offers possibilities to develop smart drug delivery systems and to discover novel bioactive molecules. For instance, curcumin nanoparticles have exhibited higher solubility and anti-cancer and anti-inflammatory effects which signify their future potential in cancer and inflammatory disease treatment. Another opportunity is the incorporation of these botanicals into multi-target approaches. The combined action of their phytochemicals could be

exploited to manage poly-etiological diseases like metabolic syndrome and neurodegenerative diseases where single molecular targeted drugs are ineffective. Also, the increasing focus on the concept of pharmacogenomics creates new opportunities for developing individualized botanical drugs that will improve therapeutic efficacy.

CONCLUSION

The review also emphasizes the great curative prospect of Amla (*Embllica officinalis*), Turmeric (*Curcuma longa*), and Ginger (*Embllica officinalis*) as sources of bioactive compounds. These botanicals display diverse pharmacological effects such as antioxidant, anti-inflammatory, anticancer, neuroprotective, and improved metabolic profile. Their high phytochemical content—vitamin C, curcuminoids, gingerols—which serve as the primary pharmacological agents give both products their therapeutic value. Furthermore, the interaction between these botanicals when used together improves their pharmacology, which presents potential therapeutic and preventative approaches to chronic diseases. The study also highlights the numerous problems that are related to the process of standardization and quality assurance of these botanicals. The variations in phytochemicals, poor absorption, and lack of standard quality assurance measures are all included in these. Nevertheless, formulation development trends, such as nanoencapsulation and liposomal delivery systems, have demonstrated the potential to overcome these challenges. Other bio-enhancers like Piperine enhance the bioavailability and effectiveness of these ingredients when combined with these natural compounds. The findings of this review are extremely important to the pharmaceutical and nutraceutical businesses. Better quality formulations of Amla, Turmeric, and Ginger could fill the increasing gap of natural plant-based remedies and the need to standardize better quality formulations of Amla, Turmeric, and Ginger could be met. They present an option for developing polypharmacological agents for complicated diseases including cancer, diabetes, and neurodegenerative diseases. Besides, their incorporation into functional foods and dietary supplements can meet the growing demand for health and nutritional prevention. In conclusion, Amla, Turmeric, and Ginger represent a robust medicine platform—however, to advance medicine to patient care potential the existing science gaps must be addressed, formulations improved, and efficacy tested in systematic randomized clinical trials. These efforts will set the stage for the expansion of these botanicals into other forms of medicine and the inclusion of these botanicals into standard protocols of treatment.

REFERENCES

1. Abd El-Hack, M. E., El-Saadony, M. T., Swelum, A. A., Arif, M., Abo Ghanima, M. M., Shukry, M., ... & El-Tarabily, K. A. (2021). Curcumin, the active substance of turmeric: its effects on health and ways to improve its bioavailability. *Journal of the Science of Food and Agriculture*, 101(14), 5747-5762.
2. Alhazmi, H. A., Najmi, A., Javed, S. A., Sultana, S., Al Bratty, M., Makeen, H. A., Meraya, A. M., Ahsan, W., Mohan, S., Taha, M. M. E., & Khalid, A. (2021). Medicinal Plants and Isolated Molecules Demonstrating Immunomodulation Activity as Potential Alternative Therapies for Viral Diseases Including COVID-19. *Frontiers in immunology*, 12, 637553. <https://doi.org/10.3389/fimmu.2021.637553>
3. Holga, R. N., Wang, F., Zhang, X., Li, J., Tran, L. S. P., & Yin, X. (2022). Bioactive compounds from the Zingiberaceae Family with known antioxidant activities for possible therapeutic uses. *Antioxidants*, 11(7), 1281.
4. Ameer, S. F., Mohamed, M. Y., Elzubair, Q. A., Sharif, E. A. M., & Ibrahim, W. N. (2024). Curcumin as a novel therapeutic candidate for cancer: can this natural compound revolutionize cancer treatment? *Frontiers in oncology*, 14, 1438040. <https://doi.org/10.3389/fonc.2024.1438040>
5. Ballester, P., Cerdá, B., Arcusa, R., Marhuenda, J., Yamedjeu, K., & Zafrilla, P. (2022). Effect of Ginger on Inflammatory Diseases. *Molecules (Basel, Switzerland)*, 27(21), 7223. <https://doi.org/10.3390/molecules27217223>
6. Bertocini-Silva, C., Vlad, A., Ricciarelli, R., Giacomo Fassini, P., Suen, V. M. M., & Zingg, J. M. (2024). Enhancing the Bioavailability and Bioactivity of Curcumin for Disease Prevention and Treatment. *Antioxidants (Basel, Switzerland)*, 13(3), 331. <https://doi.org/10.3390/antiox13030331>
7. Boskabady, M. H., Shakeri, F., & Naghdi, F. (2020). The effects of *Curcuma longa* L. and its constituents in respiratory disorders and molecular mechanisms of their action. *Studies in Natural Products Chemistry*, 65, 239-269.
8. Cacciola, N. A., Cuciniello, R., Petillo, G. D., Piccioni, M., Filosa, S., & Crispi, S. (2023). An Overview of the Enhanced Effects of Curcumin and Chemotherapeutic Agents in Combined Cancer Treatments. *International journal of molecular sciences*, 24(16), 12587. <https://doi.org/10.3390/ijms241612587>
9. Chainoglou, E., & Hadjipavlou-Litina, D. (2019). Curcumin analogs and derivatives with anti-proliferative and anti-inflammatory activity: Structural characteristics and molecular targets. *Expert opinion on drug discovery*, 14(8), 821-842.
10. Chavda, Bhumika & Sharma, Anil. (2017). Efficacy of Combination of Fenugreek, Amla and Turmeric Powder to Reduce Blood Glucose Level among Diabetics-Literature Review. *International Journal of Nursing Care*. 5. 55. 10.5958/2320-8651.2017.00012.6.
11. Crichton, M., Marshall, S., Isenring, E., Lohning, A., McCarthy, A. L., Molassiotis, A., Bird, R., Shannon, C., Koh, A., McPherson, I., & Marx, W. (2024). Effect of a Standardized Ginger Root Powder Regimen on Chemotherapy-Induced Nausea and Vomiting: A Multicenter, Double-Blind, Placebo-controlled Randomized Trial.

- Journal of the Academy of Nutrition and Dietetics, 124(3), 313–330.e6. <https://doi.org/10.1016/j.jand.2023.09.003>
12. Deshmukh, Chinmay & Choudhari, Shradha. (2021). Phytochemical and Pharmacological Profile of *Emblca Officinalis* Linn, Journal of Medical pharmaceutical and allied sciences. 10. 10.22270/jmpas.V10I2.1054.
13. Dosoky, N. S., & Setzer, W. N. (2018). Chemical Composition and Biological Activities of Essential Oils of *Curcuma* Species. *Nutrients*, 10(9), 1196. <https://doi.org/10.3390/nu10091196>
14. Fitriyanti, D., & Sulung, R. (2020). Effectiveness of ginger to overcome nausea and vomiting caused by chemotherapy in breast cancer patients. *Canadian oncology nursing journal = Revue canadienne de nursing oncologique*, 30(1), 3–5. <https://doi.org/10.5737/2368807630135>
15. Fuloria, S., Mehta, J., Chandel, A., Sekar, M., Rani, N. N. I. M., Begum, M. Y., Subramanian, V., Chidambaram, K., Thangavelu, L., Nordin, R., Wu, Y. S., Sathasivam, K. V., Lum, P. T., Meenakshi, D. U., Kumarasamy, V., Azad, A. K., & Fuloria, N. K. (2022). A Comprehensive Review on the Therapeutic Potential of *Curcuma longa* Linn. about its Major Active Constituent Curcumin. *Frontiers in pharmacology*, 13, 820806. <https://doi.org/10.3389/fphar.2022.820806>
16. G Singh, P., S Jain, A., B Sridhara Setty, P., Bv, S., S Patil, S., P, A., Ts, G., Suresh, K. P., Dugganaboyana, G. K., Murugesan, K., Gnanasekaran, A., Shivamallu, C., Kollur, S. P., Srinivasa, C., Hl, R., Rudrapathy, P., & M Basalingappa, K. (2022). Antimicrobial, antioxidant, and anti-inflammatory activities of seeds from *emblca officinalis* (Gaertn.). *Bioinformation*, 18(8), 683–691. <https://doi.org/10.6026/97320630018683>
17. Gandhi, Yashika & Grewal, Jyotika & Jain, Vipin & Rawat, Hemant & Kumar, Vijay & Kumar, Ravi & Shakya, Santosh & Sharma, Preeti & Dhanjal, Daljeet & Prasad, Dr. Shyam & Charde, Vaibhav & Arya, Jagdish Chandra & Narasimhaji, Venkata & Singh, Arjun & Singh, Ravindra & Srikanth, Naryanam & Acharya, Rabinarayan. (2023). *Emblca officinalis*: A promising herb confining versatile applications. *South African Journal of Botany*. 159. 519-531. 10.1016/j.sajb.2023.06.041.
18. Garg, A., Agrawal, R., Chauhan, C. S., & Deshmukh, R. (2024). In-situ gel: a smart carrier for drug delivery. *International Journal of Pharmaceutics*, 123819.
19. Gęgotek, A., & Skrzydlewska, E. (2022). Antioxidative and Anti-Inflammatory Activity of Ascorbic Acid. *Antioxidants* (Basel, Switzerland), 11(10), 1993. <https://doi.org/10.3390/antiox11101993>
20. Genchi, G., Lauria, G., Catalano, A., Carocci, A., & Sinicropi, M. S. (2024). Neuroprotective Effects of Curcumin in Neurodegenerative Diseases. *Foods* (Basel, Switzerland), 13(11), 1774. <https://doi.org/10.3390/foods13111774>
21. Giordano, A., & Tommonaro, G. (2019). Curcumin and Cancer. *Nutrients*, 11(10), 2376. <https://doi.org/10.3390/nu11102376>
22. Gul, M., Liu, Z. W., Iahitsham-Ul-Haq, Rabail, R., Faheem, F., Walayat, N., Nawaz, A., Shabbir, M. A., Muneekata, P. E. S., Lorenzo, J. M., & Aadil, R. M. (2022). Functional and Nutraceutical Significance of Amla (*Phyllanthus emblica* L.): A Review. *Antioxidants* (Basel, Switzerland), 11(5), 816. <https://doi.org/10.3390/antiox11050816>
23. Gupta, R., Aggarwal, A., Upreti, K., Rawat, S., & Tiwari, B. (2023). Efficacy of phytotherapeutics in allergic rhinitis: A review. *Indian Journal of Natural Products and Resources (IJNPR)[Formerly Natural Product Radiance (NPR)]*, 13(4), 451-459.
24. Haftcheshmeh, S. M., Mirhafez, S. R., Abedi, M., Heydarlou, H., Shakeri, A., Mohammadi, A., & Sahebkar, A. (2022). Therapeutic potency of curcumin for allergic diseases: A focus on immunomodulatory actions. *Biomedicine & Pharmacotherapy*, 154, 113646.
25. Halder, S., Anand, U., Nandy, S., Oleksak, P., Qusti, S., Alshammari, E. M., El-Saber Batiha, G., Koshy, E. P., & Dey, A. (2021). Herbal drugs and natural bioactive products as potential therapeutics: A review on pro-cognitive and brain boosters perspectives. *Saudi Pharmaceutical Journal: SPJ: the official publication of the Saudi Pharmaceutical Society*, 29(8), 879–907. <https://doi.org/10.1016/j.jsps.2021.07.003>
26. Hassan, Syeda & Mughal, Asma & Farman, Muhammad. (2020). *Emblca Officinalis* (Amla): A Prospective Review On Distinctive Properties And Therapeutic Applications Of Amla. 22-30. 10.7537/marsbnj060220.03.
27. Hewlings, S. J., & Kalman, D. S. (2017). Curcumin: A review of its effects on human health. *Foods*, 6(10), 92. <https://doi.org/10.3390/foods6100092>
28. Hussain, Y., Alam, W., Ullah, H., Dacrema, M., Daglia, M., Khan, H., & Arciola, C. R. (2022). Antimicrobial Potential of Curcumin: Therapeutic Potential and Challenges to Clinical Applications. *Antibiotics* (Basel, Switzerland), 11(3), 322. <https://doi.org/10.3390/antibiotics11030322>
29. Ishfaq, M. & Hu, Wanying & Hu, Zhihua & Guan, Yurong & Zhang, Ruihong. (2022). A review of nutritional implications of bioactive compounds of Ginger (*Emblca officinalis* Roscoe), their biological activities, and nano-formulations. *Italian Journal of Food Science*. 34. 1-12. 10.15586/ijfs.v34i3.2212.
30. Jafarzadeh, E., Shoeibi, S., Bahramvand, Y., Nasrollahi, E., Maghsoudi, A. S., Yazdi, F., KarkonShayan, S., & Hassani, S. (2022). Turmeric for Treatment of Irritable Bowel Syndrome: A Systematic Review of Population-Based Evidence. *Iranian journal of public health*, 51(6), 1223–1231. <https://doi.org/10.18502/ijph.v51i6.9656>
31. Jain, Pushpendra & Das, Debajyoti & Pandey, Nalini & Jain, Prachi. (2016). TRADITIONAL INDIAN HERB *EMBLCA OFFICINALIS* AND ITS MEDICINAL IMPORTANCE. *International Journal of Pharmacy and Pharmaceutical Sciences*. 4. 1-15. .

32. Jantrapirom, S., Hirunsatitpron, P., Potikanond, S., Nimlamool, W., & Hanprasertpong, N. (2021). Pharmacological Benefits of Triphala: A Perspective for Allergic Rhinitis. *Frontiers in pharmacology*, 12, 628198. <https://doi.org/10.3389/fphar.2021.628198>
33. KAB., Fernando & P., Perera. (2023). Review on Ayurveda and Modern aspects of Spices for Cancer Prevention. *International Journal of Scientific and Research Publications*. 13. 288-296. 10.29322/IJSRP.13.08 2023.p14039.
34. Kahkhaie, K. R., Mirhosseini, A., Aliabadi, A., Mohammadi, A., Mousavi, M. J., Haftcheshmeh, S. M., Sathyapalan, T., & Sahebkar, A. (2019). Curcumin: a modulator of inflammatory signaling pathways in the immune system. *Inflammopharmacology*, 27(5), 885–900. <https://doi.org/10.1007/s10787-019-00607-3>
35. Kapoor, M. P., Suzuki, K., Derek, T., Ozeki, M., & Okubo, T. (2019). Clinical evaluation of *Emblia officinalis* Gaertn (Amla) in healthy human subjects: Health benefits and safety results from a randomized, double-blind, crossover placebo-controlled study. *Contemporary clinical trials communications*, 17, 100499. <https://doi.org/10.1016/j.conctc.2019.100499>
36. Kawamoto, Y., Ueno, Y., Nakahashi, E., Obayashi, M., Sugihara, K., Qiao, S., Iida, M., Kumasaka, M. Y., Yajima, I., Goto, Y., Ohgami, N., Kato, M., & Takeda, K. (2016). Prevention of allergic rhinitis by ginger and the molecular basis of immunosuppression by 6-gingerol through T cell inactivation. *The Journal of Nutritional Biochemistry*, 27, 112–122. <https://doi.org/10.1016/j.jnutbio.2015.08.025>
37. Kulsum, Safeena & Suresh, Amritha & Mehta, Alka. (2018). Correlation of antioxidant and antiproliferative activity of amla and ginger. *Asian Journal of Pharmaceutical and Clinical Research*. 11. 263. 10.22159/ajpcr 2018.v11i8.26073.
38. Kumar, A., P, N., Kumar, M., Jose, A., Tomer, V., Oz, E., Proestos, C., Zeng, M., Elobeid, T., K, S., & Oz, F. (2023). Major Phytochemicals: Recent Advances in Health Benefits and Extraction Method. *Molecules* (Basel, Switzerland), 28(2), 887. <https://doi.org/10.3390/molecules28020887>
39. Kunnumakkara, A. B., Hegde, M., Parama, D., Girisai, S., Kumar, A., Daimary, U. D., Garodia, P., Yeniseti, S. C., Oommen, O. V., & Aggarwal, B. B. (2023). Role of Turmeric and Curcumin in Prevention and Treatment of Chronic Diseases: Lessons Learned from Clinical Trials. *ACS pharmacology & translational science*, 6(4), 447–518. <https://doi.org/10.1021/acsptsci.2c00012>
40. Lechner, J. F., & Stoner, G. D. (2019). Gingers and Their Purified Components as Cancer Chemopreventative Agents. *Molecules* (Basel, Switzerland), 24(16), 2859. <https://doi.org/10.3390/molecules24162859>
41. Lete, I., & Allué, J. (2016). The Effectiveness of Ginger in the Prevention of Nausea and Vomiting during Pregnancy and Chemotherapy. *Integrative medicine insights*, 11, 11–17. <https://doi.org/10.4137/IMI.S36273>
42. Li, G., Yu, Q., Li, M., Zhang, D., Yu, J., Yu, X., ... & Huang, H. (2023). *Phyllanthus emblica* fruits: A polyphenol-rich fruit with potential benefits for oral management. *Food & Function*, 14(17), 7738–7759.
43. Maheshwari, S., Kumar, V., Bhadauria, G., & Mishra, A. (2022). Immunomodulatory potential of phytochemicals and other bioactive compounds of fruits: A review. *Food Frontiers*, 3(2), 221–238.
44. Mahomoodally, M. F., Aumeeruddy, M. Z., Rengasamy, K. R. R., Roshan, S., Hammad, S., Pandohee, J., Hu, X., & Zengin, G. (2021). Ginger and its active compounds in cancer therapy: From folk use to nano-therapeutic applications. *Seminars in cancer biology*, 69, 140–149. <https://doi.org/10.1016/j.semcancer.2019.08.009>
45. Malik, S., Suchal, K., Bhatia, J., Khan, S. I., Vasisth, S., Tomar, A., Goyal, S., Kumar, R., Arya, D. S., & Ojha, S. K. (2016). Therapeutic Potential and Molecular Mechanisms of *Emblia officinalis* Gaertn in Countering Nephrotoxicity in Rats Induced by the Chemotherapeutic Agent Cisplatin. *Frontiers in pharmacology*, 7, 350. <https://doi.org/10.3389/fphar.2016.00350>
46. Manasa, P. & Kamble, Alka & Chilakamarthi, Ushasri. (2023). Various Extraction Techniques of Curcumin—A Comprehensive Review. *ACS Omega*. 8. 10.1021/acsomega.3c04205.
47. Mao, Q. Q., Xu, X. Y., Cao, S. Y., Gan, R. Y., Corke, H., Beta, T., & Li, H. B. (2019). Bioactive Compounds and Bioactivities of Ginger (*Emblia officinalis* Roscoe). *Foods* (Basel, Switzerland), 8(6), 185. <https://doi.org/10.3390/foods8060185>
48. Marfori, Eufrocínio. (2016). A Simple HPLC Method for the Analysis of [6]-Gingerol Produced by Multiple Shoot Culture of Ginger (*Emblia officinalis*). 8.
49. Mohaddese, Mahboubi. (2019). *Emblia officinalis* Rosc. essential oil, a review of its composition and bioactivity. *Clinical Phytoscience*. 5. 10.1186/s40816-018-0097-4.
50. Nagaraju, M., Kalahasti, K. K., Reddy, K. P., Addi, U. R., Satyavani, M., Reddy, G. B., & Reddy, S. S. (2023). Anti-inflammatory potential of turmeric, amla, and black pepper mixture against sepsis-induced acute lung injury in rats. *Journal of Food Science and Technology*, 60(1), 252–261.
51. Nikkhah Bodagh, M., Maleki, I., & Hekmatdoost, A. (2018). Ginger in gastrointestinal disorders: A systematic review of clinical trials. *Food science & nutrition*, 7(1), 96–108. <https://doi.org/10.1002/fsn3.807>
52. Pathak, S., Pratap, A., Sharma, R., & Jha, M. K. (2024). The *Phyllanthus emblica* Fruits: A Review on Phytochemistry Traditional Uses, Bioactive Composition and Pharmacological Activities. *Current topics in medicinal chemistry*, 24(22), 1917–1939. <https://doi.org/10.2174/0115680266321320240708060327>
53. Peng, Y., Ao, M., Dong, B., Jiang, Y., Yu, L., Chen, Z., Hu, C., & Xu, R. (2021). Anti-Inflammatory

- Effects of Curcumin in the Inflammatory Diseases: Status, Limitations and Countermeasures. *Drug design, development, and therapy*, 15, 4503–4525. <https://doi.org/10.2147/DDDT.S327378>
54. Pourhabibi-Zarandi, F., Shojaei-Zarghani, S., & Rafrat, M. (2021). Curcumin and rheumatoid arthritis: A systematic review of the literature. *International journal of clinical practice*, 75(10), e14280. <https://doi.org/10.1111/ijcp.14280>
 55. Prananda, A. T., Dalimunthe, A., Harahap, U., Simanjuntak, Y., Peronika, E., Karosekali, N. E., Hasibuan, P. A. Z., Syahputra, R. A., Situmorang, P. C., & Nurkolis, F. (2023). *Phyllanthus emblica*: a comprehensive review of its phytochemical composition and pharmacological properties. *Frontiers in pharmacology*, 14, 1288618. <https://doi.org/10.3389/fphar.2023.1288618>
 56. Promdam, N., & Panichayupakaranant, P. (2022). [6]-Gingerol: a narrative review of its beneficial effect on human health. *Food Chemistry Advances*, 1, 100043.
 57. Reddy, P. H., Manczak, M., Yin, X., Grady, M. C., Mitchell, A., Tonk, S., Kuruva, C. S., Bhatti, J. S., Kandimalla, R., Vijayan, M., Kumar, S., Wang, R., Pradeepkiran, J. A., Ogunmokun, G., Thamarai, K., Quesada, K., Boles, A., & Reddy, A. P. (2018). Protective Effects of Indian Spice Curcumin Against Amyloid- β in Alzheimer's Disease. *Journal of Alzheimer's disease: JAD*, 61(3), 843–866. <https://doi.org/10.3233/JAD-170512>
 58. Rudrapal, M., Eltayeb, W. A., Rakshit, G., El-Arabey, A. A., Khan, J., Aldosari, S. M., ... & Abdalla, M. (2023). Dual synergistic inhibition of COX and LOX by potential chemicals from Indian daily spices investigated through detailed computational studies. *Scientific Reports*, 13(1), 8656.
 59. Saini, R., & Dhiman, N. K. (2022). Natural anti-inflammatory and anti-allergy agents: herbs and botanical ingredients. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry* (Current Medicinal Chemistry-Anti-Inflammatory and Anti-Allergy Agents), 21(2), 90-114.
 60. Salaramoli, S., Mehri, S., Yarmohammadi, F., Hashemy, S. I., & Hosseinzadeh, H. (2022). The effects of ginger and its constituents in the prevention of metabolic syndrome: A review. *Iranian journal of basic medical sciences*, 25(6), 664–674. <https://doi.org/10.22038/IJBMS.2022.59627.13231>
 61. Setyaningsih, D., Santoso, Y. A., Hartini, Y. S., Murti, Y. B., Hinrichs, W. L. J., & Patramurti, C. (2021). Isocratic high-performance liquid chromatography (HPLC) for simultaneous quantification of curcumin and piperine in a microparticle formulation containing *Curcuma longa* and *Piper nigrum*. *Heliyon*, 7(3), e06541. <https://doi.org/10.1016/j.heliyon.2021.e06541>
 62. Shahrajabian, M., Hesam & Sun, Wenli & Cheng, Qi. (2019). Clinical aspects and health benefits of ginger (*Emblica officinalis*) in both traditional Chinese medicine and modern industry. *Acta Agriculturae Scandinavica, Section B - Soil & Plant Science*. 69. 546-556. 10.1080/09064710.2019.1606930.
 63. Shanmugam, M. K., Rane, G., Kanchi, M. M., Arfuso, F., Chinnathambi, A., Zayed, M. E., Alharbi, S. A., Tan, B. K., Kumar, A. P., & Sethi, G. (2015). The multifaceted role of curcumin in cancer prevention and treatment. *Molecules (Basel, Switzerland)*, 20(2), 2728–2769. <https://doi.org/10.3390/molecules20022728>
 64. Sharma, S., Shukla, M. K., Sharma, K. C., Tirath, Kumar, L., Anal, J. M. H., Upadhyay, S. K., Bhattacharyya, S., & Kumar, D. (2023). Revisiting the therapeutic potential of gingerols against different pharmacological activities. *Naunyn-Schmiedeberg's archives of pharmacology*, 396(4), 633–647. <https://doi.org/10.1007/s00210-022-02372-7>
 65. Silvestro, S., Sindona, C., Bramanti, P., & Mazzon, E. (2021). A State of the Art of Antioxidant Properties of Curcuminoids in Neurodegenerative Diseases. *International journal of molecular sciences*, 22(6), 3168. <https://doi.org/10.3390/ijms22063168>
 66. Talreja, Shreya & Kumari, Sonam & Srivastava, Prateek & Pandey, Swarnima. (2020). A COMPLETE PHARMACOGNOSTIC REVIEW ON AMLA. 622-637 10.20959/wjpps20194-13486.
 67. Tomeh, M. A., Hadianamrei, R., & Zhao, X. (2019). A Review of Curcumin and Its Derivatives as Anticancer Agents. *International journal of molecular sciences*, 20(5), 1033. <https://doi.org/10.3390/ijms20051033>
 68. Ullah, A., Munir, S., Badshah, S. L., Khan, N., Ghani, L., Poulson, B. G., Emwas, A. H., & Jaremko, M. (2020). Important Flavonoids and Their Role as a Therapeutic Agent. *Molecules (Basel, Switzerland)*, 25(22), 5243. <https://doi.org/10.3390/molecules25225243>
 69. Variya, B. C., Bakrania, A. K., & Patel, S. S. (2016). *Emblica Officinalis* (Amla): A review of its phytochemistry, ethnomedicinal uses, and medicinal potentials concerning molecular mechanisms. *Pharmacological Research*, 111, 180–200. <https://doi.org/10.1016/j.phrs.2016.06.013>
 70. Voulgaropoulou, S. D., van Amelsvoort, T. A. M. J., Prickaerts, J., & Vingerhoets, C. (2019). The effect of curcumin on cognition in Alzheimer's disease and healthy aging: A systematic review of pre-clinical and clinical studies. *Brain Research*, 1725, 146476. <https://doi.org/10.1016/j.brainres.2019.146476>
 71. Wakchaure, Rajesh & Ganguly, Subha. (2018). Phytochemistry and Pharmacological Properties of Ginger (*Emblica officinalis*).
 72. Walayat, R.N & Rabail, Roshina. (2022). Functional and Nutraceutical Significance of Amla (*Phyllanthus emblica* L.): A Review. *Antioxidants*. 11 10.3390/antiox11050816.
 73. Wang, Y., Cui, C., & Sun, H. (2020). Anti-inflammatory effect of mangiferin on an experimental model of allergic rhinitis through the inhibition of nf-kb signaling pathways. *Journal of Environmental Pathology, Toxicology and Oncology*, 39(4).

74. Widmann, A. K., Wahl, M. A., Kammerer, D. R., & Daniels, R. (2022). Supercritical Fluid Extraction with CO₂ of *Curcuma longa* L. in Comparison to Conventional Solvent Extraction. *Pharmaceutics*, 14(9), 1943. <https://doi.org/10.3390/pharmaceutics14091943>
75. Wu S, Xiao D. Effect of curcumin on nasal symptoms and airflow in patients with perennial allergic rhinitis. *Ann Allergy Asthma Immunol.* 2016 Dec;117(6):697-702.e1. doi: 10.1016/j.anai.2016.09.427. Epub 2016 Oct 24. PMID: 27789120.
76. Zhou, X., Afzal, S., Wohlmuth, H., Münch, G., Leach, D., Low, M., & Li, C. G. (2022). Synergistic Anti-Inflammatory Activity of Ginger and Turmeric Extracts in Inhibiting Lipopolysaccharide and Interferon- γ -Induced Proinflammatory Mediators. *Molecules* (Basel, Switzerland), 27(12), 3877. <https://doi.org/10.3390/molecules27123877>