Research Square

Preprints are preliminary reports that have not undergone peer review.
They should not be considered conclusive, used to inform clinical practice,
or referenced by the media as validated information.

Initial Silvicultural Performance of Pilocarpus microphyllus in Agroforestry systems in the Amazon

Érick Martins Nieri

ericknieri@utfpr.edu.br

Federal University of Technology – Paraná

Camila Pereira Silva

Universidade Federal do Sul e Sudeste do Pará

Luana Maria dos Santos

Federal University of Technology – Paraná

Eleandro José Brun

Federal University of Technology – Paraná

Research Article

Keywords: pilocarpine, silvicultural treatments, jaborandi

Posted Date: April 6th, 2026

DOI: https://doi.org/10.21203/rs.3.rs-9235597/v1

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
Read Full License

Additional Declarations: No competing interests reported.

Initial Silvicultural Performance of *Pilocarpus microphyllus* in Agroforestry systems in the Amazon

Erick Martins Nieri[1]*, Camila Pereira Silva[2], Luana Maria dos Santos[1], Eleandro José Brun[1]

[1] Federal University of Technology – Paraná (UTFPR), Dois Vizinhos, PR, Brazil
[2] Forest Engineer, Federal University of South and Southeast of Pará (UNIFESSPA), São Félix do Xingu, PA, Brazil

*Corresponding author: Erick Martins Nieri
Email: ericknieri@utfpr.edu.br

Abstract

The Amazon region faces significant pressure from deforestation and cattle ranching, threatening the natural habitat of *Pilocarpus microphyllus*. This study evaluated the initial silvicultural performance of *P. microphyllus* under different treatments in agroforestry systems (AFS) in São Félix do Xingu, Pará, Brazil. The experiment followed a randomized complete block design in a 2 \times 6 factorial scheme with three replicates. Variables measured included collar diameter, seedling height, crown area, number of shoots, and mean branch length. Results indicated that the AFS presented distinct light regimes (17,445.5; 3,787.5; and 2,954.5 Lux). Significant differences were observed among systems, with AFS 1 (intermediate light) showing the best growth responses for all variables. A significant interaction between pruning and fertilization was found only for seedling height. In the absence of pruning, NPK doses of 175 g and 75 g produced the highest growth. Conversely, when pruning was applied, no significant differences were found among fertilization treatments, although the combination of 45 g NPK + 45 g MAP showed the best performance at 20 months. Pruning significantly enhanced the number of shoots and mean branch length from the 19 th month onward. Higher NPK doses were effective in increasing crown area. We recommend planting *P. microphyllus* in AFS with moderate solar radiation, combined with formative pruning and targeted fertilization (175 g NPK without pruning; 45 g NPK + 45 g MAP with pruning).
Keywords: pilocarpine; silvicultural treatments; jaborandi

Introduction

The jaborandi, *Pilocarpus microphyllus*, has been economically exploited since the 1990s in the Alto Xingu region, with extractivists collecting its leaves for commercialization (Amaral et al., 2022). However, increasing demand together with deforestation activities have rendered the species vulnerable, which highlights the potential role of *P. microphyllus* in agroforestry systems (AFS) as an alternative for species conservation and for understanding its performance under cultivation (Ministério do Meio Ambiente, 2022).

The species has economic potential because it is robust, naturally occurs in the region, and is the sole natural source of pilocarpine. Caldeira et al. (2017) demonstrated that the species can develop in its native occurrence areas and can provide a relatively rapid economic return through leaf harvesting; leaves can be managed from 50 cm in height to promote crown formation.

Leaf collection may be carried out by extractives or by harvesting from cultivated stands. Nevertheless, limits to supply capacity, the onset of extraction decline, depletion of reserves, and rising demand characterize the need to initiate cultivation—an issue already evident in the Amazon region (Homma, 2014).

Unmanaged leaf extraction underscores the need to understand the behavior and performance of *P. microphyllus* when integrated into alternative production systems, with emphasis on its productive potential and pilocarpine content. Differences in pilocarpine concentration have been observed; Gumier-Costa (2016) reported concentrations of 1.0% in native areas versus 0.5% in monocultures. Early cultivation efforts were undertaken by Merck in monoculture plantations in Maranhão (Homma, 2012). However, studies on the species in other cultivation systems, such as agroforestry systems, remain scarce.

Plant growth and development depend on factors such as light, water, and carbon dioxide, as well as essential chemical elements—commonly referred to as nutrients—that are integral to plant physiology. The primary macronutrients include nitrogen (N), phosphorus (P), and potassium (K) (Teixeira et al., 2019; Nieri et al., 2019). These nutrients are fundamental for healthy seedling growth in the field, promoting vigorous development and increasing survival probability. Amazonian soils generally exhibit low natural fertility, which increases nutrient demand and makes fertilization necessary to enhance productivity (Costa et al., 2023).

The Amazon region faces strong pressures from deforestation, fire, and extensive cattle ranching; moreover, cattle raising by smallholders has become increasingly unviable economically and environmentally. Consequently, agroforestry systems have gained traction within family farming as a more sustainable activity that diversifies household income. Therefore, knowledge of the potential of species to be used in these systems, and their associated economic value, is essential (Silva et al., 2023).


In agroforestry systems, *P. microphyllus* is an important component because it can develop under adverse conditions and because of the demand for pilocarpine, given that it is the only natural source of this alkaloid and has broad pharmacological applications (Pinheiro, 2001; Grandi, 2021; De Sá et al., 2021). Additionally, the state of Pará reported a production of 50 thousand tonnes in 2022 and sustained demand for the product, with commercialization supported and stimulated by cooperatives such as the Cooperativa Alternativa Mista dos Pequenos Produtores do Alto Xingu (CAMPPAX) and the Cooperativa dos Extrativistas da Flona de Carajás (COEX-CARAJÁS), which purchase leaves and assist in production and marketing (IBGE, 2022).

Few studies have examined the behavior and development of *P. microphyllus* in agroforestry systems, where pilocarpine concentrations required by industry, species consociation, and silvicultural treatments must be considered. Therefore, this study aimed to evaluate the initial silvicultural performance of *P. microphyllus* in relation to light regime and silvicultural treatments in agroforestry systems established in Alto Xingu, Pará.

Material and Methods

The study was conducted on three smallholder properties in the municipality of São Félix do Xingu, state of Pará, Brazil. According to Köppen classification, the climate is Aw, with a rainy winter and a well-defined dry season, mean annual precipitation of approximately 2,100 mm, and mean annual temperature of 26.5 °C. Vegetation is predominantly dense ombrophilous forest, with smaller patches of Cerrado and other formations. Figure 1 shows monthly precipitation and temperature for 2021.

Figure 1. Monthly precipitation (blue bars) and temperature (red line) in 2021. Precipitation is measured in millimeters (mm) on the left vertical axis, and temperature in degrees Celsius (°C) on the right vertical axis



Source: author elaboration from INMET data (INMET, 2024).

The agroforestry systems (AFSs) were established on dystrophic red-yellow Argisols, one of the most abundant soil classes in Brazil, typically occurring on more rugged relief and characterized by low natural fertility. *P. microphyllus* was planted in three distinct AFS compositions; each AFS was treated as a replicate. Soil characterization for the 0–20 cm layer is summarized in Table 1.

Table 1 – Summary of soil analyses (0–20 cm depth) for agroforestry systems

AFS	depth (cm)	pH	OM	K	Fe	P	Ca	Mg	Al	H+Al	T	V	M
			%	mg dm^{-3}			cmol(c) dm^{-3}					%	
1	0 - 20	4,83	4,89	67,40	0,03	5,38	1,60	1,00	0,10	3,31	6,08	45,58	3,48
2	0 - 20	4,80	19,00	37,60	86,30	3,00	1,90	0,40	0,10	3,40	5,80	41,38	4,00

3	0 - 20	5,10	20,00	35,70	46,60	2,30	2,10	0,40	0,00	1,80	4,39	59,00	0,00

Em que: OM = organic matter; T = cation exchange capacity at pH 7.0; V = base saturation index; m = aluminum saturation index.

The three AFS compositions are multi-stratified silvopastoral/ silviagricultural systems that provide different light regimes for planted specimens. Each composition was treated as a replicate because the study's focal point was the initial silvicultural performance of *P. microphyllus* under the applied silvicultural treatments.

- **AFS 1:** *Musa* spp., *Ananas comosus*, *Manihot esculenta*, *Tithonia diversifolia*, *Theobroma cacao*, *Handroanthus albus*.
- **AFS 2:** *Malpighia emarginata*, *Plinia cauliflora*, *Anacardium occidentale*, *Citrus sinensis*, *Eugenia uniflora*, *Mangifera indica*, *Citrus limon*, *Spondias purpurea*, *Lafoensia glyptocarpa*, *Musa* spp., *Passiflora edulis*, *Theobroma cacao*, *Khaya ivorensis*, *Euterpe oleracea*, *Handroanthus albus*, *Spondias mombin*, *Inga edulis*, *Maximiliana maripa*, *Guazuma ulmifolia*, *Psidium guajava*, *Anadenanthera colubrina*.
- **AFS 3:** characterized by greater shading; tree species include *Mauritia flexuosa*, *Theobroma cacao*, *Euterpe oleracea*, *Theobroma grandiflorum*, *Artocarpus heterophyllus*, *Persea americana*, *Mangifera indica*, *Averrhoa carambola*, *Syzygium jambos*, *Maximiliana maripa*, *Genipa americana*, *Ipomoea carnea*, *Calycophyllum spruceanum*, *Inga edulis*, *Oenocarpus bacaba*, *Sclerolobium paniculatum*, *Croton urucurana*.

P. microphyllus seedlings were planted in 2022 at 50 cm × 50 cm spacing. Seedlings were donated by the Centroflora nursery and originated from local seed sources. The experiment used a randomized complete block design (RCBD) in a 2 x 6 factorial arrangement (two pruning regimes × six fertilization combinations), with each AFS serving as a block (replicate), Table 2. Each plot contained three plants, totaling 108 sampling units.

Table 2 — Treatments (pruning and fertilization combinations applied to *Pilocarpus microphyllus* in the different AFS compositions)

Treatment	
1	Pruning
2	Pruning + 75 g NPK*
3	Pruning + 45 g NPK*
4	Pruning + 175 g NPK*
5	Pruning + 45 g MAP**
6	Pruning + 45 g NPK* + 45 g MAP**
7	No pruning
8	No pruning + 75 g NPK*
9	No pruning + 45 g NPK*
10	No pruning + 175 g NPK*
11	No pruning + 45 g MAP**
12	No pruning + 45 g NPK* + 45 g MAP**

NPK*: the source used was the 20:0:20 formulation; MAP**: monoammonium phosphate with 61% P_2O_5 and 12% N.

Formative pruning was applied at experiment establishment at 30 cm height or when specimens reached an equivalent stem diameter of 4 mm. Fertilizers were applied once in lateral side holes, except for treatments 2 and 4 (higher fertilizer rates), which received two split applications: one at establishment and a second dose 30 days later to reduce nutrient losses.

Pruning was performed at experiment establishment at a height of 30 cm or when specimens reached a stem diameter of 4 mm. Fertilization was applied once in lateral planting pits for all treatments except treatments 2 and 4, which received higher fertilizer rates and were therefore applied in two split doses: the first at experiment establishment and the second 30 days after the first application to reduce nutrient losses.

Results were obtained by measuring root collar diameter, total seedling height, crown area, number of shoots > 30 cm, and mean shoot length. Measurements were taken every 30 days after experiment installation.

To better characterize treatment responses, the light regime in the three AFS compositions was recorded using a thermo-hygro-anemometer to assess whether illumination significantly affected the analyzed variables. For subsequent analyses, the mean of the three plants per plot was calculated for root collar diameter, total seedling height, crown area, number of shoots > 30 cm, and mean shoot length.
Crown area (CA) was calculated using the following equation:

$$CA = \frac{(CWI)x(CWR)x\,\pi}{4} \quad (1)$$

Where,
CA: is crown area per plant (m²)
CWI: is crown width in the interrow direction (m);
CWR: is crown width in the row direction (m);
π: constant (3.141592654).

Data were tabulated and subjected to analysis of variance (ANOVA). When effects were significant, mean comparisons were performed using Tukey's test and/or the Scott-Knott test at a 5% significance level, implemented in R statistical software.

5 Results

The measurement of light intensity showed that AFS 2 had the highest mean irradiance (17,445.5 lux), followed by AFS 1 (3,787.5 lux) and AFS 3 (2,954.5 lux). Comparison among the systems revealed a significant effect on root collar diameter (DC), total height (HT), number of shoots (NS), and crown area (CA). For DC and HT, AFS 1 exhibited significantly higher means than the other systems (Table 3).

Table 3 – Comparison among agroforestry systems (AFS) for root collar diameter (DC), total height (HT), number of shoots (NS), and crown area (CA) at 17, 18, 19, and 20 months after planting

Root collar diameter (mm)				
AFS	**Age at fertilization**			
	17 months	18 months	19 months	20 months
1	7.56 a	8.28 a	8.55 a	9.20 a
2	4.10 b	4.61 b	5.03 b	5.84 b
3	4.35 b	4.59 b	4.88 b	5.22 b
CV (%)	21.34	21.59	20.75	18.9
Mean (%)	5.34	5.83	6.15	6.75
Total height (cm)				
AFS	**Age at fertilization**			
	17 months	18 months	19 months	20 months
1	44.25 a	46.92 a	50.67 a	60.92 a
2	33.90 b	36.93 b	39.90 b	44.58 b
3	34.31 b	35.81 b	38.16 b	43.58 b
CV (%)	21.55	20.64	18.34	20.65
Mean (%)	37.49	39.88	42.91	49.69
Number of shoots (unit.)				
AFS	**Age at fertilization**			
	17 months	18 months	19 months	20 months
1	3.00 a	3.58 a	4.08 a	4.25 a
2	2.70 a	2.97 ab	3.10 ab	3.14 ab
3	1.83 a	2.11 b	2.11 b	2.22 b
CV (%)	46.86	46.74	49.51	51.43
Mean (%)	2.51	2.89	3.10	3.20

Crown area (m²)				
AFS	**Age at fertilization**			
	17 months	18 months	19 months	20 months
1	0.03 a	0.05 a	0.07 a	0.09 a
2	0.02 a	0.03 b	0.03 b	0.04 b
3	0.03 a	0.04 ab	0.04 b	0.05 b
CV (%)	48.41	41.56	42.66	47.99
Mean (%)	0.03	0.04	0.05	0.06

* Means followed by at least one identical letter within a column do not differ significantly by Tukey's test at the 5% probability level.

Over the monitoring period, AFS 1 consistently exhibited higher means for root collar diameter (DC) and total height (HT) at all evaluated ages, while AFS 2 and AFS 3 showed statistically lower means. For number of shoots (NS) and crown area (CA), AFS 1 demonstrated an advantage beginning at the 19th month. AFS 3 presented the poorest overall performance, which was associated with greater shading, whereas AFS 2, despite having the highest measured irradiance, did not show superior growth, indicating that factors other than light availability influenced seedling performance.

Analysis of the effect of silvicultural treatments on the agroforestry systems revealed a significant interaction between pruning (P vs. NP) and fertilization for the variable total height (HT). When plants were pruned, no statistically significant differences were observed among the fertilization doses tested. In contrast, under no pruning (NP) conditions, the higher NPK doses (175 g NPK and 75 g NPK) produced the most pronounced height growth, particularly from the 19th month onward (Table 4).

Table 4 – Evaluation of the effect of pruning in combination with fertilization on the variable Total Height (HT) in cm at 17, 18, 19, and 20 months after planting.

Age at fertilizer application: 17 month						
Treatment	0 g NPK	45 g NPK	75 g NPK	175 g NPK	45 g MAP	45 g NPK and 45 g MAP
P	38.2 A a	36.0 A a	38.00 A a	37.72 A a	33.78 A a	37.56 A a
NP	33.67 A a	35.3 A a	40.89 A a	51.11 A a	39.67 A a	27.89 A a
CV (%)	51.71					
Mean (%)	16.81					
Age at fertilizer application: 18 month						
Treatment	0 g NPK	45 g NPK	75 g NPK	175 g NPK	45 g MAP	45 g NPK and 45 g MAP
P	44.2 A a	37.9 A a	42.89 A a	38.06 A a	35.78 A a	39.00 A a
NP	35.00 B a	35.9 B a	43.67 B a	57.11 A a	41.00 B a	28.11 B a
CV (%)	20.64					
Média (%)	39.9					
Age at fertilizer application: 19 month						
Treatment	0 g NPK	45 g NPK	75 g NPK	175 g NPK	45 g MAP	45 g NPK and 45 g MAP
P	46.6 A a	42.8 A a	46.11 A a	40.39 A b	40.33 A a	43.11 A a
NP	36.72 B a	38 B a	50.89 A a	57.94 A a	42.33 B a	30.22 B a
CV (%)	18.34					
Mean (%)	42.9					
Age at fertilizer application: 17 month						
Tratamento	0 g NPK	45 g NPK	75 g NPK	175 g NPK	45 g MAP	45 g NPK and 45 g MAP
P	54.8 A a	53.0 A a	55.44 A a	46.1 A b	54.11 A a	52.89 A a
NP	38.00 B a	39.9 B a	56.67 A a	66.22 A a	46.11 B a	33.11 B b
CV (%)	20.65					

Mean (%)	49.69

Lowercase letters indicate comparisons among rows; means followed by at least one identical lowercase letter within the same column do not differ significantly according to Tukey's test at the 5% significance level. Uppercase letters indicate comparisons among columns; means followed by at least one identical uppercase letter within the same row do not differ significantly according to the Scott-Knott test at the 5% significance level

Multiple comparison tests indicated differences both among rows and among columns (Table 4). In the absence of pruning, the largest NPK doses (175 g and 75 g) promoted the greatest increases in height, while under pruning the combined fertilization treatment 45 g NPK + 45 g MAP stood out at the 20th month.

These results suggest that pruning modifies the seedlings' response to fertilization, reducing the relative effect of fertilizer dose on height growth, whereas in unpruned plants fertilizer dose is a key driver of vertical growth. There was no interaction between the factors for number of shoots (NS) and mean shoot length (MSL); however, a significant isolated effect of the pruning factor was observed. Pruning resulted in higher mean values for both NS and MSL in all evaluated months (17, 18, 19, and 20 months after planting) (see Table 5).

Table 5 – Evaluation of the effect of pruning on Number of shoots (NS) and Mean shoot length (MSL) at 17, 18, 19, and 20 months after planting

Number of shoots (NS)				
	Age at fertilizer			
Pruning	17 months	18 months	19 months	20 months
P	3.96 a	4.24 a	4.24 a	4.31 a
NP	1.06 b	1.54 b	1.95 b	2.09 b
CV (%)	46.86	46.74	49.51	51.43
Média (%)	2.51	2.89	3.10	3.20
Mean shoot length (cm)				
	Age at fertilizer			
Pruning	17 months	18 months	19 months	20 months
P	8.05 a	11.43 a	12.76 a	16.86 a
NP	4.05 b	5.41 b	6.96 b	8.22 b
CV (%)	67.48	55.77	59.57	60.05
Média (%)	6.06	8.42	9.86	12.54

* Means followed by at least one identical letter in the column do not differ significantly according to the Scott-Knott test at the 5% probability level.

At 20 months (Table 6), crown area (CA) showed a significant effect of fertilization, with the 75 g NPK and 175 g NPK doses standing out relative to the others. The Scott-Knott analysis at the 5% probability level indicated that these two doses formed the group of superior means at the end of the evaluation period.

Table 6 – Effect of fertilization on Crown Area (CA) at 17, 18, 19 and 20 months after planting

Crown area (m²)				
Fertilization	**Age at fertilization**			
	17 months	18 months	19 months	20 months
0 g of NPK	0.02 a	0.04 a	0.04 a	0.04 b
45 g of NPK	0.03 a	0.03 a	0.04 a	0.05 b
75 g of NPK	0.03 a	0.05 a	0.06 a	0.09 a
175 g of NPK	0.03 a	0.05 a	0.06 a	0.08 a
45 g of MAP	0.02 a	0.04 a	0.04 a	0.04 b
45 g of NPK e 45 g and MAP	0.02 a	0.05 a	0.06 a	0.06 b
CV (%)	48.41	41.56	42.66	47.99

Mean (%)	0.03	0.04	0.05	0.06

* Means followed by at least one identical letter in the column do not differ significantly according to the Scott-Knott test at the 5% probability level.

Discussion

The results indicate that, under the evaluated conditions, intermediate lighting (AFS 1, ~3,787.5 Lux) favored the initial vegetative development of *Pilocarpus microphyllus*, as reflected in greater root collar diameters, total heights, number of shoots, and crown area. This pattern suggests that the species, during the seedling and establishment phase, may exhibit better performance under moderate light intensity rather than under extreme full sun or intense shading.

The observed discrepancy between the higher irradiance in AFS 2 and its inferior performance can be explained by complementary mechanisms: stress due to excessive radiation (possible photoinhibition or increased water demand) and edaphic limitations associated with the pasture history (reduced fertility, soil compaction, and low organic matter content), which compromise the conversion of available light into growth. Conversely, the reduced performance in AFS 3 is consistent with the classic effects of shading on photosynthesis and the accumulation of photoassimilates, resulting in a lower resource allocation to the stem and crown (Fonseca, 2002; Engel et al., 1990; Viana et al., 2022).

Compared to the literature, some studies report greater sprouting and leaf number under higher radiation (Assis et al., 2016; Ribeiro et al., 2011; Cardoso et al., 2012), whereas other works on analogous species indicate optimal responses under half-shade conditions (Mazuchowski et al., 2007). Thus, the findings of this study do not contradict the literature but highlight that the response of *P. microphyllus* is dependent on the environmental context and soil management; the optimal light intensity may vary according to water availability, nutrients, and land-use history (Siebeneichler et al., 2008; Rayol, 2018).

For practical applications, the data suggest that management practices promoting intermediate lighting (controlled partial shading or intercropping that reduces excessive direct radiation) can optimize the initial growth of jaborandi seedlings in Agroforestry Systems (AFS). In areas recovered from pasture, soil evaluation and correction prior to planting are recommended to minimize edaphic limitations that could negate the benefits of higher light availability (Mizobata et al., 2017; Pinho et al., 2021).

Upon evaluating the application of silvicultural treatments, the findings indicate that the height response of *P. microphyllus* depends on the interaction between pruning management and fertilization. The superiority of the 175 g and 75 g NPK doses in the absence of pruning suggests that when the plant maintains a larger leaf area, the higher availability of nutrients (especially nitrogen) is effectively utilized for vertical growth, corroborating previous observations regarding the importance of N for the initial development of jaborandi (Marques and Costa, 1994; Brasil and Viégas, 1999).

The observation that there were no differences among fertilization treatments under pruning indicates that the removal of part of the aboveground biomass alters resource demand and allocation, reducing the immediate response to the different doses tested. In this context, the better performance of the 45 g NPK + 45 g MAP combination in the presence of pruning at the 20th month reinforces the need for balanced formulations that supply nitrogen and phosphorus for recovery and sprouting after pruning management, aligning with the recommendations for split fertilization in jaborandi (Marques and Costa, 1994).

The results observed in *P. microphyllus* also converge with evidence from other forest species, in which fertilization with N, P, and K promotes height increments (Vieira et al., 2017; Souza et al., 2013). Studies such as those by Vieira et al. (2017), Souza et al. (2013), Nieri et al. (2019), and Silva et al. (2022) demonstrate that adequate doses of nitrogen and phosphorus increase height growth, which supports the interpretation that nitrogen was a determining factor in the responses observed in this experiment.

From a physiological standpoint, nitrogen is fundamental in the synthesis of chlorophyll, proteins, and enzymes, favoring photosynthesis and growth, while phosphorus participates in energy transfer and essential metabolic processes; the deficiency of these nutrients at the beginning of the cycle can irreversibly compromise development (Souza et al., 2013). Thus, higher NPK doses can accelerate initial growth when the plant has sufficient leaf area to capture light and convert nutrients into biomass, which explains the greater height response in the absence of pruning.

For practical management, the data suggest that in plantations without intensive pruning, it is advantageous to apply higher doses of NPK (up to 175 g) to promote height growth, whereas in systems with scheduled pruning, the NPK + MAP combination at moderate doses can favor recovery and sprouting. These recommendations resonate with historical guidelines for jaborandi (Marques and Costa, 1994) and

with evidence that nitrogen is one of the main limiting factors for the species' initial growth (Brasil and Viégas, 1999).

The findings indicate that fertilization with higher NPK doses (75 g and 175 g) favored the development of the *P. microphyllus* crown area up to the 20th month, suggesting a positive effect of fertilization on leaf mass expansion. This result can be attributed, in particular, to the nitrogen and potassium present in the 20-0-20 formulation used: nitrogen is an essential component of the protoplasm and chlorophyll, influencing cell division and photosynthetic capacity; potassium regulates the osmotic potential, cell turgor, and enzymatic processes that affect transpiration and CO2 diffusion, contributing to a higher photosynthetic rate and, consequently, greater crown growth (Santos et al., 2010).

The positive influence of nitrogen fertilization on shoot production is corroborated by Cruz et al. (2006), who observed greater aboveground dry matter gains in *Samanea inopinata* seedlings with higher applications of ammonium sulfate. Similarly, the results of this study reinforce the role of N in promoting vegetative growth in forest species and commercial seedlings.

It is noted, however, that the significance of the differences emerged mainly in the final months of evaluation, indicating that responses to fertilization may require time to fully manifest in terms of crown area. This pattern reinforces the need for medium-term monitoring, as effects on morphological attributes and foliar biomass accumulation tend to consolidate with plant development.

From a management perspective, the data suggest that NPK applications at higher doses can be recommended to promote greater crown development during the initial stages of establishment, which can accelerate canopy closure and foliar biomass production. However, it is recommended that such guidelines be accompanied by soil analyses and fertilization programs tailored to the actual nutrient availability and the phenological needs of the species.

Finally, considering that pilocarpine levels and other phytochemical attributes of jaborandi typically stabilize only at more advanced ages (approximately four years in field experiments), it is imperative to extend the experimental monitoring to evaluate whether the fertilization regimes that maximize crown area also positively influence phytochemical quality and long-term productivity. Future studies should include foliar analyses, physiological measurements, and monitoring of N and K dynamics in the soil to elucidate mechanisms and optimize fertilization recommendations for *P. microphyllus* (Santos et al., 2010; Cruz et al., 2006).

Conclusion

The different agroforestry systems demonstrated that jaborandi develops better under intermediate light incidence. In the presence of pruning, fertilization did not affect height growth. Pruning positively influenced both the number of shoots and their length.

Fertilization with 175 g and 75 g NPK promoted increases in crown area and seedling height, with pronounced effects observed only by the twentieth month, indicating the need for prolonged monitoring to better characterize responses. When pruning is performed, to maximize height growth the recommended fertilization is 45 g NPK combined with 45 g MAP.

Silvicultural treatments benefited the development of *P. microphyllus* seedlings; planting the species in agroforestry systems that provide moderate solar radiation is recommended. Among silvicultural practices, formative pruning is advised to stimulate new shoot development and crown expansion, thereby enhancing leaf production.

References

Amaral, G. C., Pezzopane, J. E. M., Nóia Júnior, R. S., Fernández Martínez, M., Fonseca, M. D. S., Gibson, E. L., Toledo, J. V., & Pezzopane, J. R. M. (2022). *Pilocarpus microphyllus* seedling growth threatened by climate change: an ecophysiological approach. *Theoretical and Applied Climatology*, 147(1), 347–361. https://doi.org/10.1007/s00704-021-03831-6

Assis, R. M. A. de, Lameira, O. A., Moura, R. C., & Costa, K. J. A. (2016). Effect of pruning and growing environment on branch development of jaborandi (*Pilocarpus microphyllus* Stapf). *Revista de Ciências Agrárias*, 59(2), 123–130.

Brasil, E. C., & Viégas, I. de J. M. (1999). *Influence of fertilization with nitrogen, phosphorus and potassium on the production of jaborandi seedlings* (Technical Communication No. 100). Embrapa Eastern Amazon.

Caldeira, C. F., Giannini, T. C., Ramos, S. J., Vasconcelos, S., Mitre, S. K., Pires, J. P. de A., ... & Furtini, A. E. (2017). Sustainability of jaborandi in the eastern Brazilian Amazon. *Perspectives in Ecology and Conservation*, 15, 161–171.

Cardoso, L. G., Lameira, O. A., Ribeiro, F. N., & Rocha, T. T. (2012). Effect of different cutting distances on branches of jaborandi (*Pilocarpus microphyllus* Stapf ex Wardleworth). *Scientia Plena*, 8(4).
Costa, L. R. de J., et al. (2023). Soil fertility in oil palm agroforestry systems in the Eastern Amazon, Brazil. *Agroforestry Systems*, 97, 865–881. https://doi.org/10.1007/s10457-023-00832-4

Cruz, C. A. F., Paiva, H. N., & Guerrero, C. R. A. (2006). Effect of nitrogen fertilization on the production of sete-cascas (*Samanea inopinata* (Harms) Ducke) seedlings. *Revista Árvore*, 30(4), 537–546.

De Sá, É. R. A., Costa, A. N., Costa, R. K. M., Souza, J. L., Ramos, R. M., & Lima, F. D. C. A. (2021). In silico study of the interactions of *Pilocarpus microphyllus* imidazolic alkaloids with the main protease (Mpro) of SARS-CoV-2. *Molecular Simulation*, 47(12), 1–14. https://doi.org/10.1080/08927022.2021.1873321

Engel, V. L., & Poggiani, F. (1990). Influence of shading on the growth of seedlings of some native species and their ecological and silvicultural implications. *IPEF*, (43/44), 1–10.

Fonseca, É. de P., Valéri, S. V., Miglioranza, É., Fonseca, N. A. N., & Couto, L. (2002). Quality pattern of *Trema micrantha* (L.) Blume seedlings produced under different shading periods. *Revista Árvore*, 26(4), 515–523. https://doi.org/10.1590/S0100-67622002000400015

Grandi, M. C. F. (2021). *Study of the biosynthetic route of pilocarpine in jaborandi through biochemical and molecular analysis* [Doctoral dissertation, State University of Campinas]. UNICAMP Repository.

Gumier-Costa, F., McGrath, D. G., Pezzuti, J. C. B., & Homma, A. K. O. (2016). Institutional partnerships and the evolution of jaborandi extraction in the Carajás National Forest, Pará, Brazil. *Sustainability in Debate*, 7(3), 91–111. https://doi.org/10.18472/SustDeb.v7n3.2016.18955

Homma, A. K. O. (2012). Plant extraction or cultivation: which is the option for the Amazon? *Estudos Avançados*, 26(74), 167–186.

Homma, A. K. O. (2014). *Plant extraction in the Amazon: history, ecology, economy and domestication*. Embrapa.

IBGE — Brazilian Institute of Geography and Statistics. (2022). *Production of vegetal extraction and forestry 2022*. https://sidra.ibge.gov.br/pesquisa/pevs/tabelas

INMET — National Institute of Meteorology. (2024). *Climatic monitoring (annual historical data)*. https://portal.inmet.gov.br/

Marques, M. E. T., & Costa, J. P. C. (1994). *Jaborandi (Pilocarpus microphyllus)* (Recommendation Basics No. 27). Eastern Amazon Agroforestry Research Center.

Mazuchowski, J. Z., Silva, E. T. da, & Maccari Junior, A. (2007). Effect of light intensity and nitrogen addition on the growth of *Ilex paraguariensis* plants. *Revista Árvore*, 31(4), 619–627. https://doi.org/10.1590/S0100-67622007000400006

Ministry of the Environment. (2022). *Ministerial Ordinance MMA No. 148, 7 June 2022*. Brasília, Brazil.

Mizobata, K. K. G. da S., Cassiolato, A. M. R., & Maltoni, K. L. (2017). Growth of Baru and Gonçalo-Alves seedlings on degraded soil supplemented with residue, Ilha Solteira – SP. *Ciência Florestal*, 27(2), 429–444. https://doi.org/10.5902/1980509827726

Nieri, E. M., Teixeira, C. E. S., Melo, L. A., Silva, D. S. N., Luz, M. S., & Botelho, S. A. (2019).

Hydroretentive polymer and fertilization in the establishment of *Toona ciliata* M. Roemer. *Scientia Forestalis*, 47(122), 317–325. https://doi.org/10.18671/scifor.v47n122.14

Pinheiro, C. U. B. (2001). Extraction, cultivation and privatization of jaborandi (*Pilocarpus microphyllus* Stapf ex Holm.; Rutaceae) in Maranhão, Brazil. *Acta Botanica Brasilica*, 16(2), 141–150.

Pinho, L. L., de Lacerda, C. F., de Sousa, J. A., Bezerra, A. M. E., Cavalcante, E. S., & de Oliveira, A. C. (2021). Leaf temperature and water-use efficiency in *Anadenanthera colubrina* seedlings under artificial shading and irrigation with brackish water. *[Adicionar Nome da Revista e Páginas]*.

Rayol, B. P., & Alvino-Rayol, F. de O. (2018). Initial development of tree species in agroforestry systems in the Lower Amazon, Pará, Brazil. *Revista de Ciências Agroveterinárias*, 18(1), 59–64. https://doi.org/10.5965/223811711812019059

Ribeiro, F. N., Lameira, O. A., Rocha, T. T., & Azulay, L. B. O. (2011). Evaluation of branch regeneration of jaborandi (*Pilocarpus microphyllus* Stapf ex Wardleworth). *[Adicionar Nome da Revista e Páginas]*.

Santos, D. R. dos, & Silva, L. S. da. (2010). *Fertilidade do solo e nutrição de plantas* [Soil fertility and plant nutrition]. Universidade Federal de Santa Maria.

Siebeneichler, S. C., Freitas, G. A. de, Silva, R. R. da, Adorian, G. C., & Capellari, D. (2008). Morphophysiological characteristics of *Tabebuia heptaphylla* under different light conditions. *Acta Amazonica*, 38(3), 467–472. https://doi.org/10.1590/S0044-59672008000300011

Silva, C. P., Santos, L. M., Faria, J. C. T., Bruzinga, J. S. C., Melo, L. A., & Nieri, É. M. (2023). Silvopastoral system: scenario in the municipality of São Félix do Xingu-PA. *Ciência Florestal*, 33(3), e71818. https://doi.org/10.5902/1980509871818

Silva, O. M. C., Nieri, E. M., Santana, L. S., Almeida, R. S., Araújo, G. C. R., Botelho, S. A., & Melo, L. A. (2022). Phosphated fertilization in the initial growth of seven native forest species for recovery of a degraded area. *Ciência Florestal*, 32(1), 371–394. https://doi.org/10.5902/1980509861339

Souza, C. B. G., Campos, A. S., Santos, A. M. A. dos, Sousa, F. B. B. de, Souza, N. H. de, ... & Silva, E. F. da. (2013). Nutritional study of canafístula (I): growth and seedling quality in response to nitrogen and phosphorus fertilization. *Revista Árvore*, 37(4), 717–724. https://doi.org/10.1590/S0100-67622013000400015

Teixeira, C. E. S., Torres, A. Q. A., Nieri, E. M., Melo, L. A. de, Santos, L. V. dos, & Botelho, S. A. (2019). Hydroretentive polymer and mineral fertilization in the establishment of a *Eucalyptus urophylla* × *Eucalyptus grandis* hybrid. *Ciência Florestal*, 29(3), 1060–1071. https://doi.org/10.5902/1980509834950

Viana, E. de O., Bezerra, A. M. E., Pereira, D. de S., Lacerda, C. F. de, & Medeiros Filho, S. (2022). Growth and quality of seedlings of three *Copernicia* species under different shading conditions. *Ciência Florestal*, 32(4), 2094–2112. https://doi.org/10.5902/1980509866763

Vieira, C. R., Weber, O. L. dos S., & Scaramuzza, J. F. (2017). Influence of NPK fertilization on the initial growth of *Tectona grandis* L. F. in the field. *Revista do Instituto Florestal*, 29(1), 57–69. https://doi.org/10.24278/2178-5031.201729104