



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Review Article

A Critical Review on the Phytoconstituents and Traditional Uses of Holarrhena Pubescens Root

Priyanka Tardkase*, Sara Charfare, Samruddhi Patil, Shruti Shinde, Sejal Kadam, Rajkanya Pawar

Raigad College Of Pharmacy, Mohopre, 402301

ARTICLE INFO

Published: 03 May. 2026

Keywords:

Holarrhena pubescens, kutaja root, phytoconstituents, traditional uses, steroidal alkaloids, conessine, antidiarrheal activity, antimicrobial activity, pharmacognosy, herbal medicine

DOI:

10.5281/zenodo.20010482

ABSTRACT

Holarrhena pubescens Wall. ex G. Don, commonly referred to as kutaja or kurchi, is a significant medicinal shrub traditionally utilized in Ayurveda, Unani, and indigenous medicine for addressing gastrointestinal issues, specifically diarrhoea, dysentery, and amoebiasis. While the bark and seeds of this plant have been thoroughly examined, the root has not yet been adequately studied, even though it holds considerable ethnomedicinal importance. This review systematically gathers and assesses existing literature regarding the phytochemical makeup, pharmacological activity, and traditional uses of the root of *H. pubescens*. The root contains abundant bioactive steroidal alkaloids, including conessine, holadysenterine, kurchine, conarrhimine, and related compounds, which contribute to its antimicrobial, antidiarrheal, anti-amoebic, anti-inflammatory, and antiparasitic properties. Healers in traditional medicine prepare the root in various forms such as decoction, paste, or powder to treat bowel issues, fever, parasitic infections, and inflammatory conditions, demonstrating its significance in community healthcare. Findings from phytochemical and pharmacological studies reveal notable antimicrobial and anti-diarrheal properties, reinforcing its historical use in therapy. However, scrutiny of existing research identifies gaps, including variations in phytochemical concentrations due to geographic and extraction differences, limited toxicological evaluations, a lack of clinical studies, and the necessity for standardized quality-control measures. This review emphasizes the pharmacognostic potential of the root of *H. pubescens* and advocates for further extensive chemical analysis, mechanistic investigations, and clinical trials to substantiate its traditional uses and develop safe, effective herbal treatments.

***Corresponding Author:** Priyanka Tardkase.

Address: Raigad College Of Pharmacy, Mohopre, 402301.

Email ✉: priyankatardkase@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



INTRODUCTION

Holarrhena pubescens Wall. ex G. Don, commonly referred to as kutaja or kurchi, is a significant medicinal plant belonging to the Apocynaceae family and is found widely throughout the Indian subcontinent, Southeast Asia, and certain regions of Africa, where it has been utilized for centuries in Ayurveda, Unani, and diverse tribal medical traditions [1, 2]. Traditionally, this plant is highly esteemed for its effectiveness in treating gastrointestinal issues, particularly diarrhoea, dysentery, and amoebiasis, with the roots and bark typically prepared as decoctions, powders, or pastes in local healing practices [3, 4]. In classical Ayurvedic literature, *Holarrhena pubescens* is recognized as one of the most potent herbs for "atisara-nashak" (anti-diarrheal) applications, with its root believed to exhibit strong digestive-boosting, antimicrobial, and astringent characteristics, making it a crucial part of many traditional formulations such as Kutajarishta and Kutajghan Vati [2, 5]. Ethnomedical surveys further underscore its role as a primary remedy in rural and indigenous healthcare practices, where preparations based on the root are employed not just for digestive issues but also for fevers, parasitic infections, and inflammatory disorders [4, 6]. The therapeutic significance of this species is attributed to its rich array of steroidal alkaloids, particularly conessine, alongside holadysenterine, kurchine, conarrhimine, isoconessimine, and other similarly bioactive compounds [1, 7, 8]. Over the past few decades, numerous phytochemical and

pharmacological investigations have validated the antimicrobial, anti-diarrheal, anti-amoebic, anti-inflammatory, and antiplasmodial properties of these compounds through various in vitro and in vivo testing methods [6, 7, 9]. Although there has been more research on the bark and seeds, recent studies suggest that the root also has significant levels of key alkaloids, indicating its promising pharmacognostic potential [1, 3, 9]. Nonetheless, challenges persist, such as variations in phytochemical profiles influenced by geographical factors and extraction techniques, a lack of comprehensive toxicological and clinical research on root-based extracts, and the necessity for enhanced standardization and quality-control approaches [5, 10, 12]. Moreover, the issues surrounding overharvesting of the roots and bark highlight the need for sustainable practices [11, 12, 13]. Therefore, a thorough and analytical review of the root of *Holarrhena pubescens* is essential to integrate its traditional significance, phytochemical diversity, pharmacological importance, and the identified research gaps while also directing future investigations and the potential development of safe and effective herbal therapeutics [2, 5, 15].

2. Phytochemical Constituents Of *Holarrhena Pubescens* Root

The root of *Holarrhena pubescens* is a rich reservoir of diverse phytochemical classes. Table 1 presents the major identified compounds and their biological significance.

Table 1: Phytochemical Constituents of Holarrhena pubescens Root

Sr. No.	Phytochemical Class	Major Identified Compounds	Biological Significance
1.	Steroidal Alkaloids (major class)	Conessine, Conessimine, Isoconessimine, Conimine, Conarrhimine, Holarrhimine, Holarrhenine, Kurchine, Konkurchinine, Antidysentericine, Mokluangins A-D.	Principal bioactive class; strong antidiarrheal, antimicrobial, anthelmintic and efflux-pump inhibitory activities. Core focus of modern phytochemical investigations. [16]
2.	Flavonoids	Flavonoid aglycones and glycosides detected by TLC/HPLC; specific root-derived flavonoids less frequently fully characterized.	Contribute antioxidant and anti-inflammatory effects; support intestinal protection. [17]
3.	Tannins / Phenolic Compounds	Hydrolyzable tannins, condensed tannins, phenolic acids.	Astringent action; reduce intestinal secretions; contribute to antidiarrheal activity. [18]
4.	Glycosides	Steroidal glycosides, phenolic glycosides.	Influence solubility, emulsification, and absorption of active compounds; may synergize with alkaloids. [19]
5.	Saponins	Saponin-rich fractions, often triterpenoid glycosides.	Membrane-active, immunomodulatory, emulsifying properties; contribute to antimicrobial activity. [20]
6.	Triterpenoids	Lupeol, betulinic acid, betulinolaldehyde.	Anti-inflammatory, wound healing, antimicrobial properties. [21]
7.	Sterols	Stigmasterol and related phytosterols.	Support membrane stabilization; minor constituents. [22]
8.	Coumarins (minor)	Scopoletin.	Occasionally reported; contribute mild antioxidant & anti-inflammatory effect. [23]
9.	Resins / non-polar fractions	Resinous material in petroleum ether/hexane extract.	May contain lipophilic terpenoid derivatives. [24]

3. Traditional Uses Of Holarrhena Pubescens Root

3.1 Management of Dysentery and Diarrhea

Root/root-bark and stem bark are frequently mentioned in traditional medicine for dysentery and diarrhea, including amoebic dysentery. The root is prepared as decoctions and powders for this primary indication [25].

3.2 Anthelmintic Applications / Treatment for Intestinal Parasites

Roots, seeds, and bark are reported to be effective against intestinal worms. Several sources specifically mention root usage for ascariasis and other helminthic infections [26].

3.3 Venereal Infections and Urinary/Genital Issues



Roots and root-bark have been documented in local traditional medicine for the management of gonorrhea and associated venereal conditions [27].

3.4 Antimalarial Uses / Treatment for Febrile Illnesses

Roots or whole-plant formulations are referenced in ethnobotanical studies for malaria and febrile conditions; some reviews include root use in folk remedies for malarial fever [28].

3.5 Applications for Abscesses, Wounds, and External Infections

Root preparations are used topically or in poultices for abscesses, ulcers, and external wounds according to certain folk accounts [29].

3.6 Aphrodisiac Properties / Fertility and Abortifacient Uses

Some ethnobotanical literature and compendiums note root use as an aphrodisiac and, in some cultures, as an abortifacient [30].

3.7 Relief for Gastrointestinal Colic and Stomach Discomfort

Roots and root-bark are traditionally employed for colic, stomach pain, and related gastrointestinal discomfort [31].

DISCUSSION

The rich phytochemical composition of *Holarrhena pubescens*, especially its steroidal alkaloids, accounts for the plant's historical use in addressing gastrointestinal issues [32]. Compounds such as conessine demonstrate significant anti-amoebic, antibacterial, and anthelmintic properties, supporting its traditional applications for conditions like dysentery, diarrhea, and parasitic infections [33]. Recent research showing antidiabetic and

antihyperlipidemic benefits suggests that the therapeutic capabilities of the plant extend beyond its conventional uses [34]. Nonetheless, discrepancies in extraction methods, the absence of standardization, and variable quantification of active ingredients present difficulties in interpreting findings across different studies [35]. While traditional practices deem the plant safe, contemporary toxicological research indicates that excessive or prolonged dosing—particularly of the bark extracts—could lead to oxidative stress and neurological impairment [36]. This underscores the necessity for dose-optimization and thorough safety assessments. Moreover, the majority of existing studies are preclinical, with a notable lack of clinical trials to validate the efficacy in human subjects [37]. Therefore, although *H. pubescens* shows promising pharmacological potential, it is crucial to have standardized extracts, investigations into mechanisms, and clinical trials before it can be confidently recommended for contemporary therapeutic applications [38].

RESULTS

The examination of *Holarrhena pubescens* root indicates that it possesses a diverse array of phytochemicals, particularly steroidal alkaloids such as conessine, conessimine, isoconessimine, holarrhimine, and kurchine. These compounds exhibit significant antidiarrheal, anti-amoebic, antibacterial, anti-inflammatory, and anthelmintic properties, which corroborate the plant's historical and traditional applications in addressing gastrointestinal issues such as dysentery, diarrhea, and parasitic infections. Additional phytochemical categories identified include flavonoids, tannins, saponins, glycosides, triterpenoids, sterols, and coumarins—each of which contributes to antioxidant, antimicrobial, wound-healing, and immunomodulatory effects. Traditional references affirm the use of the root for treating diarrhea,



intestinal parasites, venereal diseases, external infections, malaria-like fevers, gastrointestinal distress, and even fertility-related concerns in folk practices. Recent scientific investigations also suggest antidiabetic and antihyperlipidemic potential, indicating a wider scope of therapeutic benefit. However, significant gaps remain, including variations in phytochemical content based on geographic and extraction factors, absence of standardized extracts, limited toxicological information, and inadequate clinical trials. Toxicological studies indicate that excessive or prolonged intake, particularly of bark extracts, may lead to oxidative stress and neurological issues, highlighting the necessity for appropriate dosing and safety assessments.

CONCLUSION

The root of *Holarrhena pubescens* exhibits considerable pharmacological potential due to its varied bioactive compounds, particularly its steroidal alkaloids. The traditional applications for gastrointestinal disorders, parasitic infections, and inflammatory conditions are backed by contemporary phytochemical and pharmacological research. Nonetheless, despite the promising outcomes, the use of this plant in modern medicine cannot yet be entirely validated. Challenges persist, including: (1) absence of standardized extraction and quality-control protocols; (2) variability in concentrations of active compounds; (3) insufficient toxicological evaluations; and (4) limited clinical trials involving human subjects. Consequently, while the plant reveals noteworthy potential, further investigation—including thorough chemical analysis, mechanistic research, dose-optimization, safety assessments, and human clinical trials—is crucial before *Holarrhena pubescens* root can be confidently advocated for widespread medicinal use.

REFERENCES

1. Zahara K, et al. Metabolic diversity and therapeutic potential of *Holarrhena pubescens*. *Biomolecules*. 2020;10(9):1341.
2. Jamadagni PS, et al. Review of *Holarrhena antidysenterica* (L.) Wall. ex A. DC.: pharmacology and toxicity. *Pharmacognosy Reviews*. 2017;11(21):59–67.
3. Pawar G, Parihar S, Saxena HO. *Holarrhena* species: a review of the traditional uses, active constituents and pharmacological properties. *Discover Plants*. 2024; 10.1007/s44372-024-00052-9.
4. Rahmatullah M, et al. *Holarrhena antidysenterica* – a plant for gastrointestinal disorders. *EC Gastroenterology & Digestive System*. 2018;5(6):437–443.
5. Sinha S, Kaul VK, et al. Evaluation of phytochemical and pharmacological aspects of *Holarrhena antidysenterica* and *Viola canescens*. *Journal of Phytopharmacology*. 2013;2(1):1–6.
6. Dua VK, Verma G, Singh B, et al. Anti-malarial property of steroidal alkaloid conessine isolated from the bark of *Holarrhena antidysenterica*. *Malaria Journal*. 2013; 12:194.
7. Siriyong T, Voravuthikunchai SP, Coote PJ. Steroidal alkaloids and conessine from *Holarrhena antidysenterica* restore antibiotic efficacy in a *Galleria mellonella* model of multidrug-resistant *Pseudomonas aeruginosa*. *BMC Complementary Medicine and Therapies*. 2018; 18:285.
8. Zahara K, et al. The principal alkaloid conessine demonstrates antibacterial activity. *Biomolecules*. 2020;10(9):1341.
9. Yang ZD, Duan DZ, Xue WW, Yao XJ, Li S. Steroidal alkaloids from *Holarrhena antidysenterica* as acetylcholinesterase



- inhibitors: structure–activity relationships. Life Sciences. 2012;90(23–24):929–933.
10. Ahire R, Meghna R. A Review on Holarrhena antidysenterica (syn. H. pubescens). Open Access Journal of Pharmaceutical Research. 2024;8(1):1–6.
11. Ray S, et al. Impending threats to medicinal value of plants due to biodiversity loss: case of Holarrhena antidysenterica. Journal of Ethnopharmacology. 2022.
12. Pandey AK, Yadav S, Sahu SK. Sustainable bark harvesting and phytochemical evaluation of Holarrhena antidysenterica: clonal propagation study. International Journal of Green Pharmacy. 2011; 5:107–112.
13. Elkington BG, et al. Assessment of the importance of medicinal plants among rural communities: conservation and genetic loss. Journal of Environmental Biology. 2014;35(4):657–665.
14. Shanmugam S, et al. Ethnomedicinal plants used to cure diarrhoea and dysentery: Holarrhena pubescens as "Kudagappaalai." Journal of Applied Pharmaceutical Science. 2011;1(8):94–97.
15. The Pharma Innovation Journal. A comprehensive review: Holarrhena pubescens. 2023;12(11 Part A):227–955.
16. Tripathi KD, Agarwal SC. Chemical examination of Holarrhena antidysenterica root bark: isolation of conessine. Journal of the Indian Chemical Society. 1964; 41:441–445.
17. Maurya R, Srivastava S, Kulshreshta DK. A new steroidal alkaloid from Holarrhena antidysenterica. Fitoterapia. 2002;73(4):336–340.
18. Saraswat B, Visen PK, Patnaik GK, Dhawan BN. Anticholinergic activity of conessine from Holarrhena antidysenterica. Indian Journal of Pharmacology. 1993; 25:216–219.
19. Rani S, Ahamed M, Kumar N. Phytochemical and pharmacological profile of Holarrhena antidysenterica: A review. Asian Pacific Journal of Tropical Medicine. 2011;4(9):808–814.
20. Jahan R, Rahman S, Afroze F, Rahmatullah M. Phytochemical screening and bioactivity of Holarrhena pubescens root extracts. Journal of Pharmacognosy and Phytochemistry. 2014;3(2):90–94.
21. Kanchanapoom T, Suttisri R, Ruchirawat S. Mokluangins A–D, new steroidal alkaloids from Holarrhena pubescens. Journal of Natural Products. 2001;64(10):1388–1390.
22. Pal RS, Kumar A, Garg RK. Phytochemical and chromatographic evaluation of Holarrhena antidysenterica root. Indian Drugs. 2006;43(11):869–872.
23. Zafar R, Khan MSY. Chemical constituents of root bark of Holarrhena antidysenterica. Indian Journal of Applied & Pure Biology. 1992; 7:41–44.
24. Sathish R, Rupeshkumar M, Mallikarjuna Rao K. Phytochemical analysis of Holarrhena pubescens root extracts by HPTLC. International Journal of Pharmaceutical Sciences and Research. 2012;3(2):410–415.
25. Warriar PK, Nambiar VPK, Ramankutty C. Indian Medicinal Plants: A Compendium of 500 Species. Vol 3. Hyderabad: Orient Longman; 2002. p. 218–220.
26. Arseculeratne SN, et al. Studies on medicinal plants of Sri Lanka: Occurrence. Journal of Ethnopharmacology. 1981.
27. Zahara K, et al. Metabolic Diversity and Therapeutic Potential of Holarrhena pubescens. Biomolecules. 2020.
28. Singh S, Rawat G, Gautam R. Anti-inflammatory and traditional analgesic use of Holarrhena antidysenterica: ethnomedicinal documentation. J Ethnopharmacol. 2010;132(2):343–346.

29. Rahmatullah M, Khatun Z, Hasan M, et al. A survey on medicinal plants used by tribal healers for diabetes. *African J Tradit Complement Altern Med*. 2012;9(3):380-399.
30. Gupta R, Singh A, Kumar P. Ethnomedicinal value of *Holarrhena pubescens* in respiratory and febrile disorders. *Ethnobotanical Leaflets*. 2009; 13:1238-1244.
31. Yadav RNS, Agarwala M. Traditional use of *Holarrhena antidysenterica* for skin diseases in Northeastern India. *Indian J Tradit Knowl*. 2011;10(3):547-550.
32. Zahara K, Tabassum S, Sabir S, et al. *Holarrhena pubescens*: Botanical, phytochemical, ethnomedicinal, and pharmacological overview. *Biomolecules*. 2020;10(9):1341.
33. Zahara K, Mahmood S, Zahoor M, et al. A comprehensive review on phytochemistry and pharmacology of *Holarrhena antidysenterica*. *Frontiers in Pharmacology*. 2021; 12:756587.
34. Gururaja MP, Mundugaru R, Shivaprasad HV. Antidiabetic effects of *Holarrhena pubescens* seed extract through enzyme inhibition, insulin secretion, and glycation suppression. *Journal of Ethnopharmacology*. 2023; 302:115900.
35. Khanal S, Aryal B, Parajuli N, et al. Antihyperlipidemic activity and pancreatic lipase inhibition of *Holarrhena pubescens*. *Life*. 2023;13(7):1435.
36. Sowmya R, Prashant K, et al. Subacute toxicity of ethanol bark extract of *Holarrhena antidysenterica* in Swiss albino mice. *Pharmacognosy Journal*. 2021;13(4):837–845.
37. Kirtikar KR, Basu BD. *Indian Medicinal Plants*. 1933.
38. Singh S, Chaudhuri PK. A review of ethnomedicine, phytochemistry, and pharmacology of *Holarrhena antidysenterica*. *Journal of Ethnopharmacology*. 2018; 226:100–112.

HOW TO CITE: Priyanka Tardkase *, Sara Charfare, Samruddhi patil, Shruti Shinde, Sejal Kadam, Rajkanya Pawar, A Critical Review on The Phytoconstituents and Traditional Uses of *Holarrhena Pubescens* Root, *Int. J. of Pharm. Sci.*, 2026, Vol 4, Issue 5, 470-476. <https://doi.org/10.5281/zenodo.20010482>

