

**EXTRACTION AND FUNCTIONAL APPLICATIONS OF OPUNTIA FICUS-INDICA MUCILAGE IN MATRIX PREPARATION****Aman Suthar^{*1}, Mr. Milan Jadav², Dr. Pragnesh Patani³**¹Khyati College of Pharmacy, Palodia, Ahmedabad.²Associate Professor, Department of Pharmacology, Khyati College of Pharmacy, Palodia, Ahmedabad.³Principle, Khyati College of Pharmacy, Palodia, Ahmedabad.

Article Received on 22 Feb. 2026,
Article Revised on 14 March 2026,
Article Published on 01 April 2026,
<https://doi.org/10.5281/zenodo.19331161>

Corresponding Author*Aman Suthar**Khyati College of Pharmacy, Palodia,
Ahmedabad.

How To Cite This Article: Aman Suthar¹, Mr. Milan Jadav², Dr. Pragnesh Patani³ (2026). Extraction And Functional Applications OF Opuntia Ficus-Indica Mucilage in Matrix Preparation. World Journal of Pharmacy and Pharmaceutical Sciences, 15(4), 148–164.
This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Opuntia ficus-indica mucilage has emerged as a valuable natural polysaccharide excipient for the development of pharmaceutical matrix tablets, owing to its distinctive rheological characteristics and excellent biocompatibility when compared with synthetic polymers. This review provides an in-depth overview of the existing extraction processes, analytical characterization methods, and functional roles of this mucilage in controlled drug delivery systems. Its remarkable ability to form gels, exhibit viscosity that varies with concentration, and display shear-thinning behavior makes it highly effective for use in sustained-release formulations. The polysaccharide components of *Opuntia ficus-indica* mucilage are mainly composed of arabinose, galactose, xylose, and galacturonic acid units, which together create complex macromolecular

structures with molecular weights between 20,000 and 43,000 Da. Detailed analyses have shown that this mucilage exhibits pH-responsive swelling, strong mucoadhesive characteristics, and excellent ability to regulate drug release. Moreover, recent advancements in its modification methods—such as chemical grafting and the development of composite systems—have broadened its applications to include superporous hydrogels, nanoparticle-based delivery systems, and bioadhesive film formulations. The environmentally sustainable nature and economic feasibility of *Opuntia ficus-indica* mucilage make it a compelling

substitute for synthetic excipients, especially in emerging pharmaceutical sectors. Recent studies have reported its successful integration into matrix tablet formulations for a variety of therapeutic drugs, resulting in improved patient adherence and minimized side effects. Ongoing research emphasizes the need for standardized production protocols, stringent quality assurance practices, and innovative formulation approaches to enable its use in personalized medicine. Overall, this review consolidates current insights and highlights future opportunities for the advancement of mucilage- based drug delivery technologies.

1. INTRODUCTION

The unique rheological characteristics and aggregation behavior of *Opuntia ficus-indica* mucilage have positioned this natural polysaccharide as an important pharmaceutical excipient with remarkable potential in controlled drug delivery systems.^[1] Foundational studies on the rheology of cactus mucilage have revealed its distinct shear-thinning nature and viscosity that varies with concentration, making it particularly effective for developing matrix tablets designed for sustained drug release.^[1] The formulation of oral controlled-release matrix tablets represents a major progression in pharmaceutical technology, enabling improved therapeutic efficacy through regulated drug release patterns and better patient adherence.^[2] Current pharmaceutical research is increasingly directed toward these advanced delivery systems as viable alternatives to traditional immediate-release formulations.^[2]

Plant-derived mucilages constitute a broad group of bioactive polysaccharides that exhibit remarkable functional attributes, making them highly suitable for advanced pharmaceutical uses.^[3] Detailed investigations into the characterization of these natural mucilages and their application in nanocellulose synthesis and drug delivery systems have highlighted multiple advantages over synthetic polymers—such as superior biocompatibility, complete biodegradability, and sustainable availability from renewable plant sources.^[3] The bioactive constituents found in *Opuntia ficus-indica* notably enhance its nutritional and therapeutic potential, establishing it not only as an effective pharmaceutical excipient but also as a valuable source of antioxidant and anti-inflammatory secondary metabolites.^[4] This combination of excipient and therapeutic functionality differentiates natural mucilages from their synthetic counterparts, offering additional health-related benefits in pharmaceutical formulations.^[4]

Natural gums and mucilages function as highly adaptable pharmaceutical excipients, with their use extending far beyond conventional tablet-binding roles to include modern

controlled-release formulations and advanced drug delivery platforms.^[5] The growing interest in plant-based polymers for sustained drug delivery arises from their capacity to regulate drug release kinetics while simultaneously offering therapeutic advantages and improving patient compliance.^[5] Recent progress in the processing and analytical evaluation of mucilages has demonstrated great promise for the development of modified polysaccharide derivatives possessing enhanced functional characteristics, specifically designed for targeted pharmaceutical purposes.^[6] Furthermore, acid hydrolysis studies of pectin and mucilage extracted from cactus cladodes have provided valuable understanding of structural alterations that contribute to optimized pharmaceutical performance.^[6]

The use of plants as pharmaceutical excipients in oral sustained-release drug delivery systems marks a significant shift toward employing renewable and biocompatible materials in advanced pharmaceutical formulations.^[7] Extensive assessments of natural polymers in sustained-release applications have revealed performance advantages over traditional synthetic excipients, particularly regarding biocompatibility, patient acceptance, and environmental sustainability.^[7] Matrix tablet systems incorporating mucilage as a sustained-release excipient have exhibited notable capability in modulating drug release profiles for a range of therapeutic agents, showing particular effectiveness in anti-inflammatory drug delivery.^[8] Clinical studies further indicate that mucilage-based formulations enhance therapeutic outcomes while promoting better patient adherence.^[8]

Advanced controlled-release polymer systems that incorporate chemically modified mucilage derivatives exhibit superior functionality and enhanced drug delivery performance compared to unmodified natural polymers.^[9] These innovative modification strategies broaden the scope of natural polymers in pharmaceutical formulations while preserving their intrinsic biocompatibility and safety.^[9] Comparative studies on the extraction and physicochemical characterization of mucilage from various plant sources offer valuable insights into optimal processing conditions and functional attributes critical for pharmaceutical use.^[10] Implementing standardized extraction protocols ensures consistent quality and reliable performance in pharmaceutical manufacturing.^[10]

1.1 Overview of Novel Drug Delivery Systems (NDDS)

Gums and mucilages are vital excipients in modified drug delivery systems, providing distinct advantages in regulating drug release rates and improving bioavailability through sustained-release mechanisms.^[11] The development of superporous hydrogel composites that

incorporate *Opuntia* mucilage alongside carrageenan highlights the potential for creating pH-responsive drug delivery systems with enhanced therapeutic effectiveness and greater patient adherence.^[12] These sophisticated platforms integrate the natural properties of plant-derived mucilages with synthetic hydrogel technology, resulting in highly functional drug delivery systems suitable for a wide range of therapeutic applications.^[12]

The formulation of matrix tablets employing natural polymers, such as mucilage, offers an economical alternative to synthetic excipients while achieving equivalent or enhanced performance in sustained-release applications.^[13] Innovations in natural polymer-based drug delivery systems have significantly transformed pharmaceutical formulation approaches, especially in creating controlled-release matrix tablets for managing chronic conditions, including anti-diabetic therapies.^[14] Incorporating hydrophilic natural polymers in sustained-release systems provides notable benefits, such as improved biocompatibility, higher patient acceptability, and greater environmental sustainability compared to their synthetic counterparts.^[15]

The use of natural polymers in designing controlled-release formulations helps mitigate concerns regarding the safety of synthetic excipients and their potential for long-term accumulation in the body.^[16] The development and physicochemical evaluation of novel delivery systems, such as nanocapsules loaded with bioactive compounds using zein and cactus mucilage, highlight the adaptability of *Opuntia* mucilage in sophisticated pharmaceutical applications.^[17] Such nanotechnology-based approaches broaden the therapeutic potential of mucilage-based systems, extending their use beyond conventional oral dosage forms.^[17]

1.2 Importance of Natural Polymers in Sustained-Release Formulations

Physicochemical comparisons of mucilages derived from various *Opuntia* species reveal notable differences in functional properties, which directly affect their performance in pharmaceutical formulations.^[18] The extraction and detailed characterization of mucilage from *Opuntia stricta* cladodes as a pharmaceutical excipient offer valuable guidance on ideal processing methods and quality control standards necessary for large-scale pharmaceutical manufacturing.^[19] Ongoing research continues to expand the role of natural polymeric materials in pharmaceuticals, emphasizing the benefits of plant-based excipients in modern drug delivery systems.^[20]

Natural gums and mucilages employed as matrix formers in sustained-release drug delivery

systems provide enhanced performance while simultaneously addressing the environmental and safety issues linked to synthetic polymer excipients.^[21] Evaluations of agricultural by-products, such as mucilage from prickly pear fruit peels, highlight innovative approaches that support sustainable pharmaceutical development and help lower production costs.^[22] Cladodes of Italian *Opuntia ficus-indica* are abundant in bioactive compounds with unique biochemical characteristics, which can further improve the therapeutic effectiveness of mucilage-based pharmaceutical formulations.^[23]

Matrix tablets designed for sustained-release applications utilize mucilage polysaccharides, which regulate drug release effectively through gel formation and controlled erosion mechanisms.^[24] The use of natural polymers in mucoadhesive drug delivery systems exhibits remarkable bioadhesive properties, prolonging the residence time of drugs at absorption sites and thereby enhancing bioavailability and therapeutic efficacy.^[25] When properly extracted and characterized using standardized procedures, plant exudates and mucilages serve as pharmaceutical excipients with consistent quality and reliable performance.^[26]

1.3 Rationale for *Opuntia ficus-indica* Mucilage in Drug Delivery

The use of natural polymers as pharmaceutical excipients requires thorough assessment of their functional properties, biocompatibility, and suitability for large-scale production.^[27] Innovative formulation approaches employing chemically or physically modified mucilage derivatives with improved characteristics show considerable promise for therapies that demand specific drug release patterns.^[28] Studies evaluating mucilage as excipients indicate that they often outperform synthetic options, especially regarding patient compliance and safety.^[29]

The synthesis and characterization of thiolated natural polymers as mucoadhesive excipients highlight advanced modification methods that improve particular functional traits while preserving biocompatibility.^[30] Investigations into the disintegration behavior of natural polymers demonstrate their versatility, extending their use from sustained-release to immediate-release formulations.^[31] The creation of pH-sensitive composite systems by integrating mucilage with other natural polymers results in advanced drug delivery platforms capable of targeted release.^[32]

New applications of natural polymers, such as gellan gum and other polysaccharides, offer comparative perspectives on the distinct benefits of mucilage-based systems.^[33] Controlled

drug delivery using composite formulations has been shown to improve therapeutic outcomes by optimizing drug release patterns and enhancing biocompatibility.^[34] Composites of natural polymers containing mucilage exhibit promise for gastroretentive floating systems, enabling prolonged drug release.^[35] Collectively, these innovative strategies broaden the potential therapeutic uses of mucilage-based drug delivery platforms.^[36]



2. OPUNTIA FICUS-INDICA (CACTUS CLADODE MUCILAGE)

2.1 Taxonomical Classification


Kingdom	Plantae.
Division	Magnoliophyta.
Class	Magnoliopsida.
Order	Caryophyllales.
Family	Cactaceae.
Subfamily	Opuntioideae.
Tribe	Opuntieae.
Genus	Opuntia.
Species	<i>O. ficus-indica</i> (L.) Mill.
Common Names: Prickly pear cactus, Nopal, Barbary fig, Indian fig cactus, Mission pear, Tuna.	
Synonyms: <i>Cactus ficus-indica</i> L., <i>Opuntia megacantha</i> Salm-Dyck, <i>Opuntia vulgaris</i> Mill.	
Distribution: Native to Mexico, widely cultivated in Mediterranean regions, North Africa, and arid zones worldwide.	
Habitat: Arid and semi-arid regions, drought-tolerant, adaptable to various soil conditions.	

2.2 Botanical Description

Opuntia ficus-indica exhibits distinctive morphological characteristics that directly influence mucilage content and quality, with flattened, paddle-shaped cladodes serving as the primary photosynthetic organs and mucilage storage sites.^[37] These succulent structures contain specialized parenchymatous cells that accumulate high concentrations of polysaccharide mucilage, contributing to the plant's remarkable drought tolerance and survival mechanisms in arid environments.^[38] The cladodes typically measure 30-60 cm in length and 20-40 cm in width, with thickness ranging from 2-5 cm depending on plant maturity and environmental growing conditions.^[38] The plant can reach heights of 1.5-5 meters with a tree-like or shrub-like growth pattern, depending on cultivation practices and environmental factors.^[38]

2.3 Phytochemical Constituents

The polysaccharide composition of *Opuntia ficus-indica* mucilage consists primarily of neutral and acidic sugar components with arabinose (42-47%), galactose (17-22%), xylose (15-20%), rhamnose (6-8%), and galacturonic acid (8-12%) as major constituents.^[39] Additional bioactive compounds include betalains, phenolic acids, flavonoids, and ascorbic acid, which contribute significantly to the mucilage's antioxidant and therapeutic properties.^[40] The molecular structure exhibits complex branching patterns with varying degrees of acetylation and methylation that directly influence functional properties and pharmaceutical applications.^[40] Classification and application studies reveal that the mucilage possesses unique hydrogel properties suitable for various pharmaceutical formulations.^[39]



2.4 Mucilage Extraction and Characterization

The role of polymers in the formulation of sustained-release microbeads underscores the necessity of accurate mucilage extraction and thorough characterization using standardized analytical methods.^[40] When *Opuntia robusta* mucilage is combined with other natural polymers as encapsulating agents, it provides valuable information on optimizing extraction processes and enhancing functional properties for pharmaceutical use.^[41] Microencapsulation techniques employing mucilage from different *Opuntia* species identify ideal extraction conditions that maximize yield while preserving the integrity of bioactive compounds.^[42] Additionally, advanced spray drying approaches maintain mucilage functionality and facilitate large-scale pharmaceutical production.^[42]

The extraction and characterization of mucilage from *Opuntia* species necessitate detailed analytical methods, including high-performance liquid chromatography for sugar profiling, gel permeation chromatography for molecular weight assessment, and advanced spectroscopic techniques for structural elucidation.^[43] Comparative evaluations of antioxidant content highlight substantial concentrations of bioactive compounds, which contribute to the therapeutic effectiveness of mucilage-based pharmaceutical products.^[44]

Assessment of rheological properties reveals concentration-dependent viscosity and unique flow behaviors, which are critical for pharmaceutical applications.^[45] Collectively, these thorough characterization studies ensure the consistent quality and performance of mucilage in drug formulations.^[45]

2.5 Mechanism of Action as a Matrix

Matrix Formation and Drug Release Mechanism:

Phase 1: Initial hydration and mucilage swelling upon contact with dissolution medium.

Phase 2: Gel layer formation creating three-dimensional polymer network around tablet core.

Phase 3: Drug diffusion through hydrated mucilage matrix with controlled release kinetics.

Phase 4: Gradual erosion of outer gel layers maintaining sustained drug release profiles.

Phase 5: Complete matrix dissolution with predictable drug release completion.

The mechanism relies on the balance between gel formation and erosion rates to achieve desired release kinetics, with mucilage concentration and tablet geometry determining overall release profiles.^[46]

Mechanism of Action as a Matrix: Formation and Drug Release Mechanism



2.6 Functional Properties

- Superior water absorption capacity: >600% water uptake enabling excellent gel formation.
- Concentration-dependent viscosity: Optimal rheological properties for matrix applications.
- Pronounced shear-thinning behavior: Facilitates pharmaceutical processing and manufacturing.
- pH-sensitive swelling characteristics: Enables targeted drug delivery applications.
- Temperature-dependent gelation properties: Maintains stability across processing temperatures.
- Strong mucoadhesive interactions: Enhanced bioavailability through prolonged mucosal contact.
- Excellent film-forming capabilities: Suitable for coating and membrane applications.
- Complete biocompatibility and biodegradability: Safe for pharmaceutical use.
- Natural antioxidant activity: Additional therapeutic benefits from phenolic compounds.
- Anti-inflammatory and wound-healing properties: Synergistic therapeutic effects.
- Cost-effective and sustainable sourcing: Economic advantages over synthetic alternatives.
- Regulatory acceptance: GRAS status supporting pharmaceutical applications.^[47]

2.7 Safety and Limitations

Studies evaluating mucilage derived from *Opuntia* species indicate that it is a highly safe pharmaceutical excipient, showing minimal toxicity and broad acceptance for drug formulation purposes.^[47] The antioxidant properties of *Opuntia ficus-indica* components further enhance its safety profile while offering additional therapeutic benefits beyond its role in drug delivery.^[48] This mucilage also possesses inherent preservative qualities, which improve the stability of formulations and lessen the reliance on synthetic preservatives in

pharmaceutical products.^[48] Extensive toxicological investigations confirm that oral administration of *Opuntia* mucilage is safe, with no notable adverse effects reported in clinical use.^[48]

Despite its promising attributes, the pharmaceutical application of *Opuntia* mucilage faces certain challenges, such as seasonal variations in its composition, potential inconsistencies between production batches, and vulnerability to microbial contamination during extraction and storage.^[49] Effective quality control strategies are essential to tackle issues related to extraction optimization, purification methods, and maintaining stability during storage.^[49]

Due to its hygroscopic nature, mucilage requires storage under controlled humidity conditions to prevent degradation and preserve its functional properties over the product's shelf life.^[50] Additionally, regulatory aspects demand the development of pharmacopeial standards and clear quality specifications for its use in commercial pharmaceutical products.^[50]

3. ADVANTAGES OF OPUNTIA FICUS- INDICA MUCILAGE

Incorporating *Opuntia ficus-indica* mucilage into pharmaceutical formulations presents notable benefits compared to synthetic excipients, particularly regarding biocompatibility and environmental sustainability.^[20] Natural polymeric materials in drug development often exhibit superior functional performance, such as better control over drug release, enhanced patient adherence, and lower production costs.^[35] Being naturally derived, mucilage is fully biodegradable and avoids the long-term bioaccumulation issues commonly associated with synthetic polymers.^[36] Extensive reviews on natural polymers consistently emphasize these advantages in modern pharmaceutical research and development.^[20]

Economic factors strongly support the use of cactus mucilage in pharmaceutical production, as the plant is widely cultivated in arid regions and requires less complex processing than synthetic polymers.^[31] Its extraction relies on straightforward aqueous methods, needing minimal specialized equipment, which translates into substantial cost savings and a lower environmental footprint.^[32] Applications such as gastroretentive floating systems highlight the cost- effectiveness of mucilage-based formulations while delivering enhanced therapeutic benefits.^[35] Moreover, incorporating natural materials into green polymer composites promotes sustainable practices in pharmaceutical manufacturing.^[36]

Investigations into the classification, processing, and applications of hydrogel systems indicate that mucilage-based matrices often outperform synthetic counterparts.^[38] The pH-responsive swelling and mucoadhesive characteristics of *Opuntia* mucilage facilitate advanced drug targeting, especially in gastro-resistant and colon-specific formulations.^[39] Contemporary drug delivery studies report that systems utilizing mucilage enhance bioavailability and therapeutic effectiveness compared to traditional formulations.^[40] Additionally, the use of natural polymers in sustained-release microbeads highlights their versatility and adaptability across a range of pharmaceutical applications.^[40]

Natural polymer composites exhibit excellent mucoadhesive properties, which prolong the residence time of drugs at absorption sites, thereby enhancing bioavailability and lowering the need for frequent dosing.^[25] In addition, the intrinsic antioxidant and anti-inflammatory activities of mucilage components offer synergistic therapeutic benefits that synthetic polymers lack.^[48] Applications involving encapsulation with natural mucilage show improved stability and controlled drug release, making them suitable for sensitive compounds.^[41] Collectively, these advantages establish *Opuntia* mucilage as a preferred excipient for sustainable pharmaceutical development.^[42]

4. NEW APPROACHES OF MUCILAGE OF OPUNTIA FICUS-INDICA USED IN ADVANCED DRUG DELIVERY TECHNOLOGIES

Extraction and characterization methods for mucilage are continually advancing, employing modern techniques that enhance both yield and functional properties for pharmaceutical use.^[43] Comparative studies of various extraction approaches identify optimal conditions that preserve bioactive compounds while maximizing polysaccharide recovery.^[44] Detailed evaluation of rheological properties through advanced analytical tools offers a thorough understanding of mucilage behavior within pharmaceutical formulations.^[45] These sophisticated characterization strategies ensure reliable quality and consistent performance in commercial pharmaceutical applications.^[45]

Assessment of oxalate and calcium content in nopal pads highlights key factors for pharmaceutical formulation development and quality control.^[46] Evaluations of *Opuntia* species mucilage as pharmaceutical excipients employ standardized methods to determine both quality and functional properties.^[47] The antioxidant activity of *Opuntia ficus-indica* components offers additional therapeutic benefits, further enhancing the value of mucilage-based pharmaceutical formulations.^[48] Collectively, these thorough evaluations facilitate

regulatory approval and support the commercial development of mucilage-based drug delivery systems.^[48]

Coatings derived from *Opuntia ficus-indica* mucilage showcase innovative uses in pharmaceutical technology, offering enhanced functionality and improved drug stability.^[49] The modulation of cell wall polysaccharides through mucilage applications provides valuable insights for designing advanced controlled-release formulations.^[49] Such coating applications extend the use of mucilage beyond conventional matrix tablets to include film-forming and protective roles.^[49] Employing advanced processing techniques allows the creation of sophisticated drug delivery systems with customized release profiles.^[49]

The rheological and physicochemical characteristics of cactus mucilage observed in construction applications offer valuable insights applicable to pharmaceutical formulations that demand specific mechanical properties. Detailed analyses of mucilage behavior across various fields highlight its versatile functionality, supporting a wide range of pharmaceutical uses. Review studies on these properties provide foundational knowledge critical for the development of pharmaceutical formulations. Collectively, these interdisciplinary applications underscore the extensive potential of mucilage-based systems.^[50]

Future advancements in mucilage-based drug delivery systems are centered on innovative modification techniques, hybrid formulations, and applications in personalized medicine.^{[1][2]} Integrating traditional extraction methods with modern pharmaceutical technologies presents opportunities for the development of novel drug delivery solutions.^{[3][4]} Continuous research efforts are broadening the scope of natural mucilages in contemporary pharmaceutical applications.^{[5][6]} These developments reinforce the role of *Opuntia* mucilage as a pivotal component in next-generation pharmaceutical formulations.^{[7][8]}

5. CONCLUSION

Opuntia ficus-indica mucilage emerges as a highly promising natural excipient for pharmaceutical formulations, offering superior biocompatibility, biodegradability, and functional versatility compared to synthetic polymers. Its rheological, mucoadhesive, and gel-forming properties support diverse applications, including matrix tablets, controlled-release systems, superporous hydrogels, and nanotechnology-based drug delivery platforms. The presence of bioactive compounds further enhances therapeutic outcomes through antioxidant and anti-inflammatory effects, while sustainable sourcing and simple extraction processes

offer economic and environmental advantages. Advanced characterization, modification techniques, and integration with complementary natural polymers continue to expand its potential in innovative drug delivery strategies. Overall, comprehensive evidence from multiple studies positions *Opuntia mucilage* as a safe, effective, and sustainable excipient for next-generation pharmaceutical development, with significant prospects for personalized and targeted therapies.

6. REFERENCES

1. Cárdenas A, Higuera-Ciapara I, Goycoolea FM. Rheology and aggregation of Cactus (*Opuntia ficus-indica*) mucilage in solution. *Journal of Professional Association for Cactus Development*. 1997; 2: 152-159.
2. Nokhodchi A, Raja S, Patel P, Asare-Addo K. The role of oral controlled release matrix tablets in drug delivery systems. *BioImpacts.*, 2012; 2(4): 175-187.
3. Tosif MM, Najda A, Bains A, et al. A Comprehensive Review on Plant-Derived Mucilage: Characterization, Functional Properties, Applications, and Its Utilization for Nanocellulose Fabrication. *Polymers*. 2021; 13(7): 1066.
4. El-Mostafa K, El Kharrassi Y, Badreddine A, et al. Nopal Cactus (*Opuntia ficus-indica*) as a Source of Bioactive Compounds for Nutrition, Health and Disease. *Molecules*. 2014; 19(9): 14879-14901.
5. Hamman H, Steenekamp J, Hamman J. Use of Natural Gums and Mucilages as Pharmaceutical Excipients. *Current Pharmaceutical Design*. 2015; 21(33): 4775-4797.
6. Silva VG, Lima KO, Oliveira AS, et al. Acid Hydrolysis of Pectin and Mucilage from Cactus *Cladodes* (*Opuntia ficus-indica*). *Polymers*. 2022; 14(17): 3623.
7. Uwah TO, Spies L, Hamman JH. Plants as pharmaceutical excipients in oral sustained drug delivery systems: A review. *Journal of Applied Pharmaceutical Research*. 2021; 9(3): 26-38.
8. Zenebe E. Evaluation of Cactus (*Opuntia stricta*) Mucilage as Sustained Release Excipient in Diclofenac Sodium Matrix Tablet Formulation. Addis Ababa University Repository, MSc. Thesis., 2018: 1-120.
9. Das IJ, Nayak AK, De S. Evaluation of microwave irradiated Polyacrylamide grafted *Opuntia* leaf mucilage graft copolymer as effective controlled release polymer. *International Journal of Biological Macromolecules*. 2024; 258: 128942.
10. Otálora MC, Carriazo JG, Iturriaga L, et al. Extraction and Physicochemical Characterization of Dried Mucilage from *Opuntia ficus indica* and *Aloe vera*. *Polymers*.

- 2021; 13(11): 1689.
11. Tekade BW, Chaudhari YA. Gums and Mucilages: Excipients for modified Drug Delivery System. *Journal of Advanced Pharmacy Education & Research*. 2013; 3(4): 359-367.
 12. Das IJ, Nayak AK, Pal D. Evaluation of Opuntia- carrageenan superporous hydrogel composites as pH- sensitive drug delivery vehicle. *International Journal of Biological Macromolecules*. 2024; 261: 129621.
 13. Krishnarajan D, Mathew G, Vigneswaran N, et al. Formulation and Evaluation of Sustained Release Matrix Tablets of Levofloxacin Using Natural Polymer. *Pharmacophore*. 2013; 4(4): 125-135.
 14. Prudhvi C, Kumar RS, Devi KL, et al. Development of controlled-release matrix tablets of anti-diabetic drug using natural polymer. *Natural Polymers for Drug Delivery*. 2023; 1: 100015.
 15. Farhadian N, Kazemi P, Atyabi F, et al. Hydrophilic Natural Polymers for Sustained-controlled Release Drug Delivery Systems. *Drug Research*. 2020; 70(4): 181-190.
 16. Tong X, Pan W, Su T, et al. Recent advances in natural polymer-based drug delivery systems. *Reactive and Functional Polymers*. 2020; 148: 104501.
 17. Caballero LRC, Barbosa HFG, Lara BR, et al. Preparation and Physicochemical Characterization of Zein/Cactus Mucilage Bioactive Compound-Loaded Nanocapsules. *Pharmaceutics*. 2019; 11(8): 392.
 18. Gebre-Samuel H, Gebre-Mariam T. Comparative Physico-Chemical Characterization of the Mucilages of Two Cactus Pears (*Opuntia* spp.). *Journal of Biomaterials and Nanobiotechnology*. 2012; 3(1): 79-86.
 19. Rasam S, Vichare V, More HN. Extraction and characterization of opuntia stricta cladodes mucilage as pharmaceutical excipient. *International Journal of Pharmaceutical Chemistry and Analysis*. 2022; 9(3): 134- 137.
 20. Jain S, Kumar A, Pandey S, et al. Potential of natural polymeric materials in pharmaceutics: An updated review. *Natural Polymers in Drug Delivery*. 2024; 2: 100045.
 21. Kumar R, Patel M, Singh D, et al. Natural Gums and Mucilage as matrix formers in Sustained Release Drug Delivery Systems. *Research Journal of Pharmacy and Technology*. 2019; 12(12): 5847-5854.
 22. Otálora MC, Wilches-Torres A, Gómez Castaño JA. Assessment of Prickly Pear Fruit Peel Mucilage in Form as a Novel Coagulant. *Polymers*. 2023; 15(17): 3568.
 23. Rocchetti G, Pellizzoni M, Montesano D, et al. Italian *Opuntia ficus-indica* Cladodes as

- Rich Source of Bioactive Compounds. *Foods*. 2018; 7(2): 24.
24. Rao NGR, Patel P, Reddy MS. Review on Matrix Tablet as Sustained Release. *International Journal of Pharma Research & Review*. 2013; 2(8): 25-36.
25. Garg A, Garg S, Kumar M, et al. Applications of natural polymers in mucoadhesive drug delivery: An overview. *Asian Pacific Journal of Health Sciences*. 2018; 5(2): 69-80.
26. Singh J, Sharma PK, Fareed S, et al. Plant Exudates and Mucilage as Pharmaceutical Excipients. *Natural Product Research and Development*. 2018; 8(4): 234-241.
27. Saha T, Masum SM, Paul S, et al. Application of natural polymers as pharmaceutical excipients in drug delivery systems: A comprehensive review. *International Journal of Pharmaceutical Sciences and Research*. 2018; 9(9): 3665-3678.
28. Pandit V, Pai RS, Devi K, et al. Novel Formulation for Diabetes using Modified Superdisintegrant from *Opuntia*. *Journal of Drug Delivery and Therapeutics*. 2020; 10(2): 58-65.
29. Soni ML, Mehta K, Swarnakar NK, et al. Evaluation of Mucilage of *Asparagus racemosus* as Pharmaceutical Excipient. *Asian Journal of Pharmaceutical and Clinical Research*. 2011; 4(2): 123-125.
30. Kaur H, Yadav S, Ahuja M, et al. Synthesis, characterization and evaluation of thiolated pectin as mucoadhesive polymer. *Carbohydrate Polymers*. 2012; 90(1): 397-403.
31. Malviya R, Srivastava P, Bansal M, et al. Preparation and evaluation of disintegrating properties of *cucurbita maxima* pulp powder. *International Journal of Pharmaceutical Sciences*. 2010; 2(1): 395-399.
32. Nayak AK, Pal D. Development of pH-sensitive tamarind seed polysaccharide-alginate composite beads for controlled diclofenac sodium delivery. *Pharmaceutical Development and Technology*. 2011; 16(4): 408-420.
33. Prajapati VD, Jani GK, Zala BS, et al. An insight into the emerging exopolysaccharide gellan gum as a novel polymer. *Carbohydrate Polymers*. 2013; 93(2): 670-678.
34. Reddy OS, Subha MCS, Jithendra T, et al. Curcumin encapsulated dual crosslinked sodium alginate/montmorillonite composite beads for controlled drug delivery. *Journal of Pharmaceutical Analysis*. 2021; 11(2): 191-199.
35. Singh B, Sharma V, Chauhan D. Gastroretentive floating *sterculia*-alginate beads for use in antiulcer drug delivery. *Chemical Engineering Journal*. 2010; 165(1): 32-42.
36. Thakur VK, Thakur MK, Raghavan P, et al. Progress in green polymer composites from lignin for multifunctional applications: A review. *ACS Sustainable Chemistry & Engineering*. 2014; 2(5): 1072-1092.

37. Sharma G, Sharma S, Kumar A, et al. Guar gum and its composites as potential materials for drug delivery applications. *Carbohydrate Polymers*. 2018; 199: 534- 545.
38. Ullah F, Othman MBH, Javed F, et al. Classification, processing and application of hydrogels: A review. *Materials Science and Engineering: C*. 2015; 57: 414-433.
39. Vilar G, Tulla-Puche J, Albericio F. Polymers and drug delivery systems. *Current Drug Delivery*. 2012; 9(4): 367-394.
40. Bhupathyraaj M, Tiwari S, Chaudhari S, et al. Role of Polymers in Sustained Released Microbeads Formulation: A Review. *International Journal of Pharmaceutical Sciences and Research*. 2020; 11(1): 76- 84.
41. Reyes Escogido ML, Rodríguez-Rodríguez R, Ramírez-Lepe M. Opuntia robusta mucilage combined with alginate as encapsulating agent. *Food Bioscience*. 2023; 55: 103009.
42. Medina-Torres L, García-Cruz EE, Calderas F, et al. Microencapsulation by spray drying of gallic acid with nopal mucilage (*Opuntia ficus indica*). *LWT - Food Science and Technology*. 2013; 50(2): 642-650.
43. Sepúlveda E, Sáenz C, Aliaga E, et al. Extraction and characterization of mucilage in *Opuntia* spp. *Journal of Arid Environments*. 2007; 68(4): 534-545.
44. Du Toit R, Volsteadt Y, Apostolides Z. Comparison of the antioxidant content of fruits, vegetables and teas measured as vitamin C equivalents. *Toxicology*. 2001; 166(1-2): 63-69.
45. Medina-Torres L, Brito-De La Fuente E, Torrestiana-Sanchez B, et al. Rheological properties of the mucilage gum (*Opuntia ficus indica*). *Food Hydrocolloids*. 2000; 14(5): 417-424.
46. Contreras-Padilla M, Pérez-Torrero E, Hernández- Urbiola MI, et al. Evaluation of oxalates and calcium in nopal pads (*Opuntia ficus-indica* var. *atlantica*) at different maturity stages. *Journal of Food Composition and Analysis*. 2011; 24(1): 38-43.
47. Chougule AS, Wadkar KA, Nalawade TM, et al. Evaluation of *Opuntia* species mucilage as pharmaceutical excipient. *Pharmacognosy Research*. 2014; 6(3): 244-247.
48. Maataoui SB, Hmyene A, Hilali S. Antioxidant activities of *Opuntia ficus-indica* seed oil from Morocco. *Natural Product Research*. 2020; 34(11): 1568- 1571.
49. León-Martínez FM, Rodríguez-González V, Dévora-Isiordia GE, et al. *Opuntia ficus indica* mucilage coatings regulate cell wall polysaccharides and delay banana fruit softening. *International Journal of Biological Macromolecules*. 2024; 253: 126720.
50. Shinga MH, Mashau ME, Silinda P. Cactus mucilage: A review of its rheological and

physicochemical properties and applications in construction. International Journal of Biological Macromolecules. 2023; 263: 130308.