Mutation Breeding in Coriander: Mutagenic Agents and Application Techniques

Salve K. M.

Department of Botany, Pemraj Sarda College, Ahilyanagar

Email: skbotany@gmail.com

Article DOI Link: https://zenodo.org/uploads/19661194

DOI: 10.5281/zenodo.19661194

Abstract

A significant spice crop, *Coriandrum sativum* L. (coriander) is grown extensively for its leaves, seeds, and essential oils. Mutation breeding is a useful technique for increasing coriander yield, its quality, and disease resistance. The use of physical and chemical mutagens in coriander breeding, as well as the ensuing genetic variability and enhanced features.

Keywords: *Coriandrum sativum* L., Mutation breeding, Physical and chemical mutagens, genetic variability.

Introduction

The genetic improvement of coriander (*Coriandrum sativum* L.) relies heavily on the presence of genetic variability. When natural variation is limited, mutation breeding serves as a powerful tool to induce novel traits. By employing physical and chemical agents, breeders can disrupt the genetic code to create "hopeful monsters"- variants that may possess superior yield, higher essential oil content, or better disease resistance.

Classification of Mutagenic Agents

Mutagens are broadly categorized into physical and chemical agents. In coriander research, these tools are selected based on their ability to penetrate seed tissues and their efficiency in inducing chromosomal or point mutations.

A. Physical Mutagens (Ionizing Radiation)

Physical mutagens typically involve ionizing radiation, which strips electrons from atoms, leading to DNA strand breaks.

- **Gamma Rays:** These are the most widely used physical mutagens in coriander breeding. Emitted from sources like Cobalt-60 (^ {60} Co), they have high penetrating power.

- **Dosage Dynamics:** Research indicates that the dosage of radiation is critical. For coriander, doses ranging from 10 kR to 25 kR (kilo-Roentgens) are standard.
- **Biological Impact:** While higher doses increase the likelihood of mutation, they often lead to a "biological cost" in the M_1 generation, such as reduced pollen fertility and stunted growth. Moderate doses are generally preferred to balance mutation frequency with plant survival.

Chemical Mutagens

Chemical mutagens are often preferred for inducing point mutations (single nucleotide changes) rather than large-scale chromosomal aberrations.

- **Alkylating Agents:** This group includes Ethyl Methane Sulphonate (EMS) and Methyl Methane Sulphonate (MMS). They work by adding ethyl or methyl groups to DNA bases, leading to mispairing during replication.
- **Sodium Azide (SA):** A potent chemical mutagen known for its high efficiency and low toxicity to the plant's overall physiology. It is frequently used to create variations in seed size and aromatic properties.
- **Comparison:** Some studies suggest that MMS can be particularly effective because it can induce desired mutations at lower concentrations with less biological damage to the embryo compared to harsher chemicals.

Synergy through Combination Treatments

To maximize the “mutation spectrum” – the variety of different types of mutations produced – breeders often employ combination treatments. By exposing coriander seeds to a physical mutagen (like Gamma rays) followed by a chemical one (like EMS), the cumulative effect can increase mutation frequency. This approach aims to hit different parts of the genome or utilize different repair mechanisms within the cell to lock in beneficial traits.

The Relationship between Dose and Response

A fundamental principle in mutation breeding is the inverse relationship between the mutagen dose and plant vitality.

- **Germination and Growth:** As the concentration of a chemical or the intensity of radiation increases, the germination rate typically decreases.
- **Seedling Vigor:** High doses often result in slower seedling growth and higher lethality (the LD_ {50} or Lethal Dose 50% is a common benchmark used to determine the ideal treatment level).
- **Optimal Selection:** The goal is to find the "sweet spot" where the dose is high enough to cause genetic change but low enough to allow the M_1 generation to survive and produce seeds (M_2) for further selection.

Application Techniques and the M_1 Generation

The process of inducing mutations is a precise laboratory to field procedure. For chemical mutagens, the following protocol is generally standard:

- **Pre-soaking:** Seeds are soaked in distilled water (usually for 6 – 12 hours). This softens the seed coat and initiates metabolic activity, making the DNA more susceptible to the chemical agent.
- **Chemical Treatment:** The pre – soaked seeds are immersed in the mutagenic solution (EMS, SA, or MMS) at a specific concentration and temperature.
- **Washing and Drying:** After the treatment period, seeds are thoroughly washed to remove residual chemicals and then air-dried.
- **Sowing the M_1 Generation:** The treated seeds are sown in the field to raise the M_1 generation. This generation is primarily used to observe the direct effects of the mutagen and to harvest seeds for the M_2 generation, where recessive mutations finally become visible.

Screening and Selection in the M_2 Generation

While the M_1 generation shows the immediate physiological impact of the mutagens, the M_2 generation is where the real genetic "treasure hunt" begins. Because most induced mutations are recessive, they only manifest visually when the plant inherits the mutated gene from both parents, which typically occurs during the segregation that happens in the second generation.

Visual Macro – Mutation Screening

The first step in the M_2 generation is the identification of macro – mutations – changes that are easily visible to the naked eye. In coriander, breeders look for:

- **Morphological Variants:** Changes in leaf shape (e.g., feathery vs. broad), plant height (dwarfing), or branching patterns.
- **Chlorophyll Mutations:** Variations in leaf color (albino, xantha, or chlorina), which serve as indicators of the mutagen's effectiveness, even if these specific plants aren't commercially viable.
- **Phenological Changes:** Plants that flower significantly earlier or later than the control group.

Quantitative Micro-Mutation Analysis

Not all improvements are visible. Micro – mutations involve small, cumulative changes in polygenic traits. These require statistical analysis and precise measurement:

- **Yield Components:** Measuring the number of umbels per plant, the number of seeds per umbellate, and the 1000 seed weight.

- **Essential Oil Content:** Using techniques like Hydro – distillation or Gas Chromatography (GC) to determine if the mutation has enhanced the concentration of Linalool, the primary aromatic compound in coriander.

Biotic and Abiotic Stress Screening

Mutant populations are often subjected to "stress tests" to identify resilient lines:

- **Disease Resistance:** Screening for resistance to common coriander pathogens like powdery mildew (Erysiphe polygoni) or stem gall.
- **Drought and Salinity:** Testing the M_2 and M_3 lines in controlled environments to find variants that can thrive in marginal soils or with less water.

Verification in the M_3 and M_4 Generations

Once a promising M_2 plant is identified, its seeds are harvested and sown in "plant to grow" progenies in the M_3 generation. This step is crucial to confirm that the observed trait is truly heritable and not just a temporary environmental fluke. If the trait remains stable and uniform through the M_4 generation, it is officially considered a "mutant line" and may be fast – tracked for variety release.

Comparison and Conclusion

To wrap up this chapter on coriander mutation breeding, it is essential to compare the primary tools at a breeder's disposal and summarize the strategic importance of these techniques.

1. Comparative Analysis: EMS vs. Gamma Radiation

Choosing between a physical and chemical mutagen depends on the breeder's specific goals. Below is a comparison of the two most common agents used in coriander research.

Feature	Gamma Radiation (Physical)	Ethyl Methane Sulphonate (EMS)
Primary Action	Physical energy (Ionizing radiation)	Chemical alkylation
Genetic Change	Large chromosomal aberrations/deletions	Precise point mutations (SNPs)
Ease of Use	Requires specialized facility (Gamma cell)	Can be done in a standard lab
Penetration	Uniform and high	Dependent on seed soaking/permeability
Coriander Impact	Effective for yield and structural traits	Excellent for oil quality and leaf traits
Handling Risk	High (Radiation safety protocols)	High (Carcinogenic; requires fume hood)

2. Summary and Future Outlook

The application of mutagenic agents like Gamma rays, EMS, and Sodium Azide has revolutionized our ability to "engineer" coriander. By carefully controlling the dosage and following rigorous pre – soaking and screening protocols, researchers can bypass the limitations of natural selection.

As we move forward, the integration of traditional mutation breeding with modern molecular markers will allow for even faster identification of desirable mutants. Instead of waiting for the M_2 generation to grow to maturity, breeders may soon be able to scan the DNA of young seedlings for specific genetic markers associated with high Linalool content or disease resistance.

3. Practical Takeaways for the Breeder

- **Start with the LD_ {50}:** Always determine the lethal dose for your specific coriander variety before starting a large-scale project.
- **Prioritize the M_2:** Devote the most resources to the second generation, as this is where recessive traits emerge.
- **Quality over Quantity:** A few stable, high-quality mutants in the M_3 generation are more valuable than hundreds of unstable variants in the M_1.

References

1. Andersson, L. (2008) Swedish University of Agricultural Sciences, Uppsala, Sweden.
2. Bhosale R. S. (2014) Genetic improvement in Withania somnifera Dunal, Ph.D Thesis, SPPU. Pune.
3. Gottschalk W and Wolf G (1983) Induced mutations in plant breeding. Monographs on theoretical and applied genetics, Berlin, Springer Verlag.7: 238.
4. Salve K. M. (2021) Improvement in seed germination percentage and productivity of Coriandrum sativum L. through seed priming. International Journal of Researches in Bioscience, Agriculture and Technology, issue (special – 17), 67-70.
5. Salve K. M. and More A. D. (2018), Biochemical study of Dark Green and Luxuriant Mutants in Coriandrum sativum Linn., International Journal of Researches in Bioscience, Agriculture and Technology, 6(special issue 2), 100-103.
6. Salve K. M. and More A. D. (2019) Biochemical investigation of Tall and Dwarf mutants in Coriander. International Journal of Research and Analytical Reviews; 6(2): 643i-644i.
7. Verma, A., Pandeya, S. N., Yadav, S. K., Singh, S. and Soni, P. (2011) A Review on Coriandrum sativum (Linn.): An Ayurvedic Medicinal Herb of Happines. Journal of Advances in Pharmacy and Healthcare Research 1(3): 28-48.
8. Wallis, T. E., (1985). Text book of Pharmacognosy, (Fifth edition), Published by S.K. Jain for CBS publishers and Distributors, Delhi 246 -248.