

RESEARCH ARTICLE

Transcriptome-Wide Identification of Salt-Responsive Members of the *WRKY* Gene Family in *Gossypium aridum*

Xinqi Fan^{1,2}, Qi Guo², Peng Xu², YuanYong Gong², Hongmei Shu², Yang Yang², Wanchao Ni², Xianggui Zhang², Xinlian Shen^{2*}

1 State Key Laboratory of Crop Genetics & Germplasm Enhancement, Nanjing Agricultural University, Nanjing, China, **2** Key Laboratory of Cotton and Rapeseed (Nanjing), Ministry of Agriculture, P. R. China, The Institute of Industrial Crops, Jiangsu Academy of Agricultural Sciences, Nanjing, China

* xlshen68@126.com



OPEN ACCESS

Citation: Fan X, Guo Q, Xu P, Gong Y, Shu H, Yang Y, et al. (2015) Transcriptome-Wide Identification of Salt-Responsive Members of the *WRKY* Gene Family in *Gossypium aridum*. PLoS ONE 10(5): e0126148. doi:10.1371/journal.pone.0126148

Academic Editor: Zonghua Wang, Fujian Agriculture and Forestry University, CHINA

Received: November 11, 2014

Accepted: March 30, 2015

Published: May 7, 2015

Copyright: © 2015 Fan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by grants from the Major Project of the National Transgene (2014ZX08005-004), the National Natural Science Foundation of China (NSFC) (31171595) and the Jiangsu Independent Innovation Funds of Agricultural Technology (CX (13)3059).

Competing Interests: The authors have declared that no competing interests exist.

Abstract

WRKY transcription factors are plant-specific, zinc finger-type transcription factors. The WRKY superfamily is involved in abiotic stress responses in many crops including cotton, a major fiber crop that is widely cultivated and consumed throughout the world. Salinity is an important abiotic stress that results in considerable yield losses. In this study, we identified 109 WRKY genes (*GarWRKYs*) in a salt-tolerant wild cotton species *Gossypium aridum* from transcriptome sequencing data to elucidate the roles of these factors in cotton salt tolerance. According to their structural features, the predicted members were divided into three groups (Groups I–III), as previously described for *Arabidopsis*. Furthermore, 28 salt-responsive *GarWRKY* genes were identified from digital gene expression data and subjected to real-time quantitative RT-PCR analysis. The expression patterns of most *GarWRKY* genes revealed by this analysis are in good agreement with those revealed by RNA-Seq analysis. RT-PCR analysis revealed that 27 *GarWRKY* genes were expressed in roots and one was exclusively expressed in roots. Analysis of gene orthology and motif compositions indicated that WRKY members from *Arabidopsis*, rice and soybean generally shared the similar motifs within the same subgroup, suggesting they have the similar function. Overexpression-*GarWRKY17* and –*GarWRKY104* in *Arabidopsis* revealed that they could positively regulate salt tolerance of transgenic *Arabidopsis* during different development stages. The comprehensive data generated in this study provide a platform for elucidating the functions of WRKY transcription factors in salt tolerance of *G. aridum*. In addition, *GarWRKYs* related to salt tolerance identified in this study will be potential candidates for genetic improvement of cultivated cotton salt stress tolerance.

Introduction

Plants have developed a series of complex, effective systems to protect themselves from a variety of adverse environmental conditions during growth over the long course of evolution [1].

Studies have shown that these coping mechanisms mainly function via transcriptional activation or inhibition of transcription-related genes. Transcription factors play an important role in this process by activating or inhibiting the expression of target genes alone or via interactions with other proteins [2–4].

WRKY transcription factors are recently identified plant-specific transcription factors comprising one or two WRKY domains and a zinc-finger motif. The name WRKY is derived from the highly conserved WRKY domain, which consists of 60 amino acids. The WRKY domain comprises the highly conserved WRKYGQK sequence followed by a novel zinc-finger motif (C_2H_2 or C_2HC) located in the N-terminus [5]. Based on the number of WRKY domains and the pattern of zinc-finger motifs, WRKY proteins can be classified into three main groups and eight subgroups [5]. Group I WRKYs typically have two WRKY domains containing a C_2H_2 zinc-finger motif. The C-terminal WRKY domain has DNA binding activity. However, the N-terminal WRKY domain cannot bind to DNA alone, but it can assist the C-terminal WRKY domain in its binding to DNA, increasing its DNA binding affinity and specificity. In addition, the zinc-finger motif followed by the N-terminal WRKY domain may provide a protein–protein interaction interface; the zinc-finger motif's structure is C_2H_2 ($C-X_{4-5}-C-X_{22-23}-H-X-H$, X is any amino acid). Group II WRKYs have a single WRKY domain with a C_2H_2 zinc-finger motif, which is the same as the C-terminal domain of Group I. Based on the phylogeny of the WRKY domains, group II WRKYs can be further divided into five subgroups: IIa ($C-X_5-CX_{23}-HXH$), IIb ($C-X_5-CX_{23}-HXH$), IIc ($C-X_4-CX_{23}-HXH$), IId ($C-X_5-CX_{23}-HXH$) and IIe ($C-X_5-CX_{23}-HXH$). Group III WRKYs also have a single WRKY domain, but their zinc-finger motif structure is C_2HC ($C-X_7-C-X_{23}-H-X-C$). Based on the zinc finger structure, Group III can be further divided into two subgroups: IIIa and IIIb. The structure of subgroup IIIa zinc fingers is $C-X_7-C-X_{23}-HXC$, while that of subgroup IIIb is $C-X_7C-Xn-HXC$ ($n \geq 24$) [6]. WRKYs specifically bind to the W-box, i.e., (T) (T) TGAC (C/T), in the promoter regions of target genes to regulate their transcription levels, enabling them to perform their biological functions [5].

Since the first report of *SPF1* WRKY transcription factors from sweet potato in 1994 [7], numerous WRKY transcription factors have been experimentally identified from many other plant species including *Arabidopsis*, rice, tobacco, potato, cotton, barley, wheat, chamomile, soybean, cacao, grape, tomato, cucumber and so on [8–20]. Except for plant development, WRKY family genes also play important roles in plant biotic and abiotic stress responses including pathogen-induced defense programs and responses to drought, salt, stress et al. [20–22]. One of the most important functions of WRKYs appears to be regulation of the salt stress response. For example, *AtWRKY25* and *AtWRKY33* play a role in the salt stress response and improve salt tolerance in *Arabidopsis* [23]. *OsWRKY11* and *OsWRKY45* play a regulatory role in salt tolerance in transgenic rice [24,25]. *GmWRKY54*-overexpressing plants are more salt and drought tolerant than the wild type, and *GmWRKY13* overexpression results in increased sensitivity to salt and mannitol stress [26]. Finally, overexpression of *TaWRKY10* enhances drought and salt tolerance in transgenic tobacco plants [27].

In recent years, some WRKY transcription factors have also been reported in cotton. Guo et al. isolated *GhWRKY3* from upland cotton (*G. hirsutum*), which is upregulated by the application of various phytohormones but not by cytokinin, auxin analog, drought, NaCl or cold [18]. Yu et al. identified *GhWRKY15* from *G. hirsutum*, which is involved in disease resistance and plant development [28]. Shi et al. isolated a putative IId WRKY gene, *GhWRKY39*, from *G. hirsutum* and showed that it is activated by pathogens and salt stress. *GhWRKY39*-overexpressing *Arabidopsis* plants display enhanced tolerance to salt [29]. Zhou et al. identified 26 genes encoding putative WRKY proteins in *G. hirsutum*; five *GhWRKY* genes are upregulated in roots treated with NaCl [30]. With the release of the *G. raimondii* genome sequence, two comprehensive studies of *G. raimondii* WRKY family members were conducted, revealing 120

and 116 *GrWRKY* genes and 109 WRKY genes in *G. arboreum* [31–33], respectively; the expression of these genes under abiotic stress was analyzed in *G. hirsutum* [31,32]. All of these results suggest that WRKY genes are involved in the response to salinity stress in cotton.

High salinity is a major abiotic stress in cotton production worldwide. However, as described in detail previously [34], modern cotton cultivars, which are often grown under unstressed conditions, are the result of intensive selection to produce large amounts of specific types of fiber. Selection has unintentionally narrowed the genetic variability for salt tolerance. *G. aridum* is a D-genome diploid species from the Pacific coastal states of Mexico that shows remarkably tolerance to salt stress. Understanding how *G. aridum* responds to and develops tolerance to salt stress is the first step towards improving the adaptation of cotton to salt stress. In a previous study, we performed deep sequencing analysis of the *G. aridum* transcriptome in response to salt stress. A total of 281 unigenes were annotated as WRKY family members. Differentially expressed genes encoding transcription factors belonging to the WRKY family were predominant during different stages of salt stress [34]. These results suggest that WRKY family members are important regulators of the response to salt stress in *G. aridum*. In the current study, we focused on whole transcriptome-wide identification of salt-responsive members of the WRKY family in *G. aridum*. The results of this study increase our understanding of the *GarWRKY* gene family, which will facilitate the genetic improvement of salt stress tolerance in cotton.

Materials and Methods

Plant materials and treatment conditions

Seeds from the wild *Gossypium* species *G. aridum* were kindly supplied by the National Wild Cotton Plantation in Hainan Island, China. The same treatment procedure was used as described in Xu et al. [34]. *G. aridum* plants were treated with 200 mM NaCl for 0, 1, 3, 6, 12, 24 and 72 h. Root and leaf tissues were collected at every stage of stress treatment. All tissues were immediately frozen in liquid nitrogen and stored at -70°C .

GarWRKY gene identification and chromosomal location

The annotated genome sequences of *G. raimondii* were downloaded from <http://www.phytozome.net/cn/>. The WRKY domain (PF03106) downloaded from PFAM protein family database was used as a query for identifying WRKY transcription factors by HMMER software version 3.0 [35]. The annotated *G. raimondii* WRKY genes were used to establish a local nucleic acid database. Based on de novo transcriptome sequencing data with a pooled RNA sample from roots and leaves of 12 h salt-stressed and unstressed plants [34], more than 90,000 transcripts were identified and used to BLAST the *G. raimondii* WRKY gene database using a local BLAST program with E-value $< 10^{-10}$. Redundant sequences were removed. The *GarWRKY* protein sequences were further analyzed to confirm the presence of WRKY domains using the InterPro program (protein sequence analysis & classification, <http://www.ebi.ac.uk/interpro/>) [36].

To determine the location of *GarWRKY* genes on chromosomes, the *GarWRKY* sequences were further used as query sequences for a BLASTN search against *G. raimondii* whole-genome scaffold data (<http://www.phytozome.net>). Finally, the locations of all 109 *GarWRKY*s in the *G. raimondii* genome were detected. Mapping of *GarWRKY* genes was performed using MapInspect (http://www.plantbreeding.wur.nl/UK/software_mapinspect.html). To better identify and classify the WRKY transcription factors (TFs) in *Gossypium*, a uniform nomenclature was assigned to the WRKY genes in *G. aridum* as in *G. raimondii* [31].

GarWRKY gene classification and phylogenetic analysis

To classify *GarWRKY* genes into different groups and subgroups, a total of 14 *AtWRKY* genes representing three main groups and seven subgroups were selected, including Group I (*AtWRKY7* and *AtWRKY28*), Group IIa (*AtWRKY25* and *AtWRKY27*), Group IIb (*AtWRKY28* and *AtWRKY36*), Group IIc (*AtWRKY40* and *AtWRKY41*), Group IId (*AtWRKY42* and *AtWRKY49*), Group IIE (*AtWRKY58* and *AtWRKY60*) and Group III (*AtWRKY69* and *AtWRKY70*). Complete WRKY domains from all *GarWRKY* proteins and select *AtWRKY* proteins were used to align WRKY domains with ClustalW [37]. The consequential alignment was used to create a phylogenetic tree in MEGA v4.0 using the neighbor-joining (NJ) method with 1,000 bootstrap replications [38].

To assess structural divergence of WRKY genes in different species, the Multiple Expectation Maximization for Motif Elicitation program (MEME, <http://meme.nbcr.net/meme/tools/meme>) was used to identify conserved motifs in the encoded proteins. Parameters employed in the analysis were: minimum motif width, 6; maximum motif width, 50; and maximum number of motifs, 10.

Isolation of salt-induced WRKY family genes

In a previous study, digital gene expression (DGE) analysis of *G. aridum* under salt stress revealed that the expression levels of 28 of 109 WRKY genes were significantly altered under salt stress. Using gene-specific primers (S2 Table) designed based on corresponding homologous gene in the *G. raimondii* genome, WRKY genes were cloned by PCR and their transcripts were amplified from *G. aridum* roots treated with 200 mM NaCl for 12 hours. PCR was performed using EasyPfu DNA Polymerase (TransGen Biotech, China). The PCR products were cloned into the PTG19-T vector (TransGen Biotech, China) and sequencing was performed by Invitrogen (Shanghai, China). The subcellular localizations of proteins were predicted by CELLO v.2.5 (subcellular localization predictor, <http://cello.life.nctu.edu.tw>) [39].

Expression analysis of salt-induced *GarWRKY* members in *G. aridum*

RNA isolation. Total RNA from *G. aridum* leaves and roots was extracted using an improved CTAB method [40]. For RT-PCR, first-strand cDNA was synthesized with a BU-SuperScript RT Kit (Biouniquer, China) according to the manufacturer's protocol. For quantitative real-time PCR, first-strand cDNA was synthesized with PrimeScript RT Master Mix (Perfect Real Time) from TaKaRa (Dalian, China).

Semi-quantitative RT-PCR. Semi-quantitative RT-PCR was performed to detect the differential expression of *GarWRKY* genes in different tissues including root, stem, leaf, petal, stigma, stamen and calyx tissue under normal conditions. The RT-PCR condition were as follow: preheating at 94°C for 3 min, followed by 36 cycles of 95°C for 45 s, 60°C for 45 s and 72°C for 1 min, with a final extension at 72°C for 10 min to complete the reaction. The PCR products were separated on a 1.2% agarose gel and quantified using an Imaging System (Bioshine GelX 1520, China). The specific primers used for RT-PCR are listed in S3 Table. The cotton actin gene was used as an internal reference gene (S3 Table).

Quantitative real-time PCR. The qRT-PCR was performed on an ABI PRISM 7500 Real-Time PCR System. The qRT-PCR reaction system (20 µl) contained 10 µl 2×SYBR Premix Ex Taq, 1 µl cDNA template, 0.4 µl each of forward and reverse primer (10 µM) and 0.4 µl ROX Reference Dye II (50×). The thermal cycling conditions were as follow: pre-denaturation at 95°C for 30 s, followed by 40 cycles at 95°C for 5 s and 60°C for 34 s. After the cycle was complete, melting curves analysis was performed at 60–95°C to verify the specificity of the amplicon for each primer pair. Quantitative real-time PCR was carried out with three biological replicates. The cotton actin gene was used as an internal reference gene. The specific primers used for

qRT-PCR are listed in [S3 Table](#). The relative gene expression values were analyzed by the $2^{-\Delta\Delta t}$ method.

Generation of *GarWRKY17* and *GarWRKY104* transgenic *Arabidopsis* plants. The cDNA sequence containing the full-length coding sequence of *GarWRKY17* and *GarWRKY104* were cloned into pCAMBIA2301 expression vector separately under the control of CaMV35S promoter between the *Xba* I and *Kpn* I sites. The recombinant plasmid pCAMBIA2301-*GarWRKY17* and pCAMBIA2301-*GarWRKY104* were transformed into *Agrobacterium* EH105 with electroporation method ($V = 2.4\text{kv}$). The *Arabidopsis* plants were transformed by the floral dip method [41]. Their T₂ transgenic lines were obtained after screening of Kanamycin resistance, PCR and RT-PCR analysis.

Analysis of salt tolerance in transgenic *Arabidopsis* plants. To evaluate salt tolerance of *GarWRKY17* and *GarWRKY104* transgenic *Arabidopsis* plants during seed germination stage, 50 sterilized seeds of wide type (WT) and T₂ generation transgenic lines (3 lines for *GarWRKY17* and 2 lines for *GarWRKY104*) were sowed in MS medium with and without 150 mM NaCl. The experiment was conducted with three biological replicates. Germination rate was calculated after 10 days.

To obtain further evidence that whether the overexpression of *GarWRKY17* and *GarWRKY104* could confer resistance to salt stress during vegetative growth, sterilized seeds of WT and T₂ transgenic *Arabidopsis* were sowed in soil and grew for 20 days. Plants were growing in a pot supplemented with 150ml water or 150mM NaCl. Phenotype symptoms were observed after 4 weeks.

To determine the activities of antioxidant enzymes, three-week-old WT seedlings and T₂ seedlings of 2 *GarWRKY17*-overexpressing (*GarWRKY17*-10 and *GarWRKY17*-25) were exposed to 150 mM NaCl for 24h. Leaves were collected from 10 plants of wild type and two transgenic lines. The activities of superoxide dismutase (SOD) and peroxidase (POD) was assayed using the procedure described by Liu et al. [42]. The tests were performed in triplicate.

Results

Identification of WRKY genes in *G. aridum* and their chromosomal locations

In a previous study, we identified more than 90,000 transcripts from de novo transcriptome sequencing of *G. aridum* under salt stress [34]. To perform genome-wide analysis of WRKY genes in *G. aridum*, all of these transcripts were used to BLAST the *G. raimondii* WRKY gene database using a local BLAST program. A total of 109 genes in *G. aridum* were ultimately identified as possible members of the WRKY superfamily and named according to their position from the top to the bottom of cotton chromosomes 1–13 (Fig 1). To facilitate communication, a uniform nomenclature for the 109 *GarWRKY* genes identified in this work was adopted based on that used in *G. raimondii* by Cai et al. [31]. The authors identified 120 candidate WRKY genes in the *G. raimondii* genome. By comparison, 11 WRKY genes (*WRKY7*, 16, 33, 42, 44, 49, 89, 96, 102, 109 and 112) were not identified in the current study.

GarWRKY genes are distributed on all 13 *G. aridum* chromosomes. Of these, chromosome 7 has the highest number of *GarWRKY* genes (13) and chromosome 5 has the lowest (3). The distribution patterns of these genes on individual chromosomes reveal some regions with numerous *GarWRKY* gene clusters. For example, *GarWRKY* genes located on chromosomes 4, 8, 9 and 11 appear to be concentrated on the lower or upper ends of the arms of chromosomes, respectively (Fig 1).




Fig 1. Chromosomal locations of *GarWRKY* genes in the *G. raimondii* genome. The candidate *WRKY* genes in *G. aridum* were designated *GarWRKY1* to *GarWRKY120* based on their orders on chromosomes. Eleven *WRKY* genes in *G. aridum* (*GarWRKY7*, 16, 33, 42, 44, 49, 89, 96, 102, 109 and 112) not identified in the current study were replaced by corresponding *WRKY* genes in the *G. raimondii* genome based on a report by Cai et al. (2014). Salt-responsive *GarWRKYs* are indicated in red. The scale is in megabases (Mb).

doi:10.1371/journal.pone.0126148.g001

Classification and phylogenetic analysis of *GarWRKY* genes

The *WRKY* domain and the zinc finger motif are the most prominent structural features of *WRKYs*. The *WRKY* domain contains the highly conserved heptapeptide stretch *WRKYGQK* at its N-terminus followed by a zinc-finger motif. A total of 126 *WRKY* domains were found in the 109 *GarWRKY* protein sequences.

The amino acid sequences of 126 *WRKY* domains were aligned using ClustalW with default settings. Based on *AtWRKY* classification and the *WRKY* domain features of *GarWRKYs*, the *GarWRKYs* were mainly classified into three groups, designated Group I to Group III (Fig 2). Seventeen *GarWRKYs* were assigned to Group I; these proteins contain two *WRKY* domains, an N-terminal WD (NTWD) and a C-terminal WD (CTWD). The NTWD zinc-finger type is C-X₄-C-X₂₂-HXX and the CTWD zinc-finger type is C-X₄-CX₂₃-HXX. Eighty *GarWRKYs* were assigned to Group II, which contain a single *WRKY* domain. The zinc-finger type is also C₂H₂, but its structure is C-X₄₋₅-C-X₂₃-HXX, which is structurally different from the Group I zinc finger motif. These 80 Group II *GarWRKYs* were further divided into five subgroups: subgroup IIa (7), IIb (15) IIc (30), IId (15) and IIE (13). Twelve *GarWRKYs* were assigned to Group III; these proteins contain a single *WRKY* domain with a C₂HC zinc-finger motif (C-X₇-C-X₂₃-HXC). Detailed information about the *GarWRKYs* can be found in S1 Table.

Cloning and classification of salt-induced *WRKY* family genes

In a previous study, we performed digital gene expression profile analysis of *G. aridum* during different stages of salt stress and identified 28 *WRKY* ESTs that were differentially expressed in roots or leaves during at least one stage [34]. In the current study, using gene-specific primers designed based on the corresponding homologous genes in the *G. raimondii* genome, we performed PCR cloning of *WRKY* genes and amplified the transcripts from roots of *G. aridum* treated with 200 mM NaCl for 12 h. We obtained 28 cDNA sequences of *WRKY* genes with complete open reading frames (ORFs; GenBank accession numbers KM438453–KM438480; Table 1). The ORF length ranged from 474 bp to 1,848 bp, with an average length of 1,048 bp. The identified *GarWRKY* genes encode proteins ranging from 157 to 615 amino acids (aa) in length with an average of 348 aa. The similarities between *GarWRKY* and *GrWRKY* based on their full-length nucleotide sequences ranged from 93.5% to 99.88%. All 28 salt-responsive *GarWRKY* proteins were predicted to be located in the nucleus.

Expression patterns of *GarWRKY* genes in various tissues revealed by semi-quantitative RT-PCR analysis

Using RT-PCR, we examined the expression patterns of *GarWRKY* genes in plants grown under normal growth conditions in seven different tissues including root, stem, leaf, petal, stigma, stamen and calyx tissue. As shown in Fig 3, 27 *GarWRKY* genes were expressed in roots, while 6 genes (*GarWRKY5*, *GarWRKY17*, *GarWRKY27*, *GarWRKY75*, *GarWRKY90* and *GarWRKY104*) were expressed in all tissues (with varying expression levels). Moreover, 21 genes had different tissue-specific expression profiles and *GarWRKY51* had very low transcript abundance in all tissues. Notably, *GarWRKY52* was exclusively expressed in roots.




Fig 2. Phylogenetic analysis of WRKY domains in *G. aridum*. WRKY protein name with the suffix 'N' or 'C' indicates the N-terminal WRKY domains or the C-terminal WRKY domains. The black arcs indicate different groups of WRKY domains. ● represent *AtWRKY* proteins, ▲ represent salt-responsive *GarWRKY* proteins.

doi:10.1371/journal.pone.0126148.g002

Validation of significant changes in *GarWRKY* expression in response to salt stress using real-time PCR

Gene expression patterns can provide important clues for determining gene function. To validate the results obtained from RNA-Seq, we performed quantitative RT-PCR analysis

Table 1. Characterization of 28 salt-responsive *GarWRKY* members.

GarWRKY	Group	ORF (bp)	Polypeptide (aa)	GrWRKY	Similarity (%)	Subcellular localization	GenBank accession number
GarWRKY5	III	921	306	WRKY5	98.05	Nuclear	KM438453
GarWRKY6	IIc	1077	358	WRKY6	99.07	Nuclear	KM438454
GarWRKY9	IId	951	316	WRKY9	98.74	Nuclear	KM438455
GarWRKY17	IIc	957	318	WRKY17	98.03	Nuclear	KM438456
GarWRKY22	I	1530	509	WRKY22	99.22	Nuclear	KM438457
GarWRKY27	IIe	1386	461	WRKY27	99.49	Nuclear	KM438458
GarWRKY28	IIa	933	310	WRKY28	99.41	Nuclear	KM438459
GarWRKY29	IIc	960	319	WRKY29	99.06	Nuclear	KM438460
GarWRKY31	IIb	1698	565	WRKY31	99.04	Nuclear	KM438461
GarWRKY38	IIc	474	157	WRKY38	99.37	Nuclear	KM438462
GarWRKY43	IIe	828	275	WRKY43	98.56	Nuclear	KM438463
GarWRKY51	IIa	876	291	WRKY51	93.5	Nuclear	KM438464
GarWRKY52	IIa	786	261	WRKY52	98.85	Nuclear	KM438465
GarWRKY54	IIb	978	325	WRKY54	98.56	Nuclear	KM438466
GarWRKY56	III	1008	335	WRKY56	99.01	Nuclear	KM438467
GarWRKY65	IIc	867	288	WRKY65	99.88	Nuclear	KM438468
GarWRKY67	IIe	978	325	WRKY67	99.39	Nuclear	KM438469
GarWRKY72	IId	1020	339	WRKY72	97.65	Nuclear	KM438470
GarWRKY75	IIb	1443	480	WRKY75	98.83	Nuclear	KM438471
GarWRKY78	IIc	885	294	WRKY78	98.8	Nuclear	KM438472
GarWRKY90	III	999	332	WRKY90	98.8	Nuclear	KM438473
GarWRKY95	I	1215	404	WRKY95	99.42	Nuclear	KM438474
GarWRKY104	IIc	966	321	WRKY104	98.65	Nuclear	KM438475
GarWRKY105	IIe	717	238	WRKY105	99.02	Nuclear	KM438476
GarWRKY107	IIb	1848	615	WRKY107	98.11	Nuclear	KM438477
GarWRKY113	IIe	1194	397	WRKY113	98.49	Nuclear	KM438478
GarWRKY114	III	876	291	WRKY114	95.97	Nuclear	KM438479
GarWRKY117	IIc	786	261	WRKY117	99.46	Nuclear	KM438480

doi:10.1371/journal.pone.0126148.t001

(Q-PCR) of the differentially expressed *GarWRKY* genes to confirm their levels of expression in roots and leaves after salt stress treatment. Quantitative expression analysis of these genes was performed using samples obtained from independent experiments carried out under the same conditions used to obtain RNA samples for RNA-Seq analysis. RNA was extracted from roots and leaves at 0, 1, 3, 6, 12, 24 and 72 h post salt stress, while RNA-Seq analysis was performed with fewer time points (0, 3, 12 and 72 h). As shown in Fig 4, the *GarWRKY* genes displayed distinct expression patterns in response to salt stress. Real-time PCR generally produced similar results to those obtained by differential expression profiling by RNA-Seq. However, in contrast to the RNA-Seq results, qRT-PCR revealed reduced expression of *GarWRKY95* and *GarWRKY114* in leaves.

Most *GarWRKY* genes in roots were activated within 1 h following salt treatment, with peak expression during the middle period and a return to basal levels during the latest period (72 h; Fig 4). However, in leaves, the majority of the analyzed *WRKY* genes were activated later than 6 h after salt treatment, with peak expression during the latest period (72 h). Expression analysis of *WRKY* genes during salt stress revealed delayed induction kinetics for most of the analyzed genes in leaves compared with roots, indicating that plants perceive salt stress and activate their defense machinery rapidly in roots.




Fig 3. Expression patterns of *GarWRKY* genes in various tissues. Semi-quantitative RT-PCR was conducted under normal growth condition. The cotton actin gene was used as an internal reference.

doi:10.1371/journal.pone.0126148.g003

Based on hierarchical clustering, the expression patterns of the *GarWRKY* genes were divided into five groups, i.e., Cluster A, Cluster B, Cluster C, Cluster D and Cluster E (Fig 5). Cluster A includes three genes (*GarWRKY9*, *GarWRKY27* and *GarWRKY107*) that were slightly downregulated in roots but upregulated in leaves. Cluster B includes one gene (*GarWRKY22*) that was downregulated in roots and leaves. Cluster C contains nine genes (*GarWRKY90*, *GarWRKY72*, *GarWRKY113*, *GarWRKY75*, *GarWRKY31*, *GarWRKY78*, *GarWRKY104*, *GarWRKY67* and *GarWRKY29*) that were upregulated in roots and downregulated in leaves at early stages (before 24 h) but upregulated at later stage (72 h). Cluster D contains two genes (*GarWRKY95* and *GarWRKY114*) that were upregulated in roots but downregulated in leaves. Finally, Cluster E contains 13 genes (*GarWRKY56*, *GarWRKY5*, *GarWRKY6*, *GarWRKY105*, *GarWRKY17*, *GarWRKY38*, *GarWRKY51*, *GarWRKY52*, *GarWRKY54*, *GarWRKY65*, *GarWRKY117*, *GarWRKY28* and *GarWRKY43*) that were upregulated in roots and leaves.

Comparison of orthologous *GarWRKY* genes from *Arabidopsis*, rice and soybean

To further implicate the functions of *GarWRKYs* in plant defenses to stresses, the orthologs in *Arabidopsis*, rice and soybean of 28 salt-responsive *WRKY* members were identified using BLASTP ($E < 1e^{-20}$), and the 58 top hit (17 in *Arabidopsis*, 19 in rice and 22 in soybean, respectively) were collected. A phylogenetic tree was constructed with the MEGA 4.0 software and conserved motif compositions were analyzed by MEME program.




Fig 4. Expression profiles of *GarWRKY* genes under NaCl stress. A: roots; B: leaves. The cotton actin gene was used as an internal reference. The data are mean $\pm$ SE of three biological replicates. * and ** indicate statistical significance at the 0.05 and 0.01 probability level, respectively.

doi:10.1371/journal.pone.0126148.g004

Our results showed that all 86 *WRKY* members from different species could be unambiguously classified as Group I, GroupIIa, IIb, IIc, IId, IId, IId and Group III (Fig 6). In total, 10 motifs, named, motif 1–10, were identified. Motifs 1, 2, 3, 4, were located in the *WRKY* domains, while the other motifs were dispersed around the *WRKY* domains. Motifs 1 and 2 were conserved in all *WRKY* family members; motif 5 was conserved in GroupIIa and IIb except for *GarWRKY54*; motif 4 was specific to Group I *WRKY* members. In Group I, only *GarWRKY22* and *OsWRKY78* contained motif 10. In Group IId and IId, motif 10 was found only in *GarWRKY9*, *GarWRKY27*, *GarWRKY113*, *OsWRKY14* and *OsWRKY31*. Overall, similar motif compositions were found in *WRKY* members of *Arabidopsis*, rice and soybean within the same subgroup, indicating the *WRKY* members have the similar function in these plants. It is worthwhile to note that 20 orthologous *WRKY* members in *Arabidopsis*, rice and/or soybean were also induced under salt stress according to previous reports [20–26, 42–44] (Fig 6), further implying these 28 *GarWRKY* members may play an important role in underlying mechanism of gene regulation under salt stress.

Salt tolerance validation of *GarWRKY17* and *GarWRKY104* in transgenic *Arabidopsis*

GarWRKY17 and *GarWRKY104* showed relatively higher differentially expression level in qRT-PCR analysis (Fig 4). Therefore, they were selected to generate overexpression transgenic *Arabidopsis* plants to further evaluate their function in response to salt stress. When these lines germinated on the medium adding 150 mM NaCl, the germination rates of two *GarWRKY104*-overexpressing lines (*GarWRKY104-1* and *GarWRKY104-2*) were significantly higher than that of WT (80.0% and 81.3% vs 22.0%). However, no significant difference was observed between WT and three *GarWRKY17*-overexpressing lines (*GarWRKY17-4*, -10, -25) (Fig 7).

To further validate the salt tolerance of *GarWRKY17* and *GarWRKY104*, 20 day-old overexpressing transgenic plants were treated with 150 mM NaCl solution using water as the control. Four weeks later, growth of the WT seedlings was completely inhibited while the *GarWRKY17*-overexpressing seedlings remained green and continued to grow, especially for *GarWRKY17-10* and *GarWRKY17-25* lines (Fig 8). Similar to WT, the *GarWRKY104*-overexpressing lines displayed the inhibited growth (data not shown). These data indicate that *GarWRKY17* positively regulate salt tolerance in a wide range of vegetative growth, while *GarWRKY104* exhibited the salt tolerance only during seed germination.

To study the physiological response of overexpressing-*GarWRKY17* *Arabidopsis* plants to salt stress, two transgenic lines (*GarWRKY17-10* and *GarWRKY17-25*) which showed better growth than WT plants during vegetative stage under salt stress were selected for further enzyme activities analysis. The activities of SOD and POD were determined in leaves of the control and the salt-treated *Arabidopsis* plants (Fig 8). The activity of SOD was much higher in the transgenic plants as compared to the WT plants with or without salt stress. In particular, the SOD activity in the two transgenic lines increased 2.6 times (*GarWRKY17-10*) and 2.8 times (*GarWRKY17-25*) compared with that in the wild type plants under salt stress. The activity of POD was significant higher in the transgenic plants as compared to the WT plants with or without salt stress. The activity of POD in the two transgenic lines increased 41% (*GarWRKY17-10*) and 36% (*GarWRKY17-25*) compared with that in WT plants under salt




Fig 5. Heat map of the expression patterns of 28 salt-responsive *GarWRKYs* from RNA-Seq and qRT-PCR data. Changes in expression levels are displayed from green (downregulated) to red (upregulated), as shown in the color gradient at the bottom right corner (color figure online). Heat maps were generated and hierarchical clustering was performed using Cluster 3.0 based on log2 fold-change data in response to NaCl stress. The normalized expression values from RNA-Seq data were provided in [S4 Table](#).

doi:10.1371/journal.pone.0126148.g005




Fig 6. Phylogenetic relationships and motif compositions of 28 salt-responsive GarWRKY members and their orthologs of *Arabidopsis*, rice and soybean. Unrooted phylogenetic tree was constructed by using MEGA4.0. The motifs identified by MEME software are represented by colored boxes. * indicated the genes induced under salt stress.

doi:10.1371/journal.pone.0126148.g006




Fig 7. Overexpressing *GarWRKYs* *Arabidopsis* plants at germination stage under salt stress. (A) Seedlings of the WT and 2 *GarWRKY104*-overexpressing lines germinated on MS medium adding 150 mM NaCl; (B) Seeds from the WT and 3 *GarWRKY17*-overexpressing lines were germinated on MS medium; (C) Germination rate of WT and 2 *GarWRKY104*-overexpressing lines; (D) Germination rate of WT and 3 *GarWRKY17*-overexpressing lines. The data are mean $\pm$ SE of three biological replicates. * and ** indicate statistical significance at the 0.05 and 0.01 probability level, respectively.

doi:10.1371/journal.pone.0126148.g007

stress. These data indicate that overexpression of *GarWRKY17* confer a more efficient antioxidant system to counteract the oxidative stress caused by salt stress.

Discussion

A few studies have revealed WRKY family members in *G. hirsutum* (cultivated species) and *G. raimondii* (for which complete genome sequence information is available), and the expression patterns of various WRKY TFs in response to abiotic stress (including salt stress) have been analyzed in upland cotton. As described in detail previously [34], modern cotton cultivars are the result of intensive selection to produce large amounts of specific types of fiber under unstressed




Fig 8. Effect of salt treatment on GarWRKY17-overexpressing Arabidopsis plants at vegetative growth stage. (A) Plants of WT and 3 *GarWRKY17*-overexpressing *Arabidopsis* lines were watered with 150 mM NaCl solution; (B) The expression level of *GarWRKY17* in WT and transgenic T₂ lines; (C) SOD activity; (D) POD activity. The data are mean $\pm$ SE of three biological replicates. * and ** indicate statistical significance at the 0.05 and 0.01 probability level, respectively.

doi:10.1371/journal.pone.0126148.g008

conditions. Selection has unintentionally narrowed the genetic variability for salt tolerance. Therefore, improving salt tolerance in upland cotton requires the use of valuable alleles from other *Gossypium* species. In this study, we selected *G. aridum*, a D-genome diploid species that grows in the Pacific coastal states of Mexico, to conduct transcriptome-wide identification of WRKY family members. We identified 28 salt-responsive *GarWRKY* genes and validated their expression patterns in *G. aridum* using RNA-Seq and qRT-PCR technology. In a recent study,

Cai et al. (2014) identified 120 candidate WRKY genes from *G. raimondii* and revealed that 17 WRKY genes are significantly induced by salt treatment in *G. hirsutum* [31]. Using uniform nomenclature, we found that WRKY22 in *G. hirsutum* and GarWRKY22 in *G. aridum* had similar expression patterns in *G. hirsutum* and *G. aridum* (significantly downregulated after salt treatment). However, WRKY6 in *G. hirsutum* and GarWRKY6 in *G. aridum* exhibited opposite expression patterns (downregulated in *G. hirsutum* and upregulated in *G. aridum*), indicating that some orthologous WRKY genes between *G. hirsutum* and *G. aridum* may have similar functions in regulating the stress response, while some may not.

In this study, we identified a total of 109 WRKY genes in *G. aridum* via de novo transcriptome sequencing of *G. aridum* under salt stress. In a recent report, Cai et al. identified 120 candidate WRKY genes from the *G. raimondii* genome, while the 11 remaining WRKY genes (WRKY7, 16, 33, 42, 44, 49, 89, 96, 102, 109 and 112) were not identified in the current study. Since *G. aridum* and *G. raimondii* belong to the same subgenome (D genome), they are expected to have the same number of WRKY genes in their genomes. The transcript abundance of these 11 WRKY genes was too low for them to be detected in the RNA-Seq libraries. These genes may be pseudogenes, or they may be expressed at specific developmental stages, under specific conditions or in specific tissues. In Cai's study, the expression patterns of WRKY16 and WRKY44 were examined in different tissues. WRKY16 was not expressed in roots and was only slightly expressed in leaves, but it was preferentially expressed in fibers. WRKY44 exhibited very low transcript abundance in roots and leaves but preferential expression in anthers. These WRKY genes, which are expressed under specific conditions or in specific plant tissues, may be involved in the growth and development of different organs.

A number of studies have shown that WRKY genes are induced by salt stress and that overexpression of some WRKY genes alters stress tolerance in plants. In *Arabidopsis*, 18 out of 35 WRKY genes from roots are induced by salt stress, such as AtWRKY8, 25, 33, 46, 57 and so on [43]. In rice, 27 WRKY transcription factor genes are induced in response to salt stress, 26 of which are upregulated [44]. In soybean, 22 of 64 WRKY genes are differentially expressed under salt stress [26]. In transgenic rice, overexpression of OsWRKY11/01g43650 and OsWRKY45/05g2577 results in enhanced salt, heat and drought tolerance [25]. Overexpression of OsWRKY45 results in enhanced salt and drought tolerance, in addition to increased disease resistance [24]. In *Arabidopsis*, overexpression of either AtWRKY25 or AtWRKY33 increases salt tolerance [23], while overexpression of AtWRKY18 or AtWRKY60 increases plant sensitivity to salt [45]. *GmWRKY54*-overexpressing plants are more salt and drought tolerant than the control, and *GmWRKY13* overexpression results in increased sensitivity to salt and mannitol stress [26]. In the current study, integrated transcriptome analysis indicated that 28 WRKY genes in *G. aridum* were differentially regulated in response to salt stress conditions. To further explore the functions of GarWRKYs that may be involved in plant defense responses to salt stress, we identified orthologous pairs among GarWRKYs, AtWRKYs, OsWRKYs and GmWRKYs based on full-length protein sequence similarities. Of the 28 salt-induced GarWRKY genes, 20 orthologs in *Arabidopsis*, rice and soybean also exhibit significant induction under salt stress. Among these, GarWRKY22, GarWRKY65 and GarWRKY5 are orthologous to AtWRKY33, GmWRKY54 and OsWRKY45, respectively. Motif compositions analysis showed that similar motif compositions were generally shared by WRKY members from *Arabidopsis*, rice and soybean within the same subgroup. These findings suggest that these members are important regulators of the response to salt stress. We also observed there was a possible relation between motif composition and gene function. GarWRKY22 contained motif 10, which was downregulated under salt stress. However, its orthologous gene in *Arabidopsis* (AtWRKY33) didn't possess motif 10, which was upregulated under salt stress [23]. Further

study is required to clarify whether motif 10 is a key composition in regulating alteration of gene expression and functional divergence in different species.

Recent studies have identified and revealed the expression patterns of various salt-responsive WRKY TFs in cotton [18, 28–34]. However, functional analysis of these TFs in an important crop such as cotton has rarely been performed. To date, only one WRKY gene in cotton has been functionally characterized [29], while the biological functions of most cotton WRKY genes remain largely unknown. In this study, we performed the first transcriptome-wide analysis of salt-responsive members of the WRKY family in *G. aridum*, a highly salt-tolerant wild species. Functional analysis by overexpression of *GarWRKY17* and *GarWRKY104* in *Arabidopsis* indicated they could positively regulate salt tolerance in *Arabidopsis* different development stages. This study demonstrates that a number of WRKY genes might be involved in the response to salt stress, and it provides clues for the selection of candidate genes for use in improving the stress tolerance of upland cotton.

Supporting Information

S1 Table. Characterization of 109 *GarWRKY* members.

(DOC)

S2 Table. Primers used for *GarWRKY* gene cloning.

(DOC)

S3 Table. Primers used for qRT-PCR.

(DOC)

S4 Table. The normalized expression values of 28 salt-responsive *GarWRKYs* from RNA-Seq.

(DOC)

Author Contributions

Conceived and designed the experiments: XLS. Performed the experiments: XQF QG YYG HMS XGZ. Analyzed the data: XQF PX YY. Contributed reagents/materials/analysis tools: WCN. Wrote the paper: XQF XLS.

References

1. Glazebrook J (2001) Genes controlling expression of defense responses in *Arabidopsis*—2001 status. *Current Opinion in Plant Biology* 4: 301–308. PMID: [11418339](#)
2. Chen W, Provart NJ, Glazebrook J, Katagiri F, Chang H, Eulgem T, et al. (2002) Expression profile matrix of *Arabidopsis* transcription factor genes suggests their putative functions in response to environmental stresses. *The Plant Cell* 14: 559–574. PMID: [11910004](#)
3. Kalde M, Barth M, Somssich IE, Lippok B (2003) Members of the *Arabidopsis* WRKY group III transcription factors are part of different plant defense signaling pathways. *Molecular Plant-Microbe Interactions* 16: 295–305. PMID: [12744458](#)
4. Pabo CO, Sauer RT (1992) Transcription factors: structural families and principles of DNA recognition. *Annual Review of Biochemistry* 61: 1053–1095. PMID: [1497306](#)
5. Rushton PJ, Somssich IE, Ringler P, Shen QJ (2010) WRKY transcription factors. *Trends in Plant Science* 15: 247–258. doi: [10.1016/j.tplants.2010.02.006](#) PMID: [20304701](#)
6. Zhang Y, Wang L (2005) The WRKY transcription factor superfamily: its origin in eukaryotes and expansion in plants. *BMC Evolutionary Biology* 5: 1. PMID: [15629062](#)
7. Ishiguro S, Nakamura K (1994) Characterization of a cDNA encoding a novel DNA-binding protein, *SPF1*, that recognizes *SP8* sequences in the 5' upstream regions of genes coding for sporamin and β -amylase from sweet potato. *Molecular and General Genetics* 244: 563–571. PMID: [7969025](#)

8. Sun C, Palmqvist S, Olsson H, Borén M, Ahlandsberg S, Jansson C (2003) A novel WRKY transcription factor, *SUSIBA2*, participates in sugar signaling in barley by binding to the sugar-responsive elements of the *iso1* promoter. *The Plant Cell* 15: 2076–2092. PMID: [12953112](#)
9. Chen C, Chen Z (2000) Isolation and characterization of two pathogen- and salicylic acid-induced genes encoding WRKY DNA-binding proteins from tobacco. *Plant Molecular Biology* 42: 387–396. PMID: [10794538](#)
10. Zhang Z, Xie Z, Zou X, Casaretto J, Ho TD, Shen Q (2004) A rice WRKY gene encodes a transcriptional repressor of the gibberellin signaling pathway in aleurone cells. *Plant Physiology* 134: 1500–1513. PMID: [15047897](#)
11. Lambais MR (2001) In silico differential display of defense-related expressed sequence tags from *sugarcane* tissues infected with diazotrophic endophytes. *Genetics and Molecular Biology* 24: 103–111.
12. Borrone JW, Kuhn DN, Schnell RJ (2004) Isolation, characterization, and development of WRKY genes as useful genetic markers in *Theobroma cacao*. *Theoretical and Applied Genetics* 109: 495–507. PMID: [15148571](#)
13. Pnueli L, Hallak-Herr E, Rozenberg M, Cohen M, Goloubinoff P, Kaplan A, et al. (2002) Molecular and biochemical mechanisms associated with dormancy and drought tolerance in the desert legume *Retama raetam*. *The Plant Journal* 31: 319–330. PMID: [12164811](#)
14. Ashida Y, Nishimoto M, Matsushima A, Watanabe J, Hirata T (2002) Molecular cloning and mRNA expression of geraniol-inducible genes in cultured shoot primordia of *Matricaria chamomilla*. *Bioscience, Biotechnology, and Biochemistry* 66: 2511–2514. PMID: [12506999](#)
15. Li H, Xu Y, Xiao Y, Zhu Z, Xie X, Zhao H, et al. (2010) Expression and functional analysis of two genes encoding transcription factors, *VpWRKY1* and *VpWRKY2*, isolated from Chinese wild *Vitis pseudoreticulata*. *Planta* 232: 1325–1337. doi: [10.1007/s00425-010-1258-y](#) PMID: [20811906](#)
16. Xu Y, Wang J, Wang S, Wang J, Chen X (2004) Characterization of *GaWRKY1*, a cotton transcription factor that regulates the sesquiterpene synthase gene (+)- δ -cadinene synthase-A. *Plant Physiology* 135: 507–515. PMID: [15133151](#)
17. Ling J, Jiang W, Zhang Y, Yu H, Mao Z, Gu X, et al. (2011) Genome-wide analysis of WRKY gene family in *Cucumis sativus*. *BMC Genomics* 12: 471. doi: [10.1186/1471-2164-12-471](#) PMID: [21955985](#)
18. Guo R, Yu F, Gao Z, An H, Cao X, Guo X (2011) *GhWRKY3*, a novel cotton (*Gossypium hirsutum* L.) WRKY gene, is involved in diverse stress responses. *Molecular Biology Reports* 38: 49–58. doi: [10.1007/s11033-010-0076-4](#) PMID: [20238169](#)
19. Huang S, Gao Y, Liu J, Peng X, Niu X, Fei Z, et al. (2012) Genome-wide analysis of WRKY transcription factors in *Solanum lycopersicum*. *Molecular Genetics and Genomics* 287: 495–513. doi: [10.1007/s00438-012-0696-6](#) PMID: [22570076](#)
20. Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, et al. (2002) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *The Plant Journal* 31: 279–292. PMID: [12164808](#)
21. Chen L, Song Y, Li S, Zhang L, Zou C, Yu D (2012) The role of WRKY transcription factors in plant abiotic stresses. *Biochimica et Biophysica Acta—Gene Regulatory Mechanisms* 1819: 120–128.
22. Cheong Y, Chang H, Gupta R, Wang X, Zhu T, Luan S (2002) Transcriptional profiling reveals novel interactions between wounding, pathogen, abiotic stress, and hormonal responses in *Arabidopsis*. *Plant Physiology* 129: 661–677. PMID: [12068110](#)
23. Jiang Y, Deyholos MK (2009) Functional characterization of *Arabidopsis* NaCl-inducible *WRKY25* and *WRKY33* transcription factors in abiotic stresses. *Plant Molecular Biology* 69: 91–105. doi: [10.1007/s11103-008-9408-3](#) PMID: [18839316](#)
24. Qiu Y, Yu D (2009) Over-expression of the stress-induced *OsWRKY45* enhances disease resistance and drought tolerance in *Arabidopsis*. *Environmental and Experimental Botany* 65: 35–47.
25. Wu X, Kishitani S, Ito Y, Toriyama K (2009) Accumulation of raffinose in rice seedlings overexpressing *OsWRKY11* in relation to desiccation tolerance. *Plant Biotechnology* 26: 431–434.
26. Zhou Q, Tian A, Zou H, Xie Z, Lei G, Huang J, et al. (2008) Soybean WRKY-type transcription factor genes, *GmWRKY13*, *GmWRKY21*, and *GmWRKY54*, confer differential tolerance to abiotic stresses in transgenic *Arabidopsis* plants. *Plant Biotechnology Journal* 6: 486–503. doi: [10.1111/j.1467-7652.2008.00336.x](#) PMID: [18384508](#)
27. Wang C, Deng P, Chen L, Wang X, Ma H, Hu W, et al. (2013) A wheat WRKY transcription factor *TaWRKY10* confers tolerance to multiple abiotic stresses in transgenic tobacco. *PloS one* 8: e65120. doi: [10.1371/journal.pone.0065120](#) PMID: [23762295](#)
28. Yu F, Hua X, Lu W, Wu C, Cao X, Guo X (2012) *GhWRKY15*, a member of the WRKY transcription factor family identified from cotton (*Gossypium hirsutum* L.), is involved in disease resistance and plant development. *BMC Plant Biology* 12: 144. doi: [10.1186/1471-2229-12-144](#) PMID: [22883108](#)

29. Shi W, Liu D, Hao L, Wu C, Guo X, Li H (2014) *GhWRKY39*, a member of the WRKY transcription factor family in cotton, has a positive role in disease resistance and salt stress tolerance. *Plant Cell, Tissue and Organ Culture* 18: 1–16.
30. Zhou L, Wang N, Kong L, Gong S, Li Y, Li X (2014) Molecular characterization of 26 cotton WRKY genes that are expressed differentially in tissues and are induced in seedlings under high salinity and osmotic stress. *Plant Cell, Tissue and Organ Culture* 1–16.
31. Cai C, Niu E, Du H, Zhao L, Feng Y, Guo W (2014) Genome-wide analysis of the WRKY transcription factor gene family in *Gossypium raimondii* and the expression of orthologs in cultivated tetraploid cotton. *The Crop Journal*.
32. Dou L, Zhang X, Pang C, Song M, Wei H, Fan S, et al. (2014) Genome-wide analysis of the WRKY gene family in cotton. *Molecular Genetics and Genomics*: 1–19.
33. Ding M, Chen J, Jiang Y, Lin L, Cao Y, Wang M, et al. (2015) Genome-wide investigation and transcriptome analysis of the WRKY gene family in *Gossypium*, *Mol Genet Genomics*, 290:151–171. doi: [10.1007/s00438-014-0904-7](https://doi.org/10.1007/s00438-014-0904-7) PMID: [25190108](https://pubmed.ncbi.nlm.nih.gov/25190108/)
34. Xu P, Liu Z, Fan X, Gao J, Zhang X, Zhang X, et al. (2013) *De novo* transcriptome sequencing and comparative analysis of differentially expressed genes in *Gossypium aridum* under salt stress. *Gene* 525: 26–34. doi: [10.1016/j.gene.2013.04.066](https://doi.org/10.1016/j.gene.2013.04.066) PMID: [23651590](https://pubmed.ncbi.nlm.nih.gov/23651590/)
35. Punta M, Coggill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, et al. (2012) The Pfam protein families database. *Nucleic Acids Research* 40: D290–D301. doi: [10.1093/nar/gkr1065](https://doi.org/10.1093/nar/gkr1065) PMID: [22127870](https://pubmed.ncbi.nlm.nih.gov/22127870/)
36. Jones P, Binns D, Chang H, Fraser M, Li W, McAnulla C, et al. (2014) InterProScan 5: genome-scale protein function classification. *Bioinformatics*: btu031.
37. Li KB (2003) ClustalW-MPI: ClustalW analysis using distributed and parallel computing. *Bioinformatics* 19: 1585–1586. PMID: [12912844](https://pubmed.ncbi.nlm.nih.gov/12912844/)
38. Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24: 1596–1599. PMID: [17488738](https://pubmed.ncbi.nlm.nih.gov/17488738/)
39. Yu C, Chen Y, Lu C, Hwang J (2006) Prediction of protein subcellular localization. *Proteins: Structure, Function, and Bioinformatics* 64: 643–651. PMID: [16752418](https://pubmed.ncbi.nlm.nih.gov/16752418/)
40. Hu G, Yu S (2007) Extraction of high-quality total RNA in cotton leaf with improved CTAB Method. *Cotton science* 19: 69–70.
41. Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium* mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16, 735–743. PMID: [10069079](https://pubmed.ncbi.nlm.nih.gov/10069079/)
42. Liu RX, Guo WQ, Chen BL, Zhou ZG (2008) Effects of N fertilization on root development and activity of water-stressed cotton plants, *Agricultural Water Management*, 95:1261–1270.
43. Jiang Y, Deyholos MK (2006) Comprehensive transcriptional profiling of NaCl-stressed *Arabidopsis* roots reveals novel classes of responsive genes. *BMC Plant Biology* 6: 25. PMID: [17038189](https://pubmed.ncbi.nlm.nih.gov/17038189/)
44. Ramamoorthy R, Jiang S, Kumar N, Venkatesh PN, Ramachandran S (2008) A comprehensive transcriptional profiling of the WRKY gene family in rice under various abiotic and phytohormone treatments. *Plant and Cell Physiology* 49: 865–879. doi: [10.1093/pcp/pcn061](https://doi.org/10.1093/pcp/pcn061) PMID: [18413358](https://pubmed.ncbi.nlm.nih.gov/18413358/)
45. Chen H, Lai Z, Shi J, Xiao Y, Chen Z, Xu X (2010) Roles of *Arabidopsis WRKY18*, *WRKY40* and *WRKY60* transcription factors in plant responses to abscisic acid and abiotic stress. *BMC Plant Biology* 10: 281. doi: [10.1186/1471-2229-10-281](https://doi.org/10.1186/1471-2229-10-281) PMID: [21167067](https://pubmed.ncbi.nlm.nih.gov/21167067/)