

Sonerila lundinii, a new species of Melastomataceae from southern Western Ghats with notes on *Sonerila pedunculosa*, a less known taxon from Sri Lanka

Resmi S.¹, Nampy S.^{1*}, Cellinese N.² & M.P. Krishnapriya¹

¹Department of Botany, University of Calicut, Malappuram District, Kerala – 673 635, India

²Florida Museum of Natural History, University of Florida, 1659 Museum Road, Gainesville, FL 32611-7800, U.S.A.

E-mail: santhoshnampy2019@gmail.com

Abstract: *Sonerila lundinii* Resmi & Nampy, a new species of Melastomataceae from the southern Western Ghats is here described. The new species is morphologically similar to *Sonerila pedunculosa* Thwaites by its prostrate or creeping habit, but differs by its pubescence, length of the peduncle, shape of the petals, anthers and capsules. Specimens so far collected from India wrongly identified as *S. pedunculosa*, which is an endemic species with a restricted distribution in Sri Lanka. Therefore, these specimens are considered as novelty and remarks on the occurrence of *S. pedunculosa* in South Indian Floras are clarified in the present paper.

Keywords: India, Endemism, Sonerileae, *Sonerila*, Sri Lanka, Taxonomy.

Introduction

Sonerila Roxb., a taxonomically complex, species-rich, tropical Old World genus in the family Melastomataceae (tribe Sonerileae), is represented by around 180 taxa (Cellinese, 1997; Resmi & Nampy, 2021a, b; Resmi *et al.*, 2018; 2021). It is widely distributed spanning an area from Sri Lanka, India, Nepal, Bhutan, South China, Taiwan, Indo-China and the Malay Archipelago. It is the largest genus in the Indian Sonerileae with 49 species and one variety, and exhibits a high percentage of endemism (Resmi, 2022). The greatest diversity of *Sonerila* in India occurs in the Western Ghats, where 43 species and one variety are found with 86% of endemics.

As a part of an ongoing taxonomic revision, the authors collected specimens from across India, and noticed that at Chemunji, in the

Thiruvananthapuram district of Kerala, a few populations exhibited a prostrate or creeping habit, a densely villous stem, leaves with finely serrulate laminae, acuminate anthers, and 3-ribbed capsules. Detailed comparative studies with the morphologically allied *S. pedunculosa* Thwaites, and consultation of types and protologues, showed that these specimens are morphologically distinct from any of the previously described taxa, and thus represent a new species.

Materials and Methods

Materials for this study were collected through extensive field trips to different parts of the southern Western Ghats during 2018–2020. Herbarium specimens were prepared and flowers and fruits were preserved for further study in a solution containing 4% formalin and 70% ethanol (De Vogel, 1987; Forman & Bridson, 1989). Living plants were maintained in the Calicut University Botanical Garden (CUBG) in order to observe variation under cultivation. Detailed morphological observations were recorded and described using terms of Stearn (1992) and Beentje (2016). Photographs were taken with an EZ4 HD stereo microscope (Leica, Heerbrugg, Switzerland) with a 3.0 megapixel digital camera (Leica) and Stemi 508 stereo microscope (Zeiss, Oberkochen, Germany) outfitted with an Axiocam 105 color camera (Zeiss). Relevant taxonomic accounts (Thwaites, 1864; Triana, 1871; Clarke, 1879; Cogniaux, 1891; Stapf, 1892; Triménis, 1894; Willis, 1911; Yoganarasimhan *et al.*, 1976; Lundin, 1983; Sharma *et al.*, 1984; Giri & Nayar, 1986; Bremer, 1987; Mohanan & Henry, 1994; Saldanha, 1996; Sasidharan, 2004; Nayar *et al.*, 2006; Manickam *et al.*, 2008; Nayar *et al.*, 2014; Murthy

Received: 31.11.2022; Revised & Accepted: 03.02.2023

Published Online: 02.03.2023

& Nair, 2016), herbarium specimens (BM, K, CAL, CALI, TBGT, XCH), and digital images available from BR, G, HUH, P and PDA were consulted for this study. Acronyms of herbaria follow the Index Herbariorum (Thiers, updated continuously). We generated a distribution map using QGIS ver. 3.28.2 (QGIS, 2022), based on coordinates and geo-referenced estimates of collection localities. The Area of Occupancy (AOO) and Extent of Occurrence (EOO) were estimated using GeoCAT (Bachman *et al.*, 2011), and a provisional conservation status was assessed according to the IUCN Red List Categories and Criteria (IUCN, 2012; IUCN Standards and Petitions Committee, 2022).

***Sonerila lundinii* Resmi & Nampy, sp. nov. Figs. 1–3**

Sonerila lundinii is morphologically close to *S. pedunculosa* sharing a prostrate or creeping habit rooting at nodes, but is distinguished by its densely villous stem (*vs.* glabrous in *S. pedunculosa*), finely serrulate laminae (*vs.* crenate or finely serrate), 2–4 cm long peduncles (*vs.* 3–7 cm), broadly ovate, pale rose petals (*vs.* ovate, dark pink), lanceolate acuminate anthers (*vs.* narrowly ovate, obtuse to acute), and campanulate, 3-ribbed capsules (*vs.* obconic, 6-ribbed).

Type: INDIA, Kerala, Thiruvananthapuram district, Agasthyamala Biosphere Reserve, Chemunji, way to hill top, 1200 m, 05.01.2018, *S. Resmi, M.P. Krishnapriya & S. Nampy* 151257 (holo CALI!; iso MH!).

Annual, caulescent, prostrate or creeping herbs. Root stock thick; roots arising from all nodes, 5–8 cm long. Stems terete, 0.2–0.3 cm thick, green or vinaceous, repeatedly branched from a common node and ramify in all directions, densely covered with eglandular trichomes; branches 5–15 cm long; internodes 1–2.5 cm long, short towards apex. Leaves decussate, whorled towards apex forming a rosette in young stage; petioles adaxially grooved or canaliculated, 0.5–3 cm long, green, densely covered with eglandular trichomes; laminae ovate to broadly ovate, 2–5 × 1.7–3 cm, dark green or with a purple tinge adaxially, equal and rounded or sub-cordate at base with non-overlapping margins, finely serrulate at margins with each tooth ending in a terminal eglandular trichome, acute to obtuse at apex, membranous, densely villous adaxially with eglandular trichomes, but only on veins abaxially; veins pinnate, 1–2 pairs from the base and 1–2 pairs from midrib above, less branched, impressed

adaxially, conspicuous abaxially. Cymes scorpioid, unbranched, 3–7-flowered, terminal; peduncles sub-terete to quadrangular, 2–4 × 0.10–0.15 cm, green or green with a vinaceous or claret tinge; bracts sometimes present. Flowers 1–1.5 × 1–1.5 cm; pedicels sub-angular, 1–3 × 1–2 mm, longer in fruit, vinaceous or claret, sparsely covered by gland-tipped trichomes. Hypanthia campanulate, 5–7 × 2–2.5 mm, 3-lobed, 3-ribbed, green with a vinaceous or claret tinge, sparsely covered by gland tipped trichomes; lobes triangular, 1–1.5 × 1–1.5 mm, acute at apex. Petals 3, broadly ovate, 6–9 × 4–5 mm, pale rose with a darker midrib, rounded at base, mucronate at apex, glabrous adaxially, with gland-tipped trichomes on midrib abaxially. Stamens 3; filaments 4–6 mm long, pale pink, glabrous; anthers lanceolate, 4–6 mm long, acuminate at apex. Ovary 2–3 × 2–3 mm; style 8–9 mm long, pale pink, dark towards apex, equal to the stamen; stigma capitate, pink, rugose, glabrous. Capsules campanulate, 5–7 × 3–3.5 mm, green with a vinaceous or claret tinge, 3-ribbed, sparsely covered with gland-tipped trichomes, less so with age. Seeds numerous, obovoid, 0.5–0.6 × 0.2–0.3 mm, pale brown; raphe prominent, excurrent or remaining out into a lateral appendage; testa densely pusticulate throughout, tuberculate at dorsal angle; tubercles pyramidal, covered all over by small pusticles.

Flowering & fruiting: Flowering from December to January and fruiting from January to March.

Habitat: *Sonerila lundinii* occurs in shady habitats with a relatively thick mossy substratum. It grows along the edges of evergreen forests, at elevations between 750–1200 m above sea level, in association with *S. tinneveli* C.E.C.Fisch., *S. brunonis* Wight & Arn. (Melastomataceae), *Cyanotis villosa* (Spreng.) Schult. & Schult.f. (Commelinaceae), *Henckelia repens* (Bedd.) A.Weber & B.L.Burt (Gesneriaceae), *Justicia betonica* L. (Acanthaceae) and some mosses.

Distribution: Endemic to southern Western Ghats, India (Fig. 4).

Etymology: The epithet ‘lundinii’ honours the late Roger Lundin for his valuable contribution to the taxonomy of the genus *Sonerila*.

Specimens examined: INDIA, Kerala, Thiruvananthapuram district, Chemunji,




Fig. 1. *Sonerila lundinii* Resmi & Nampy: **a–e.** Plants in its natural habitat at Chemunji hills in the Thiruvananthapuram district; **f.** Habit from above and **h.** from below, note the branching from nodes; **g.** Portion of stem showing dense trichomes (from S. Resmi, M.P. Krishnapriya & S. Nampy 151257; photos by S. Resmi).




Fig. 2. *Sonerila lundinii* Resmi & Nampy: **a–c.** Cymes in different views; **d.** Leaves, adaxial view; **e.** Leaves, abaxial view; **f.** Flower, front view; **g.** Flower, side view; **h.** Petals, adaxial view; **i.** Petals, abaxial view; **j.** Hypanthium, side view; **k.** Hypanthium with stamens and pistil; **l.** Stamens; **m.** Pistil; **n.** Immature capsules; **o.** Seeds (from *S. Resmi*, *M.P. Krishnapriya* & *S. Nampy* 151257; photos by *S. Resmi*).




Fig. 3. *Sonerila lundinii* Resmi & Nampy: **a.** Habit; **b.** Flower; **c.** Hypanthium with stamens and pistil; **d.** Petals, adaxial view; **e.** Petal, abaxial view, note the glandular trichomes; **f.** Stamens; **g.** Immature capsules; **h.** Seeds (from *S. Resmi*, *M.P. Krishnapriya* & *S. Nampy* 151257; drawn a by V.V. Drisya; b-h by S. Resmi).

21.03.1996, E.S. Santhosh Kumar 25412 (TBGT); *Ibid.*, 23.12.2020, M.K. Akhil, T.P. Krishnaraj, S. Resmi & S. Nampy 164469 (CALI); Koviltherimala, 750 m, 09.10.2009, K.P. Deepthikumari & A.G. Pandurangan 60599 (in vegetative state); *Ibid.*, 18.10.2010, K.P. Deepthikumari & Usha 67987 (TBGT).

Conservation status: The new species is currently known only from Chemunji and Koviltherimala in the Agasthyamala Biosphere Reserve. We were able to observe one population of approximately 90 mature individuals in the Chemunji hills. The habitat is severely affected by controlled burning activities by the Forest Department. It is likely that there may be more populations in similar habitats and these areas will be negatively impacted by further disturbances in the future due to local human activities involving the harvesting of wood, honey and forage grasses. The lack of collections from additional locations precludes the accurate estimation of the Extent of Occurrence (EOO), but the Area of Occupancy (AOO) is c. 8 km². This species is thus assessed here as Critically Endangered (CR), B2ab (ii,iii,v); D according to IUCN Red List Categories and Criteria (IUCN, 2012; IUCN Standards and Petitions Committee, 2022).

Notes: *Sonerila lundinii* appears to be closely related to *S. pedunculosa* in its creeping habit. Nevertheless, it differs by characteristics of the stem, leaf margins, and the morphology of petals, anthers and capsules. According to Thwaites (1864), *S. pedunculosa* is a distinct, creeping plant, rooting at nodes with rounded or ovate, obtuse, crenate leaves, sparsely hairy adaxially and, nearly so, abaxially. It is also characterised by elongated, few-flowered cymes, cordate-acute anthers, and 6-ribbed capsules. Trimen (1894) and Lundin (1983) added that these plants have glabrous stems with long internodes, which was evident in Lundin's (*l.c.*) illustration (based on Fagerlind 1229 in S). Clarke (1879), Cogniaux (1891) and Trimen (1894) recorded the extended distribution of *S. pedunculosa* in Travancore and Quilon by citing specimen Wight 1111. However, Stapf (1892), Willis (1911), Bremer (1987) and Lundin (1983)

followed Triana's treatment and considered this taxon only to occur in Southwest Sri Lanka (Ceylon).

According to Yoganarasimhan *et al.* (1976), Wight 1111 should be identified as *S. rheedei* sensu Wight & Arn. and not *S. pedunculosa* as stated by Clarke (1879). We studied Wight 1111 at K[K001325401] and confirm its identity as *S. rheedei*. Yoganarasimhan *et al.* (*l.c.*) cited a collection (Yoganarasimhan 1331) from Charmadi Ghats, Chikamagalur district, Karnataka as *S. pedunculosa* and according to Giri and Nayar (1986), this should also be identified as *S. rheedei*. We tried to locate specimen Yoganarasimhan 1331 at K, CAL, and PDA, but to no avail.

Several recent authors also wrongly recorded *S. pedunculosa* from the Western Ghats (Sharma *et al.*, 1984; Mohanan & Henry, 1994; Saldanha, 1996; Nayar *et al.*, 2006; Sasidharan, 2004; Manickam *et al.*, 2008; Nayar *et al.*, 2014). Manickam *et al.* (*l.c.*) mentioned its distribution in Tamil Nadu, but we confirmed that the voucher specimen deposited at XCH are *Sonerila travancorica* Bedd., not *S. pedunculosa*. Additionally, we noticed that some recent collections from Chemunji and Koviltherimala in TBGT were also identified as *S. pedunculosa*. According to Murthy and Nair (2016), *S. pedunculosa* reported from India by various authors is actually *S. rheedei* Wight & Arn. This misidentification created ambiguity in the identity and distribution of *S. pedunculosa* rather than clarifying the confusion. Detailed herbarium studies showed that Thwaites specimens of *S. pedunculosa* available at different herbaria (BM, CAL, HUH [HUH00073221], BR [BR000000522366, BR000000522398], K [K001325399], [K001325400], G [G00319944], P [P04888228, P05282776, P05282775] and PDA [PDA00002846]) are very distinct from *S. lundinii*. Further, no collection of this species from other parts of tropical Asia could be located in other Indian and foreign herbaria consulted by us. This study sheds light on the erroneous identification of specimens recorded from Chemunji and nearby areas as reported in local floras, and confirmed the endemism of *S. pedunculosa* in Sri Lanka.




Fig. 4. Distribution points of *Sonerila lundinii* Resmi & Nampy in Kerala, India (drawn using QGIS ver. 3.28.2).

Acknowledgements

The authors are thankful to the Head of the Department of Botany, University of Calicut, for access to the facilities used in this study; the curators of the herbaria at K, BM, CALI, MH, CAL, TBGT and XCH for access to the specimens consulted; Kerala and Tamil Nadu Forest Departments for permission to conduct field studies; Mr. Akhil M.K., Mr. Krishnaraj T.P., Dr. Soumya P. and Dr. Anto Mathew for various help during fieldwork. SN is thankful to the Science Engineering Research Board (SERB), New Delhi (EMR/2016/007346, dated 20/11/2018) for funding. RS is grateful to the International Association for Plant Taxonomy for the IAPT Research Grant 2020 and to the Government of Kerala for financial support.

Literature Cited

BACHMAN S., MOAT J., HILL A.W., DE LA TORRE J. & B. SCOTT 2011. Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. In: SMITH V. & L. PENEV (eds.), e-infrastructures for data publishing in biodiversity science. *ZooKeys* 150: 117–126. <https://doi.org/10.3897/zookeys.150.2109>

- BEENTJE H. 2016. *The Kew plant glossary: an illustrated dictionary of plant terms*, Second edition. Kew Publishing, Royal Botanic Gardens, Kew.
- BREMER K. 1987. Melastomataceae. In: DASSANAYAKE M.D. & F.R. FOSBERG (eds.), *A revised handbook to the Flora of Ceylon*. Volume 4. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi. pp. 157–240.
- CELLINESE N. 1997. Notes on the systematic and biogeography of the *Sonerila* generic alliance (Melastomataceae) with special focus on fruit characters. *Tropical Biodiversity* 4: 83–93.
- CLARKE C.B. 1879. Melastomataceae. In: HOOKER J.D. (ed.), *The Flora of British India*. Volume 2. L. Reeve & Co., London. pp. 512–565.
- COGNIAUX C.A. 1891. *Sonerila*. In: DE CANDOLLE A.L.P.P. & A.C.P. DE CANDOLLE (eds.), *Monographiae Phanerogamarum*. Volume 7. G. Masson, Paris. 7: 490–516.
- DE VOGEL E.F. 1987. *Guidelines for the preparation of revisions. Manual of herbarium taxonomy theory and practice*. UNESCO, Jakarta.

- FORMAN L. & D. BRIDSON 1989. *The herbarium handbook, First edition*. Royal Botanic Gardens, Kew.
- GIRI G.S. & M.P. NAYAR 1986. A new species and notes on two species of *Sonerila* Roxb. (Melastomataceae) from South India. *Bulletin of the Botanical Survey of India* 26(3–4): 174–180.
- IUCN 2012. *IUCN Red List Categories and Criteria: Version 3.1*. Second edition. IUCN Species Survival Commission, Gland and Cambridge.
- IUCN STANDARDS AND PETITIONS COMMITTEE 2022. Guidelines for using the IUCN Red list Categories and Criteria. Version 15. Available at: <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> (Accessed on 04.10 2022).
- LUNDIN R. 1983. Taxonomy of *Sonerila* (Melastomataceae) in Ceylon. *Nordic Journal of Botany* 3: 633–656. <https://doi.org/10.1111/j.1756-1051.1983.tb01475.x>
- MANICKAM V.S., MURUGAN C. & G.J. JOTHI 2008. *Flora of Tirunelveli hills*. Volume 1. Polypetalae. Bishen Singh Mahendra Pal Singh, Dehra Dun.
- MOHANAN M. & A.N. HENRY 1994. *Flora of Thiruvananthapuram*. Botanical Survey of India, Kolkata.
- MURTHY G.V.S. & V.J. NAIR 2016. *The Flora of Kerala*. Volume 2. Fabaceae–Cornaceae. Botanical Survey of India, Kolkata.
- NAYAR T.S., RASIYA BEEGAM A. & M. SIBI 2014. *Flowering plants of the Western Ghats, India*. Jawaharlal Nehru Tropical Botanical Garden and Research Institute, Thiruvananthapuram.
- NAYAR T.S., RASIYA BEEGAM A., MOHANAN N. & G. RAJKUMAR 2006. *Flowering plants of Kerala: a handbook*. Tropical Botanic Garden and Research Institute, Thiruvananthapuram.
- QGIS (Quantum GIS Development Team) 2022. Quantum GIS geographic information system. Version 3.28.2.
- RESMI S. 2022. *Taxonomic revision of the genus Sonerila Roxb. (Melastomataceae) in India*. Ph.D. Thesis (unpublished). University of Calicut, Malappuram.
- RESMI S. & S. NAMPY 2021a. A new addition to the stoloniferous *Sonerila* (Sonerileae, Melastomataceae) from southern Western Ghats, India. *Brittonia* 73(2): 195–202. <https://doi.org/10.1007/s12228-021-09657-w>
- RESMI S. & S. NAMPY 2021b. *Sonerila longipedunculata*, a new species of Melastomataceae from southern Western Ghats, India, with additional notes and lectotypification of *Sonerila travancorica*. *European Journal of Taxonomy* 733: 160–174. <https://doi.org/10.5852/ejt.2021.733.1231>
- RESMI S., MANUDEV K.M., PRAMOD C. & S. NAMPY 2018. A new species of *Sonerila* (Melastomataceae: Sonerileae) from southern Western Ghats, India. *Phytotaxa* 333 (2): 235–242. <https://doi.org/10.11646/phytotaxa.333.2.7>
- RESMI S., NAMPY S. & P.F. AKSHATRA 2021. *Sonerila konkanensis* (Melastomataceae), a new species from south Goa, India. *Candollea* 76(1): 139–143. <https://doi.org/10.15553/c2021v761a14>
- SALDANHA C.J. 1996. *Flora of Karnataka*. Oxford and IBH Publishing Co., New Delhi.
- SASIDHARAN N. 2004. *Biodiversity documentation for Kerala, Part 6: Flowering plants*. Kerala Forest Research Institute, Peechi.
- SHARMA B.D., SINGH N.P., RAGHAVAN R.S. & U.R. DESHPANDE 1984. *Flora of Karnataka analysis*, Series 2. Botanical Survey of India, Kolkata.
- STAPF O. 1892. On the Sonerileae of Asia. *Annals of Botany* 6(3): 291–323. <https://doi.org/10.1093/oxfordjournals.aob.a090681>
- STEARN W.T. 1992. *Latín Botánico: Historia, Gramática, Sintaxis, Terminología y Vocabulario*. Omega, David & Charles, London.
- THIERS B. 2022 (continuously updated). Index Herbariorum: a global directory of public Herbaria and associated staff. Available at: <http://sweetgum.nybg.org/ih> (Accessed on 31.11.2022).
- THWAITES G.H.K. 1864. *Enumeratio Plantarum Zeylaniae: an enumeration of Ceylon plants*. Dulau & Co., London.
- TRIANA J. 1871[1872]. Les Melastomacees. *Transactions of the Linnean Society of London* 28: 1–188. <https://doi.org/10.1111/j.1096-3642.1871.tb00222.x>
- TRIMEN H. 1894. *A hand-book to the flora of Ceylon: containing descriptions of all the species of flowering plants indigenous to the island, and notes on their history, distribution, and uses: with an atlas of plates illustrating some of the more interesting species*. Volume 2. Dulau & Co. London.
- WILLIS J.C. 1911. *Revised catalogue of the indigenous flowering plants and ferns of Ceylon*. Colombo: H. C. Cottle, Government Printer Government Press, Ceylon.
- YOGANARASIMHAN Y., SUBRAMANYAM K. & B.A. RAZI 1976. *Sonerila pedunculosa* Thw. (Melastomataceae) – A new record for India. *Current Science* 45(14): 527.