

Utricularia neottiioides

Fernando Rivadavia, Rua Inacio Pedroso 230, C.E.P.,
05612-050, Sao Paulo, S.P. Brazil

Utricularia neottiioides is a bizarre species native to the South American countries of Colombia, Venezuela, Bolivia, and Brazil (Taylor 1989). It grows anchored to rocks submerged by flowing water in cold mountain streams, a habit called rheophytic (Taylor 1989). I've observed on my trips around Brazil that *U. neottiioides* prefers the acidic, Coca-Cola-colored waters common of streams flowing down from sandstone highlands. As I've been told, this acidity and bizarre color are a result of the tannic acid present in the water. The water may look dirty to most people, but is actually almost sterile and excellent for drinking, especially since it's usually extremely cold!

I've found *U. neottiioides* growing at the Caraça Range, Cipó Range, around the town of Diamantina (all three in central Minas Gerais state), Canastra Range (southwestern M.Gerais), around the villages of Grão Mogol and Itacambira (both in northern M.Gerais), Ibitipoca Range (southeastern M.Gerais), Chapada dos Veadeiros (northern Goiás state, bordering the new state of Tocantins, which covers the top half of the old Goiás), Chapada dos Guimarães (southern Mato Grosso state), and have seen herbarium from many other localities in Brazil.

U. neottiioides flowers more profusely in shallow, slow-flowing water, most often during the dry season, though it can be found flowering all year long. During the wet season, the bare, stiff stolons are often almost invisible, fixed tightly in web-like patterns to the submerged rock surfaces. I've been fortunate enough to discover sites where *U. neottiioides* formed beautiful mats with thousands of flower scapes bearing tens of thousands of tiny greenish-white, yellowish-white, or cream flowers. Each peduncle bears 20 or more flowers and may reach 30cm in height (Taylor 1989), always flowering above the water level. Once at the Chapada dos Guimarães I observed *U. neottiioides* right after sunset and it seemed to me as if the flowers were slightly closed, with the upper lip bending together with the three lobes of the lower lip.

The leaves of *U. neottiioides* are unique in that they grow not only from the stolons, but also from the submerged, lowermost parts of the peduncles (Taylor 1989). Some kind of mucilage is produced by the leaves, which are long and feathery. At the Chapada dos Veadeiros I collected plants with leaves up to 20cm in length. The traps are extremely small and not very abundant either on the leaves or on the stolons, though at the Chapada dos Guimarães I've found leaves bearing numerous large red traps.

U. neottiioides forms Section Avesicaria together with *U. oliveriana*, a species native to Brazil, Colombia, and Venezuela (Taylor 1989). In Brazil, *U. oliveriana* has only been collected in the states of Pará, Rondônia, and Goiás (Taylor 1989). A few main differences between *U. oliveriana* and *U. neottiioides* are that the former has unbranched stolons, longer pedicels, and small, "normal-looking" *Utricularia* leaves (Taylor 1989). Unfortunately, I still haven't come across *U. oliveriana* in the wild. *U. neottiioides* is often found growing in spots of strong current in streams. It's a mystery to me how seeds are able to fasten themselves to and germinate on rocks in swift-flowing water. In this case, I imagine the seeds could have originally germinated in shallow water near the streamsides and then the plants could have spread slowly over the rocks to the central parts of the stream. The seeds in Section Avesicaria are said to be sticky (Taylor 1989), which would help in anchoring themselves to rocks. They probably only become sticky after contact with water, since the ones I've collected from dead peduncles are always dry.

Maybe the seeds are not truly shed and actually germinate directly from the fruit

while still on the flower scape. This would be an advantage since loose seeds in the water will surely be carried further downstream than a dead peduncle which will get stuck somewhere among the rocks sooner. Or maybe the flower scapes could bend over into the water once dead, while remaining attached to the stolons. That would allow the seeds to germinate directly from the fruit onto the submerged rocks.

Of course these would only work if the seeds maintain their viability after drying out, which is usually not the case with aquatic *Utricularia*. I'm not sure how it works for *U. neottiioides*, but I've seen dead plants covering the dry bed of an annual stream at the Chapada dos Guimarães. I don't think it's likely that a few fresh seeds were brought over by birds, wind, or whatever, and to have then developed into a fair-sized colony in a single season. Most likely, this population regenerates each year from a good quantity of seeds which just sit on the streambed until the wet season arrives. Maybe the dead scapes only release the seeds after the dry season is over, with the advent of rain.

Only on two occasions have I seen *U. neottiioides* growing in a habitat other than bare submerged rocks. At the Chapada dos Guimarães I found plants growing in gravel mixed with sand in a shallow stream, which is not too strange, but was surprised when I discovered plants at the Chapada dos Veadeiros growing in reddish dirt covering a streambed. Just to make sure, I even dug below the plants with my hands to be certain that the dirt wasn't only a thin layer above bare rock.

I've brought back live plants from my trips, but these never last very long in cultivation, rotting off in stagnant water. *U. neottiioides* grows in a very specific habitat and to cultivate it, you'd probably have to have a stream in your backyard. Live plants would have to be brought from the wild and quickly secured to rocks in your stream. Maybe gravel, sand or some type of soil mix might work too. But even if you did get live *U. neottiioides* to a stream near your house, there's the problem of water pH and chemical composition, which are probably very important for the survival of this exotic *Utricularia*.

References:

Taylor, Peter. 1989. The Genus *Utricularia* - A Taxonomic Monograph. Kew Bulletin Additional Series XIV, Royal Botanic Gardens, Kew.



Figure 1. *Utricularia neottiioides* growing in a shallow, rocky stream near Chapada dos Veadeiros, Brazil. Photo by Fernando Rivadavia.

CARNIVOROUS PLANT NEWSLETTER

Official Journal of the
International Carnivorous
Plant Society

Volume 25, Number 2
June 1996



Front cover: *Utricularia neottiioides* covering a streambed at the Chapada dos Veadeiros, Brazil. Photo by Fernando Rivadavia.

Rear Cover: *Utricularia tricolor* flower. Photo by R.B. Roth of Scottsville, Natal, South Africa.

The coeditors of CPN would like everyone to pay particular attention to the following policies regarding your dues to the ICPS.

All Correspondence regarding dues, address changes and missing issues should be sent to ICPS C/O Kevin Snively, P.O. Box 1013, Everett Wa. 98206-1013, U.S.A. or ICPS c/o Fullerton Arboretum, CSUF, Fullerton, CA 92634. DO NOT SEND TO THE COEDITORS. Checks for subscription and reprints should be made payable to ICPS.

All material for publication, comments and general correspondence about your plants, field trips or special noteworthy events relating to CP should be directed to one of the coeditors. We are interested in all news related to carnivorous plants and rely on the membership to supply us with this information so that we can share it with others.

Views expressed in this publication are those of the authors, not necessarily the editorial staff.

- President - Rick Walker, MS 26U4, P.O. Box 10350, Palo Alto, CA, 94303-0867
Phone: (415) 856-2354 Email: walker@opus.hpl.hp.com
V. Pres. - Christoph A. Belanger, 88 Strathmore Rd., Apt#4, Brookline, MA 02146
Phone: (617) 739-9795 Email: cbelan9630@aol.com
Sec/Treas - Kevin Snively, P.O. Box 1013, Everett Wa. 98206-1013
Phone: (206) 252-2911 Email: ksnive@premier1.net

Coeditors

- D.E. Schnell, Rt. 1, Box 145C, Pulaski, VA 24301
J.A. Mazrimas, 329 Helen Way, Livermore, CA 94550
Steve Baker, Rt. 1, Box 540-19AB, Conover, NC 28613-(flytrap@twave.net)
Seed Bank: Tom Johnson, P.O. Box 12281, Glendale, CA 91224-0981., (Carnipla@aol.com)
Business Manager: Leo C. Song, Jr. (LeoSong@fullerton.edu)

PUBLISHER: The International Carnivorous Plant Society by the Fullerton Arboretum, California State University, Fullerton, CA 92634. Published quarterly with one volume annually. Desktop Publishing: Steve Baker, Rt. 1, Box 540-19AB, Conover, NC 28613. Printer: Kandid Litho, 1077 East Enda Place, Covina, CA 91724. Masthead Art: Paul Milauskas, 38 Manchester Court, Fox River Grove, IL 60021. Dues: \$15.00 annually. \$20.00 foreign. Reprints available by volume only © 1996 Carnivorous Plant Newsletter. All rights reserved. ISSN #0190-9215. Circulation 865 yearly.

CARNIVOROUS PLANT NEWSLETTER

VOLUME 25, NUMBER 2

JUNE 1996

