

Screening tropical and sub-tropical maize germplasm for resistance to *Striga hermonthica* and *S. asiatica* and yield-related traits

Emeline N. Dossa

dossaemeline@gmail.com

University of KwaZulu-Natal

Hussein Shimelis

University of KwaZulu-Natal

Admire I. T. Shayanowako

University of KwaZulu-Natal

Mark D. Laing

University of KwaZulu-Natal

Research Article

Keywords: Dual resistance, *Striga asiatica*, *Striga hermonthica*, maize breeding

Posted Date: November 1st, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3507906/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Euphytica on March 15th, 2024. See the published version at <https://doi.org/10.1007/s10681-024-03309-2>.

Abstract

Identification of maize germplasm with dual resistance to *Striga hermonthica* (*Sh*) and *S. asiatica* (*Sa*), could lead to the development of cultivars with stable resistance. 130 tropical and sub-tropical maize germplasms were evaluated in a controlled environment for their reaction to *Sh* and *Sa* infestations using a 13x10 alpha lattice design with two replications over two seasons. Significant differences ($P < 0.05$) were detected among the assessed genotypes for all the recorded traits in *Sh* and *Sa*-infested treatments. Under *Sa*-infested conditions, mean *Striga* emergence counts 8 weeks after planting (SEC8) and 10 weeks after planting (SEC10) were 5.00 and 45.50 respectively, while the mean *Striga* damage rate 8 weeks after planting (SDR8) and 10 weeks after planting (SDR10) were 3.35 and 3.07, respectively. Under *Sh*-infested conditions, SEC8 and SEC10 mean values were 3.66 and 3.77, respectively, while the SDR8 and SDR10 values were 5.25 and 2.75 respectively. Positive and significant ($P < 0.05$) correlations were found between anthesis-silking interval (ASI) and SDR8 ($r = 0.18$) and SDR10 (0.32) under *Sa*-infested conditions. Negative and significant correlations were recorded between ear per plant (EPP) and SEC8, SDR8, and SDR10, with $r = -0.18$, $r = -0.27$, and $r = -0.24$, respectively. Under *Sh*-infested conditions, significant and negative correlations were recorded between SDR8 and EPP ($r = -0.20$), EHT and SEC8 ($r = -0.22$), EHT and SDR8 ($r = -0.36$), PLHT and SDR8 (-0.48), and PLHT and SDR10 (-0.22). The results suggest that dual resistance to the two *Striga* species exists in some tropical and sub-tropical maize lines. The following genotypes have dual resistance to *Sa* and *Sh*: CML440, CML566, CML540, CML539, CLHP0343, CLHP0326, TZISTR1248, TZSTR115, TZISTR25, TZISTR1205, TZSTR113, TZISTR1119, TZISTR1174 and the OPVs B.King/1421, Shesha/1421, ZM1421, DTSTR-W SYN13, DTSTR-Y SYN14, and 2*TZECOMP3DT/WhiteDTSTRSYN) C2. The identified genotypes are suitable for use as parents in developing and deploying high-performing maize varieties with *Striga* resistance and improved grain yield.

1. Introduction

Maize (*Zea mays* L., $2n = 2x = 20$) is the world's third most widely cultivated cereal crop, after wheat and rice. It is a vital food security crop in sub-Saharan Africa (SSA), constituting 85–95% of the region's carbohydrate intake (Johnmark et al., 2022). In SSA, maize is a source of livelihood for more than 300 million Africans (Regassa et al., 2021; Kansiime et al., 2023). It is a raw material for manufacturing industrial products, including livestock feed (Dabija et al., 2021). Despite the importance of the cereal, one out of five people living in communal and small-scale maize farming systems are at risk of starvation (Arndt et al., 2023). This is because the crop's is vulnerable to many stress factors, one major biotic stress being parasitic weeds of the genus *Striga*.

Striga, popularly known as witchweed, is endemic and widespread throughout SSA. The species *S. asiatica* (red flower type) is predominant in Southern Africa, and *S. hermonthica* (purple flower) is widely distributed in Western, Central, and Eastern Africa (Dossa et al., 2023). The two holoparasites cause severe crop damage through stunting and leaf chlorosis, leading to yield losses ranging from 30–100% under severe infestation (Mutsvanga et al., 2022). Maize suffers yield losses caused by these parasites due to the paucity of *Striga* resistance sources in the maize gene pool, and only partial resistance has been reported. Over 50 million hectares of agricultural land under cereal cultivations, including maize, have been infested by *Striga* spp (Dafaallah, 2019; David et al., 2022). The annual losses caused by *Striga* have been estimated to USD 10 billion across SSA (Dafaallah, 2019; Samejima and Sugimoto, 2022). Under heavy *Striga* infestations farmers are often forced to abandon their farms.

Cultural practices, crop protection chemicals, biological control, host plant resistance, and integrated *Striga* management approach are the main control strategies recommended for *Striga* management. Germination stimulants such as ethylene and ethephon, when applied on infested croplands before sowing, can deplete *Striga* seed banks by inducing suicidal germination without a host (Samejima et al., 2016). Imidazolinone herbicides are also effective against *Striga* in the field (Kanampiu et al., 2003). Maize seeds with imazapyr resistance can be coated with small doses of the herbicide, significantly reducing *Striga* emergence (David et al., 2011; Kamara et al., 2020). However, using imazapyr-coated maize in a smallholder maize production system is difficult since the chemical is toxic to other crops (Souto et al., 2020). Most of the proposed control methods are impractical to implement under communal and small-scale farming systems. Most farmers do not have enough land to crop rotate, nor do they have access to large quantities of organic matter for effective cultural control practices (Shayanowako et al., 2020). Furthermore, they do not have access to the money required to buy agrochemicals and sprayers. Hence, the development and use of *Striga*-resistant cultivars is the most feasible management option (Gasura et al., 2021; Dossa et al., 2023).

Striga-resistant cultivars can reduce *Striga* seed production and the *Striga* seed bank in infested soils (Badu-Apraku et al., 2020a). Resistant cultivars can induce the germination of *Striga* seeds but prevent the parasite from attaching to the maize plant. Concerted efforts have been made by the International Institute of Tropical Agriculture (IITA) in Nigeria and national maize research programs to develop tropical maize genotypes with resistance to *Sh* (Badu-Apraku et al., 2020b; Yacoubou et al., 2021a). However, the performance of their germplasm against *Sa* is unknown. High-yielding and *Striga*-resistant sub-tropical maize varieties are yet to be developed (Shayanowako et al., 2018a). Therefore, the IITA genetic resources can serve as breeding parents with resistance to *Sh*, and need to be evaluated for their levels of resistance to *Sa*, and for yield improvements under *Sa*-infestation. Likewise, sub-tropical maize varieties should be screened for *Striga* resistance and yield performance under both *Sa* and *Sh* infestation. This can also benefit West and Central Africa with a diversity of resistance genes that could be useful for accumulating improved levels of partial resistance.

Striga emergence count, *Striga* damage rating, and grain yield under *Striga* infestation are the major selection indices used in resistance breeding (Menkir et al., 2007). The choice of the selection method to be used in the genetic improvement of maize depends on the type of gene action controlling *Striga* resistance in maize. *Striga* resistance in maize is quantitatively inherited with many minor genes with small additive effects and is significantly influenced by the environment (Lane et al., 1997; Badu-Apraku et al., 2020a), making breeding complex and challenging. Hybrid breeding, backcrossing, and recurrent selection are common methods widely used in incorporating *Striga* tolerance/resistance genes into well-adapted maize varieties. However, the initial steps of these methods include the collection and evaluation of maize germplasm with different genetic backgrounds to identify potential sources of resistance (Yacoubou et al., 2021b). Screening for *Striga* resistance in maize includes field, greenhouse, and laboratory conditions (Shayanowako et al., 2018b). However, the use of greenhouse conditions is the most efficient for managing the level of *Striga* infestation and environmental conditions (Kountche et al., 2019; Yacoubou et al., 2021b).

The existing *Striga* resistant cultivars of maize in SSA are bred for *Sh* resistance, while no commercially grown maize varieties are resistant to *Sa*. In most of the East African countries, the two species occur in tandem (Gethi and Smith, 2004). Germplasm with dual resistance to both parasites would be extremely valuable across the continent. This study aimed to evaluate 130 tropical and sub-tropical maize germplasms in a controlled environment for their reaction to *Sh* and *Sa* infestations and for resistance breeding. This study is one of the few attempts to report on the performance of tropical and sub-tropical maize germplasm under both *Sa* and *Sh* infestation to select inbred lines with dual resistance to the two dominant *Striga* species.

2. Materials and Methods

2.1 Plant material and study sites

The study screened maize genotypes for *Striga* resistance at the University of Kwazulu-Natal Controlled Environment Facilities (UKZN-CEF) in two cropping seasons of 2022. The UKZN CEF is situated at the UKZN College of Agriculture, Engineering, and Science (29.62° S, 30.40° E). The study used 130 maize genotypes, comprising 74 acquired from the International Institute of Tropical Agriculture (IITA)/Nigeria, 45 from the International Maize and Wheat Improvement Centre, Zimbabwe (CIMMYT)/Zimbabwe, and 10 from the National Plant Genetic Resources Centre, South Africa (NPGRC)/South Africa (Table 1). The germplasm from IITA/Nigeria comprised 55 inbred lines (genotypes number 1 to 55 in Table 1), 4 single cross hybrids (genotypes number 127 to 130 in Table 1), and 15 open-pollinated varieties (OPV) (genotypes number 112 to 126 in Table 1). The CIMMYT/Zimbabwe germplasm included 43 inbred lines (genotype number 56 to 98 in Table 1), and 1 OPV (genotype number 110 in Table 1). The NPGRC/South Africa germplasm included 5 OPVs (genotype 99 to 103 Table 1) and 6 hybrids (genotype 104 to 109 Table 1). The Nigerian accessions were tropical varieties developed for *Sh* resistance, generated from multi-parent populations (Menkir, 2006; Simon et al., 2018; Gasura et al., 2019). CIMMYT provided sub-tropical maize germplasm. Their material was developed for drought tolerance. The OPVs and hybrids germplasms from NPGRC/South Africa and CIMMYT/Zimbabwe were used as local checks, while the OPVs and hybrids from IITA were *Sh*-resistant checks. The *Sa* and *Sh* seeds were collected from sorghum and maize-infested fields in Tanzania and Kenya, respectively.

Table 1
List and source of maize genotypes evaluated in the present study.

N°	Germplasm name/designation	Source/Origin	<i>Striga</i> resistance /breed description
1	TZISTR1154	IITA/Nigeria	Resistant/inbred line
2	TZISTR1261	IITA/Nigeria	Resistant/inbred line
3	TZISTR1248	IITA/Nigeria	Resistant/inbred line
4	TZISTR1263	IITA/Nigeria	Resistant/inbred line
5	TZISTR1275	IITA/Nigeria	Resistant/inbred line
6	TZISTR1157	IITA/Nigeria	Resistant/inbred line
7	TZISTR1160	IITA/Nigeria	Resistant/inbred line
8	TZISTR1162	IITA/Nigeria	Resistant/inbred line
9	TZISTR1165	IITA/Nigeria	Resistant/inbred line
10	TZISTR1175	IITA/Nigeria	Resistant/inbred line
11	TZISTR1178	IITA/Nigeria	Resistant/inbred line
12	TZISTR1163	IITA/Nigeria	Resistant/inbred line
13	TZISTR1166	IITA/Nigeria	Resistant/inbred line
14	TZISTR1190	IITA/Nigeria	Resistant/inbred line
15	TZISTR1199	IITA/Nigeria	Resistant/inbred line
16	TZISTR1231	IITA/Nigeria	Resistant/inbred line
17	TZISTR1232	IITA/Nigeria	Resistant/inbred line
18	TZISTR1259	IITA/Nigeria	Resistant/inbred line
19	TZISTR1262	IITA/Nigeria	Resistant/inbred line
20	TZISTR1159	IITA/Nigeria	Resistant/inbred line
21	TZISTR1223	IITA/Nigeria	Resistant/inbred line
22	TZISTR1225	IITA/Nigeria	Resistant/inbred line
23	TZISTR1244	IITA/Nigeria	Resistant/inbred line
24	TZSTRI101	IITA/Nigeria	Resistant/inbred line
25	TZSTRI102	IITA/Nigeria	Resistant/inbred line
26	TZSTRI104	IITA/Nigeria	Resistant/inbred line
27	TZSTRI107	IITA/Nigeria	Resistant/inbred line
28	TZSTRI108	IITA/Nigeria	Resistant/inbred line
29	TZSTRI109	IITA/Nigeria	Resistant/inbred line
30	TZSTRI110	IITA/Nigeria	Resistant/inbred line
31	TZSTRI112	IITA/Nigeria	Resistant/inbred line
32	TZSTRI114	IITA/Nigeria	Resistant/inbred line
33	TZSTRI115	IITA/Nigeria	Resistant/inbred line
34	TZISTR25	IITA/Nigeria	Resistant/inbred line
35	TZISTR1001	IITA/Nigeria	Resistant/inbred line
36	TZISTR1003	IITA/Nigeria	Resistant/inbred line
37	TZISTR1004	IITA/Nigeria	Resistant/inbred line
38	TZISTR1008	IITA/Nigeria	Resistant/inbred line
39	TZISTR1011	IITA/Nigeria	Resistant/inbred line
40	TZISTR1018	IITA/Nigeria	Resistant/inbred line
41	TZEEI21	IITA/Nigeria	Resistant/inbred line

N°	Germplasm name/designation	Source/Origin	<i>Striga</i> resistance /breed description
42	TZEEI13	IITA/Nigeria	Resistant/inbred line
43	TZEEI14	IITA/Nigeria	Resistant/inbred line
44	TZEEI49	IITA/Nigeria	Resistant/inbred line
45	TZDEEI55	IITA/Nigeria	Resistant/inbred line
46	TZDEEI50	IITA/Nigeria	Resistant/inbred line
47	TZEEI34	IITA/Nigeria	Resistant/inbred line
48	TZISTR1174	IITA/Nigeria	Resistant/inbred line
49	TZISTR1205	IITA/Nigeria	Resistant/inbred line
50	TZSTRI113	IITA/Nigeria	Resistant/inbred line
51	TZISTR1119	IITA/Nigeria	Resistant/inbred line
52	TZISTR1015	IITA/Nigeria	Resistant/inbred line
53	TZDEEI64	IITA/Nigeria	Resistant/inbred line
54	TZDEEI54	IITA/Nigeria	Resistant/inbred line
55	TZEEI10	IITA/Nigeria	Resistant/inbred line
56	CML312	CIMMYT/Zimbabwe	Unknown/Inbred line
57	CML444	CIMMYT/Zimbabwe	Unknown/Inbred line
58	CML442	CIMMYT/Zimbabwe	Unknown/Inbred line
59	CML550	CIMMYT/Zimbabwe	Unknown/Inbred line
60	CML547	CIMMYT/Zimbabwe	Unknown/Inbred line
61	CML539	CIMMYT/Zimbabwe	Unknown/Inbred line
62	CML440	CIMMYT/Zimbabwe	Unknown/Inbred line
63	CML566	CIMMYT/Zimbabwe	Unknown/Inbred line
64	CML540	CIMMYT/Zimbabwe	Unknown/Inbred line
65	CML545	CIMMYT/Zimbabwe	Unknown/Inbred line
66	CML571	CIMMYT/Zimbabwe	Unknown/Inbred line
67	CML390	CIMMYT/Zimbabwe	Unknown/Inbred line
68	CLHP0352	CIMMYT/Zimbabwe	Unknown/Inbred line
69	HA04A-2107-36	CIMMYT/Zimbabwe	Unknown/Inbred line
70	CLHP0303	CIMMYT/Zimbabwe	Unknown/Inbred line
71	CLHP0221	CIMMYT/Zimbabwe	Unknown/Inbred line
72	CLHP0020	CIMMYT/Zimbabwe	Unknown/Inbred line
73	CLHP0058	CIMMYT/Zimbabwe	Unknown/Inbred line
74	CKDHL0378	CIMMYT/Zimbabwe	Unknown/Inbred line
75	CLHP0312	CIMMYT/Zimbabwe	Unknown/Inbred line
76	CLHP0310	CIMMYT/Zimbabwe	Unknown/Inbred line
77	CLHP0003	CIMMYT/Zimbabwe	Unknown/Inbred line
78	CKDHL0467	CIMMYT/Zimbabwe	Unknown/Inbred line
79	CLHP00378	CIMMYT/Zimbabwe	Unknown/Inbred line
80	CLHP0156	CIMMYT/Zimbabwe	Unknown/Inbred line
81	CLHP0113	CIMMYT/Zimbabwe	Unknown/Inbred line
82	CLHP03302	CIMMYT/Zimbabwe	Unknown/Inbred line
83	CLHP0404	CIMMYT/Zimbabwe	Unknown/Inbred line
84	CLHP0343	CIMMYT/Zimbabwe	Unknown/Inbred line

N°	Germplasm name/designation	Source/Origin	<i>Striga</i> resistance /breed description
85	CZL1380	CIMMYT/Zimbabwe	Unknown/Inbred line
86	CLHP0326	CIMMYT/Zimbabwe	Unknown/Inbred line
87	CZL99017	CIMMYT/Zimbabwe	Unknown/Inbred line
88	CLHP0049	CIMMYT/Zimbabwe	Unknown/Inbred line
89	CLHP00478	CIMMYT/Zimbabwe	Unknown/Inbred line
90	CLHP00286	CIMMYT/Zimbabwe	Unknown/Inbred line
91	CML451	CIMMYT/Zimbabwe	Unknown/Inbred line
92	CLHP0302	CIMMYT/Zimbabwe	Unknown/Inbred line
93	CLHP0364	CIMMYT/Zimbabwe	Unknown/Inbred line
94	CLHP0350	CIMMYT/Zimbabwe	Unknown/Inbred line
95	CLHP00294	CIMMYT/Zimbabwe	Unknown/Inbred line
96	CLHP0005	CIMMYT/Zimbabwe	Unknown/Inbred line
97	CLHP0022	CIMMYT/Zimbabwe	Unknown/Inbred line
98	CML304	CIMMYT/Zimbabwe	Unknown/Inbred line
99	ZM1423/Z.DLO	NPGRG/South Africa	Unknown/local OPV
100	NC.QPM/Z.DPLO	NPGRG/South Africa	Unknown/local OPV
101	M.Pearl/DT-STR	NPGRG/South Africa	Unknown/local OPV
102	NC.QPM/DT-STR	NPGRG/South Africa	Unknown/local OPV
103	ZM1421/DT-STR	NPGRG/South Africa	Unknown/local OPV
104	N.Choice/1421	NPGRG/South Africa	Unknown/local hybrid
105	B.King/1421	NPGRG/South Africa	Unknown/local hybrid
106	Colorado/1421	NPGRG/South Africa	Unknown/local hybrid
107	Hickory/1421	NPGRG/South Africa	Unknown/local hybrid
108	Kep/1421	NPGRG/South Africa	Unknown/local hybrid
109	Shesha/1421	NPGRG/South Africa	Unknown/local hybrid
110	ZM1423	CIMMYT/Zimbabwe	Unknown/local OPV
111	ZM1421	CIMMYT/Zimbabwe	Unknown/local hybrid
112	STR-SYN-Y2	IITA/Nigeria	Resistant/OPV
113	Z.diplo-BC4-C3-W/DOGONA-1/Z.diplo-BC4-C3-W	IITA/Nigeria	Resistant/OPV
114	Z. Diplo.BC4C3-W-DT C1	IITA/Nigeria	Resistant/OPV
115	TZBSTR (Susceptible)(RE)	IITA/Nigeria	Resistant/OPV
116	STR-SYN-W1	IITA/Nigeria	Resistant/OPV
117	DTSTR-W SYN13	IITA/Nigeria	Resistant/OPV
118	DTSTR-Y SYN15	IITA/Nigeria	Resistant/OPV
119	((IWD C3 SYN*2/(White DT STR Syn))-DT C1	IITA/Nigeria	Resistant/OPV
120	DTSTR-W SYN11	IITA/Nigeria	Resistant/OPV
121	SAMMMZ16	IITA/Nigeria	Resistant/OPV
122	(TZEOMP5C7/TZECOMP3DTC2) C2	IITA/Nigeria	Resistant/OPV
123	((TZL COMP1-W C6*2/(White DT STR Syn))-DT C1	IITA/Nigeria	Resistant/OPV
124	TZCOM1/ZDPSYN	IITA/Nigeria	Resistant/OPV
125	DTSTR-Y SYN14	IITA/Nigeria	Resistant/OPV
126	(2*TZECOMP3DT/WhiteDTSTRSYN) C2	IITA/Nigeria	Resistant/OPV
127	TZSTR1137/TZSTR1132	IITA/Nigeria	Resistant/hybrid

N°	Germplasm name/designation	Source/Origin	<i>Striga</i> resistance /breed description
128	TZSTR1159/TZSTR1132	IITA/Nigeria	Resistant/hybrid
129	TZSTR1160/TZSTR1132	IITA/Nigeria	Resistant/hybrid
130	TZSTR1166/TZSTR1132	IITA/Nigeria	Resistant/hybrid

CIMMYT, International Maize and Wheat Improvement Centre; IITA, International Institute of Tropical Agriculture; NPGRC/SA, National Plant Genetic Resources Centre/South Africa, OPV, open-pollinated variety

2.2 Experimental design and trial management

The 130 genotypes were evaluated under two *Striga* treatments using a 13 x 10 alpha lattice design with two replications in each *Striga*-infested environment. The maize genotypes were evaluated under *Sa* and *Sh* infestations. The experimental unit consisted of 4 plastic pots of 15-L capacity, filled with a composted pine bark potting mix for each *Striga*-infested environment. Two weeks before planting, each pot was infested with a scoop of sand mixed with 0.03 g of two years old *Sa* or *Sh* seed. Standard agronomic practices recommended for maize production were followed. Hand weeding was routinely done to remove all weeds except *Striga*. Figure 1 shows the experimental setup with artificial infestations of maize with *Sa* (A, B, and C), and *Sh* (D and E) in the greenhouse condition at the University of Kwazulu-Natal, South Africa.

2.3 Data collection

Data were collected on maize phenotypic traits and *Striga* parameters in the *Sa* and *Sh*-infested environments. The following phenotypic traits were evaluated on maize: Days to 50% silking (DS), recorded as the number of days taken by 50% of the plants to silk in each plot; Days to anthesis (DA), recorded as the number of days from planting until 50% of the plants have emerged silks and shed pollen, respectively; Anthesis-silking interval (ASI), measured as the difference between days to 50% silking and 50% anthesis; Plant height (PLHT) and ear height (EHT), measured as the distance from the base of the plant to the height of the first tassel branch and the node bearing the upper ear, respectively; Root lodging (RL) was recorded as a percentage of plants leaning more than 30° from the vertical; Stalk lodging (SLG) (percentage broken at or below the highest ear node); and Number of rotten ears (EROT). The number of ears per plant (EPP) was obtained by dividing the total number of ears per plot by the number of plants harvested. Husk cover (HUSK) was rated on a scale of 1 to 5, where 1 = husks tightly arranged and extended beyond the ear tip and 5 = ear tips exposed. Ear aspect (EASP) was recorded based on a scale of 1 to 9, where 1 = clean, uniform, large, and well-filled ears and 9 = ears with undesirable features. Grain yield per plant (GY/plant) was determined as the weight (g) of the grain from the ears of individual plants after shelling, adjusted to a constant moisture of 12.5%.

The *Striga* parameters were recorded, including the number of emerged *Sa* and *Sh* plants 8 and 10 weeks after planting, denoted as SEC8 and SEC10. A rating of host plant damage 8 and 10 weeks after planting, designated as SDR8 and SDR10, was done using a visual rating score of 1 to 9 where 1 = no damage, indicating normal plant growth and a high level of tolerance, and 9 = complete collapse or death of the maize plant, i.e., highly susceptible (Kim, 1994).

2.4 Data-analysis

The collected data from *Sa* and *Sh*-infested environments were subjected to analysis of variance using a lattice procedure, using the package *agricolae* in RStudio version 2023. 06.1 (R Core Team, 2023). The mean values of the test genotypes for the assessed traits were compared at the 5% significance level using Fisher's least significance difference (LSD). Broad sense heritability (H^2) (hereafter referred to as heritability) was computed using DeltaGen (Jahufer and Luo, 2018) with the following formula:

$$(H^2) = \frac{\sigma^2 g}{\sigma^2 g + \frac{\sigma^2 s}{ns} + \frac{\sigma^2 r}{nr} + \frac{\sigma^2 b}{nb} + \frac{\sigma^2}{ns+nr+nb}}$$

where $\sigma^2 g$, $\sigma^2 s$, $\sigma^2 r$, $\sigma^2 b$, and σ^2 are the variance components for genotypes, season, replication, block, and the pooled error, respectively, and ns, nr, and nb are the number of seasons, replications, and blocks, respectively.

Pearson's correlation coefficients (r) were calculated separately for *Sa* and *Sh*-infested conditions using RStudio version 4.3.1 (Team, 2010). The rotated component matrix and principal component analysis biplots (PCA) were generated separately for the assessed traits under *Sa*, and *Sh*-infested conditions using the packages *ggplot2*, *factoextra*, and *FactoMiner* (Alboukadel, 2017) in RStudio version 4.3.1. Cluster heatmap plots were generated based on the mean values of the traits recorded in both *Sa* and *Sh* environments to establish the Clustering of the genotypes using Deltagen (Jahufer and Luo, 2018).

3. Results

3.1 Analysis of variance (ANOVA)

The analysis of variance revealed significant differences among the evaluated genotypes for all the recorded traits ($P < 0.001$) (Table 2) under both *Sa* and *Sh*-infested environments. Testing seasons exerted significant effects ($P < 0.001$) on all the traits under *Sa*-infested conditions except for EPP, PLHT, HUSK, and SEC10, and under *Sh*-conditions except for EPP, PLHT, EHT, and HUSK.

Table 2 Analyse of variance for yield components and *Striga* parameters of 126 maize genotypes evaluated under *Sa* and *Sh* infestation conditions.

<i>S. asiatica</i>												
Source of variation	Df	DA	DS	ASI	EPP	PLHT	EHT	HUSK	CL	EASP	GY	SEC8
Genotypes (G)	125	9.34***	7.27***	4.06***	6.49***	2.10***	5.82***	3.65***	9.95***	2.47***	4.05***	2.79***
Seasons (S)	1.00	273.78***	100.64***	33.01***	0.00	0.27	20.82***	0.00	29.74***	43.79***	54.71***	1518.79***
G x S	125	0.0666	0.00	0.084	0.00	0.00	0.00	0.06	0.00	0.036	0.00	0.54
Replications in seasons	1.00	0.35	0.00	0.44	0.00	0.00	0.00	0.00	0.00	0.39	0.00	51.62***
Blocks in seasons	12.00	1.20	0.78	3.34***	0.00	2.01*	3.73***	0.40	4.37***	2.65**	0.85	24.71***
Error	238.00	15.40	20.03	12.28	0.03	106.99	0.05	0.28	2.08	3.47	1439.00	1.51
<i>S. hermonthica</i>												
Source of variation	df	DA	DS	ASI	EPP	PLHT	EHT	HUSK	CL	EASP	GY	SEC8
Genotypes (G)	125	2.43***	2.29***	1.76***	3.01***	2.17***	1.84***	1.48**	1.47**	2.76***	2.45***	4.97***
Seasons (S)	1.00	125.84***	158.08***	32.01***	0.00	3.14	23.37	1.03	17.77***	18.89***	72.91***	3255.33***
G x S	125	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.00	0.00
Replications in seasons	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.26	0.00	0.00	0.00	0.00
Blocks in seasons	18.00	0.73	1.33	3.18***	0.00	1.68	0.21	0.10	0.08	0.27	1.50	0.54
Error	233.00	36.00	51.00	15.75	0.03	3.61	0.08	0.49	7.09	5.40	1473.00	1.20

* denote significant difference at $P < 0.05$, ** at $P < 0.01$ and *** at $P < 0.001$, Df = degrees of freedom, DA = days to 50% anthesis, DS = days to 50% silking, ASI = anthesis-silking interval, EPP = ear per plant, PLHT = Plant height, EHT = ear height, HUSK = husk cover, CL = cob length, EASP = ear aspect, GY = grain yield, SEC8 = *Striga* emergence counts eight days after sowing, SEC10 = *Striga* emergence counts ten days after sowing, SDR8 = *Striga* damage rating eight days after sowing, and SDR10 = *Striga* damage rating 10 days after sowing.

3.2 Mean performance and statistical summary

3.2.1 Under *Sa*-infested conditions

Table S1 summarizes the mean performance and the statistics of the 126 genotypes evaluated under *Sa* infestation. The coefficient of variation (CV%) of all the traits under *Sa* infestation ranged from 4.89–426.82% (Table 3). Smaller variations were obtained for DA followed by DS, with CV values of 4.89% and 5.60%, respectively. The highest variation was exhibited by PLHT followed by ASI, with CV values of 426.82% and 268.88%, respectively. Inbred lines had a mean ASI of 2.77, while the OPV and hybrid checks had mean ASI values of 1.86 and 1.77, respectively. The EPP ranged from 1.00 to 2.00. The mean yield ranged from 0.00 g/plant to 277.50 g/plant for TZISTR1262 and CML540, respectively, with a mean of 62.77 g/plant for the inbred lines, while ranging from 00.00 g/plant to 214.00 g/plant for Hickory/1421 and N.Choice/1421, respectively for the hybrids checks, and from 35.00 g/plant to 169.50 g/plant (((IWD C3 SYN*2/(White DT STR Syn)) -DT C1 and NC.QPM/Z.DPLO respectively) for the OPVs checks. The top inbred lines and checks are shown in Table 3. The best-yielding genotypes were generally taller than the poorer-performing genotypes, with the longest cobs and moderate or high EASP. The greatest reduction in *Striga* emergence in SEC8 occurred with relatively high yielding genotypes, e.g., TZISTR1154 (2.00), TZISTR1263 (2.00), TZISTR1261 (2.50), TZISTR1015 (2.50), TZISTR1174 (3.00), TZISTR1113 (3.00), TZISTR1119 (3.00), TZISTR1205 (3.50), TZISTR1248 (3.50), the local OPVs M.Pearl/DT-STR (2.00), ZM1421/DT-STR (2.50), and the *Striga*-resistant checks DTSTR-Y SYN14 (2.50), and NC.QPM/DT-STR (3.00). The SEC10 mean was 45.50, 10.32, and 9.07 for the inbred lines, the OPVs, and the hybrids, respectively. Genotypes such as inbred line TZISTR1174 and *Striga*-resistant check OPV DTSTR-W SYN11 exhibited high numbers for SEC10, and are still relatively high yielding. The inbred lines exhibited an SDR8 mean value of 3.35, while the OPVs and hybrids showed an SDR8 of 3.14 and 2.36, respectively. SDR10 mean scores were 3.07, 4.30, and 2.71 for the inbred lines, the OPVs, and the hybrids checks, respectively. Genotypes with high yields generally showed moderate or large reductions in SDR8 and SDR10 values. The following high-yielding genotypes displayed relatively high and moderate reductions in scores for SDR8 and SDR10: CML566, CML440, TZISTR1248, ZM1421, and local check (B.King/1421). None of the genotypes exhibited SLG, RL, and EROT under *Sa* infestation. High broad-sense heritability was recorded for CL (0.97), EHT (0.96), EASP (0.96), DS (0.94), ASI (0.94), DA (0.90), and GY (0.88) under *Sa*-infested conditions. However, the low heritability of *Sa* resistance is worth noting, reflected in the *Striga* parameters SEC10, SDR8, and SDR10 under *Sa* infested conditions. In contrast, under *Sh*-infested conditions, high heritability values were estimated for all the traits except for GY (0.02).

Table 3
Mean responses for 14 traits of 126 maize genotypes evaluated under *Sa* infestation, showing the top 10 inbred lines and the top 4 hybrids and 6 OPVs.

Top 10 inbred lines														
Genotype	DA	DS	ASI	EPP	PLHT	EHT	HUSKCL	EASPGY	SEC8	SEC10	SDR8	SDR10		
CML540	77.00	81.50	4.50	1.00	2.03	0.77	1.00	11.00	3.50	277.50	4.00	1.50	3.50	3.00
CML566	82.50	79.00	3.50	1.00	2.22	1.15	1.00	12.00	1.50	155.50	4.00	5.50	1.50	1.50
TZISTR1001	82.00	82.00	0.00	1.00	2.10	1.28	1.00	11.00	1.50	140.00	4.50	4.50	3.00	2.50
TZISTR1205	81.50	75.50	6.00	1.00	1.85	0.91	1.00	11.00	1.00	114.25	3.50	13.00	3.00	2.50
TZSTR115	77.50	77.00	0.50	1.00	2.10	1.20	1.00	11.50	1.50	112.50	5.00	2.00	3.50	2.50
CLHP0350	75.00	76.00	1.00	1.00	2.35	0.81	3.00	14.00	3.50	102.75	5.00	3.50	2.00	3.50
CLHP0049	80.50	78.00	2.50	1.00	1.25	0.70	1.00	10.00	3.00	101.25	7.00	4.00	1.00	2.50
CLHP0302	81.00	80.50	0.50	1.00	1.76	1.00	3.00	13.25	3.00	98.00	4.50	7.00	2.00	3.50
CML440	82.50	76.00	6.50	1.00	2.36	1.08	1.00	11.00	1.50	96.25	4.50	13.50	3.00	1.50
CLHP0303	84.50	83.50	1.00	1.50	1.87	1.15	1.00	7.25	3.00	92.50	4.50	8.50	3.00	3.00
Top 4 hybrids and 6 OPVs														
N.Choice/1421	82.00	76.50	5.50	1.00	1.90	1.03	1.00	13.25	1.50	214.00	5.00	4.00	3.50	3.50
Shesha/1421	75.50	72.50	3.00	1.00	2.03	1.75	1.00	18.75	1.50	165.75	7.50	18.50	2.00	2.00
B.King/1421	80.50	78.50	2.00	1.00	2.05	1.15	2.00	23.50	1.50	157.25	5.00	4.50	1.00	2.00
ZM1421/DT-STR	77.00	76.50	0.50	1.00	2.38	1.10	1.50	10.75	3.00	93.50	2.50	19.00	2.50	2.00
NC.QPM/Z.DPLO	71.50	71.50	0.00	1.00	2.25	1.03	0.00	12.50	4.50	154.25	3.00	5.00	3.50	2.00
Z.diplo-BC4-C3-W/DOGONA-1/Z.diplo-BC4-C3-W	81.00	83.00	2.00	1.00	2.36	1.05	1.00	11.50	1.50	112.00	6.00	8.50	3.00	3.50
DTSTR-W SYN13	85.50	85.50	0.00	1.00	1.25	0.85	1.50	13.00	3.50	107.50	4.50	3.50	1.50	1.00
TZBSTR	83.00	83.50	0.50	1.00	2.65	1.30	1.50	14.50	1.00	103.00	6.50	3.00	3.00	2.50
ZM1423	69.00	69.50	0.50	1.00	0.85	1.39	1.00	10.50	1.50	99.25	4.50	16.50	5.00	3.00
(2*TZECOMP3DT/WhiteDTSTRSYN) C2	69.00	78.00	9.00	1.00	1.75	0.85	1.50	12.25	2.50	89.00	6.00	0.50	2.00	1.50
Trial statistics														
LSD (5%)	3.94	4.44	3.69	0.15	10.57	0.23	0.52	1.55	1.93	37.73	1.87	12.44	1.64	1.61
SEM	7.64	8.75	7.27	0.21	20.82	0.46	1.02	3.05	3.80	74.31	3.69	24.51	3.23	3.17
%CV	4.89	5.60	268.88	14.65	426.82	25.36	41.21	13.63	58.57	56.12	40.91	129.74	50.29	49.50
Heritability	0.90	0.94	0.94	0.11	0.11	0.96	0.96	0.97	0.96	0.88	0.34	0.01	0.11	0.16

DA = days to 50% anthesis, DS = days to 50% silking, ASI = anthesis-silking interval, EPP = ear per plant, PLHT = Plant height, EHT = ear height, HUSK = husk cover, CL = cob length, EASP = ear aspect, GY = grain yield, SEC8 = *Striga* emergence counts eight days after sowing, SEC10 = *Striga* emergence counts ten days after sowing, SDR8 = *Striga* damage rating eight days after sowing, and SDR10 = *Striga* damage rating 10 days after sowing.

3.2.2 Under *Sh*-infested conditions

Table S2 shows the mean performances and the summary statistics of the evaluated genotypes for all traits under *Sh*-infested conditions. The CV% of the traits ranged from 7.15–597.49% (Table 4). The DA exhibited the lowest variation of 7.15%, while PLHT exhibited the highest variation of 597.49%, as was the case under *Sa* infested environment. The mean ASI was 1.52 for the inbred lines, 1.26, and 2.03 for the OPVs and the hybrids, respectively. The mean yield ranged from 10.05 g/plant to 151 g/plant, with a mean of 63.83 g/plant for the inbred lines, from 34.75 g/plant to 133.25 g/plant with a mean of 79.79 g/plant for the hybrids, and 33.60 g/plant to 144.25 g/plant with a mean of 70.81 g/plant for the OPVs. The sub-tropical inbred line CML304, which exhibited the lowest SDR10 under *Sa*, showed the highest GY under *Sh*. The lowest GY among the inbred lines was exhibited by the sub-tropical line HA04A-2107-36. The local check OPV ZM1423 exhibited the highest GY, whereas the *Striga* susceptible check TZBSTR showed the lowest GY. The top 10 inbred lines, as well as the top 10 check genotypes showing high GY under *Sh*, are presented in Table 4. The SEC8 mean was 63.89, 70.81, and 79.79, whereas the SEC10 mean was 3.25, 4.43, and 3.28 for the inbred lines, the OPVs, and the hybrids, respectively. The following high-yielding genotypes displayed the greatest reduction in SEC10 scores: TZDEEI50 (1.70) CML550 (2.63), TZISTR1001 (2.70), and B.King/1421 (2.63). The mean SDR8 was 5.25, 3.86, and 3.11 for the inbred lines, the OPVs, and the hybrids checks, respectively while the SDR10 means were 2.75, 3.05, and 3.11 in the same order. The high-yielding genotypes CML539 and ZM1421 resulted in the greatest reduction in SDR8. The genotype TZDEEI54 resulted in the greatest reduction of 0.75 in SDR10, while the least reductions were displayed by the low-yielding genotypes CKDHL0378 (5.75) and CZL1380 (5.50).

Table 4

Mean responses for 14 traits of 126 maize genotypes evaluated under *Sh* infestation, showing the top 10 inbred lines and the top 4 hybrids and 6 OPVs

Top 10 lines														
Genotypes	DA	DS	ASI	EPP	PLHT	EHT	HUSK	CL	EASPGY	SEC8	SEC10	SDR8	SDR10	
CML304	79.25	79.63	0.38	1.00	1.44	0.75	1.00	12.00	4.75	151.00	3.18	2.00	4.75	2.75
TZSTR1101	90.00	87.50	2.50	1.00	1.45	0.75	1.00	12.46	3.25	144.00	3.63	6.50	3.75	3.00
CLHP0404	74.75	74.25	0.50	1.00	2.07	0.75	1.00	10.00	6.25	137.35	3.18	5.00	6.00	2.00
TZISTR1119	78.75	77.00	1.75	1.00	1.81	0.95	1.00	10.50	3.75	135.75	3.68	4.50	5.50	3.50
TZISTR25	75.75	75.50	0.25	1.00	2.25	1.05	1.00	12.00	1.25	131.00	3.20	2.00	3.75	2.00
TZISTR1205	81.00	83.00	2.00	1.00	2.21	1.00	1.00	9.50	1.75	129.00	4.18	1.00	3.75	2.50
CML566	78.00	76.00	2.00	1.00	2.07	1.20	1.00	12.00	1.25	127.00	2.25	4.50	1.75	2.75
TZISTR1001	79.63	78.25	1.38	1.00	2.10	1.03	1.00	11.50	1.75	120.00	2.70	1.50	3.50	2.50
TZISTR1174	79.25	79.63	0.38	1.00	1.44	0.75	1.00	12.00	4.75	151.00	3.18	2.00	4.75	2.75
TZSTR1113	74.50	73.00	1.50	1.00	1.41	0.90	1.00	9.00	1.75	111.75	2.68	3.50	3.75	2.75
Top 4 hybrids and 6 OPVs														
N.Choice/1421	81.25	75.25	6.00	1.00	1.62	0.85	1.50	10.96	1.75	133.25	3.13	3.50	4.00	3.00
Shesha/1421	71.50	70.75	0.75	1.00	1.82	0.88	1.50	10.71	1.75	112.25	4.63	2.50	4.00	2.75
B.King/1421	81.00	77.75	3.25	1.00	2.35	1.05	1.50	11.71	3.25	91.75	2.63	4.50	2.25	2.25
ZM1421	82.38	80.75	1.63	1.00	2.10	0.95	1.50	10.71	2.25	88.00	2.63	2.00	1.75	3.25
ZM1423	70.25	71.88	1.63	1.00	2.17	0.94	1.00	13.71	1.25	144.25	4.63	2.50	1.75	2.50
STR-SYN-Y2	85.25	85.25	0.00	1.00	1.60	0.80	1.00	11.25	3.25	126.85	8.18	3.50	3.25	2.50
DTSTR-W SYN13	89.25	88.50	0.75	1.00	0.98	0.75	1.50	10.00	3.75	115.35	4.68	5.50	3.50	2.50
ZM1423/Z.DLO	81.25	83.25	2.00	1.00	12.41	1.03	1.00	10.75	4.75	96.75	2.68	5.00	3.75	2.75
DTSTR-Y SYN14	80.13	79.75	0.38	1.00	1.36	0.75	1.50	11.50	1.75	93.25	3.68	1.00	3.75	3.75
DTSTR-Y SYN15	83.75	84.25	0.50	1.00	1.78	0.65	1.00	9.00	6.25	87.35	3.18	4.50	4.00	2.75
Trial statistics														
LSD (5%)	5.71	7.10	2.58	0.09	1.58	1.15	0.62	2.18	1.39	34.87	34.87	2.37	1.62	1.17
SEM	8.19	15.88	5.08	0.18	2.58	2.26	1.39	4.87	4.20	77.95	77.95	5.31	3.62	2.61
%CV	7.15	8.95	597.49	12.24	87.15	28.59	49.30	21.39	43.80	52.81	52.81	64.59	47.58	41.79
Heritability	0.42	0.89	0.89	0.88	0.92	0.92	0.88	0.92	0.88	0.002	0.87	0.92	0.82	0.91

DA = days to 50% anthesis, DS = days to 50% silking, ASI = anthesis-silking interval, EPP = ear per plant, PLHT = Plant height, EHT = ear height, HUSK = husk cover, CL = cob length, EASP = ear aspect, GY = grain yield, SEC8 = *Striga* emergence counts eight days after sowing, SEC10 = *Striga* emergence counts ten days after sowing, SDR8 = *Striga* damage rating eight days after sowing, and SDR10 = *Striga* damage rating 10 days after sowing.

3.3 Principal component and biplot analyses

The rotated components matrix showing the percentages of variances of different principal components (PC) and the respective loadings of recorded traits are shown in Table 5. The first four PCs under *Sa*-infested conditions had a cumulative variance of 50.80%. The first PC had the highest variation of 17.38% followed by PC2 with 13.32%. DS, DA, EHT, CL, GY, and SDR8 made the highest contributions to PC1, while EPP, EASP, SDR8, and SDR10 contributed strongly to PC2. The highest loadings for PC3 were DS, GY, and DA, followed by SEC10, while the highest loadings for PC4 were SEC8, and CL, followed by HUSK, EPP, and SEC10. Under *Sh*-infested conditions, DS, DA, EHT, SDR8, and SEC8 had high positive loadings into the first PC, explaining 17.06% of the total variance. PC2 was highly influenced by GY, SEC8, CL, and EHT, which had a high loading, explaining 31.82% of the total variance. SDR8, SDR10, DA, and DS had the highest loadings into PC3, while PC4 was most influenced by HUSK, PLHT, and ASI.

Table 5
Rotated component matrix for 10 yield components and 4 *Striga* parameters in 126 maize genotypes under *Sa* and *Sh*-infested conditions.

Traits	<i>Sa</i>				<i>Sh</i>			
	PC1	PC2	PC3	PC4	PC1	PC2	PC3	PC4
DA	17.17	9.90	16.59	0.90	18.58	2.32	18.85	0.93
DS	22.02	1.17	21.56	0.80	26.88	3.70	14.19	1.26
ASI	1.44	8.42	1.31	0.44	8.96	1.55	0.00	16.19
EPP	1.77	18.61	0.42	10.40	3.65	0.08	3.75	14.48
PLHT	2.44	4.09	0.11	15.37	2.42	3.41	0.48	21.22
EHT	12.49	2.32	2.57	0.05	11.45	12.91	0.22	0.00
HUSK	3.26	3.76	4.41	13.84	0.39	1.21	0.37	34.65
CL	11.47	0.00	0.91	17.26	0.00	12.91	1.83	3.15
EASP	5.33	15.95	12.88	0.05	0.35	25.29	4.19	0.65
GY	9.46	4.52	20.73	5.10	0.03	25.52	7.97	1.43
SEC8	1.10	0.54	1.10	21.64	10.09	0.02	2.58	0.12
SEC10	0.05	2.77	14.31	9.02	0.10	5.48	0.06	0.94
SDR8	8.64	12.92	2.99	3.64	11.85	3.51	18.57	0.09
SDR10	3.35	15.03	0.10	1.50	5.25	2.09	26.94	4.89
Eigenvalue	2.43	1.91	1.53	1.24	2.39	2.07	1.66	1.29
Variance percentage (%)	17.38	13.63	10.96	8.82	17.06	14.77	11.85	9.24
Cumulative variance percentage (%)	17.38	31.01	41.97	50.80	17.06	31.82	43.67	52.91

PC: principal component, DA = days to 50% anthesis, DS = days to 50% silking, ASI = anthesis-silking interval, EPP = ear per plant, PLHT = Plant height, EHT = ear height, HUSK = husk cover, CL = cob length, EASP = ear aspect, GY = grain yield, SEC8 = *Striga* emergence counts eight days after sowing, SEC10 = *Striga* emergence counts ten days after sowing, SDR8 = *Striga* damage rating eight days after sowing, and SDR10 = *Striga* damage rating 10 days after sowing. Biplot based on principal components are presented to decipher the performance of the maize germplasm with *Sa* and *Sh* infestations using Fig. 2 and Fig. 3, respectively. Smaller angles between vectors of recorded traits indicate a high correlation between the traits in discriminating genotypes. Genotypes plotted closer to and further along a vector line scored highly in that trait. Under *Sa*-infested conditions, the genotypes were evenly scattered across both PC1 and PC2 (Fig. 2). The angles between vectors of GY and those of PLHT and HUSK, were acute, indicating a strong positive correlation between the traits and with high-yielding genotypes TZSTR1115 (34), TZISTR1001 (36), CML451 (86), TZISTR1015 (98), TZBSTR (110), ZM1423 (104), TZISTR1205 (95), and NC.QPM/Z.DPLO (106). The same analysis can be made with all the *Striga* parameters SEC8, SEC10, SDR8, and SDR10, which showed small angles with each other and with ASI, pointing to a high positive correlation between the traits. Genotypes CML312 (49), CML571 (60), CML539 (55), CLHP0049 (83), CLHP00286 (85), CML304 (93), STR-SYN-Y2 (107), and SAMMMZ16 (115) showed more susceptibility to *Sa* because they were closely associated with

vectors of SEC8, SEC10, SDR8, and SDR10. However, the angles formed between the vectors of GY, SEC8, SEC10, SDR8, and SDR10 were close to 90°, which means that the association between the traits is weak. EASP, DA, DS, and EPP showed a negative correlation with poor yielding including TZISTR1154 (1), TZISTR1275 (5), TZISTR1165 (9), TZISTR1159 (20), and TZSTR1109 (30).

Under *Sh*-infested conditions (Fig. 3), GY had a strong correlation with EHT, PLHT, and EPP and with the high-yielding genotypes, including CML304 (93) and ZM1423 (104), which showed a high score in these traits. The *Striga* parameters SEC8, SEC10, SDR8, and SDR10 positively correlated with EASP, and all plotted far from the vector of GY, indicating a negative correlation between the traits and low *Striga* emergence counts genotypes, including TZISTR1263 and Colorado/1421. The same summary is made with low SDR8 and SDR10 reduction genotypes CLHP0404, CKDHL0378, CZL1380, M.Pearl/DT-STR, and Colorado/1421, which were negatively correlated with GY.

3.4 Correlation of maize yield components and *Striga* parameters

Phenotypic correlation coefficients showing the relationship between GY and agronomic traits, within agronomic traits, within *Striga* parameters, and between GY and *Striga* parameters under both *Sa* and *Sh* infested conditions are shown in Fig. 4A and Fig. 4B, respectively. Under *Sa* conditions (Fig. 4A), GY had a positive and significant correlation with CL and PLHT, with correlation coefficients of $r = 0.33$ and $r = 0.18$, respectively. Negative and significant correlations were found between GY and EASP ($r = -0.50$). The agronomic traits PLHT and EHT showed positive and significant correlations of $r = 0.28$ and $r = 0.29$ with CL, respectively. A positive and significant correlation was also recorded between DA and DS ($r = 0.83$). The correlation between EPP and EASP, DS and ASI, and PLHT and EHT were also positive and significant ($r = 0.18$, $r = 0.30$, and $r = 0.54$, respectively). However, negative and significant correlations were recorded between the following agronomic traits: DA and ASI ($r = -0.29$), EPP and ASI ($r = -0.22$), EHT and DS ($r = -0.18$), CL and DA ($r = -0.20$), CL and DS ($r = -0.21$), PLHT and DA ($r = -0.18$), PLHT and DS ($r = -0.20$). ASI showed a positive and significant correlation between SDR8 ($r = 0.18$) and SDR10 (0.32). PLHT and SDR10 had a positive and significant correlation of $r = 0.18$. EPP exhibited a negative and significant correlation with SEC8, SDR8, and SDR10 ($r = -0.18$, $r = -0.27$, and $r = -0.24$, respectively). SDR8 exhibited negative and significant correlations with EHT ($r = -0.32$) and CL ($r = -0.22$). The correlation between SDR8 and SDR10 was positive and significant ($r = 0.54$).

Under *Sh* conditions (Fig. 4B), GY was positively correlated with EHT ($r = 0.20$), CL ($r = 0.29$), and PLHT (0.20), whereas with EASP and CL, GY exhibited a negative correlation with $r = -0.57$ and $r = -0.21$, respectively. Positive and significant correlations were revealed between the agronomic parameters DS and DA ($r = 0.93$), ASI and DS ($r = 0.30$), and PLHT and EHT ($r = 0.61$). However, negative and significant correlations were found between EHT and DS ($r = -0.19$), PLHT and DA ($r = -0.26$), and PLHT and DS ($r = -0.26$). Positive and significant correlations were observed between maize agronomic traits and *Striga* parameters. These include the positive correlation recorded between DS and SEC8 ($r = 0.21$), EASP and SEC10 ($r = 0.18$), and ASI and SEC8 ($r = 0.20$). However, negative, and significant correlations were recorded for SDR8 with EPP ($r = -0.20$) and EHT ($r = -0.20$). A negative correlation was recorded for EHT with SEC8 ($r = -0.22$) and SDR8 ($r = -0.36$), and PLHT with SDR10 ($r = -0.22$), and SDR8 ($r = -0.48$). The *Striga* parameter SDR8 was positively correlated with SEC8 ($r = 0.26$) and SDR10 ($r = 0.52$).

3.5 Cluster analysis based on yield components and *Striga* parameters

Cluster heatmap analysis based on eight maize and *Striga* phenotypic traits is presented for both *Sa* (Fig. 5 and Table S3) and *Sh*-infested environments (Fig. 6 and Table S4). The heatmap shows clusters based on the mean performances of each trait from the lowest performance (blue colour) to the highest performance (red colour). In an *Sa*-infested environment, the heatmap revealed six clusters (Fig. 5) where genotypes in the first cluster (I) had the highest scores in EASP, ASI, and SDR8 reduction. For instance, the IITA lines TZISTR1154, TZISTR1225, TZISTR1018, TZISTR1178, TZISTR1163, and hybrid ZM1421, which exhibited high SDR8 reductions and high scores in EASP, are classified in that cluster. Genotypes in the second cluster (II) had low ASI and high SEC10 values, including some IITA lines TZISTR1166, TZISTR1244, TZSTR1101, TZSTR1104, some CIMMYT lines CLHP0326, CLHP00378, CLHP0049, CLHP0022, and IITA OPVs DTSTR-W SYN13, and TZL COMP1-W C6*2/(White DT STR Syn-DT C1, which all exhibited low SEC10 reductions. Genotypes from the third cluster (III) had moderate EASP and moderate SEC8 values and included some CIMMYT lines CML312, CML444, CML442, CML571, CML390, IITA lines TZSTR1108, TZEEI14, TZEEI49, TZISTR1011, and TZDEEI50 and OPVs STR-SYN-Y2, and Z. Diplo.BC4C3-W-DT C1. Cluster IV comprised genotypes exhibiting the lowest SEC10 reduction, moderate ASI, moderate PLHT, and moderate SDR10 reduction. The genotype CIMMYT CML304, which showed a high SEC10 number, was clustered in that cluster together with the IITA OPVs (2*TZECOMP3DT/WhiteDTSTRSYN) C2, NC.QPM/DT-STR, ZM1421/DT-STR, and DTSTR-Y SYN14. Cluster V consisted of genotypes with high SDR10 reduction, moderate PLHT, and moderate SDR10, and comprised some good-yielding genotypes, including CKDHL0378, CML451, CLHP0350, CLHP0005, ZM1423/Z.DLO, TZISTR1174, (TZEOMP5C7/TZECOMP3DTC2) C2, and Colorado/1421. Cluster VI comprised genotypes that exhibited high GY, moderate PLHT, and high SDR10 reduction. The cluster had the high-yielding CIMMYT lines CML440, CML566, CML540, CML545, CLHP0156, and the IITA OPVs TZBSTR, Z.diplo-BC4-C3-W/DOGONA-1/Z.diplo-BC4-C3-W, and showed high GY under *Sa* infestation.

Under *Sh*-infested conditions (Fig. 6 and Supplemental Table 4), the genotypes clustered into four, with Cluster I consisting of genotypes exhibiting low reductions in SDR8, SDR10, SEC8, and SEC10. This cluster included TZBSTR, CZL1380, and Colorado/1421, which were characterized by low GY, low SDR10, and lowest SEC10 reduction under *Sh*. Cluster II was composed of two types of genotypes. The first group comprised genotypes that showed low GY, low SEC8, and SEC10 reduction, and included the genotype HA04A-2107-36, which showed the lowest GY in *Sh*-infested environment. The second group comprised genotypes that showed average GY and moderate resistance to *Sh*. Cluster III genotypes had high values in GY, EASP, and had moderate reduction in all *Sh* parameters. These included the CIMMYT lines CML540, CML566, CML304, CML550, CML539, CML440, CML545, and the IITA genotypes TZISTR25, TZISTR1174, TZISTR1119, TZISTR1166, TZSTR1113, B.King/1421, Shesha/1421, ZM1423, N.Choice/1421, DTSTR-W SYN13 all exhibited high mean yields under *Sh* infestation. Cluster IV comprised the NPGRC/SA OPV ZM1423/Z.DLO and the CIMMYT line CML571 showed exceptional scores in PLHT and GY.

4. Discussion

Breeding highly *Striga*-resistant maize genotypes adaptable to wide agro-ecological areas in SSA would be a sustainable solution to manage *Striga* infestations in maize crops. The mean values, genotypic and phenotypic variances, heritability, and the correlation coefficients of agronomic traits are key parameters used in breeding programs to select superior genotypes (Nzuve et al., 2014). The present study discerned genotypic differences for maize and *Striga* resistance traits, indicating that the target traits are amenable to selection. The genotype-by-season interaction effect was significant for any of the evaluated traits in both *Striga* species environments, suggesting the strong influence of environment on *Striga* traits.

The rotated component matrix showed that DA, DS, EHT, CL, GY, and SDR8 were the most discriminating traits under both *Sa* and *Sh* environments. This indicates that these traits are important selection indices. In addition to these traits, EPP, EASP, SDR8, and SDR10, which were loaded in the second PC in an *Sa*-infested environment, can also be considered during selection. Badu-Apraku et al. (2010) reported that SDR8, SDR10, EASP, and EPP were the best traits for selection under *Striga*-infested conditions. SDR was also reported to be the best selection criterion for improving GY performance in maize under *Striga* infestation by Dossa et al. (Dossa et al., 2023). All the traits exhibited high heritability values in both the *Sa* and *Sh* environments, except for GY under *Sh*-infested conditions. This indicates a large contribution of the genotypes to the phenotypic variance (Ngugi, 2013). Traits having high heritability are improved by selection based on phenotypic traits (Shekhawat and Singhania, 2005; Boghara et al., 2016). However, it is important to note the low heritability of the *Striga* parameters SEC10, SDR8, and SDR10, with values of only 0.008, 0.11, and 0.16, respectively, under *Sa* infested conditions, and of GY, with a value of 0.02 under *Sh* infested conditions. Notably, maize selection parameters recorded high heritabilities compared to the heritability of *Sa* parameters. This suggests that the genotypic variance of the *Striga* resistance traits was low compared to the phenotypic variance, making direct selection difficult for these traits. This is also indicative of the complex nature of the *Striga* resistance traits. Badu-Apraku et al. (2007) reported in a previous study low heritability values for SDR and SEC.

The PC biplots confirmed the positive and significant correlation between GY and secondary traits, including PLHT, EHT, and CL, under both *Sa*- and *Sh*-infested conditions, indicating the importance of these traits in improving maize GY under *Striga* infestation. Previous studies reported similar findings (Golam et al., 2011; Nzuve et al., 2014; Yahaya et al., 2021).

The ASI showed a positive correlation between SDR8 ($r = 0.18$) and SDR10 (0.32) under *Sa*-infested conditions, indicating that severe *Striga* damage prolongs the ASI. This confirms the stunted growth symptom of *Striga*-infested plants (Waweru et al., 2019). This is also supported by the positive correlation between EASP and SEC10, indicating that higher numbers of *Striga* plants increased the EASP score. The negative and significant associations between yield components, including EPP, CL, PLHT and EHT, and *Striga* parameters, including SEC8, SEC10, SDR8, and SDR10, under both *Sa* and *Sh* infestation, were expected because when the parasitic weeds proliferate, damage to maize yield components increase (Shayanowako et al., 2020). Therefore, reducing *Striga* damage scores and the parasitic plant counts at 8 and 10 days after planting will have a positive impact on yield component and enhance GY under *Striga* infestation (Menkir and Meseka, 2019; Badu-Apraku et al., 2021).

With *Sa* infestation, the yield of the evaluated inbred lines was higher than the yield of the OPVs checks. This was not expected, but it can be explained by the fact that most of these OPVs were bred for resistance or tolerance to *Sh*, with no screening for resistance or tolerance to *Sa*. These results indicate that these OPVs that have been bred for *Sh* resistance are not resistant to *Sa*. In the present study, some genotypes, e.g. TZISTR1174, showed excellent GY performance and relatively high SEC under *Sa*-infested conditions compared to others, and could be used as a source of resistance. Resistant genotypes support fewer *Striga* plants and produce a higher grain yield than susceptible ones. Conversely, in the case of tolerance, the host supports an equally severe level of the parasitic weed without losses in GY (Rodenburg et al., 2005). The following lines showed resistance to both *Sa* and *Sh*: sub-tropical lines CML440, CML566, CML540, CML539, CML451, CLHP0343, CLHP0326, and the tropical lines TZISTR1248, TZSTR115, TZISTR25, TZISTR1205, TZSTR113, TZISTR1119. The local hybrids B.King/1421, Shesha/1421, ZM1421, and the *Striga*-resistant check OPV DTSTR-W SYN13.

The phenotypic clustering analysis based on eight morphological traits clustered 126 maize genotypes into six clusters under *Sa* infestation and four clusters under *Sh* infestation. This indicates that the studied genotypes exhibited genetic divergence regarding morphological traits. The formation of different clusters using phenotypic traits in maize genotypes agreed with other reports (Saiyad and Kumar, 2017; Han et al., 2018; Kasoma et al., 2020; Shayanowako et al., 2020). All clusters consisted of a mixture of genotypes from different sources (IITA, CIMMYT, and NPGRC). Cluster mean values of evaluated genotypes in *Sa* environment suggested that Cluster VI was the best with outstanding trait values, especially for GY and SDR. In a *Sh*-infested environment, cluster mean values showed that Cluster IV was best, with outstanding values for all the evaluated traits. Therefore, these clusters containing tropical and sub-tropical maize genotypes should be good sources for heterosis breeding in *Striga* resistance breeding programs.

5. Conclusions

This study identified IITA, CIMMYT, and NPGRC/SA conserved maize genetic resources that could be used as breeding parents to improve for both *Sa*, *Sh*, and dual resistance. The top five most resistant to *Sa* included tropical and sub-tropical maize inbred lines: CML540 (277.50 g/plant), CML566 (155.50 g/plant), TZISTR1001 (140.00 g/plant), TZISTR1205 (114.25 g/plant) and TZSTR115 (112.50 g/plant). On the other hand, the top five yielding inbred lines under *Sh* were CML304 (151.00 g/plant), TZSTR101 (144.00 g/plant), CLHP0404 (137.35 g/plant), TZISTR1119 (135.75 g/plant), and TZISTR25 (131.00 g/plant). The genotypes CML440, CML566, CML540, CML539, CLHP0343, CLHP0326, TZISTR1248, TZSTR115, TZISTR25, TZISTR1205, TZSTR113, and TZISTR1119, showed resistance to both *Striga* species, whereas TZISTR1174 showed tolerance to *Sa*. The identified tolerant and resistant inbred lines could be used to produce new hybrids combining tolerance and resistance to both *Sa* and *Sh* and market-preferred agronomic traits.

Declarations

Supplementary Materials: The following supporting information can be downloaded: Table S1: Mean values for 10 yield components and 4 *Striga* parameters measured for 126 maize genotypes evaluated under *Sa* infestation. Table S2. Mean values for 10 yield components and 4 *Striga* parameters measured for 126 maize genotypes evaluated under *Sh* infestation. Table S3. Clusters and their member genotypes of 126 maize genotypes evaluated under *Sa* infestation. Table S4: Clusters and their member genotypes of 126 maize genotypes evaluated under *Sh* infestation.

Author Contributions: Each author participated sufficiently in the completion of this work. ED: Conceptualization; Investigation; Methodology; Resources; Writing original draft; Writing – Review & Editing. HS: Conceptualization; Funding acquisition; Methodology; Project administration; Resources; Supervision;

Validation; Visualization; Writing – Review & Editing. AS: Methodology; Visualization; Validation; Writing – Review & Editing. ML: Funding acquisition; Resources; Supervision; Validation; Visualization; Writing – Review & Editing; All authors have read and agreed to the published version of the manuscript.

Acknowledgements

We are grateful to the University of KwaZulu-Natal (UKZN) and the Organisation for Women in Science for the Developing World (OWSD) for the financial support of this study.

Conflicts of Interest:

The authors declare that they have no competing financial interests or personal relationships that could have appeared to influence this work.

References

1. Alboukadel, K. (2017). Practical guide to principal component methods in R: PCA, M (CA), FAMD, MFA, HCPC, and factoextra. *Statistical Tools For High-Throughput Data Analysis* 2, 1–154.
2. Arndt, C., Diao, X., Dorosh, P., Pauw, K., and Thurlow, J. (2023). The Ukraine war and rising commodity prices: Implications for developing countries. *Global Food Security* 36, 1–9. doi:10.1016/j.gfs.2023.100680
3. Badu-Apraku, B., Adewale, S., Paterne, A., Gedil, M., and Asiedu, R. (2020a). Identification of QTLs controlling resistance/tolerance to *Striga hermonthica* in an extra-early maturing yellow maize population. *Agronomy-Basel* 10, 1–18. doi:10.3390/agronomy10081168
4. Badu-Apraku, B., Adu, G. B., Yacoubou, A. M., Toyinbo, J., and Adewale, S. (2020b). Gains in genetic enhancement of early maturing maize hybrids developed during three breeding periods under *Striga*-infested and *Striga*-free environments. *Agronomy-Basel* 10, 1–19. doi:10.3390/agronomy10081188
5. Badu-Apraku, B., Akinwale, R. O., and Fakorede, M. A. B. (2010). Selection of early maturing maize inbred lines for hybrid production using multiple traits under *Striga*-infested and *Striga*-free environments. *Maydica* 55(3), 261–274.
6. Badu-Apraku, B., Fakorede, M. A. B., Akinwale, R. O., Adewale, S. A., and Akaogu, I. C. (2021). Developing high-yielding *Striga*-resistant maize in sub-Saharan Africa. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 16(30), 1–12. doi:10.1079/PAVSNR202116030
7. Badu-Apraku, B., Menkir, A., and Lum, A. F. (2007). Genetic variability for grain yield and its components in an early tropical yellow maize population under *Striga hermonthica* infestation. *Journal of Crop Improvement* 20, 107–122. doi:10.1300/J411v20n01_06
8. Boghara, M. C., Dhaduk, H. L., Kumar, S., Parekh, M. J., Patel, N. J., and Sharma, R. (2016). Genetic divergence, path analysis, and molecular diversity analysis in cluster bean (*Cyamopsis tetragonoloba* L. Taub.). *Industrial Crops and Products* 89, 468–477. doi:10.1016/j.indcrop.2016.05.049
9. Dabija, A., Ciocan, M. E., Chetariu, A., and Codină, G. G. (2021). Maize and sorghum as raw materials for brewing, a review. *Applied Sciences* 11(7), 1–24. doi:10.3390/app11073139
10. Dafaallah, A. B. (2019). Biology and physiology of witchweed (*Striga* spp.): A review. *International Journal of Academic Multidisciplinary Research* 3(10), 42–51.
11. David, C., Fontem, L. A., and Menkir, A. (2011). Seed coating herbicide tolerant maize hybrids with imazapyr for *Striga hermonthica* (Del.) Benth control in the West African savanna. *Journal of Food Agriculture and Environment* 9, 416–421.
12. David, O. G., Ayangbenro, A. S., Odhiambo, J. J. O., and Babalola, O. O. (2022). *Striga hermonthica*: A highly destructive pathogen in maize production. *Environmental Challenges* 8, 1–9. doi:10.1016/j.envc.2022.100590
13. Dossa, E. N., Shimelis, H., Shayanowako, A. I. T., and Laing, M. D. (2023). A meta-analysis of the effects of *Striga* control methods on maize, sorghum, and major millets production in sub-Saharan Africa. *Crop Science*, 1–20. doi:10.1002/csc2.20889
14. Gasura, E., Nyandoro, B., Mabasa, S., Setimela, P. S., Kyalo, M., and Yao, N. (2021). Breeding strategy for resistance to *Striga asiatica* (L.) Kuntze based on genetic diversity and population structure of tropical maize (*Zea mays* L.) lines. *Genetic Resources and Crop Evolution*, 69, 987–996. doi:10.1007/s10722-021-01274-6
15. Gasura, E., Setimela, P., Mabasa, S., Rwafa, R., Kageler, S., and Nyakurwa, C. (2019). Response of IITA maize inbred lines bred for *Striga hermonthica* resistance to *Striga asiatica* and associated resistance mechanisms in southern Africa. *Euphytica* 215(151), 1–15. doi:10.1007/s10681-019-2467-5
16. Gethi, J. G., and Smith, M. E. (2004). Genetic responses of single crosses of maize to *Striga hermonthica* (Del.) Benth. and *Striga asiatica* (L.) Kuntze. *Crop Science* 44, 2068–2077. doi:10.2135/cropsci2004.2068
17. Golam, F., Farhana, N., Zain, M. F., Majid, N. A., Rahman, M., Rahman, M. M., and Kadir, M. A. (2011). Grain yield and associated traits of maize (*Zea mays* L.) genotypes in Malaysian tropical environment. *African Journal of Agricultural Research* 6, 6147–6154.
18. Han, L., Yang, G., Yang, H., Xu, B., Li, Z., and Yang, X. (2018). Clustering field-based maize phenotyping of plant-height growth and canopy spectral dynamics using a UAV remote-sensing approach. *Frontiers in Plant Science* 9, 1–18. doi:10.3389/fpls.2018.01638
19. Jahufer, M. Z. Z., and Luo, D. (2018). DeltaGen: a comprehensive decision support tool for plant breeders. *Crop Science* 58, 1118–1131. doi:10.2135/cropsci2017.07.0456
20. Johnmark, O., Indieka, S., Liu, G., Gowda, M., Suresh, L. M., Zhang, W., and Gao, X. (2022). Fighting death for living: recent advances in molecular and genetic mechanisms underlying maize lethal necrosis disease resistance. *Viruses* 14(2), 1–21. doi:10.3390/v14122765
21. Kamara, A. Y., Menkir, A., Chikoye, D., Solomon, R., Tofa, A. I., and Omoigui, L. O. (2020). Seed dressing maize with imazapyr to control *Striga hermonthica* in farmers' fields in the savannas of Nigeria. *Agriculture-Basel* 10(83), 1–9. doi:10.3390/agriculture10030083

22. Kanampiu, F. K., Kabambe, V., Massawe, C., Jasi, L., Friesen, D., Ransom, J. K., and Gressel, J. (2003). Multi-site, multi-season field tests demonstrate that herbicide seed-coating herbicide-resistance maize controls *Striga* spp. and increases yields in several African countries. *Crop Protection* 22, 697–706. doi:10.1016/s0261-2194(03)00007-3
23. Kansime, M. K., Rwomushana, I., and Mugambi, I. (2023). Fall armyworm invasion in Sub-Saharan Africa and impacts on community sustainability in the wake of Coronavirus Disease 2019: reviewing the evidence. *Current Opinion in Environmental Sustainability* 62, 1–6. doi:10.1016/j.cosust.2023.101279
24. Kasoma, C., Shimelis, H., Laing, M. D., Shayanowako, A. I. T., and Mathew, I. (2020). Revealing the genetic diversity of maize (*Zea mays* L.) populations by phenotypic traits and DArTseq markers for variable resistance to fall armyworm. *Genetic Resources and Crop Evolution* 68, 243–259. doi:10.1007/s10722-020-00982-9
25. Kim, S. K. (1994). Genetics of maize tolerance of *Striga hermonthica*. *Crop Science* 34, 900–907.
26. Kountche, B. A., Jamil, M., Yonli, D., Nikiema, M. P., Blanco-Ania, D., Asami, T., Zwanenburg, B., and Al-Babili, S. (2019). Suicidal germination as a control strategy for *Striga hermonthica* (Benth.) in smallholder farms of sub-Saharan Africa. *Plants People Planet* 1, 107–118. doi:10.1002/ppp3.32
27. Lane, J., Child, D., Moore, T., Arnold, G., and Bailey, J. (1997). Phenotypic characterisation of resistance in *Zea diploperennis* to *Striga hermonthica*. *Maydica* 42, 45–51.
28. Lobulu, J., Shimelis, H., Laing, M., Mushongi, A., and Shayanowako, A. I. T. (2021). "Characterization of maize genotypes (*Zea mays* L.) for resistance to *Striga asiatica* and *S. hermonthica* and compatibility with *Fusarium oxysporum* f. sp. strigae (FOS) in Tanzania." *Agronomy* 11(5), 1–27.
29. Menkir, A. (2006). Assessment of reactions of diverse maize inbred lines to *Striga hermonthica* (Del.) Benth. *Plant Breeding* 125, 131–139. doi:10.1111/j.1439-0523.2006.01175.x
30. Menkir, A., Badu-Apraku, B., Yallou, C. G., Kamara, A. Y., and Ejeta, G. (2007). Breeding maize for broad-based resistance to *Striga Hermonthica*. In: *Integrating New Technologies for Striga Control: Towards Ending the Witch-Hunt*, World Scientific pp. 99–114.
31. Menkir, A., Kling, J. G., Badu-Apraku, B., and Ibikunle, O. (2006). Registration of 26 tropical maize germplasm lines with resistance to *Striga hermonthica*. *Crop Science* 46(2), 1007–1009.
32. Menkir, A., and Meseka, S. (2019). Genetic improvement in resistance to *Striga* in tropical maize hybrids. *Crop Science* 59, 2484–2497.
33. Mutsvanga, S., Gasura, E., Setimela, P. S., Nyakurwa, C. S., and Mabasa, S. (2022). Nutritional management and maize variety combination effectively control *Striga asiatica* in southern Africa. *CABI Agriculture and Bioscience* 3, 1–14.
34. Ngugi, K. (2013). Anthesis to silking interval usefulness in developing drought tolerant maize. *Journal of Renewable Agriculture* 1(5), 84–90. doi:10.12966/jra.08.03.2013
35. Nzuve, F., Githiri, S., Mukunya, D. M., and Gethi, J. (2014). Genetic variability and correlation studies of grain yield and related agronomic traits in maize. *Journal of Agricultural Science* 6(9), 166–176. doi:10.5539/jas.v6n9p166
36. R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. 2018. <https://www.r-project.org/>.
37. R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. 2023. <https://www.r-project.org/>.
38. Regassa, B., Wegary, D., Fininsa, C., and Abraham, A. (2021). Screening maize genotypes for resistance to maize lethal necrosis disease in Ethiopia. *Tropical Plant Pathology* 46, 583–595. doi:10.1007/s40858-021-00458-w
39. Rodenburg, J., Bastiaans, L., Weltzien, E., and Hess, D. E. (2005). How can field selection for *Striga* resistance and tolerance in sorghum be improved? *Field Crops Research* 93, 34–50. doi:10.1016/j.fcr.2004.09.004
40. Saiyad, M. M., and Kumar, S. (2017). Evaluation of maize genotypes for fodder quality traits and SSR diversity. *Journal of Plant Biochemistry and Biotechnology* 27, 78–89. doi:10.1007/s13562-017-0418-6
41. Samejima, H., Babiker, A. G., Takikawa, H., Sasaki, M., and Sugimoto, Y. (2016). Practicality of the suicidal germination approach for controlling *Striga hermonthica*. *Pest Management Science* 72, 2035–2042. doi:10.1002/ps.4215
42. Samejima, H., and Sugimoto, Y. (2022). Phenotypic diversity in pre- and post-attachment resistance to *Striga hermonthica* in a core collection of rice germplasms. *Plants (Basel)* 12(1), 1–19. doi:10.3390/plants12010019
43. Shayanowako, A. I., Shimelis, H., Laing, M. D., and Mwadzingeni, L. (2018a). Variance components and heritability of traits related to *Striga asiatica* resistance and compatibility to *Fusarium oxysporum* f. sp. strigae in maize. *Maydica* 63(1), 1–8.
44. Shayanowako, A. I. T., Shimelis, H., Laing, M. D., and Mwadzingeni, L. (2020). *Striga* resistance and compatibility of maize genotypes to a biocontrol agent, *Fusarium oxysporum* f.sp. strigae. *Journal of Crop Improvement* 34, 437–454. doi:10.1080/15427528.2020.1728599
45. Shayanowako, A. I., Shimelis, H., Laing, M. D., and Mwadzingeni, L. (2018b). Resistance breeding and biocontrol of *Striga asiatica* (L.) Kuntze in maize: a review. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science* 68, 110–120. doi:10.1080/09064710.2017.1370493
46. Shekhawat, S., and Singhania, D. (2005). Correlation and path analysis in cluster bean. *Forage Research* 30, 196–199.
47. Simon, Z., Kasozi, L. C., Patrick, R., and M. Abubaker (2018). Gene action for grain yield and agronomic traits in selected maize inbred lines with resistance to *Striga hermonthica* in Uganda. *Journal of Food Security* 6(4), 155–162. doi:10.12691/jfs-6-4-3
48. Souto, K. M., Jacques, R. J. S., Zanella, R., Machado, S. L. O., Balbinot, A., and Avila, L. A. (2020). Phytostimulation of lowland soil contaminated with imidazolinone herbicides. *International Journal of Phytoremediation* 22, 774–780. doi:10.1080/15226514.2019.1710814
49. Waweru, D. N., Kuria, E. K., Bradley, J. M., Scholes, J. D., and Runo, S. (2019). Tissue culture protocols for the obligate parasitic plant *Striga hermonthica* and implications for host-parasite co-cultivation. *Plant Cell, Tissue and Organ Culture (PCTOC)* 138, 247–256.

50. Yacoubou, A. M., Aboudou, A., Yaoitcha, A. S., Menkir, A., Badu-Apraku, B., Olanmi, B., and Zoumarou, N. W. (2021a). Screening of early hybrids for resistance to *Striga hermonthica* (del.) benth in maize African Journal of Plant Breeding ISSN 8 (11), 001–018.
51. Yacoubou, A. M., Wallis, N. Z., Menkir, A., Zinsou, V. A., Onzo, A., Garcia-Oliveira, A. L., Meseka, S., and Paterne, A. (2021b). Breeding maize (*Zea mays*) for *Striga* resistance: Past, current and prospects in sub-Saharan Africa. Plant Breeding 140, 195–210. doi:10.1111/pbr.12896
52. Yahaya, M., Bello, I., and Unguwanrimi, A. (2021). Correlation and path-coefficient analysis for grain yield and agronomic traits of maize (*Zea mays* L.). Science World Journal 16, 10–13.
53. Zziwa, S. (2018). Genetic resistance of selected maize inbred lines to *Striga hermonthica*. PhD thesis, Makerere University, p88.

Figures



Figure 1: Photographs showing the experimental setup with artificial infestations of maize with *Striga asiatica* (A, B, and C), and *S. hermonthica* (D and E) in the African Center for Crop Improvement greenhouse at the University of Kwazulu-Natal, South Africa

Figure 1

See image above for figure legend.




Figure 2: Principal component of 126 maize population under *Striga*-infested condition. Genotypes are coded with numbers as recorded in Table S1. Dim= dimension, DA= days to 50% anthesis, DS= days to 50% silking, ASI= anthesis-silking interval, EPP= ear per plant, PLHT= Plant height, EHT= ear height, HUSK= husk cover, CL= cob length, EASP= ear aspect, GY= grain yield, SEC8= *Striga* emergence counts eight days after sowing, SEC10= *Striga* emergence counts ten days after sowing, SDR8= *Striga* damage rating eight days after sowing, and SDR10= *Striga* damage rating 10 days after sowing

Figure 2

See image above for figure legend.




Figure 3: Principal component of 126 maize population under *Striga*-infested condition. Genotypes are coded with numbers as recorded in Table S2. Dim= dimension, DA= days to 50% anthesis, DS= days to 50% silking, ASI= anthesis-silking interval, EPP = ear per plant, PLHT= Plant height, EHT= ear height, HUSK= husk cover, CL= cob length, EASP= ear aspect, GY= grain yield, SEC8= *Striga* emergence counts eight days after sowing, SEC10= *Striga* emergence counts ten days after sowing, SDR8= *Striga* damage rating eight days after sowing, and SDR10= *Striga* damage rating 10 days after sowing.

Figure 3

See image above for figure legend.




Figure 4: Correlation matrix plot between yield components and *Striga* parameters in a population of 126 maize genotypes under *Sa* (A) and *Sh* (B) infestation. The colour variation indicates the magnitude of correlations; traits with deep blue are strongly and positively correlated, while deep red are strongly negatively correlated. Note *, ** and *** denote the level of significance associations of traits at 0.05, 0.01 and <0.001 probability values, respectively and ns=non-significant. EHT= ear height, CL= cob length, GY= grain yield, DA= days to 50% anthesis, DS= days to 50% silking, SEC8= *Striga* emergence counts eight days after sowing, SEC10= *Striga* emergence counts ten days after sowing, EPP = ear per plant, EASP= ear aspect, HUSK= husk cover, SDR8= *Striga* damage rating eight days after sowing, ASI= anthesis-silking interval, SDR10= *Striga* damage rating 10 days after sowing, PLHT= Plant height.

Figure 4

See image above for figure legend.




Figure 5: Six clusters (I to VI) heatmap plot showing a colour pattern of 126 maize genotypes based on 4 maize yield components and 4 *Striga* parameters recorded in *Sa*-infested environment. Blue: lowest performance; red: highest performance. The numbers at the right represent the genotype numbers as coded in Table S1. EASP= ear aspect, SEC8= *Striga* emergence counts eight days after sowing, SEC10= *Striga* emergence counts ten days after sowing, PLHT= Plant height, GY= grain yield, ASI= anthesis-silking interval, SDR10= *Striga* damage rating 10 days after sowing, SDR8= *Striga* damage rating eight days after sowing.

Figure 5

See image above for figure legend.




Figure 6: Four clusters (I to IV) heatmap plot showing a colour pattern of 126 maize genotypes based on 4 maize yield components and 4 *Striga* parameters recorded in *Sh*-infested environment. Blue: lowest performance; red: highest performance. The numbers at the right represent the genotype numbers as coded in Table S2. EASP= ear aspect, SEC10= *Striga* emergence counts ten days after sowing, SDR10= *Striga* damage rating 10 days after sowing, SDR8= *Striga* damage rating eight days after sowing, SEC8= *Striga* emergence counts eight days after sowing, ASI= anthesis-silking interval, PLHT= plant height, GY= grain yield.

Figure 6

See image above for figure legend.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalfileDossaetal2023.docx](#)