Neilreichia **12**: S1–S6 (2021)

Biogeographical classification of Austria based on hierarchical cluster analysis of vascular plant distributions

Angelika Billensteiner & Harald Niklfeld

(Neilreichia **12**: 199–218)

Electronic Supplement

DOI: https://doi.org/10.5281/zenodo.5821417




Fig. S1: Cluster dendrogram based on Ward method and Euclidean distance. Blue rectangles indicate clusters at cutlevel 25.




Fig. S2: Cluster dendrogram based on Complete Linkage method and Euclidean distance. Blue rectangles indicate clusters at cutlevel 25.




Fig. S3: Cluster dendrogram based on Complete Linkage method and Jaccard index. Blue rectangles indicate clusters at cutlevel 25.




Fig. S4: Cluster dendrogram based on Complete Linkage method and Dice index. Blue rectangles indicate clusters at cutlevel 25.




Fig. S5: Geographic delimitation of clusters identified at cutlevel 25 based on Ward method and Euclidean distance.




Fig. S6: Geographic delimitation of clusters identified at cutlevel 25 based on Complete Linkage method and Euclidean distance.




Fig. S7: Geographic delimitation of clusters identified at cutlevel 25 based on Complete Linkage method and Jaccard index.




Fig. S8: Geographic delimitation of clusters identified at cutlevel 25 based on Complete Linkage method and Dice index.




Fig. S9: NbClust diagram, Krzanowski and Lai Index (Ward method, Euclidean distance).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Neilreichia - Zeitschrift für Pflanzensystematik und Floristik Österreichs

Jahr/Year: 2021

Band/Volume: 12

Autor(en)/Author(s): Billensteiner Angelika, Niklfeld Harald

Artikel/Article: Biogeographical classification of Austria based on hierarchical cluster analysis of vascular plant distributions S1-S6