

## Species delimitation and population structure analysis in *Polygonum* species (section *Polygonum*)

Samaneh Mosaferi, Masoud Sheidai, Maryam Keshavarzi & Zahra Noormohammadi

**Summary:** The genus *Polygonum* L. (Polygonaceae) contains about 230 species that are distributed worldwide. These herbaceous annuals and shrubby perennials are usually invasive in nature and occupy different geographical regions. They show morphological and genetic plasticity and they are difficult to deal with taxonomically. In spite of many *Polygonum* species that grow in Iran, we do not have any report on their population genetic structure, mode of divergence and dispersal. Therefore, we planned molecular (ISSR markers) and morphological studies of 37 accessions from 17 species of *Polygonum* (section *Polygonum*) that were collected from different regions in Iran. We try to address the following questions: 1) can ISSR markers delimit *Polygonum* species, 2) what is the genetic structure of these taxa in Iran, and 3) what evolutionary forces are acting for *Polygonum* species divergence? MDS plot of morphological characters separated annual species from perennials. ISSR molecular markers revealed a low degree of genetic variability in some of the studied species but showed high genetic diversity in some others. AMOVA showed strong genetic differentiation among the studied species. STRUCTURE analysis and assignment test revealed some degree of genetic admixture among studied species/populations. The present study show that invasion followed by genetic divergence due to genetic drift and inbreeding, local adaptation and occasional interspecific cross-pollination are the major evolutionary forces in *Polygonum* species and populations divergence.

**Keywords:** *Polygonum*, ISSR, genetic divergence, morphological variability

*Polygonum* L. with about 230 species in the world is the largest genus of Polygonaceae (GALASSO et al. 2009; VOYLOKOVA et al. 2009). The genus as a whole has a wide distribution ranging from the tropics to the polar region and from sea level to the highest altitudes.

There has always been disagreement on classification of *Polygonum* s.l. and different subdivisions on section and genus level have been reported (MEISSNER 1826; BENTHAM & HOOKER 1880; GROSS 1913; HEDBERG 1946; HARALDSON 1978; RONSE DECRAENE & AKEROYD 1988). According to LAMB-FRYE & KRON (2003), molecular analysis on *Polygonum* species of North America showed that *Polygonum* is paraphyletic. Phylogenetic studies on European *Polygonum* confirm that Polygonaceae are monophyletic and the subfamily Polygonoideae seems to be paraphyletic (GALASSO et al. 2009). In this study, *Polygonum* s.str. (tribe Polygoneae) has been adopted. According to previous studies, 29 taxa of *Polygonum* s.str. belonging to 4 sections (*Tiniaria* Meisn., *Aconogonon* Meisn., *Pleuropterus* (Turcz.) Benth. & Hooker., *Polygonum* Tourn.) occur in Iran (RECHINGER & SCHIMAN-CZEKA 1968; MOZAFFARIAN 2012).

*Polygonum* s.str. species are annual and perennial herbs, subshrubs or shrubs with woody stocks (LI et al. 2003). Their stems are erect, prostrate or ascending, usually with conspicuously swollen nodes. All of them have membranous tubular ochreas. Their leaves are linear, lanceolate, elliptic, ovate or oblong with or without petioles. Inflorescences are axillary with solitary flowers or few flowered fascicles or lax spikes. Some of them are weeds, while some others have ornamental and medicinal importance across the world (RECHINGER & SCHIMAN-CZEKA 1968; KAISER 2001).

*Polygonum* species, particularly annual taxa, show a wide range of ecological adaptations and occupy different habitats due to their adaptability and plasticity (SULTAN 2003; GRIFFITH & SULTAN 2006). Although *Polygonum* plants are mainly autogamous, hybridization can occur between annual species (LÖVE & LÖVE 1956; YURTSEVA 2001). Based on YURTSEVA (2001) and VOYLOKOVA et al. (2009), possible hybrids between *P. aviculare* and *P. arenastrum* were found in Russia and supported by morphological and molecular studies. Morphological complexity caused by adaptation, plasticity and hybridization made identification of *Polygonum* species difficult. Extensive studies have been performed on *Polygonum*. These were mainly dealing with their taxonomy and molecular phylogenetics (MEERTS et al. 1990; RONSE DECRAENE et al. 2000; LAMB-FRYE & KRON 2003; GALASSO et al. 2009; KESHAVARZI et al. 2012), but there has been no attempt to study genetic diversity, ecological adaptation and intra- and interspecific differentiation along with morphometric studies on *Polygonum* s.str. of Iran.

In this study, we examine the morphology, genetic diversity and population structure of 17 species of section *Polygonum* in Iran. We try to answer the following questions: 1) Is there intra- and interspecific genetic diversity among studied species? 2) Is genetic distance among these species correlated with their geographical distance? 3) What is the genetic structure of populations and taxa? 4) Is there any gene exchange between *Polygonum* species in Iran?

## Materials and methods

**Plant materials.** Hundred and five plants were randomly collected from 37 populations of 17 *Polygonum* species (section *Polygonum*). These species include both annuals (numbers 1–9 in Table 1) and perennials (numbers 10–17 in Table 1). Details of localities of the studied species and populations are given in Table 1. The voucher specimens are deposited in the herbarium of Shahid Beheshti University [HSBU], Alzahra University Herbarium [ALUH], Ferdowsi University Mashhad Herbarium [FUMH], Kharazmi University Herbarium [T] and in the herbarium of Shahid Bahonar University of Kerman.

**ISSR assays and genetic analyses.** Fresh leaves were randomly collected and dried in silica gel powder. For some specimens, we used herbarium materials. Genomic DNA was extracted using CTAB activated charcoal protocol (KRIŽMAN et al. 2006). 5 ISSR primers, UBC 834, (CA)7GT, (GA)9C, UBC 807 and (GA)9T commercialized by UBC (University of British Columbia), were used. PCR reactions were performed in a 25 µl volume containing 10 mM Tris-HCl buffer at pH 8, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.2 mM of each dNTP (Bioron, Germany), 0.2 µM of a single primer, 20 ng genomic DNA and 3 U of *Taq* DNA polymerase (Bioron, Germany). The amplification reactions were performed in a Techne thermocycler (Germany) with the following program: 5 min at 94°C, 40 cycles of 1 min at 94°C, 1 min at 51–56°C and 1 min at 72°C and a final cycle of 7 min at 72°C. The amplification products were visualized by running on 1% agarose gel, followed by ethidium bromide staining. The fragment size was estimated by using a 100 bp molecular size ladder (Fermentas, Germany).

## Data analyses

### Morphological study

For morphometric studies, 15 qualitative and 17 quantitative characters were used (Table 2). The analysis of variance (ANOVA) test was performed to show significant morphological differences

**Table 1.** *Polygonum* species and populations, their localities and voucher numbers.

| Species                                          | Locality                                                          | Latitude      | Longitude     | Altitude (m) | Voucher no.                             |
|--------------------------------------------------|-------------------------------------------------------------------|---------------|---------------|--------------|-----------------------------------------|
| 1. <i>P. arenastrum</i> Boreau                   | East Azerbaijan, Sar village                                      | 38° 20' 03" N | 46° 11' 37" E | 1946         | ALUH 800                                |
|                                                  | Tehran, Darakeh                                                   | 35° 48' 34" N | 51° 22' 52" E | 1756         | ALUH 863                                |
|                                                  | Guilan, Bandar-e Anzali                                           | 37° 28' 21" N | 49° 27' 31" E | -27          | ALUH 801                                |
|                                                  | Kermanshah, Kermanshah                                            | 34° 18' 51" N | 47° 03' 54" E | 1392         | ALUH 802                                |
| 2. <i>P. patulum</i> M. Bieb.                    | Tehran, Darakeh                                                   | 35° 48' 34" N | 51° 22' 52" E | 1756         | HSBU 2014219                            |
|                                                  | Golestan, Gorgan                                                  | 36° 50' 44" N | 54° 26' 22" E | 125          | HSBU 2014212                            |
| 3. <i>P. olivascens</i> Rech.f. & Schiman-Czeika | Sistan & Baluchestan, Hirmand, Dust Mohammad village              | 31° 08' 41" N | 61° 47' 33" E | 470          | HSBU 2014213                            |
|                                                  | East Azerbaijan, Khosrowshahr                                     | 37° 56' 59" N | 46° 03' 10" E | 1356         | ALUH 804                                |
|                                                  | East Azerbaijan, Khvor Khvor village                              | 37° 51' 55" N | 45° 46' 58" E | 1278         | ALUH 805                                |
|                                                  | Tehran, Nasim Shahr                                               | 35° 33' 49" N | 51° 09' 31" E | 1073         | ALUH 906                                |
|                                                  | Markazi, Saveh, Yalabad village                                   | 34° 57' 14" N | 50° 18' 24" E | 1022         | ALUH 905                                |
| 4. <i>P. argyrocoleon</i> Steud. ex Kunze        | Khuzestan, Behbahan                                               | 30° 35' 45" N | 50° 14' 30" E | 325          | HSBU 2014214                            |
|                                                  | West Azerbaijan, Dash Band village                                | 36° 38' 45" N | 46° 10' 32" E | 1336         | ALUH 869                                |
|                                                  | Razavi Khorasan, Mashhad, Kuh sangi                               | 36° 17' 16" N | 59° 34' 12" E | 1008         | ALUH 872                                |
|                                                  | Kurdistan, 5 km to Divandarreh                                    | 35° 54' 50" N | 47° 01' 26" E | 1838         | ALUH 874                                |
|                                                  | Guilan, Rasht                                                     | 37° 16' 51" N | 49° 34' 59" E | -2           | ALUH 866                                |
| 5. <i>P. molliiforme</i> Boiss.                  | Khorasan Razavi, Saroogh mountain                                 | 36° 20' 33" N | 57° 18' 41" E | 1660         | FUMH- 42438                             |
|                                                  | Khorasan Razavi, Afchang mountains                                | 36° 24' 29" N | 57° 38' 18" E | 1750         | FUMH- 39101                             |
| 6. <i>P. polycnemoides</i> Jaub. & Spach         | Tehran, Abali                                                     | 35° 45' 45" N | 51° 57' 45" E | 2085         | ALUH 872                                |
|                                                  | Mazandaran, Savadkuh, Kachid village                              | 36° 12' 20" N | 53° 00' 38" E | 1704         | ALUH 803                                |
|                                                  | Alborz, Asara                                                     | 36° 02' 11" N | 51° 11' 39" E | 1877         | HSBU- 2014215                           |
|                                                  | Kerman, Kerman                                                    | 30° 11' 18" N | 57° 08' 11" E | 1784         | HSBU- 2014216                           |
|                                                  | Khorasan Razavi, Mashhad                                          | 36° 17' 16" N | 59° 34' 12" E | 1006         | HSBU- 2014217                           |
| 7. <i>P. kitaibelianum</i> Sadler.               | Khuzestan, Shushtar                                               | 32° 02' 44" N | 48° 51' 24" E | 61           | T- 42861                                |
| 8. <i>P. aviculare</i> L.                        | Mazandaran, Gadook                                                | 35° 50' 04" N | 52° 55' 47" E | 220          | HSBU- 2014200                           |
|                                                  | Kurdistan, Kamyaran                                               | 34° 47' 44" N | 46° 56' 08" E | 1466         | HSBU- 2014201                           |
|                                                  | Lorestan, Khorramabad                                             | 33° 30' 49" N | 48° 21' 89" E | 1257         | HSBU- 2014202                           |
|                                                  | Khorasan Razavi, Torbate heydarieh                                | 35° 17' 11" N | 59° 13' 17" E | 1372         | HSBU- 2014203                           |
| 9. <i>P. rottboellioides</i> Jaub. & Spach       | Alborz, Shahrestanak                                              | 35° 58' 12" N | 51° 21' 13" E | 2186         | HSBU- 2014218                           |
| 10. <i>P. thymifolium</i> Jaub. & Spach          | Khorasan Razavi, North west of Torbate heydarieh- Kadkan mountain | 32° 24' 00" N | 59° 19' 00" E | 1850         | FUMH-17529                              |
| 11. <i>P. paronychioides</i> C.A. Mey. ex Hohen  | Khorasan Razavi, Gonabad, between Disfan & Khanik village         | 34° 09' 33" N | 58° 24' 39" E | 1500         | FUMH-12815                              |
| 12. <i>P. salicornioides</i> Jaub. & Spach       | Kerman, Gonabad, Hranan village                                   | 29° 30' 25" N | 56° 42' 36" E | 2900         | Shahid Bahonar University of Kerman 291 |
| 13. <i>P. luzuloides</i> Jaub. & Spach           | Kermanshah, 40 km Sonqor to Gorveh                                | 35° 03' 36" N | 47° 34' 40" E | 2259         | Shahid Bahonar University of Kerman 292 |
| 14. <i>P. spinosum</i> H. Gross                  | Fars, Kazerun                                                     | 29° 37' 10" N | 51° 39' 15" E | 850          | ALUH 1000                               |
| 15. <i>P. hyrcanicum</i> Rech.f.                 | Khorasan Razavi, Neyshaboar, Mirabad village                      | 36° 06' 37" N | 58° 46' 47" E | 1750         | FUMH-26061                              |
| 16. <i>P. dumosum</i> Boiss.                     | Kerman, Sirjan                                                    | 29° 27' 31" N | 55° 40' 17" E | 1742         | Shahid Bahonar University of Kerman 293 |
| 17. <i>P. alpestre</i> C.A. Mey.                 | Tehran, Damavand, Tar lake                                        | 35° 43' 48" N | 52° 13' 39" E | 2900         | HSBU-2014220                            |

Genetic and morphological diversity in *Polygonum* species**Table 2.** Morphological characters in studied species.

|                                                              |                                                                                                                                                                 |
|--------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1. Leaf length (mm)                                          | 17. Status of achene to perianth (exserted 1, included 2)                                                                                                       |
| 2. Leaf width (mm)                                           | 18. Color of flower (green with whitish margin 1, green with pinkish margin 2, dark pink 3, white 4, green 5)                                                   |
| 3. Leaf length / leaf width (mm)                             | 19. Shape of base of ochrea (tubular 1, funnel form 2, inflated 3)                                                                                              |
| 4. Ochrea length (mm)                                        | 20. Shape of apex of ochrea (lacerate 1, acuminate 2, flat 3)                                                                                                   |
| 5. Flower length (mm)                                        | 21. Ochrea veins (conspicuous 1, inconspicuous 2)                                                                                                               |
| 6. Flower width (mm)                                         | 22. Achene apex (short 1, long 2)                                                                                                                               |
| 7. Flower length / flower width (mm)                         | 23. Color of leaf (light green 1, dark green 2)                                                                                                                 |
| 8. Achene length (mm)                                        | 24. Stem branches (branched from the base 1, erect with branches 2, dichotomously branched 3)                                                                   |
| 9. Achene width (mm)                                         | 25. vegetative form (prostrate 1, decumbent 2, erect 3, procumbent 4, fragile and drooping 5, shrubby 6, shrubby-caespitose 6)                                  |
| 10. Achene length / achene width (mm)                        | 26. Pedicel (pedicellate 1, subsessile 2)                                                                                                                       |
| 11. Length of longest internode (mm)                         | 27. Petiole (petiolate 1, sessile 2)                                                                                                                            |
| 12. Petiole length (mm)                                      | 28. Leaf shape (linear 1, lanceolate 2, oblong 3, elliptic 4, ovate-lanceolate 5, linear-lanceolate 6, oblong-elliptic 7, oblong-linear 8, oblong-lanceolate 9) |
| 13. Pedicel length (mm)                                      | 29. Shape of inflorescence (axillary with solitary flowers 1, flowers in fascicle 2)                                                                            |
| 14. Ratio of tube to perianth length (mm)                    | 30. Achene shape (ovoid 1, digonous 2, trigonous 3)                                                                                                             |
| 15. Ochrea length (mm)                                       | 31. Color of achene (light brown 1, dark brown 2, black 3)                                                                                                      |
| 16. Color of ochrea (light brown 1, hyaline 2, dark brown 3) | 32. Plant life form (annual 1, perennial 2)                                                                                                                     |

between the studied populations. Multidimensional Scaling (MDS) was performed to group studied species and Principle Component Analysis (PCA) was used to identify the most variable morphological characters among these species (PODANI 2000). Data analyses were performed using PAST ver. 2.17 (HAMMER et al. 2001).

### Genetic diversity and population structure

ISSR bands were coded as binary characters (presence = 1, absence = 0). Genetic diversity parameters like Nei's gene diversity (H), Shannon information index (I), number of effective alleles and percentage of polymorphism were determined for each species (WEISING et al. 2005; FREELAND et al. 2011). PopGene ver. 1.32 was used for these parameters.

Ward, NJ clustering and UPGMA methods (PopGene ver. 1.32 & PAST ver. 2.17) were used for grouping after 100 times bootstrapping replicates. Jaccard and Nei's genetic distance was used for clustering (FREELAND et al. 2011; HUSON & BRYANT 2006).

The Mantel test was performed to investigate the correlation between geographical and genetic distances of studied populations by GenAlEx ver. 6.4.

Pearson coefficient of correlation using PAST ver. 2.17 was determined between geographical features (altitude, longitude and latitude) and genetic diversity parameters.

AMOVA (Analysis of Molecular Variance) test (with 100 permutations) as implemented in GenAEx ver. 6.4 (PEAKALL & SMOUSE 2006) and Nei's  $G_{ST}$  analysis of GenoDive ver. 2 (2013) (MEIRMANS & VAN TIENDEREN 2004), were used to reveal significant genetic differences among the studied species (IJBARI et al. 2014).

Population genetic differentiation was studied by  $G'_{ST\_est}$  = standardized measure of genetic differentiation (HEDRICK 2005), and  $D_{est}$  = Jost measure of differentiation (JOST 2008). In order to overcome potential problems caused by the dominance of ISSR markers, a Bayesian program, Hickory ver. 1.0 (HOLSINGER & LEWIS 2003), was used to estimate parameters related to genetic structure (theta B value) (TERO et al. 2003).

Bayesian based model STRUCTURE analysis was used to study the genetic structure of populations (PRITCHARD et al. 2000). For this analysis, data were scored as dominant markers (FALUSH et al. 2007). The Evanno test was used to identify optimum number of genetic groups (K; EVANNO et al. 2005).

### Gene flow

Gene flow was determined by different approaches: 1) Calculating  $N_m$ : an estimate of gene flow from  $G_{ST}$  by PopGene ver. 1.32 as:  $N_m = 0.5 (1 - G_{ST})/G_{ST}$ . This approach considers equal amounts of gene flow among all populations. 2) STRUCTURE analysis based on the admixture model (PRITCHARD et al. 2000) and 3) Population assignment test based on maximum likelihood as performed in GenoDive ver. 2. To check if ISSR markers show correlation with environmental features of the studied populations, LFMM analysis (Latent Factor Mixed Models) was done. This method tests correlations between environmental and genetic variation while estimating the effects of hidden factors that represent background residual levels of population structure (FRICHOT et al. 2013).

## Results

### Morphometry

ANOVA showed significant differences ( $P < 0.01$ ) in quantitative morphological characters among the species studied. In order to determine the most variable characters among the taxa studied, PCA analysis has been performed. It revealed that the first three factors comprised over 61.54% of the total variation. In the first PCA axis with 41.03% of total variation, such characters as leaf length, leaf width, ochrea length, length of longest internode, length of petiole and pedicel have shown the highest correlation ( $>0.7$ ). Ratio of leaf length to width, shape of leaf, shape of stems branches and vegetative form and plant life form were characters influencing PCA axis 2 and 3 respectively.

The MDS plot of morphological characters (Fig. 1) separated the species into distinct groups. The plants of *P. olivascens* showed more similarity to *P. patulum* and *P. argyrocoleon*. Similarly, plants of *P. aviculare* and *P. arenastrum* were placed close to each other due to morphological similarity. Three perennial species of *P. spinosum*, *P. salicornioides* and *P. dumosum* (No. 12, 14 & 16) were placed close to each other and were separated from the annuals.


Genetic and morphological diversity in *Polygonum* species

Figure 1. MDS plot of *Polygonum* species based on morphological characters (Species numbers according to Table 1).

### Species delimitation and genetic diversity

All ISSR primers produced polymorphic bands. Genetic diversity parameters determined in the studied species are provided in Table 3. We used few plants for genetic diversity analysis in some of the species as we could obtain only few samples from different herbaria. Therefore, we believe that with a higher number of samples, we could get more precise information on the genetic variation within each species. However, in some cases, even with only three plants, high degree of genetic polymorphism was obtained. For instance, *P. rottboellioides* with three plants had a high degree (21.43%) of genetic polymorphism. Similarly, other species studied had only three samples included in the analysis but differed in genetic diversity parameters and also differed significantly in AMOVA test (see below). Moreover, irrespective of plant samples used in each species, they showed a very similar level of effective number of alleles (Table 3). Therefore, we can compare these values and draw some conclusions.

*Polygonum olivascens* and *P. argyrocoleon* had the highest value for genetic polymorphism (68.57% and 64.29% respectively). These two species also had high value for Nei gene diversity (0.18). However, *P. arenastrum* had the highest value for this parameter (0.19).

AMOVA test showed significant genetic difference ( $P = 0.01$ ) among studied species. It revealed that 40% of total variation was among species and 60% was within species. Pair-wise  $F_{ST}$  values showed significant difference among all studied species (Table 4). Moreover, genetic differentiation of these species was demonstrated by significant Nei's  $G_{ST}$  (0.54,  $P = 0.01$ ) and  $D_{est}$  values (0.174,  $P = 0.01$ ).


UPGMA, Ward and NJ clustering of the studied species produced similar results. Therefore, the Ward dendrogram is presented and discussed (Fig. 2). Most of the plants of each species were grouped together in a separate cluster. This indicates that ISSR molecular markers can be used in *Polygonum* species delimitation.

The perennial and annual species were separated in different clusters. The perennial cluster contained two sub-clusters. *P. dumosum* and *P. spinosum* showed genetic affinity to each other and formed the first sub-cluster. The other studied perennial species, i.e. *P. salicornioides*, *P. alpestre*, *P. thymifolium*, *P. paronychioides*, *P. luzuloides* and *P. hyrcanicum* comprised the other sub-cluster.

**Table 3.** Genetic diversity parameters in *Polygonum* species.

| Species                   | N  | Na   | Ne   | I    | He   | UHe  | %P    | Hs   |
|---------------------------|----|------|------|------|------|------|-------|------|
| <i>P. arenastrum</i>      | 12 | 1.21 | 1.33 | 0.29 | 0.19 | 0.20 | 60    | 0.22 |
| <i>P. patulum</i>         | 6  | 0.97 | 1.26 | 0.24 | 0.16 | 0.17 | 48.57 | 0.24 |
| <i>P. olivascens</i>      | 15 | 1.39 | 1.28 | 0.29 | 0.18 | 0.19 | 68.57 | 0.26 |
| <i>P. argyrocoleon</i>    | 15 | 1.29 | 1.29 | 0.28 | 0.18 | 0.19 | 64.29 | 0.23 |
| <i>P. molliiforme</i>     | 6  | 0.53 | 1.05 | 0.04 | 0.03 | 0.03 | 8.57  | 0.04 |
| <i>P. polycnemoides</i>   | 10 | 1.26 | 1.33 | 0.30 | 0.20 | 0.21 | 62.86 | 0.24 |
| <i>P. kitaibelianum</i>   | 2  | 0.29 | 1    | 0    | 0    | 0    | 0     | 0    |
| <i>P. aviculare</i>       | 12 | 1.17 | 1.25 | 0.25 | 0.16 | 0.17 | 58.57 | 0.23 |
| <i>P. rottboellioides</i> | 3  | 0.46 | 1.14 | 0.12 | 0.08 | 0.10 | 21.43 | 0.14 |
| <i>P. thymifolium</i>     | 3  | 0.19 | 1.07 | 0.05 | 0.03 | 0.04 | 7.14  | 0.04 |
| <i>P. paronychioides</i>  | 3  | 0.11 | 1.03 | 0.02 | 0.01 | 0.02 | 2.86  | 0.02 |
| <i>P. salicornioides</i>  | 3  | 0.13 | 1.05 | 0.04 | 0.03 | 0.03 | 5.71  | 0.01 |
| <i>P. luzuloides</i>      | 3  | 0.16 | 1.05 | 0.04 | 0.03 | 0.03 | 5.71  | 0.03 |
| <i>P. spinosum</i>        | 3  | 0.09 | 1    | 0    | 0    | 0    | 0     | 0.00 |
| <i>P. hyrcanicum</i>      | 3  | 0.06 | 1.03 | 0.02 | 0.01 | 0.02 | 2.86  | 0.01 |
| <i>P. dumosum</i>         | 3  | 0.37 | 1.08 | 0.06 | 0.04 | 0.05 | 8.57  | 0.05 |
| <i>P. alpestre</i>        | 3  | 0.30 | 1.06 | 0.05 | 0.03 | 0.04 | 7.14  | 0.04 |


N: number of samples, Na: mean number of alleles, Ne: number of effective alleles, I: Shannon's information index, He: gene diversity, UHe: unbiased genetic diversity, %P: Percentage of polymorphic loci, Hs: Genetic diversity due to population.

**Figure 2.** Ward clustering of *Polygonum* species based on ISSR data.

Genetic and morphological diversity in *Polygonum* species**Table 4.** Pair-wise  $F_{ST}$  values among the studied *Polygonum* species. (Above diagonal =  $F_{ST}$  value, below diagonal = P value).

|      | sp1   | sp2   | sp3   | sp4   | sp5   | sp6   | sp7   | sp8   | sp9   | sp10  | sp11  | sp12  | sp13  | sp14  | sp15  | sp16  | sp17  |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| sp1  | —     | 0.173 | 0.272 | 0.359 | 0.639 | 0.325 | 0.519 | 0.335 | 0.297 | 0.457 | 0.47  | 0.436 | 0.485 | 0.506 | 0.435 | 0.541 | 0.523 |
| sp2  | 0.001 | —     | 0.112 | 0.25  | 0.645 | 0.153 | 0.443 | 0.24  | 0.14  | 0.204 | 0.255 | 0.225 | 0.278 | 0.364 | 0.25  | 0.445 | 0.412 |
| sp3  | 0.001 | 0.014 | —     | 0.124 | 0.54  | 0.16  | 0.369 | 0.248 | 0.101 | 0.238 | 0.303 | 0.256 | 0.239 | 0.269 | 0.246 | 0.393 | 0.406 |
| sp4  | 0.001 | 0.001 | 0.001 | —     | 0.532 | 0.266 | 0.454 | 0.277 | 0.25  | 0.297 | 0.33  | 0.329 | 0.293 | 0.328 | 0.314 | 0.475 | 0.458 |
| sp5  | 0.001 | 0.005 | 0.001 | 0.001 | —     | 0.585 | 0.922 | 0.618 | 0.828 | 0.895 | 0.917 | 0.897 | 0.905 | 0.919 | 0.918 | 0.894 | 0.911 |
| sp6  | 0.001 | 0.003 | 0.001 | 0.001 | 0.001 | —     | 0.399 | 0.282 | 0.271 | 0.382 | 0.424 | 0.403 | 0.389 | 0.43  | 0.403 | 0.433 | 0.45  |
| sp7  | 0.006 | 0.041 | 0.008 | 0.012 | 0.029 | 0.018 | —     | 0.392 | 0.725 | 0.884 | 0.956 | 0.916 | 0.915 | 1     | 0.956 | 0.908 | 0.913 |
| sp8  | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.01  | —     | 0.212 | 0.282 | 0.278 | 0.226 | 0.311 | 0.362 | 0.185 | 0.488 | 0.419 |
| sp9  | 0.003 | 0.078 | 0.083 | 0.003 | 0.012 | 0.004 | 0.1   | 0.014 | —     | 0.545 | 0.585 | 0.482 | 0.544 | 0.651 | 0.452 | 0.677 | 0.686 |
| sp10 | 0.004 | 0.047 | 0.007 | 0.003 | 0.017 | 0.003 | 0.084 | 0.008 | 0.102 | —     | 0.323 | 0.289 | 0.372 | 0.762 | 0.696 | 0.827 | 0.799 |
| sp11 | 0.002 | 0.03  | 0.004 | 0.002 | 0.012 | 0.003 | 0.094 | 0.008 | 0.092 | 0.097 | —     | 0.561 | 0.679 | 0.889 | 0.8   | 0.867 | 0.841 |
| sp12 | 0.003 | 0.045 | 0.004 | 0.004 | 0.01  | 0.005 | 0.101 | 0.022 | 0.1   | 0.187 | 0.104 | —     | 0.529 | 0.789 | 0.6   | 0.836 | 0.816 |
| sp13 | 0.002 | 0.013 | 0.006 | 0.003 | 0.008 | 0.004 | 0.09  | 0.009 | 0.098 | 0.222 | 0.097 | 0.094 | —     | 0.778 | 0.7   | 0.831 | 0.815 |
| sp14 | 0.003 | 0.029 | 0.003 | 0.001 | 0.013 | 0.006 | 0.111 | 0.003 | 0.102 | 0.09  | 0.091 | 0.099 | 0.111 | —     | 0.909 | 0.885 | 0.891 |
| sp15 | 0.002 | 0.081 | 0.006 | 0.003 | 0.014 | 0.004 | 0.113 | 0.023 | 0.101 | 0.105 | 0.094 | 0.103 | 0.098 | 0.101 | —     | 0.844 | 0.838 |
| sp16 | 0.004 | 0.02  | 0.004 | 0.002 | 0.01  | 0.002 | 0.102 | 0.004 | 0.105 | 0.081 | 0.115 | 0.1   | 0.101 | 0.099 | 0.104 | —     | 0.764 |
| sp17 | 0.003 | 0.011 | 0.005 | 0.003 | 0.012 | 0.005 | 0.116 | 0.003 | 0.099 | 0.104 | 0.095 | 0.091 | 0.106 | 0.114 | 0.087 | 0.101 | —     |

The annual species revealed a higher variation and were scattered in different sub-clusters. In general, *P. olivascens* and *P. argyrocoleon* (coded 3 and 4 respectively), showed genetic affinity and were placed close to each other. *P. kitaibelianum* and *P. aviculare* (coded 7 and 8 respectively), showed closer genetic affinity and formed a separate sub-cluster. *P. molliiforme* (coded 5) was genetically different from the other annual species and was placed far from the other annual taxa studied. In general, species relationships obtained from ISSR data agrees well with species relationship obtained from morphological characters.

**Figure 3.** STRUCTURE plot of *Polygonum* species based on ISSR data.

**Table 5.** The population assignment test.

| Individual | Current | Inferred | Lik_max | Lik_home | Lik_ratio |
|------------|---------|----------|---------|----------|-----------|
| 10         | Pop1    | Pop3     | -37.296 | -42.281  | 9.971     |
| 16         | Pop2    | Pop3     | -32.485 | -39.157  | 13.344    |
| 17         | Pop2    | Pop8     | -18.506 | -31.885  | 26.758    |
| 18         | Pop2    | Pop1     | -28.084 | -35.181  | 14.194    |
| 24         | Pop3    | Pop4     | -39.388 | -42.327  | 5.878     |
| 27         | Pop3    | Pop2     | -37.287 | -37.303  | 0.033     |
| 34         | Pop4    | Pop6     | -26.177 | -38.972  | 25.589    |
| 44         | Pop4    | Pop3     | -38.096 | -38.596  | 0.999     |
| 63         | Pop6    | Pop2     | -18.183 | -46.952  | 57.539    |
| 64         | Pop6    | Pop3     | -40.926 | -53.259  | 24.667    |
| 65         | Pop7    | Pop6     | -45.856 | —        | —         |
| 66         | Pop7    | Pop6     | -45.856 | —        | —         |
| 79         | Pop9    | Pop3     | -28.717 | —        | —         |
| 80         | Pop9    | Pop3     | -31.621 | —        | —         |
| 81         | Pop9    | Pop3     | -26.513 | —        | —         |
| 82         | Pop10   | Pop2     | -30.156 | —        | —         |
| 83         | Pop10   | Pop2     | -30.156 | —        | —         |
| 84         | Pop10   | Pop2     | -18.183 | —        | —         |
| 85         | Pop11   | Pop2     | -24.862 | —        | —         |
| 86         | Pop11   | Pop2     | -24.862 | —        | —         |
| 87         | Pop11   | Pop2     | -26.472 | —        | —         |
| 88         | Pop12   | Pop8     | -25.745 | —        | —         |
| 89         | Pop12   | Pop2     | -25.085 | —        | —         |
| 90         | Pop12   | Pop8     | -27.806 | —        | —         |
| 91         | Pop13   | Pop2     | -31.765 | —        | —         |
| 92         | Pop13   | Pop3     | -29.956 | —        | —         |
| 93         | Pop13   | Pop3     | -29.323 | —        | —         |
| 94         | Pop14   | Pop4     | -30.386 | —        | —         |
| 95         | Pop14   | Pop4     | -30.386 | —        | —         |
| 96         | Pop14   | Pop4     | -30.386 | —        | —         |
| 97         | Pop15   | Pop8     | -23.799 | —        | —         |
| 98         | Pop15   | Pop8     | -18.506 | —        | —         |
| 99         | Pop15   | Pop8     | -23.799 | —        | —         |
| 100        | Pop16   | Pop6     | -59.693 | —        | —         |
| 101        | Pop16   | Pop6     | -59.693 | —        | —         |
| 102        | Pop16   | Pop3     | -53.156 | —        | —         |
| 103        | Pop17   | Pop2     | -51.557 | —        | —         |
| 104        | Pop17   | Pop8     | -50.993 | —        | —         |
| 105        | Pop17   | Pop2     | -48.561 | —        | —         |

Mantel test with 5000 permutations showed a significant correlation ( $r=0.13$ ,  $p<0.01$ ) between genetic distance and geographical distance, so isolation by distance (IBD) occurred among the *Polygonum* species studied. Altitude was negatively correlated with genetic polymorphism ( $r=-0.516$ ,  $P=0.03$ ) and gene diversity ( $r=-0.40$ ,  $P=0.11$ ). Similarly, latitude was negatively (but not significantly) correlated and the longitude was positively (but not significantly) correlated with these genetic diversity parameters.

### The species genetic STRUCTURE

We performed STRUCTURE analysis followed by the Evanno test to identify the optimal number of genetic groups. We used the admixture model to illustrate interspecific gene flow or/ and ancestrally shared alleles in the species studied.

STRUCTURE analysis followed by Evanno test produced  $\Delta K=5$ . The STRUCTURE plot (Fig. 3) produced more detailed information about the genetic structure of the species studied as well as shared ancestral alleles and/ or gene flow among *Polygonum* species. This plot revealed that the perennial species (10–15) are genetically different from the studied annual *Polygonum* species (1–9), supporting results from the Ward dendrogram. However, the species 16 and 17 had shared alleles (green colored segment) with some of the annual species (2, 3, 4 and 6). The STRUCTURE plot also revealed that the annual species 1 and 5 differed genetically from the other studied species. This plot revealed a higher degree of genetic admixture among the annual *Polygonum* species compared to the perennials.

The low  $N_m$  value (0.28) indicates limited gene flow or ancestrally shared alleles between the species studied. The population assignment test (Table 5) also supported gene flow between species, particularly between species 2, 3 and 4 and also revealed the occurrence of shared alleles between species 2 and 17, 2 and 8, 8 and 15, 8 and 16, etc.

The LFMM analysis identified 8 ISSR loci out of 70 as adaptive loci. These are ISSR loci 22, 30, 41, 55, 58, 65 and 68 (Table 6). Except ISSR locus 58 that had  $N_m$  value  $>1.0$ , the other 7 loci had low  $N_m$  values ( $<1.0$ ) and are private alleles. These loci had strong  $G_{ST}$  values (differentiating power) and low  $H_s$  values (diversity value) (Table 7).

## Discussion

Taxonomy of *Polygonum* is complicated due to species morphological similarities (HONG et al. 1998, MEERTS et al. 1990). For instance, morphological characters between *P. aviculare* and *P. arenastrum* are not clearly expressed and show some degree of overlap in Flora Iranica (RECHINGER & SCHIMAN-CZEIKA 1968; MOZAFFARIAN 2012). *Polygonum aviculare* s.l. is a polyploid complex of selfing annual weeds colonizing various man-disturbed habitats. Members of this complex share extensive phenotypic plasticity, autogamy and polyploidy. Genetic variation in morphometric and life history traits is related to the ecological specialization in members of this complex (MEERTS et al. 1998).

*Polygonum* species studied are predominantly self-pollinated and are expected to show low genetic diversity. In fact, some of the species showed low genetic polymorphism, but some others had a high degree of genetic variation (up to 68% genetic polymorphism).

*Polygonum olivascens* and *P. argyrocoleon* exhibited intraspecific genetic variation and genetic admixture in STRUCTURE plot. Intraspecific genetic variation has been reported in some other

**Table 6.** LFMM result showing ISSR loci with significant association with geographical parameters studied.

| Name    | Z score | -log <sub>10</sub> (p-value) | P-value     |
|---------|---------|------------------------------|-------------|
| ISSR 22 | 3.46859 | 3.28134                      | 0.000523192 |
| ISSR 30 | 2.11301 | 1.46092                      | 0.0346001   |
| ISSR 41 | 3.54403 | 3.40443                      | 0.000394064 |
| ISSR 55 | 1.89369 | 1.23458                      | 0.058266    |
| ISSR 58 | 2.98875 | 2.55266                      | 0.00280118  |
| ISSR 65 | 2.05601 | 1.40032                      | 0.0397817   |
| ISSR 68 | 3.80138 | 3.84196                      | 0.000143893 |

*Polygonum* species, too. For example, allozyme analysis of three alpine populations of *Polygonum viviparum*, a common herbaceous perennial with no observed sexual reproduction, revealed surprising levels of genetic diversity (DIGGLE et al. 1998). Similarly, high genetic variation was reported within and among selfing mating *Polygonum cespitosum* populations (MATESANZ et al. 2014).

Different reasons can be considered for intraspecific genetic variation of the species studied.

1) Occasional cross-pollination: Most authors have considered *P. aviculare* and related taxa to be self-pollinated (SCHOLZ 1958; STYLES 1962; MEERTS et al. 1998). Indeed, the inconspicuous inflorescences with flowers partially hidden in the ochrea and the reduction of floral nectaries seem to indicate an evolutionary tendency in this direction. YURTSEVA (1998) also reported that emasculated chasmogamous flowers did not produce fruits. However, the floral dimorphism present in *P. aviculare* complex is difficult to explain in view of the autogamy reported for these plants. Structures that make cross-pollination possible in other *Polygonum* s.l. are preserved in *P. aviculare* and there are numerous reports of flowers visited by insects (LÖVE & LÖVE 1956; SCHMID 1983; KARLSSON 2000). The possibility of limited ongoing gene flow among some of the species studied is supported by the fact that morphological similarities and overlapping characters do occur in these species. This is also supported by population assignment test presented before.

2) Phenotypic plasticity: Phenotypic plasticity is a well-known phenomenon in *Polygonum* (ZANGERL & BAZZAZ 1983; SULTAN 2003; MATESANZ et al. 2012). This plasticity can lead to genome expression differences and morphological differences.

**Table 7.** Nm analysis of ISSR loci showing loci identified as adaptive loci in LFMM analysis.

| Locus    | Hs     | Gst    | Nm     |
|----------|--------|--------|--------|
| Locus 22 | 0.086  | 0.8061 | 0.1203 |
| Locus 30 | 0.12   | 0.6315 | 0.2918 |
| Locus 41 | 0.0929 | 0.6966 | 0.2178 |
| Locus 55 | 0.1249 | 0.3576 | 0.8980 |
| Locus 58 | 0.1532 | 0.2670 | 1.3727 |
| Locus 65 | 0      | 1      | 0      |
| Locus 68 | 0.1606 | 0.3380 | 0.9795 |

STRUCTURE analysis and AMOVA indicated high interspecific genetic differentiation among the studied *Polygonum* species. These species are invasive and invasive plants form new colonies and occupy new places by few founder individuals. Therefore, genetic drift and inbreeding become important evolutionary forces in the species history. The low level of within species genetic variation observed in some of the species studied might be due to genetic drift (DEWALT & HAMRICK 2004; NOVAK & MACK 2005). This is supported by LFMM analysis that identified 8 ISSR loci out of 70 as adaptive loci. Seven loci had low Nm value ( $< 1.0$ ) and were private alleles. These private alleles may be fixed by genetic drift and inbreeding in new founders during invasion in new habitats (MATESANZ et al. 2012; MATESANZ et al. 2014).

To conclude, the present study revealed the use of ISSR molecular markers along with morphological characters in *Polygonum* species delimitation. This molecular marker revealed genetic distinctness of annuals versus perennials. Some degrees of interspecific genetic admixture occur in *Polygonum*, but the studied species are strongly differentiated during the speciation process and invasion in new habitats. Genetic drift, strong inbreeding and local adaptation are effective evolutionary forces operating in *Polygonum* species and population divergence and adaptation.

Plant species delimitation is of central importance in phylogenetic systematics, evolution, biogeography and biodiversity. It is significant to infer patterns and mechanisms of speciation and hybridization, the evolutionary process by which new biological species arise and gene flow between closely related phylogenetic species can occur (SCHLUTER 2001; DUMINIL & DI MICHELE 2009). Isolation by distance, local adaptation and gene flow are different mechanisms responsible for species differentiation and genetic diversity (FREELAND et al. 2011; FRICHOT et al. 2013). Genetic diversity of *Polygonum* species can be due to these factors. However, further investigations can help to interpret this diversification better.

## Acknowledgements

We are grateful to the curators of Herbarium of Ferdowsi University of Mashhad [FUMH], Kharazmi University Herbarium [T] and Shahid Bahonar University of Kerman for permission to extract DNA from selected specimens.

## References

- BENTHAM G. & HOOKER J. D. (1880): Genera Plantarum. Vol. 3: 88–105. – London: Reeve.
- DEWALT S. J. & HAMRICK J. L. (2004): Genetic variation of introduced Hawaiian and native Costa Rican populations of an invasive tropical shrub, *Clidemia hirta* (Melastomataceae). – Amer. J. Bot. **91**: 1155–1162.
- DIGGLE P. K., LOWER S. & RANKER T. A. (1998): Clonal diversity in alpine populations of *Polygonum viviparum* (Polygonaceae). – Int. J. Plant. Sci. **159**: 606–615.
- DUMINIL J. & DI MICHELE M. (2009): Plant species delimitation: A comparison of morphological and molecular markers. – Pl. Biosyst. **143**(3): 528–542.
- EVANNO G., REGNAUT S. & GOUDET J. (2005): Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. – Molec. Ecol. **14**: 2611–2620.
- FALUSH D., STEPHENS M. & PRITCHARD J. K. (2007): Inference of population structure using multilocus genotype data: dominant markers and null alleles. – Molec. Ecol. Notes **7**: 574–578.

- FREELAND J. R., KIRK H. & PETERSON S. D. (2011): *Molecular Ecology*. [2<sup>nd</sup> ed.] – Chichester, UK: Wiley-Blackwell.
- FRICHOT E., SCHOVILLE S. D., BOUCHARD G. & FRANCOIS O. (2013): Testing for associations between loci and environmental gradients using latent factor mixed models. – *Molec. Biol. Ecol.* **30**: 1687–1699.
- GALASSO G., BANFI E., DE MATTIA F., GRASSI F., SGORBATI S. & LABRA M. (2009): Molecular phylogeny of *Polygonum* L. s.l. (Polygonaceae), focusing on European taxa: preliminary results and systematic considerations based on *rbcL* plastidial sequence data. – *Atti Soc. Ital. Sci. Nat. Mus. Civico Storia Nat. Milano* **150** (I): 113–148.
- GRIFFITH T. M. & SULTAN S. E. (2006): Plastic and constant developmental traits contribute to adaptive differences in co-occurring *Polygonum* species. – *Oikos* **114**: 5–14.
- GROSS H. (1913): Remarques sur les *Polygonées* de l'Asie Orientale. – *Bull. Acad. Int. Géogr. Bot.* **23**: 7–32.
- HAMMER Ø., HARPER D. & RYAN P. D. (2001): PAST: Paleontological Statistics software package for education and data analysis. – *Palaeontologia Electronica* **4**(1): 1–9. – [http://palaeo-electronica.org/2001\\_1/past/past.pdf](http://palaeo-electronica.org/2001_1/past/past.pdf)
- HARALDSON K. (1978): Anatomy and taxonomy in Polygonaceae subfamily Polygonaceae Meisn. emend. Jaretsky. – *Symb. Bot. Upsal.* **22**: 1–95.
- HEDBERG O. (1946): Pollen morphology in the genus *Polygonum* L. sensu lato and its taxonomic significance. – *Svensk Bot. Tidskr.* **40**: 371–404.
- HEDRICK P. W. (2005): A standardized genetic differentiation measure. – *Evolution* **59**(8): 1633–1638.
- HOLSINGER K. E. & LEWIS P. O. (2003): Hickory: a package for analysis of population genetic data V1.0. – Department of Ecology and Evolutionary Biology, University of Connecticut – <http://www.eeb.uconn.edu>
- HONG S. P., RONSE DECRAENE L. P. & SMETS E. (1998): Systematic significance of tepal surface morphology in tribes Persicarieae and Polygoneae (Polygonaceae). – *Bot. J. Linn. Soc.* **127**: 91–116.
- HUSON D. H. & BRYANT D. (2006): Application of phylogenetic networks in evolutionary studies. – *Molec. Biol. Evol.* **23**: 254–267.
- IJBARI H., SHEIDAI M., MEHRABIAN A. R., NOORMOHAMMADI Z. & GHASEMZADEH-BARAKI S. (2014): K-means clustering and STRUCTURE analyses of genetic diversity in *Tamarix* L. accessions. – *Turk. J. Bot.* **38**: 1080–1094.
- JOST L. (2008): GST and its relatives do not measure differentiation. – *Molec. Ecol.* **17**(18): 4015–4026.
- KARLSSON T. (2000): The genus *Polygonum* L. – In: JONSELL B. [ed.]: *Flora Nordica*. Vol. 1: 255–273. – Stockholm: Bergius Foundation, The Royal Swedish Academy of Sciences.
- KESHAVARZI M., MOSAFERI S. & SHOJAI M. (2012): Leaf anatomical studies of the annual species of *Polygonum* s.l. (Polygonaceae) in Iran. – *Phytol. Balc.* **18**(2): 127–133.
- KRIŽMAN M., JAKSE J., BARICEVIC D., JAVORNIK B. & PROSEK M. (2006): Robust CTAB-activated charcoal protocol for plant DNA extraction. – *Acta. Agri. Slov.* **87**: 427–433.
- LAMB-FRYE A. S. & KRON K. A. (2003): *RbcL* Phylogeny and Character Evolution in Polygonaceae. – *Syst. Bot.* **28**(2): 326–332.
- LI A., BAO B., GRABOVSKAYA-BORODINA A. E., HONG S. P., MCNEILL J., MOSYAKIN S. L., OHBA H. & PARK C. W. (2003): Polygonaceae. – In: *Flora of China*. Vol 5: 277–350.
- LÖVE A. & LÖVE D. (1956): Chromosomes and Taxonomy of Eastern North American *Polygonum* l. – *Can. J. Bot.* **34**: 501–521.
- MATESANZ S., HORGAN-KOBELSKI T. & SULTAN S. E. (2012): Phenotypic plasticity and population differentiation in an ongoing species invasion. – *PLoS ONE* **12**: e44955.

- MATESANZ S., THEISS K. E., HOLSINGER K. E. & SULTAN S. E. (2014): Genetic Diversity and Population Structure in *Polygonum cespitosum*: Insights to an Ongoing Plant Invasion. – PLoS ONE **9**(4): e93217.
- MEERTS P., BAYA T. & LEFÉBVRE C. (1998): Allozyme variation in the annual weed species complex *Polygonum aviculare* (Polygonaceae) in relation to ploidy level and colonizing ability. – Pl. Syst. Evol. **211**: 239–256.
- MEERTS P., BRIANE J. P. & LEFÉBVRE C. (1990): A numerical taxonomic study of the *Polygonum aviculare* complex (Polygonaceae) in Belgium. – Pl. Syst. Evol. **173**: 71–89.
- MEIRMANS P. G. & VAN TIENDEREN P. H. (2004): Genotype and Genodive: two programs for the analysis of genetic diversity of asexual organisms. – Molec. Ecol. Notes **4**: 792–794.
- MEISSNER C. F. (1826): Monographie generis Polygoni prodromus. – Geneva: A. Lador.
- MOZAFFARIAN V. (2012): A revision of *Polygonum* L. sensu lato, (Polygonaceae) in Iran. – Iran. J. Bot. **18**(2): 159–174.
- NOVAK S. J. & MACK R. N. (2005): Genetic bottlenecks in alien plant species. – In: SAX D. F., STACHOWICZ J. J. & GAINES S. D. [eds]: Species invasions, insights into ecology, evolution and biogeography: 201–228. – Sunderland MA: Sinauer.
- PEAKALL R. & SMOUSE P. E. (2006): GENAIEX 6: genetic analysis in Excel. Population genetic software for teaching and research. – Molec. Ecol. Notes **6**(1): 288–295.
- PODANI J. (2000): Introduction to the exploration of multivariate data [English translation]. – Leiden: Backhuyes.
- PRITCHARD J. K., STEPHENS M. & DONNELLY P. (2000): Inference of population structure using multilocus genotype Data. – Genetics **155**(2): 945–59.
- QAISER M. (2001): Polygonaceae. – In: Flora of Pakistan, Vol. 205. – Missouri: Missouri Botanical Garden Press.
- RECHINGER K. H. & SCHIMAN-CZEIKA H. (1968): Polygonaceae. – In: Flora Iranica, Vol. 56: 46–83. – Graz: Akademische Druck- und Verlagsanstalt.
- RONSE DECRAENE L. P. & AKEROYD J. R. (1988): Generic limits in *Polygonum* and related genera (*Polygonaceae*) on the basis of floral characters. – Bot. J. Linn. Soc. **98**: 321–371.
- RONSE DECRAENE L. P., HONG S. P. & SMETS E. (2000): Systematic significance of fruit morphology and anatomy in tribes *Persicarieae* and *Polygoneae* (Polygonaceae). – Bot. J. Linn. Soc. **134**: 301–337.
- SCHLUTER D. (2001): Ecology and the origin of species. – Trends Ecol. Evol. **16**: 372–380.
- SCHMID K. (1983): Untersuchungen an *Polygonum aviculare* s.l. – Mitt. Bot. Staatssammlung München **19**: 29–149.
- SCHOLZ H. (1958): Die Systematik des europäischen *Polygonum aviculare* L. I. – Ber. Deutsch. Bot. Ges. **71**: 427–434; II *ibid.* **72**: 63–72.
- STYLES B. T. (1962): The taxonomy of *Polygonum aviculare* and its allies in Britain. – Watsonia **5**(4): 117–241.
- SULTAN S. E. (2003): Phenotypic plasticity in plants: a case study in ecological development. – Evol. Dev. **5**(1): 25–33.
- TERO N., ASPI J., SIKAMAKI P., JAKALANIEMI A. & TUOMI J. (2003): Genetic structure and gene flow in a metapopulation of an endangered plant species, *Silene tatarica*. – Molec. Ecol. **12**(8): 2073–2085.
- VOYLOKOVA V. N., YURTSEVA O. V., BOBROVA V. K. & TROITSKY A. V. (2009): Morphological and molecular evidences of hybrid origin of *Polygonum aschersonianum* and *Polygonum samarense* (Polygonaceae). – Bot. Zhurn. **94**(2): 161–187.
- WEISING K., NYBOM H., WOLFF K. & KAHL G. (2005): DNA Fingerprinting in Plants. – Principles, Methods, and Applications. [2<sup>nd</sup> ed.] – Boca Raton: CRC Press.

- YURTSEVA O. V. (1998): Self- pollination in species of *Polygonum* subsection *Polygonum*. – Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. **103** (5): 61–67. [In Russian]
- YURTSEVA O. V. (2001): Ultra sculpture of achene surface in *Polygonum* section *Polygonum* (Polygonaceae) in Russia. – Nord. J. Bot. **21**: 513–528.
- ZANGERL A. R. & BAZZAZ F. A. (1983): Responses of an early and a late successional species of *Polygonum* to variations in resource availability. – Oecologia **56**: 397–404.

Addresses of the authors:

Samaneh Mosaferi (author for correspondence)

Masoud Sheidai

Faculty of Biological Sciences

Shahid Beheshti University

Tehran

Iran

E-mail: samanehmosaferi@yahoo.com

Maryam Keshavarzi

Faculty of Biological Sciences

Alzahra University

Tehran

Iran

Zahra Noormohammadi

Department of Biology

Sciences and Research Branch

Islamic Azad University

Tehran

Iran

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Wulfenia](#)

Jahr/Year: 2016

Band/Volume: [23](#)

Autor(en)/Author(s): Mosaferi Samaneh, Sheidai Masoud, Keshavarzi Jamshidian Maryam, Noormohammadi Zahra

Artikel/Article: [Species delimitation and population structure analysis in Polygonum species \(section Polygonum\) 98-112](#)