

Phyton (Horn, Austria)	Vol. 42	Fasc. 2	225–236	20. 12. 2002
------------------------	---------	---------	---------	--------------

Gas Exchange Characteristics of *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis* (Tagasaste) Measured in Spring-Summer under Field Conditions

By

A. M. GONZÁLEZ-RODRÍGUEZ, A. MARTÍN-OLIVERA,

D. MORALES and M. S. JIMÉNEZ*)

With 5 figures

Received April 4, 2002

Accepted May 5, 2002

Key words: Carbon assimilation rate, diurnal courses, stomatal conductance, transpiration, water use efficiency.

Summary

GONZÁLEZ-RODRÍGUEZ A. M., MARTÍN-OLIVERA A., MORALES D. & JIMÉNEZ M. S. 2002. Gas exchange characteristics of *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis* (tagasaste) measured in spring-summer under field conditions. – *Phyton* (Horn, Austria) 42 (2): 225–236, with 5 figures. – English with German summary.

Carbon assimilation rate (A), stomatal behaviour and transpiration (E) of *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis* (tagasaste) were studied in spring-summer in natural conditions in the North-West slope of Tenerife. Daily courses of this species did not show midday depression of A . $A_{\max}$ measured in tagasaste ($10 \mu\text{mol m}^{-2} \text{s}^{-1}$) is in the limits given to tropical rain forest trees and trees of broadleaved evergreens of the subtropics and warm-temperate regions. Stomatal closure was shown under conditions of high air vapour pressure deficit (VPD) and leaf temperature, showing a relatively good stomatal control, better than other tree species of the Canarian laurel forest. Instantaneous water use efficiency decreased with higher evaporative demand, in contrast to intrinsic water use efficiency (A/g_s) that did not change significantly under different environmental conditions. In both

*) A.M. GONZÁLEZ-RODRÍGUEZ, A. MARTÍN-OLIVERA, D. MORALES & M. S. JIMÉNEZ
Dpto. de Biología Vegetal. Facultad de Farmacia. Universidad de La Laguna. C/
Astrofísico Sánchez, s/n. 38207 La Laguna. Tenerife. E-mail: aglerod@ull.es

cases these parameters were higher than those found in trees of the Canarian laurel forest, an ecosystem with similar distribution area to that of tagasaste. These characteristics together with a better stomatal control could make this species able to live outside of its natural distribution area and enable its use for revegetation in semiarid zones.

Zusammenfassung

GONZÁLEZ-RODRÍGUEZ A. M., MARTÍN-OLIVERA A., MORALES D. & JIMÉNEZ M. S. 2002. Die Gaswechseleigenschaften von *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis* (Tagasaste) im Frühling und im Sommer unter Freilandbedingungen. – Phytion (Horn, Austria) 42 (2): 225–236, 5 Abbildungen. – Englisch mit deutscher Zusammenfassung.

Die Kohlenstoffassimilationsrate (A), das Spaltöffungsverhalten und die Transpiration (E) von *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis* (Tagasaste) wurde im Frühling und Sommer unter Freilandbedingungen am Northwestabfall von Teneriffa untersucht. Der Tagesgang wies keine mittägliche Depression von A auf. Die bei Tagasaste gemessene $A_{\max}$ ($10 \mu\text{mol m}^{-2} \text{s}^{-1}$) liegt im Bereich der Bäume tropischer Regenwälder und immergrüner Laubbäume der Subtropen sowie warm temperierter Gebiete. Spaltöffnungsschluss konnte unter Bedingungen hoher Dampfdruckdefizite (VPD) und Blatttemperaturen festgestellt werden, was auf eine relativ gute stomatäre Kontrolle weist, welche viel besser ist als bei anderen Arten des kanarischen Lorbeerwaldes. Die momentane „water use efficiency“ sank mit größer werdender Beanspruchung der Wasserabgabe, was im Gegensatz zur wirklichen „water use efficiency“ (A/g_s) steht, welche sich nicht signifikant unter den verschiedenen Umweltbedingungen verändert. In beiden Fällen waren diese Parameter größer als bei den anderen Bäumen des kanarischen Lorbeerwaldes, einem Ökosystem mit ähnlicher Verbreitung wie von Tagasaste. Diese Eigenschaften, gemeinsam mit einer besseren Spaltöffnungskontrolle, könnten diese Art befähigen, auch außerhalb ihres natürlichen Verbreitungsgebiets vorzukommen und würde sie zur Wiederbepflanzung semiarider Zonen eignen.

Introduction

Tagasaste (*Chamaecytisus proliferus* (L. fil.) Link ssp. *proliferus* var. *palmensis* (Christ)) is a perennial leguminous shrub endemic of the Canary Islands (ACEBES GINOVÉS & al. 1991, FRANCISCO-ORTEGA & al. 1993) where it has been traditionally used as fodder for local herds. It was introduced in Australia and New Zealand in the XIX century (FRANCISCO-ORTEGA & al. 1991), and its cultivation has been greatly extended to feed livestock not only in these countries where it has gained economic importance, but also in others, such as Chile and Ethiopia (KAITHO & al. 1997, ARREDONDO & al. 1998, ASSEFA 1998). Most studies have been conducted about its biomass yield and quality (ARREDONDO & al. 1998, ASSEFA 1998), nutritive value and palatability (BORENS & POPPI 1990, MÉNDEZ 1993, TOLERA & al. 1997), however little is known about its physiological response under environmental conditions. Recent studies have been carried out on water use effi-

ciency and water balance of tagasaste in south-western Australia (LEFROY & al. 2001a, b) although nothing is known about its gas exchange characteristics under native field conditions.

In the Canary Islands, tagasaste is mainly present from 300/400 to 1300/1400 m of altitude in gaps and boundary zones of forests in N and NE slopes of the islands (PÉREZ DE PAZ & al. 1986) in a humid Mediterranean climate. Since many legume shrubs, because their ability for fixing nitrogen, are able to grow in low quality soils (VAN ANDEL & al. 1993) and have been used for revegetation of abandoned, altered or drought fields, tagasaste could be used for this purpose. This work is a preliminary study to know the behaviour of tagasaste in its native environment by evaluating photosynthetic and transpiration rates, and stomatal conductance. This will give insight to the general knowledge of the behaviour of this species in its native environment and also will constitute the basis for a possible use of this forage plant outside of its natural distribution area, not only in the Canary Island but also in other Mediterranean countries where more arid conditions occur.

Material and Methods

The study was conducted in adult plants of *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis* (tagasaste) from March to August of 2000 in Tenerife, Canary Islands. The experimental site is situated at an altitude of 950 m, facing NO and has a humid Mediterranean climate with a mean annual temperature between 10–13 °C.

Carbon assimilation rate, measured as CO₂ uptake (*A*), stomatal conductance (*g_s*) and transpiration rate, measured as water vapour efflux (*E*) were determined on attached apical non-woody shoots with a portable Infrared Gas Analyser (LCA2, Analytical Development Co. Ltd., Hoddesdon, Herts, U.K.) every hour from the early morning to the evening.

Gas exchange rates were calculated using equations developed by VON CAEMMERER & FARQUHAR 1981. Leaf temperature (*T_{leaf}*) was determined from air temperature and calculated energy fluxes (PARKINSON 1985).

Approximately 10 diurnal courses were measured from March to August 2000 on six attached apical non-woody shoots in different plants.

In the global analysis we included all data obtained during all measured days. Applying boundary lines to the resulting data clouds we obtained approximations of optimal ranges of gas exchange variation with environmental and physiological factors. This kind of results are very common in field measurements and has been evaluated in the same way by other authors (MASAROVICOVA & ELIAS 1986, CHEESEMAN & al. 1991, LARCHER 1995, CHEESEMAN & LEXA 1996).

Results

Figure 1 shows the diurnal courses of gas exchange during two sunny days with contrasting environmental conditions. On 25 April, a cloudless day, with a vapour pressure deficit (VPD) low (<0.58 kPa) and *T_{leaf}* below

26 °C, A_{max} reached 8.4 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and high values of g_s were recorded (150 $\text{mmol m}^{-2} \text{s}^{-1}$). No values of g_s below 130 $\text{mmol m}^{-2} \text{s}^{-1}$ were recorded. In contrast, during the weather conditions of 31 July, when VPD was high (mean value > 2 kPa) reaching maximum values of 5 kPa and T_{leaf} reached 39 °C, A_{max} was only 5 $\mu\text{mol m}^{-2} \text{s}^{-1}$, because it was limited by low g_s (80 $\text{mmol m}^{-2} \text{s}^{-1}$). On both days, A paralleled diurnal pattern of PPFD.

On the day with low VPD and T_{leaf} (25 April), E reached maximum mean values below 3 $\text{mmol m}^{-2} \text{s}^{-1}$. This value increased to 4 $\text{mmol m}^{-2} \text{s}^{-1}$ under high VPD and T_{leaf} conditions on 31 July.

As a result of the increase of VPD, a decrease in instantaneous water use efficiency (WUE) was shown, varying from 3.2 $\mu\text{mol mmol}^{-1}$ on 25 April to 1.3 $\mu\text{mol mmol}^{-1}$ on 31 July. Intrinsic water use efficiency (A/g_s) did not change significantly, remaining around 60 $\mu\text{mol mmol}^{-1}$ (Fig. 1).



In order to analyse the total data we used boundary-line analysis using all data of each parameter plotted against every variable, to estimate the maximal parameter values when other variables were not limiting. The relationship between A and T_{leaf} indicated that the optimum temperature was 22 °C. On the basis of measurements of CO_2 gas exchange, a 90% of the photosynthetic rate at optimum temperature were obtained between 17.5 and 28 °C and a reduction of 50% of A was reached at temperature below 12 °C and above 40 °C (Fig. 2).

Response of A to PPFD yielded an A_{max} of 10 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and indicated that A reached light saturation above 700 $\mu\text{mol m}^{-2} \text{s}^{-1}$. The plot of A versus g_s did not pass through the origin because the stomata were partly open when A was zero. Values of g_s above 150 $\text{mmol m}^{-2} \text{s}^{-1}$ were considered to be non-limiting (Fig. 3).

Response of A to VPD followed a characteristic tendency in which higher VPD values led to lower A values. The curve showed optimal photosynthetic values until VPD values of 0.6 kPa and a progressive decrease of A at higher VPD values. This species maintained 50% of its A_{max} until 5 kPa of VPD (Fig. 3).

Relationships between g_s and PPFD and VPD are shown in figure 4. Values of g_s above 200 $\text{mmol m}^{-2} \text{s}^{-1}$ were exhibited at low PPFD (34 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and g_s began to decrease at 0.5 kPa and was less than 50% of maximum above 1.5 kPa.

Fig. 1. Diurnal courses of photosynthetic photon flux density (PPFD) at leaf surface, leaf temperature (T_{leaf}), air vapour pressure deficit (VPD), carbon assimilation rate (A), transpiration (E), stomatal conductance (g_s), instantaneous water use efficiency ($\text{WUE} = A/E$) and intrinsic water use efficiency (A/g_s) of *Chamaecytisus proliiferus* subsp. *proliiferus* var. *palmensis* measured on 25 April (a day with low VPD) and on 31 July (a day with high VPD). Each point represents the mean of six measurements with their standard deviations. Upper graphics show PPFD (solid line), VPD (○) and T_{leaf} (●).






Fig. 2. Relationships between carbon assimilation rate (A) and leaf temperature (T_{leaf}), in *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis*. Each point is an individual measurement.




Maximum values of E ($5 \text{ mmol m}^{-2} \text{ s}^{-1}$) were reached at 3 kPa of VPD and A_{max} (optimum $T_{\text{leaf}} 22^{\circ}\text{C}$) was reached at values of E close to $3.5 \text{ mmol m}^{-2} \text{ s}^{-1}$ (Fig. 5).

Discussion

Daily courses of gas exchange were different depending on environmental conditions. On days with low VPD, A , E and g_s were mainly controlled by PPFD; however, on days in which the VPD was high, A was lower because g_s was also lower throughout the whole day. Midday depression of A has been reported for temperate-zone plants (HODGES 1967, DOUGHERTY & HINCKLEY 1981), Mediterranean climate plants (TENHUNEN & al. 1981, PEREIRA & al. 1986) and tropical trees (MEDINA & al. 1978, ZOTZ & al. 1995, PATHRE & al. 1998, ISHIDA & al. 1999), however, we did not find midday depression of A in all studied days.

The increase of VPD to 5 kPa produced a reduction in A about 50%, mainly due to low g_s . Despite stomatal closure, E increased, producing a

Fig. 3. Relationships between carbon assimilation rate (A) and photosynthetic photon flux density (PPFD) at the leaf surface, water vapour stomatal conductance (g_s) and air vapour pressure deficit (VPD) in *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis*. Each point is an individual measurement.






Fig. 4. Relationships between water vapour stomatal conductance (g_s) and photosynthetic photon flux density (PPFD) at the leaf surface, and air vapour pressure deficit (VPD) in *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis*. Each point is an individual measurement.

decrease in WUE. However, the values of WUE obtained were higher than those found in the Canarian laurel forest trees (GONZÁLEZ-RODRÍGUEZ & al. 2001, 2002). In the same way, values of A/g_s were also higher than in Canarian laurel forest trees (GONZÁLEZ-RODRÍGUEZ & al. 2001, 2002). A significant increase of this ratio during the summer has been shown in some Mediterranean shrubs and trees (PEÑUELAS & al. 1998, FLEXAS & al. 2001), optimising their photosynthetic rate under drought conditions, nevertheless this parameter did not change significantly under different environmental conditions in tagasaste.




Fig. 5. Relationships between transpiration rate (E) and leaf temperature (T_{leaf}), and air vapour pressure deficit (VPD) in *Chamaecytisus proliferus* subsp. *proliferus* var. *palmensis*. Each point is an individual measurement.

Leaf photosynthesis in trees is variable since its maximum value under natural conditions ranges from roughly 3 to 30 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ (CEULEMANS & SAUGIER 1991). Generally, photosynthetic rates of tropical rain forest trees are relatively higher (between 10–16 $\mu\text{mol m}^{-2} \text{ s}^{-1}$) than trees of broadleaved evergreens of the subtropics and warm-temperate regions (between 6–12 $\mu\text{mol m}^{-2} \text{ s}^{-1}$) (LARCHER 1995). The maximum value of A measured in tagasaste (10 $\mu\text{mol m}^{-2} \text{ s}^{-1}$) is in the limits of both ranges.

A_{max} is reached with values of PPFD above 700 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, this value is in the range given by sun leaves of tropical forest trees and non-tropical

evergreen broadleaved trees (LARCHER 1995). Temperature optimum of A (22 °C) is similar to those found to evergreen broadleaved trees of the tropics and subtropics.

Stomatal conductance was sensitive to increasing VPD over the range 0 to 5 kPa, decreasing exponentially to values below 100 mmol m⁻² s⁻¹. Maximum values of g_s in tagasaste were relatively high compared with Mediterranean evergreen shrubs (KÖRNER 1995) and lower than in Canarian laurel forest trees (GONZÁLEZ-RODRÍGUEZ & al. 2001, 2002).

All these characteristics fit well with the environment where this shrub lives: in a subtropical region but with humid Mediterranean climate in the limits of the laurel forest distribution area. The WUE and A/g_s values higher than Canarian laurel forest trees and the better stomatal control as a response to the increase of VPD could make this species able to live outside of its natural distribution area, although more studies focused on drought responses of this shrub are needed before its use in semiarid zones for revegetation.

Acknowledgements

This research was supported by IFD97-1049-C08-04. We express our gratitude to La Laguna University and Santander Bank for the concession of a grant to A. M. OLIVERA.

References

- ACEBES-GINOVÉS J. R., DEL ARCO-AGUILAR M. & WILDPRET DE LA TORRE W. 1991. Revisión taxonómica del género *Chamaecytisus* Link en Canarias. – *Vieraea* 20: 191–202.
- ARREDONDO S., ARONSON J., OVALEE C., DEL POZO A. & AVENDANO J. 1998. Screening multipurpose legume trees in central Chile. – *For. Ecol. Manage.* 109 (1–3): 221–229.
- ASSEFA G. 1998. Biomass yield, botanical fractions and quality of tagasaste, (*Chamaecytisus palmensis*) as affected by harvesting interval in the highlands of Ethiopia. – *Agrofor. Syst.* 42(1): 13–23.
- BORENS F. & POPPI D. P. 1990. The nutritive and feeding value for ruminants of tagasaste (*Chamaecytisus palmensis*), a leguminous tree. – *Animal Feed Science and Technology* 28: 275–292.
- CEULEMANS P. J. & SAUGIER B. 1991. Photosynthesis. – In: RAGHAVENDRA A. S. (Ed.), *Physiology of trees*, pp. 21–50. – Wiley-Interscience, New York.
- CHEESEMAN J. M. & LEXA M. 1996. Gas exchange: models and measurements. – In: BAKER N. R. (Ed.), *Photosynthesis and the environment*, pp. 223–240. – Kluwer Academic Publishers.
- , CLOUGH B. F., CARTER D. R., LOVELOCK C. E., ONG JIN EONG & SIM R. G. 1991. The analysis of photosynthetic performance in leaves under field conditions: a case study using *Bruguiera* mangroves. – *Photosyn. Res.* 29: 11–22.
- DOUGHERTY W. J. S. & HINCKLEY T. M. 1981. The influence of severe drought on net photosynthesis of white oak (*Quercus alba*). – *Can. J. Bot.* 59: 335–341.

- FLEXAS J., GULÍAS J., JONASSON S., MUS M. & MEDRANO H. 2001. Seasonal patterns and control of gas exchange in local populations of the Mediterranean evergreen shrub *Pistacia lentiscus* L. – *Acta Oecol.* 22: 33–43.
- FRANCISCO-ORTEGA J., NEWBURY H. J. & FORD-LLOYD B. V. 1993. Numerical analyses of RAPD data highlight the origin of cultivated tagasaste (*Chamaecytisus proliferus* ssp. *palmensis*) in the Canary Islands. – *Theor. Appl. Genet.* 87: 264–270.
- , JACKSON M. T., SANTOS-GUERRA A. & FERNÁNDEZ-GALVÁN M. 1991. Historical aspects of the origin and distribution of tagasaste (*Chamaecytisus proliferus* (L. fil.) Link ssp. *palmensis* (Christ) Kunkel), a fodder tree from the Canary islands. – *J. Adelaide Bot. Gard.* 14(1): 67–76.
- GONZÁLEZ-RODRÍGUEZ A. M., MORALES D. & JIMÉNEZ M. S. 2001. Gas exchange characteristics of a Canarian laurel forest tree species (*Laurus azorica*) in relation to environmental conditions and leaf canopy position. – *Tree Physiol.* 21: 1039–1045.
- , MORALES D. & JIMÉNEZ M. S. 2002. Leaf gas exchange characteristics of a Canarian laurel forest tree species (*Persea indica* (L.) K. Spreng.) under natural conditions. – *J. Plant Physiol.* (in press).
- HODGES J. D. 1967. Patterns of photosynthesis under natural environmental conditions. – *Ecology* 48: 234–242.
- ISHIDA A., TOMA T. & MARJENAH 1999. Limitation of leaf carbon gain by stomatal and photochemical processes in the top canopy of *Macaranga conifera*, a tropical pioneer tree. – *Tree Physiol.* 19: 467–473.
- KATHO R. J., NSAH LAI I. V., WILLIAMS B. A., UMUNNA N. N., TAMMINGA S. & VAN BRUCHEM J. 1997. Relationships between preference, rumen degradability, gas production and chemical composition of browses. – *Agrofor. Syst.* 39 (2): 129–144.
- KÖRNER CH. 1995. Leaf diffusive conductances in the major vegetation types of the globe. – In: SCHULZE E. D. & CALDWELL M. M. (Eds.), *Ecophysiology of photosynthesis*, pp. 463–490. – Springer, Berlin.
- LARCHER W. 1995. *Physiological Plant Ecology*. – Third Edn. Springer, 506 p.
- LEFROY E. C., PATE J. S. & STIRZAKER R. J. 2001a. Growth, water use efficiency, and adaptive features of the tree legume tagasaste (*Chamaecytisus proliferus* Link.) on deep sands in south-western Australia. – *Aust. J. Agric. Res.* 52: 221–234.
- , STIRZAKER R. J. & PATE J. S. 2001b. The influence of tagasaste (*Chamaecytisus proliferus* Link.) trees on the water balance of an alley cropping system on deep sand in south-western Australia. – *Aust. J. Agric. Res.* 52: 235–246.
- MASAROVICOVÁ E. & ELIAS P. 1986. Photosynthetic rate and water relations in some forest herbs in spring and summer. – *Photosynthetica* 20(2): 187–195.
- MEDINA E., SOBRADO M. & HERRERA R. 1978. Significance of leaf orientation for leaf temperature in an Amazonian sclerophyll vegetation. – *Radiation and Environmental Biophysics* 15: 131–140.
- MÉNDEZ P. 1993. Forage potential of Canary Islands legumes. Management of mediterranean shrublands and related forage resources. – *Reur. Technical Series* 28: 141–144.
- PARKINSON K. J. 1985. A simple method for determining the boundary layer resistance in leaf cuvettes. – *Plant, Cell and Environment* 8: 223–226.

- PATHRE U., SINHA A. K., SHIRKE P. A. & SANE P. V. 1998. Factors determining the mid-day depression of photosynthesis in trees under monsoon climate. – *Trees* 12: 472–481.
- PEREIRA J. S., TENHUNEN J. D., LANGE O. L., BEYSLAG W., MEYER A. & DAVID M. M. 1986. Seasonal and diurnal patterns in leaf gas exchange of *Eucalyptus globulus* trees growing in Portugal. – *Can. J. For. Res.* 16: 177–184.
- PÉREZ DE PAZ P. L., DEL ARCO AGUILAR M., ACEBES GINOVES J. R. & WILDPRET DE LA TORRE W. 1986. Leguminosas forrajeras de Canarias. Excmo. Cabildo Insular de Tenerife. – ACT Series Publ. Científicas. Subser. Mus. Ins. De Cienc. Nat. 2, 157 pp.
- PEÑUELAS J., FILELLA I., LLUSIÀ J., SISCART D. & PIÑOL J. 1998. Comparative field study of spring and summer leaf gas exchange and photobiology of the mediterranean trees *Quercus ilex* and *Phillyrea latifolia*. – *J. Exp. Bot.* 49: 229–238.
- TENHUNEN J. D., LANGE O. L. & BRAUN M. 1981. Midday stomatal closure in Mediterranean type sclerophylls under stimulated habitat conditions in an environmental chamber. II Effect on the complex of leaf temperature and air humidity on gas exchange of *Arbutus unedo* and *Quercus ilex*. – *Oecologia* 50: 5–11.
- TOLERA A., KHAZAAL K. & ORSKOV E. R. 1997. Nutritive evaluation of some browse species. – *Animal Feed Science & Technology* 67 (2–3): 181–185.
- VAN ANDEL J., BAKKER J. P. & GROOTJANS A. P. 1993. Mechanisms of vegetation succession: a review of concepts and perspectives. – *Acta Bot. Neerl.* 42: 413–433.
- VON CAEMMERER S. & FARQUHAR G. D. 1981. Some relations between the biochemistry of photosynthesis and the gas exchange of leaves. – *Planta* 153: 376–387.
- ZOTZ G., HARRIS G., KÖNIGER M. & WINTER K. 1995. High rates of photosynthesis in the tropical pioneer tree, *Ficus insipida* Willd. – *Flora* 190: 265–272.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Phyton, Annales Rei Botanicae, Horn](#)

Jahr/Year: 2002

Band/Volume: [42_2](#)

Autor(en)/Author(s): Gonzalez-Rodriguez A. M., Martin-Olivera A.,
Morales Domingo, Jimenez Marisol S.

Artikel/Article: [Gas Exchange Characteristics of Chamaecytisus proliferus subsp. proliferus var. palmensis \(Tagasate\) Measured in Spring-Summer under Field Condititons. 225-236](#)