

Palms

Journal of the International Palm Society

Vol. 61(4) Dec. 2017



THE INTERNATIONAL PALM SOCIETY, INC.

The International Palm Society

Founder: Dent Smith

The International Palm Society is a nonprofit corporation engaged in the study of palms. The society is international in scope with worldwide membership, and the formation of regional or local chapters affiliated with the international society is encouraged. Please address all inquiries regarding membership or information about the society to The International Palm Society Inc., 9300 Sandstone St., Austin, TX 78737-1135 USA, or by e-mail to info@palms.org, fax 512-607-6468.

OFFICERS:

President: Ray Hernandez, 4315 W. San Juan Street, Tampa, Florida 33629 USA, e-mail subtropicoofcancer@hotmail.com, tel. 1-813-832-3561.

Vice-Presidents: Jeff Brusseau, 1030 Heather Dr., Vista, California 92084 USA, e-mail jbrusseau@newportnational.biz, tel. 1-760-271-8003. Julie Cleaver-Malzone, Av. Paulista 66, Apto. 71, Bela Vista, São Paulo, SP, 01310-100, Brazil, e-mail cleaver.julie@gmail.com, tel. 55-11-99-601-1677.

Corresponding Secretary: Susan Hubbell, 690 Calle César González, Apt. 2402, San Juan, Puerto Rico 00918 USA, e-mail sushubbell@aol.com, tel. 1-787-765-5711.

Administrative Secretary: Larry Noblick, Montgomery Botanical Center, 11901 Old Cutler Road, Miami, Florida 33156 USA, e-mail lnob@montgomerybotanical.org, tel. 1-305-667-3800 ex 104.

Treasurer: Tom Jackson, P.O. Box 1102, Ross, California 94957 USA, e-mail jackstot@sutterhealth.org, tel. 1-415-923-3299.

Directors: 2014–2018: Cindy Adair, Puerto Rico; Faith Bishock, Florida; Julie Cleaver Malzone, Brazil; Kimberley Cyr, California; Walter Donovan, Australia; John Dransfield, United Kingdom; Len Geiger, California; Ray Hernandez, Florida; Horace Hobbs, Texas; Leland Lai, California; Francisco (Paco) Martí Navarro, Spain; Don Martin, California; Jill Menzel, Brazil; Fernando Roca, Peru; Tobias W. Spanner, Germany; David Tanswell, Australia. **2016–2020:** Robert Blenker, Florida; Jeff Brusseau, California; John DeMott, Florida; Gregg Hamann, California; Hareesh, India; Susan Hubbell, Puerto Rico; Tom Jackson, California; Michael Merritt, Hawaii; Romeo Montalvo, Texas; Larry Noblick, Florida; Lazaro Priegues, Florida; Michael Smukall, Florida; Grant Stephenson, Texas; Michael Valentine, Colorado; Scott Zona, Florida.

Bookstore: www.palms.org

Chapters: See www.palms.org

Website: www.palms.org

Palms (formerly PRINCIPES)

Journal of The International Palm Society

An illustrated, peer-reviewed quarterly devoted to information about palms and published in March, June, September and December by The International Palm Society Inc., 9300 Sandstone St., Austin, TX 78737-1135 USA.

Editors: John Dransfield, Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE United Kingdom, e-mail j.dransfield@kew.org, tel. 44-20-8332-5225.

Scott Zona, 6111 Twin Lake Drive, South Miami, Florida 33143 USA, e-mail scott.zona@yahoo.com, tel. 1-305-667-5522.

Guidelines for authors are available on request from the Editors, or on-line at: www.palms.org/palms_author_guidelines.cfm

Annual membership dues are US\$55.00 for Individuals (or US\$150 for three years) and include a subscription to the Journal. Donor memberships are US\$500 per year. Individual Lifetime memberships are available for a one-time fee of US\$1000. Benefactor memberships require a one-time payment of US\$2500. Subscription price is US\$55.00 per year for libraries and institutions. Dues include mailing of the Journal by airlift service to addresses outside the USA. Dues may be paid on-line at www.palms.org.

Change of Address: Send change of address, phone number or e-mail to The International Palm Society Inc., 9300 Sandstone St., Austin, TX 78737-1135 USA, or by e-mail to info@palms.org.

Claims for Missing Issues: Claims for issues not received in the USA should be made within three months of the mailing date; claims for issues outside the USA should be made within six months of the mailing date.

Periodical postage paid at Austin, TX, USA and additional mailing offices. Postmaster: Send address changes to The International Palm Society Inc., 9300 Sandstone St., Austin, TX 78737-1135 USA.

PALMS (ISSN 1523-4495)

Mailed at Lawrence, Kansas 15 Dec. 2017
© 2017 The International Palm Society

The full text of PALMS is available on EBSCO Publishing's database.

This publication is printed on acid-free paper.

CONTENTS

161 **Palms of Samar Island, Philippines**
J.T. ADORADOR & E.S. FERNANDO

197 **An Unusual Form of *Reinhardtia latisecta* in Cultivation**
J. DRANSFIELD & J. MARCUS

Features

Palm News	160
Financial Statement	196
Photo Feature	201
Index to Volume 61	202
Statement of Ownership	206
Patrons of the IPS	207



FRONT COVER

The mottled-leaved *Pinanga copelandii* with its broadly-truncate leaflets that are glaucous underneath is one of the characteristic palms in forests over limestone on Samar Island, Philippines. See article by Adorador & Fernando., p. 161. Photo by J.T. Adorador.

BACK COVER

A palm found in lowland evergreen forests on Samar Island, Philippines, is *Calamus* sp. “*tagsa-on*” – note its short, erect, sparsely-branched inflorescence. See article by Adorador & Fernando., p. 161. Photo by J.T. Adorador.



The colorful infructescence of the “compact form” of *Reinhardtia latisecta*, are held above the foliage. Here, the fruits are not yet mature; they will turn black when ripe. See article by Dransfield and Marcus p. 197. Photo by J. Dransfield.

PALM NEWS

The major news stories as this issue of PALMS goes to press concern the damage wrought to palm collections by Hurricanes Harvey, Irma and Maria. The areas affected by these storms are prime areas for growing palms in the Caribbean and the continental USA, and many of the public and private gardens in these regions were hard-hit by the storms.

Mercer Botanical Gardens (Humble, TX) does not have a large palm collection, but the entire garden, including its Tropical Garden, where the palms were prominent, was so damaged by Harvey that it is closed until further notice.

In Florida, Hurricane Irma was a strong, slow-moving storm that battered gardens, some of which IPS members visited during the 2014 Biennial. Harry P. Leu Gardens (Orlando), Fairchild Tropical Botanic Garden (Miami), Key West Botanical Garden (Key West), Montgomery Botanical Center (Miami), Naples Botanical Garden (Naples), National Tropical Botanical Garden's The Kampong (Coconut Grove), and Pinecrest Gardens (Pinecrest) all report significant damage to collections and tree canopies. Casual observation suggests that palms were often the least damaged directly by strong winds. Leaves were often twisted and torn, but the palms usually survived. Damage to palm collections most often came from falling trees or tree branches. Understory palms that were not buried under debris often suffered sunburn and desiccation in the days after the storm following the loss of the tree canopy. A few palm species seem highly susceptible to wind damage: *Caryota mitis* stems toppled like bowling pins and *Hyphaene* species lost entire branches. Readers may remember the interesting study, published in the pages of this journal by P. Griffith et al. (57: 115-122. 2013), that examined the correlation between the storm resistance of palms and storm frequency in their natural habitat. The authors concluded that palms native to Florida and the Caribbean are more resistant to wind damage than palms from South America.

Hurricane Irma also caused some damage to gardens in the Caribbean, but that damage paled when compared to the havoc wrought by Maria. As humanitarian efforts are rightly focused on saving lives and restoring basic services to the islands, we have only limited information on the palm collections. We have no word from the Dominica Botanical Gardens in the capital of Roseau. Past visitors may remember the large group of *Corypha umbraculifera*, many of which flowered over the years. We are hopeful that the garden survived. Dominica suffered a direct hit by the eye of the storm, so the recovery for the island and its people will be long and difficult. The St. George Village Botanical Garden (St. Croix, U.S. Virgin Islands) reports extensive damage to its collections, although garden buildings were undamaged. Video posted on-line of the Jardín Botánico of the Universidad de Puerto Rico – Río Piedras shows catastrophic damage to trees and shrubs. Many palms, however, can be seen in the video, bruised but unbowed. We expect that exotic palms were more likely to be damaged by high winds than palms native to the islands.

Other botanical gardens and the personal palm collections of residents surely experienced some damage from these three storms. We remain hopeful that as the region recovers, so too will the palm collections.

Palms of Samar Island, Philippines

JIRO T. ADORADOR
*Plant Biology Division,
Institute of Biological Sciences,
College of Arts and Sciences,
The University of the Philippines -
Los Baños,
College, 4031 Laguna, Philippines
jtadorador@up.edu.ph*

AND

EDWINO S. FERNANDO
*Department of Forest Biological
Sciences,
College of Forestry and Natural
Resources,
The University of the Philippines -
Los Baños,
College, 4031 Laguna, Philippines*


Surveys of palms (Arecaceae) on Samar Island in the Philippines were conducted in view of the island's diverse but extremely threatened forest ecosystems, its old conglomerate geology, its significant biogeography and its poor palm collection history. Inventories of palms were carried out in selected sites between 2013 and 2016. The inventories yielded 50 species of which about 70% are Philippine endemics. Twelve are new island-records, and there are 11 probable novel taxa (five in *Calamus*, and one each for *Areca*, *Oncosperma*, *Heterospatha*, *Orania*, *Pinanga* and the Ptychospermatinae) while one is a recollection. Overall, the palm flora of Samar reflects its biogeographic affinity and diverse habitat types. Notes on the observed morphological variations of most species will be utilized in the future taxonomic re-evaluation of some Philippine palm species and their complexes. The wealth of information generated in this study will ultimately be used for the conservation of palms and their habitats on the island.

Samar Island is the third largest island in the Philippines archipelago with an approximate land area of 13,107 square kilometers, extending between 10°45' and 12°45' N latitude and between 124°15' and 125°45' E longitude (PhilGIS 2016). The island's regional geomorphology includes (1) the volcanic hills

and mountains at the northwest tip, (2) the heavily dissected plateau in the north, (3) the limestone area that rims the central portion, (4) the basement area in the central and southern parts, (5) the rolling hills in the southwest and (6) the Guiuan peninsula (Travaglia et al. 1978). Based on the modified

Corona’s climate classification, the island is divided into two regions. The northeastern part manifests the Type II climate, which has no dry season and has a very pronounced rain period particularly during December and January, while the southeastern portion

1. Various sites visited and surveyed from 2013–2016. The site numbers correspond with the enumeration in Appendix 1. Geologically, limestone areas approximate the Oligocene-Miocene, whereas ultramafic sites fit in the Cretaceous-Palaeogene. The inset map shows the present-day Philippines archipelago with 100 meters below sea level bathymetry line (gray shade) where several Pleistocene Aggregated Island Complexes (PAIC) can be reconstructed.





2 (top). The vast unexplored intact lowland forests viewed atop Mt. Huraw, San Jose de Buan, Samar.

3 (bottom). The contiguous lowland forests over rugged terrain towards the central mountains viewed atop a karst formation in Paranas, Samar. Photos by J.T. Adorador.

manifests the Type IV climate, which has rainfall more or less evenly distributed throughout the year.

Samar is part of the Greater Mindanao biological region, which corresponds to the Mindanao Pleistocene Aggregate Island



4 (top). Secondary forests adjacent to local gold mining excavations in Guinmayuhan, Balangiga, E. Samar.
5 (bottom). A low karst forest along Sohoton River, Basey, Samar where *Heterospathe intermedia* abounds.
Photos by J.T. Adorador.

Complex (PAIC) that also includes Leyte, Bohol, Mindanao, Dinagat and Siargao (Heaney 1986) (see inset map in Fig. 1). Together with the eastern portion of Luzon (Sierra Madre range and Bicol peninsula), this

PAIC formed a continuous island arc into Halmahera and New Guinea in the late Oligocene, about 25 Ma (Heaney 1986, Brown & Diesmos 2009, Vallejo 2011). This island chain has been used by plants and animals to

migrate north and southwards (Lam 1945, Hall 2001, Heads 2003, Vallejo 2011). The island is one of the centers of botanical diversity in the country and in the Malesian Region (Madulid 2000). According to him there are at least six forest formations found in the island, which include (following the scheme of Fernando et al. 2008) lowland evergreen rainforest, lower montane forest, forest over limestone, forest over ultramafic rocks, mangrove forest and beach forest. Additionally, peat swamp forest was hypothesized to occur in Dolongan, Basey (Davies 2006), whereas freshwater swamp forests were observed by us along the seaward-flooded plains of Balangiga-Giporlos-Quinapondan. These two poorly documented vegetation types in the island were found to be degraded and converted into cultivated areas or fishponds.

The contiguous forests found in central portions of the island inevitably provide its inhabitants with abundant water supply and renewable forest products; likewise, it offers sanctuary to its rich flora and fauna. It was declared a natural park (Samar Island Natural Park or SINP) in 2003 through Presidential

Proclamation No. 422. The park comprises 333,300 hectares of protected area and 125,400 hectares of buffer zone – the largest terrestrial protected area in the Philippines (UNDP-GEF 2014). Despite this effort, the island's environment has been extensively degraded by logging and forest clearing for agriculture in the past 70 years (UNDP-GEF 2014). Likewise, the proliferation of small-scale illegal logging significantly contributes to forest destruction. Coal and chromite mining activities also pose a major threat to the island's biodiversity, while other threats include unregulated quarrying of limestone, charcoal-making, over-harvesting of non-timber forest products (including rattans), pollution by industries and invasion of alien species (SEARCA 2004).

Although Samar Island is considered among the under-collected places for palms in the country (Fernando 1990a, c), an extensive review of literature revealed about 35 species of indigenous palms listed for the island (Beccari 1919a, b, Pelsner et al. 2011-onwards, Fernando 1990a, b, c, Madulid 2000, Gruezo & Harries 1984, SEARCA 2004, Baja-Lapis

6. (left). *Calamus urdanetanus* – recollected and rediscovered in Arizona, Hinabangan, Samar since its description in 1919. 7 (right). *Calamus daemonoropoides* – note its densely bristly leaflets, sheath armed with dense partial whorls of spines, and unarmed sharply bulging knee. Photos by J.T. Adorador.





8 (left). Habit of *Calamus ochrolepis* – note the characteristic elevated whorls of spines in the leaf sheath.

9 (right). Dense infructescence of *Plectocomia elongata* var. *philippinensis*. Photos by J.T. Adorador.

2010). Definitely, more taxa remain unaccounted since no taxonomic studies focusing on palms have been undertaken on the island. Preliminary site visits and subsequent research and consultations, suggest several probable new species of *Pinanga* and *Calamus* and a possible new genus in the Ptychospermatinae.

SINP's diverse and conserved but extremely threatened forest ecosystems, its old conglomerate geology, its significant biogeography and its poor palm collection history altogether suggest the timeliness of this endeavor and the likelihood of discovering novel palm taxa. A robust taxonomic treatment of palms and knowledge of their ecology essential for conservation were the goals for this study.

Methods

Identification of potential sites for field surveys was done through Geographic Information System (GIS)-based processing techniques through the aid of ASTER-DEM and various thematic maps (e.g. forest cover, geology, administrative boundaries, and SINP boundary) (Fig. 1). All the eight forest formations (Fernando et al. 2008) that occur

in accessible and secured sites of the island were visited (Appendix 1, Figs. 2–5). The necessary gratuitous permit to carry out such surveys within SINP was granted by the Department and Natural Resources-Region 8 (DENR-8) on July 9, 2015.

Complete palm specimens were collected (following Dransfield 1986), fertile if available, on several sites. Likewise, silica-dried leaf tissue samples of most species were collected for future molecular studies. Oven-drying and mounting were executed in the Department of Forest Biological Sciences, College of Forestry and Natural Resources, University of the Philippines Los Baños (UPLB). The first set of palm specimens are deposited at the Forestry Herbarium (LBC), UPLB Museum of Natural History, while the other sets will be sent to the Philippine National Herbarium (PNH) and the Herbarium of the Royal Botanic Gardens, Kew (K).

All available and relevant palm taxonomic literatures were reviewed. Palm specimens were carefully examined at the following herbaria: (1) LBC [Forestry Herbarium] and (2) CAHUP [College of Agriculture Herbarium–UP], both managed by the UPLB Museum of Natural



10. The habit of *Saribus rotundifolius* – note its highly-branched infructescence. Photo by J.T. Adorador.



11. Habit of the stilt-rooted *Areca costulata* – note the closely-set multifold leaflets and yellowish crownshaft. Photo by J.T. Adorador.




12. The diminutive *Pinanga* sp. "*tagibungang-liitan*" – note its short coriaceous leaves and distichously-set fruits. Photo by J.T. Adorador.

History and (3) EBL [ERDB Botanical Laboratory and Herbarium (formerly known as FORI)] under the ERDB-DENR. Appropriate parts were measured in herbarium sheets and of specimens in situ.

The morphological species concept of different Philippine-occurring palm lineages provided through the works of Beccari (1919a, b), Baker (2015), Baker et al. (1999), Dransfield (1980,

1984, 1997), Dransfield et al. (2008), Fernando (1988a, b, 1990b, 2014), Baja-Lapis (1989), Madulid (1981), Lewis (2002), Hahn (unpublished), Heatubun et al. (2012, 2014), Dowe (2009), Bacon and Baker (2011), and Keim and Dransfield (2012), were circumspectly analyzed and utilized in species delimitation. The syntheses of data gathered from numerous taxonomic articles, herbarium material and field observations generated the



13. The spiny *Oncosperma* sp. "*anibong-kinis*" – note the glabrous prophyll and few-branched Infructescence. Photo by J.T. Adorador.



14 (left). The terminal portion of *Calamus siphonospathus* var. *dransfieldii* – note the dense brown sheath spines and indistinct ocrea. 15 (right). Habit of flowering *Calamus moseleyanus* – note its broad-ellipsoid dull green leaflets and diffusely yet stiff paniculate inflorescence. Photos by J.T. Adorador.

functional broad morphological species concept used herein to delimit the indigenous palms of the island.

Results and Discussions

The Palm Flora of Samar Island

The series of surveys across the island revealed a total of 50 palm species distributed in 16 genera together comprising 26 calamoids, 16 arecoids, seven coryphoids and the sole nypoid (Appendix 2). *Calamus* is the most species-rich genus on the island with 23 species, distantly followed by *Pinanga* and *Areca* (each with 4 species), then *Heterospatha* (3 species). Twenty-seven species were previously recorded from the island, while 12 are new records and 11 are probable new taxa. Overall, it represents a quarter (39/139) of the currently known palm species (not including varieties and subspecies) in the country. Meanwhile, Philippine endemism is about 70% (34/50), which is locally similar to the palm flora of the 16-ha Palanan Forest Dynamics Plot (Yap 2010) and regionally consistent with the total vascular flora of the Philippine archipelago (Myers et al.

2010). Among the recognized taxa, only one (*Pinanga samarana*) is reported endemic to the island, while most of the probable new species are believed to be similarly restricted. Despite the island's relative smaller geographic area, species richness is comparable with that of Sulawesi Java and Moluccas with 62, 46 and 40 species, respectively (Baker & Couvreur 2012). The transitory biogeographic setting of the island could partially explain the species compositional trend found there.

It is notable that *Calamus urdanetanus* (Fig. 6), previously known only from its type locality, Mt. Cabadbaran (Mt. Urdaneta) in Agusan del Norte, Mindanao island, was recollected for the first time since its description by Beccari in 1919. For the new island records, seven species were already predicted to occur since these palms were then known to be distributed across Greater Mindanao (except Samar) and Greater Luzon island complexes (e.g. *Calamus daemonoropoides*, *C. ochrolepis*, *Plectocomia elongata* var. *philippinensis* (Figs. 7–9). Meanwhile, two species are recorded for the first time outside Mindanao, *C. multinervis* and



16. The infructescence of *Pinanga insignis* – note its robust peduncle and distichously-set shortly-beaked fruits. Photo by J.T. Adorador.



17. Habit of fruiting *Orania decipiens* – note highly branched inflorescence axes. Photo by J.T. Adorador.

the recollected rattan, whereas *Saribus* cf. *merrillii* and *Livistona saribus* are now found outside Luzon (Fernando et al. 2008). This

conspectus of updated geographic distribution resolves the palm disjunction of Luzon-Mindanao pattern (Fernando 1990a).



18. *Areca* sp. "*mudulan*" showing its slender habit and immature Infructescence. Photo by J.T. Adorador.

Eleven are believed to be novel taxa but still warrant further investigation through study of all available herbarium material and/or application of molecular techniques: five in *Calamus*, and one each for *Areca*, *Oncosperma*, *Heterospatha*, *Orania*, *Pinanga* and the Ptychospermatinae. The species in the last four

genera are morphologically distinctive and are currently being formally described, while those in the first three require more in-depth studies. The discovery of these new palm taxa is a function of new collections from a poorly-known area, correspondence with several palm experts and recent taxonomic updates.

The morphological characters that were found to be very variable and diverse in herbarium specimens and in situ include leaflet venation, the shape of the fruits and leaf sheath indumentum for the arecoids, while such characters are sheath armature, inflorescence length and number of fruits scales for the calamoids. On the contrary, characters that were found to be consistent across different habitats and developmental stages for all palms comprise general stem form, leaflet bristles and covering, and overall inflorescence architecture.

Habitat and Ecology

There remain vast tracts of dense forests scattered throughout the island, especially those growing over limestone areas and in the central portions where communist rebels inexorably make camps. Among the island's forest types, only three were surveyed comprehensively on selected sites, which include the lowland evergreen, over limestone, and over ultramafic rocks. Several palms were noted to be widespread amidst the perceived ecological differences between the three vegetation types. Such rattans include *Calamus mollis*, *C. ochrolepis*, *C. ornatus* and *Plectocomia elongata* var. *philippinensis*; on the other hand,

C. aidae and *C. merrillii* were rarely found over limestone areas. Meanwhile, the widespread erect palms include *Saribus rotundifolius* (Fig. 10), *Areca caliso*, *A. costulata* (Fig. 11), *Pinanga samarana*, *P. copelandii*, *P. sp. "tagibungang-liitan,"* *Caryota cumingii* and *C. rumphiana*; *Oncosperma* sp. "*anibong-kinis*" was occasionally spotted over limestone-derived soils.

The widespread though sporadic occurrence of a single new species of *Pinanga* (Fig. 12) across different forest types is rather surprising. It was first encountered near the summit of Mt. Huraw (800 m elevation) then atop tower karsts of Tenani (400 m), ridges of Arizona (300 m) and Vigan (200 m). Its several morphotypes were then regarded to be distinctive species but were dismissed to be just ecotypic variations. Generally, this taxon could be characterized by its clustering diminutive and very slender habit, the typically pluricostulate coriaceous leaflets, and short inflorescence with few reflexed rachillae bearing two rows of ovoid-ellipsoidal fruits. This novel taxon generally prefers ridges where vegetation is intact and light is abundant.

The probable new species of *Oncosperma* (Fig. 13) shares morphological affinity with the Luzon-restricted *O. gracilipes* and the

19. Rattan sapling diversity in the lower slopes of Mt. Huraw – in the foreground (left to right) are *Plectocomia elongata* var. *philippinensis*, *Calamus mollis* and *C. filispadix*, while in the background are *C. aidae* and *C. daemonoropoides*. Photo by J.T. Adorador.



Mindanao-widespread *O. horridum* and could represent a distinctive race of hybrid origin. This clustering palm is characterized by its median leaflets abaxially bristly on the distal portions (first report for the genus), unarmed prophyll and fruits developing only on the central portions of the rachillae. It grows well in sheltered valleys but becomes spindly towards the ridges.

Lowland evergreen forests

The lowland evergreen forests occupy the greatest vegetation coverage within the park, which gradually merge with other forest types hence creating ecotonal boundaries. Those species observed to be restricted to this vegetation include the rarities (e.g. *Calamus pedicellaris*, *C. siphonospathus* var. *dransfieldii* (Fig. 14) and *Saribus* cf. *merrillii*) and a probable new *Calamus* (*C. sp. "tagsa-on"*). Likewise, several species were shared with forests over ultramafic areas, which comprise *Calamus elmerianus*, *C. filispadix*, *C. megaphyllus*, *C. moseleyanus* (Fig. 15), *C. vidalianus*, *Pinanga insignis* (Fig. 16) and *Orania decipiens* (Fig. 17). Such habitat sharing also occurs in the suspected new *Areca* sp. "*mudulan*" and *Calamus* sp. "*malubagacay-bilog*."

The inferred new *Areca* (Fig. 18) was observed to be sympatric (in some areas) with the limestone-preferring *A. caliso* and the eurytopic *A. costulata*. It is more similar to the former but is distinctive in its much more slender habit, shorter leaves with generally 2-costulate leaflets and its much shorter inflorescence with the uppermost rachillae still bearing at least one floral triad on its basal portions. Such variations in the vegetative traits could be dictated by the differences of the substrate but those of the floral characteristics need to be re-evaluated (along with other Philippine *Areca*) – if this really represents a new species.

Meanwhile, the other inferred new taxa in *Calamus* approach the "*C. mindorensis* group" (including *C. mindorensis*, *C. moseleyanus*, *C. multinervis* and *C. reyesianus*), which local people generally refer to as *malubagacay* or *lokuan*. Beccari (1919a) recognized this artificial group (except *C. reyesianus*) but did not provide a conspectus of it. Nevertheless, *C. sp. "tagsa-on"* (Back Cover) is distinctive in its glabrous coriaceous leaflets typically in pairs and shorter erect inflorescence. On the other hand, *C. sp. "malubagacay-bilog"* though only non-fertile

20 (left). *Pinanga samarana* – notice the peeled portion of the crownshaft, evidence of palm-heart extraction by monkeys. 21 (right). *Calamus filispadix* with a bird's nest hanging from its flagellum. Photos by J.T. Adorador.





22. *Calamus aidae* with its interwoven basal leaflets made as a nest by a forest rodent. Photo by J.T. Adorador.

plants were seen, is rather distinctive in its very chartaceous broad-lanceolate leaflets and fine, continuous tertiary veins. These two rattans together with the “*C. mindorensis* group” need taxonomic reappraisal since several morphological heterogeneities were observed in the field, such as leaflet groupings upon development or exposure to light and persistence and/or absence of sheath armature on mature individuals.

Among the intact lowland forests visited, the slopes of Mt. Huraw in San Jose de Buan offered the highest palm diversity with 29 species (17 rattans and 12 erect palms). Along the forest edges, there could be as much as five different species of juvenile rattans in a 3m × 3m plot (Fig. 19) – not including the seedlings of *Oncosperma* and *Areca*. The uniqueness of palm composition in this locality could be due to the dense canopy, hence very little light reaching the understory. *Pinanga samarana* was uncommon along the trails.

Meanwhile, the secondary forests of Arizona, Hinabangan and Ulot, Paranas possessed an almost similar palm floristic composition to Mt. Huraw but differed in the abundance of *Pinanga samarana*, *Calamus aidae*, and *C. filispadix*. Thus, the proliferation of these palms could be due to the increased light intensity

penetrating the forest floor of these relatively disturbed forests. Interestingly, traces of palm heart extraction by monkeys were observed in *P. samarana* (Fig. 20) and the two rattans mentioned were recorded to provide suitable shelter for forest animals (Figs. 21 & 22).

Forests over limestone

The forests over limestone are arguably the most intact vegetation type on Samar Island. This is due to the difficulty of penetrating the thickets of its abruptly variable terrain; thus, illegal logging is simply not profitable. It is locally known as the *kaigangan* with karst formations, which reach up to 400 m elevation. Across the sites surveyed, *Heterospathe intermedia* (Fig. 23), *Areca caliso*, *Pinanga copelandii* (Front Cover), *Calamus discolor* (Fig. 24), *C. multinervis* (Fig. 25) and *C. ochrolepis*, are the most characteristic species within the valleys and mid-slopes of this forest formation. The palm floristic gradient then changes towards the peaks of karst towers. There abounds *Pinanga samarana*, morphologically different from the lowland counterparts in having more slender stems, uncostulate leaflets and more reflexed rachillae with obovoid fruits. However, all these trait shifts could be attributed to environmental differences of their habitats, especially light and soil conditions.



23. The habit of *Heterospathe intermedia*. Photo by J.T. Adorador.



24 (left). *Calamus discolor* – note its dense brown spines and petiolate leaves with leaflets glaucous abaxially. 24 (right). Habit of *Calamus multinervis* – note its brown sheath indumentum and plaited leaflets grouped in pairs which are observed in individuals under dense forest cover. Photos by J.T. Adorador.

Interestingly, four new taxa were discovered atop this forest type. The novel rattans include two slender clustering *Calamus*, *pudlos-uban* (Fig. 26) and *pudlos-liitan* (Fig. 27) of San Rafael, Taft (Eastern Samar) and Tenani, Paranas (Samar), respectively. The former differs from typical *C. discolor* in having less spiny, glaucous green sheaths and epetiolate leaves with less bristly leaflets. Meanwhile, the latter species is apparently allied to *C. microsphaerion* but ultimately differs in the unarmed rachis and concave short-lanceolate apiculate leaflets, which are grouped in pairs. This rattan was recorded to produce shoots on cut stems, which could be the reason for its abundance on the summit. Aside from these traits, their inflorescences details ultimately segregate them from their respective analogous counterparts.

The new species of slender *Orania* (Fig. 28) atop the calcareous mountain is rather surprising since its counterparts thrive in areas where there is much thicker soil. Its closest match is *O. littoralis* of Papua New Guinea using the artificial key by Keim and Dransfield (2011). However, the inflorescence morpho-

logy is the most unusual among all known *Orania* species, in developing an all-staminate inflorescence (without triads along the rachillae) and a bisexual type (with floral triads). Moreover, its fruit traits (color, mesocarp thickness and embryo shape) are recorded to be different with those of *O. decipiens* when compared in fresh state – which could be used in re-annotating useful characters for the Philippine species.

Perhaps, the most thrilling discovery of the entire palm flora of the island would be the member of Ptychospermatinae, known locally as *amuring* (Fig. 29). It is morphologically embedded within the subtribe in its praemorse leaflet apices and bullet-shaped buds of staminate flowers. Moreover, after consultations with several experts on the subtribe (S. Zona, W. Baker and C. Heatubun), review of recent literature (Alapetite et al. 2014, Heatubun et al. 2012) and herbarium studies, we believe this taxon merits a unique generic placement. It deviates from other members in details of its inflorescence and fruit structure. It is restricted to the summits of karst formations in the central part of the island.



26 (left). The slender *Calamus* sp. “*pudlos-uban*” – note its epetiolate discolorous leaves and less spiny sheath.
 27 (right). *Calamus* sp. “*pudlos-liitan*” with its short lanceolate leaflets grouped in pairs. Photos by J.T. Adorador.

The observed populations of *amuring*, which comprise several fruiting individuals, few juveniles but numerous seedlings generally suggest its adaptability but slow growth and establishment to the unique habitat. Fruit doves (*Ptilonopus* spp.) were observed to feed on the thin but fleshy and bland-tasting mesocarps of this palm.

Forests over ultramafic rocks

In contrast, the remaining vegetation over ultramafic areas on southeast portions of the island were generally disturbed. Aside from the mining activities and illegal timber extraction of nearby communities, such areas were devastated by one of the history's strongest typhoons (Haiyan) in November 3, 2013 and most of them subsequently succumbed to wild-(or man-induced) fires in the summer of 2014. The forest surveys were carried-out in the most accessible intact forests outside mining concession areas due to the difficulty of acquiring permits from the companies.

Nevertheless, the fieldwork showed the general palm floristic make-up across the three metal-rich sites, which are dominated by the widespread *Pinanga samarana*, *Areca* spp., *Calamus aidae* and *Plectocomia elongata* var.

philippinensis. It is within these forest types that the extent of morphological variation of *P. samarana* was observed and verified. The encountered populations from these sites differ from the typical lowland and limestone morphotypes in having 3-pluricostulate leaflets and twisted rachillae with globose-ovoid fruits. Meanwhile, *Orania decipiens*, *Calamus elmerianus*, *C. megaphyllus* (Fig. 30), *C. vidalianus* (Fig. 31) and *C. sp. “tagda”* were locally abundant in Guinmayuhan, Balangiga, whereas the new slender *Pinanga*, *Calamus ochrolepis*, *C. sp. “malubagacay-bilog”* and *Heterospathe* sp. “*saguisi-mina*” were very abundant in Vigan, Gen. MacArthur.

The sites' unique edaphic conditions characterized by heavy metal concentration and low litter-nutrient inputs paved way for the niche specialization of certain plants such as *Nepenthes* spp. This habitat differentiation might also explain occurrence of the two new ultramafic-restricted palms, *Calamus* sp. “*tagda*” (Fig. 32) and *Heterospathe* sp. “*saguisi-mina*” (Fig. 33). The novel flagellate rattan is characteristic in its long-petiolate leaves and its sheath armature, reminiscent of cirrate *C. loherianus* in Luzon. It is unique among Philippine species in having the proximal joint of the flagella ornamented with a crown of



28. *Orania* sp. "*banga-igang*" with its very slender habit. Photo by J.T. Adorador.



29. The habit of *Ptychospermatinae* or *amuring* with its broadly-truncate, apically praemorse leaflets and whitish inflorescence axes. Photo by J.T. Adorador.

spines. Fertile collections would make such comparisons more complete. Meanwhile, the latter slender clustering erect palm has similarities with wider-ranging *H. philippinensis* but is different in the shorter leaf outline, thus having fewer leaflets and purplish inflorescence axes. Across different sites, this palm likewise exhibited noteworthy variation in inflorescence branching order and number of its peduncular bracts and leaflets. These two palms commonly prefer exposed ridges where the canopy is more open, but they are locally more abundant on reddish soil.

Other habitats (freshwater swamp, beach, and mangrove)

The freshwater swamps along seaward coastal portions of the island were mainly converted into rice fields or coconut plantations. Remnants of natural vegetation in these waterlogged sites mainly comprise discontinuous stands of *Terminalia*, *Anthocephalus* and *Pandanus*. The palms sporadically thriving here include *Metroxylon sagu* (Fig. 34) and *Livistona saribus* (Fig. 35). The latter fan palm is characterized by its heavily armed petiole, drooping leaf apices

and bluish fruits, while the former clustering palm has erect dark-green leaves and trunks die after a massive apical flowering. Sago palm stands were not communally maintained and utilized since the fragmented clusters seemed to be only established by chance. Meanwhile, *Nypa fruticans* typically borders rice fields where they border brackish water zones towards the mangroves.

Beach forests of appreciable extent can only be encountered in uninhabited coasts or smaller islands where *Heterospathe elata*, *Calamus discolor*, *C. usitatus* and *Caryota cumingii* flourish. These narrow coastal vegetation strips are separated from the central lowland forests by large expanses of cultivated lands or abandoned clearings where few plants of *Arenga pinnata* can be found.

Economic use of some palms

Among the rattans, at least five robust species (*Calamus aidae*, *C. merrillii*, *C. moseleyanus*, *C. multinervis* and *C. ornatus*) were recorded as used in furniture-making industry. Meanwhile, those slender-cane taxa used in handicrafts (such as weaving material and cordage) include

30 (left). Sapling of *Calamus megaphyllus* with its broad oblanceolate leaflets. 31 (right). Habit of clustering *Calamus vidalianus* with its irregularly arranged leaflets. Photos by J.T. Adorador.





32. The flagellate *Calamus* sp. "*tagda*" – note the dense brown spines of the sheath and short lamina outline. Photo by J.T. Adorador.



33. *Heterospathe* sp. "*saguisi-mina*" – note its several undivided leaves and purplish inflorescence axes. Photo by J.T. Adorador.



34 (top). Young stand of the clustering *Metroxylon sagu*. 35 (bottom). Remnant population of *Livistona saribus* in Basey, Samar with several cultivated *Cocos nucifera* in the background. Photos by J.T. Adorador.

C. daemonoropoides, *C. discolor*, *C. filispadix*, *C. megaphyllus*, *C. mollis*, *C. sp. "tagda"* and *C. sp. "pudlos-liitan."* Some species provide food for inhabitants such as *C. ornatus* for its sour fruits and *C. aidae* for its bitter palm heart.

As for the erect palms, the leaves of several species (*Corypha utan*, *Livistona saribus* and *Saribus rotundifolius*, *Metroxylon sagu*, *Nypa fruticans* and *Cocos nucifera*) are used for thatching and roofing materials. Aside from

the conventional uses of coconut, its fruit and fermented sap are commercially utilized for copra (coco-meat) and *lambanog*, respectively. Local people seldom utilize the durable wood of *Orania decipiens* as house posts. On a lesser scale, the three forest *Areca* species were observed to serve as substitutes for *A. catechu* for their traditional *mamâ*. Likewise, *Caryota rumphiana* and *Pinanga insignis* are known for their palatable palm heart, while those of *P. samarana* are also said to be edible though bitter to taste.

Threats to their habitat

The never-ending small-scale timber poaching and slash-and-burn (*kaingin*) practices always pose significant threats to the island's rich forest biodiversity. Several widespread palms were recorded to thrive in heavily disturbed lowland forests (such as *Pinanga samarana* and *Plectocomia elongata* var. *philippinesis*), but this does not correspond to their general ecological tolerance as a lineage. These palms are natural inhabitants of intact forests wherein their

population dynamics respond to natural disturbances such as canopy gaps and soil erosion. Thus, prolonged exposure to such man-induced environmental stresses could result to the decrease of palm species composition or even to the extirpation of their respective populations. Such is the case for the two annihilated populations of the new diminutive *Pinanga* in the carbon-mined wasteland of Arizona, Hinabangan (Samar), wherein *kaingin* is frequently practiced for cassava cultivation (Fig. 36). Moreover, the 1980s agricultural conversion of the entire Dolongan peatlands at Basey (Samar) could have inflicted the same drastic trend to its palm flora before we even knew what existed there.

Over-harvesting of the rattan canes for the furniture industry in past decades has resulted in the absence of mature rattans but abundance of juvenile plants in almost every forest type visited. In Mt. Huraw, Arizona, Marabut, and Guinmayuhan, long canes of

36 (left). The extirpated cluster of *Pinanga* sp. "*tagibungang-liitan*" in Arizona, Hinabangan, Samar – note the rolled and scorched leaves which could be due to extreme heat following *kaingin* practice. 37 (right). One of the numerous saplings of *Calamus merrillii* – a remnant of the over-collection of mature canes in Mt. Huraw, San Jose de Buan. Photos by J.T. Adorador.



Calamus merrillii (Fig. 37) and *C. aidae* were not seen since local people had eagerly harvested these prime species. The current economic-paucity of the said industry could pave the way to the re-establishment of the natural populations, which must be sustainably managed in the future.

Several sites on the entire island are declared mineral reservations (e.g. laterite, bauxite, gold) – just waiting for any legal propositions for these ores to be extracted. In several sites, fuel-wood gathering and unproductive farming practices incessantly pressurize the spoiled vegetation. These destructive activities likely jeopardize the habitats of ultramafic-restricted palms. Naturally, the rapidly transformed environment suppresses their respective ecological niches and consequent reproductive strategies. The surveyed burnt forests of the previous laterite-exhumed concession area in Carapdapan, Salcedo (Eastern Samar) revealed the sparse occurrence of *Calamus* sp. “*tagda*” and the disappearance of the new *Heterospatha*.

On the other hand, the palm species found on the inaccessible interior rugged terrains of forests over limestone (e.g. Ptychospermatinae, *Orania* sp. “*banga-igang*,” *Calamus* sp. “*pudlos-liitan*”) are projected to be among the least threatened by human extracting activities. However, natural disasters such as typhoons and prolonged droughts could bring the palm flora to its ecological tipping point in this limiting environment.

Summary and Conclusions

A survey of palms on Samar Island has revealed that its palm flora is very rich and particularly diverse and unusual, comprising 50 species with 70% Philippine endemism, 12 new records and 11 probable new taxa. Across different vegetation types and degree of disturbances, several palm species were observed to present morphological plasticity. Scrutiny of taxonomic literature, herbarium material and field observations required the application a broad morphological species concept on the island's palms. When taken in a broader context, this realization inevitably calls for a thorough review of several Philippine palm species and their complexes. Despite most of the island's remaining forests being declared “protected,” human-induced activities constantly exert significant environmental pressures to the palms found there. The results of these surveys will ultimately be used towards the conservation of palms and their habitats on the island.

Recommendations

Population surveys of the economically important rattans subject to commercial extraction should be carried out so that sustainable management can be proposed. Thorough habitat surveys of the novel palms and knowledge on their ecology and phenology are also recommended.

The interior swamps of Bermudahan in Hinabangan, Samar, the highlands of the central portions of the island (Mts. Capotoan and Concord), and the volcanic soils in the northwestern part (Victoria-Calbayog) could support unique palm flora, but these sites were not included due to security reasons. Thus, if rebel tensions dissipate and funds are available, surveys in these areas could produce further botanical novelties.

Acknowledgments

The authors are indebted to Dr. William Gruezo and Dr. Nathaniel Bantayan for their sound advices throughout the conduct of the study. To several palm experts who provided assistance in several ways, Drs. William Baker, Scott Zona, Charlie Heatubun, Fred Essig, Aida Baja-Lapis, Ary Keim, William Hahn and Christine Bacon. Dr. Pieter Pelser provided access to certain publications. We thank the keepers of three herbaria namely, LBC, CAHUP and EBL. JTA would also like to express gratitude to the DOST-ASTHRDP-NSC scholarship and the International Palm Society Endowment Fund for providing scholarship and research grants. The SINP-PAMB, CENRO and PENRO-Catbalogan, DENR-R8 are thanked for providing assistance from processing the gratuitous and transport permits, especially to the following; Zenaida Baisa (PASu), Eires Mate, Jovelyn Grovoso, Gene Tomas (EMS-Catbalogan), Marcialito Torres (CENRO-Catbalogan), and other members of the SINP-PAMB. We thank our field guides in various localities; Titu, Michael, David, Rommel, Marlito, Marlon, Using, Leonardo, to the Mamanwa tribe, Buchoy, Larry, Gloria family and Lito. Most importantly JTA thanks his dearest girlfriend, Zhereeleen Meneses, for her assistance and companionship on and off during fieldwork.

LITERATURE CITED

- ALAPETITE, E., W.J. BAKER AND S. NADOT. 2014. Evolution of stamen number in Ptychospermatinae (Arecaceae): Insights from a new molecular phylogeny of the subtribe. *Molecular Phylogenetics and Evolution* 76: 277–240.

- BACON, C.D. AND W.J. BAKER. 2011. *Saribus* resurrected. *Palms* 55: 109–116.
- BAJA-LAPIS, A.C. 1989. An account of taxa related to *Calamus siphonospathus* complex. *Sylvatrop* 2: 91–96.
- BAJA-LAPIS, A.C. 2010. A field guide to Philippine rattans. *Asia Life Sciences Supplement* 5: 1–214.
- BAKER, W.J. 2015. A revised delimitation of the rattan genus *Calamus* (Arecaceae). *Phytotaxa* 197: 139–152.
- BAKER, W.J. AND T.L.P. COUVREUR. 2012. Biogeography and distribution patterns of Southeast Asian palms. Pp. 164–190. In D.J. GOWER ET AL. (eds.). *Biotic Evolution and Environmental Change in Southeast Asia*. Cambridge University Press. .
- BAKER, W.J., J. DRANSFIELD, M.M. HARLEY AND A. BRUNEAU. 1999. Morphology and cladistic analysis of subfamily Calamoideae (Palmae). In A.J. HENDERSON AND F. BORCHSENIUS (eds.). *Evolution, Variation and Classification of Palms*. Pp. 307–324. The New York Botanical Garden Press, Bronx, NY.
- BECCARI, O. 1919a. Palms of the Philippine Islands. *Leaflets of Philippine Botany* 8: 2997–3067.
- BECCARI, O. 1919b. The palms of the Philippine Islands. *Philippine Journal of Science (Botany)* 14: 295–362.
- BROWN, R. AND A.C. DIEMOS. 2009. Philippines, biology. Pp. 723–732. In GILLESPIE, R.G AND D.A. CLAGUE (eds). *Encyclopedia of Islands*. University of California Press, Berkeley.
- DAVIES, J. 2006. National Action Plan for Sustainable Use and Protection of Philippine Peatlands. Pp. 39. Report for IFAD-GEF Project Proposal.
- DOWE, J.L. 2009. A taxonomic account of *Livistona* R.Br. (Arecaceae). *Gardens Bulletin Singapore* 60: 185–344.
- DRANSFIELD, J. 1980. On the identity of Sika in Palawan. *Kalikasan, The Philippine Journal of Biology* 9: 43–48.
- DRANSFIELD, J. 1984. The Rattans of Sabah. *Sabah Forest Record No. 13*, Forest Department, Sabah. Pp. 182.
- DRANSFIELD, J. 1986. A guide to collecting palms. *Annals of the Missouri Botanical Garden* 73: 166–176.
- DRANSFIELD, J. 1997. The Rattans of Brunei Darrussalam. *Forestry Department Brunei Darrussalam and The Royal botanic Gardens*. Pp. 200.
- DRANSFIELD, J., N.W. UHL, C.B. ASMUSSEN, W.J. BAKER, M.M. HARLEY AND C.E. LEWIS. 2008. *Genera Palmarum – the Evolution and Classification of Palms*, ed. 2. Royal Botanic Gardens, Kew, Richmond, UK. Pp. 732.
- FERNANDO, E.S. 1988a. The mottled-leaved species of *Pinanga* in the Philippines. *Principes* 32: 165–174.
- FERNANDO, E.S. 1988b. Four new taxa of Philippine rattans (Palmae Calamoideae). *Gardens' Bulletin Singapore* 41: 49–58.
- FERNANDO, E.S. 1990a. A preliminary analysis of the palm flora of the Philippine islands. *Principes* 34: 28–45.
- FERNANDO, E.S. 1990b. The genus *Heterospathes* (Palmae: Arecoideae) in the Philippines. *Kew Bulletin* 45: 219–234.
- FERNANDO, E.S. 1990c. Diversity and conservation status of the Philippine rattans. *Rattan Information Centre Bulletin (Malaysia)* 9: 7–21.
- FERNANDO, E.S. 2007–present. A Provisional Checklist of the Palms of the Philippines. Unpublished.
- FERNANDO, E.S. 2014. Three new species in *Calamus* sect. *Podocephalus* (Arecaceae: Calamoideae) from the Philippines, Indonesia, and Papua New Guinea. *Phytotaxa* 166: 069–076.
- FERNANDO, E.S., M.H. SUH, J. LEE, AND D.K. LEE. 2008. *Forest Formations of the Philippines*. ASEAN-Korea Environmental Cooperation Unit (AKECU), Seoul National University, Korea. Pp. 232.
- GRUEZO, W.S. AND H.C. HARRIES. 1984. Self-sown, wild-type coconuts in the Philippines. *Biotropica* 16: 140–147.
- HAHN, W.J. (unpublished). Revision of *Caryota*. Pp. 195.
- HALL, R. 2001. Cenozoic reconstruction of SE Asia and the SW Pacific; changing patterns of land and sea. In METCALFE, I., J.M.B. SMITH, M. MORWOOD AND I.D. DAVIDSON (eds.), *Faunal and Floral Migrations and Evolution in SE Asia – Australia*. Pp. 35–56. AA Balkema (Swets & Zeitlinger Publishers), Lisse.
- HEADS, M.J. 2003. Ericaceae in Malesia: vicariance biogeography, terrane tectonics and ecology. *Telopea* 10: 311–449.

- HEANEY, L.R. 1986. Biogeography of mammals in SE Asia: estimates of rates of colonization, extinction, and speciation. *Biological Journal of the Linnean Society* 28: 127–165.
- HEATUBUN, C.D., J. DRANSFIELD, F. THOMAS, S. TJITROSOEDIRDOJO, J.P. MOGEA AND W.J. BAKER. 2012. A monograph of the betel but palms (*Areca*: Arecaceae) of East Malesia. *Botanical Journal of the Linnean Society* 168: 147–173.
- HEATUBUN, C.D., S. ZONA AND W.J. BAKER. 2014. Three new genera of arecoid palm (Arecaceae) from eastern Malesia. *Kew Bulletin* 69: e9525.
- KEIM, A.P. AND J. DRANSFIELD. 2012. A monograph of the genus *Orania* (Arecaceae: Oranieae). *Kew Bulletin* 67: 127–190.
- LAM, H.J. 1945. Notes on the historical phyto-geography of Celebes. *Blumea* 5: 600–640.
- LEWIS, C.E. 2002. A phylogenetic analysis of the palm subtribe Oncospermatinae (Arecaceae) based on morphological characters. *Brittonia* 54: 78–91.
- MADULID, D.A. 1981. A monograph of *Plectocomia* (Palmae: Lepidocaryoideae). *Kalikasan, Philippine Journal of Biology* 10: 1–94.
- MADULID, D.A. 2000. A review and assessment of the floristic knowledge of Samar Island: based on literature, PNH Records, and current knowledge. USAID. Pp. 16.
- MYERS, N, R.A. MITTERMEIER, C.G. MITTERMEIER, G.A. DA FONSECA AND J. KENT. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- PELSE, P.B., J.F. BARCELONA AND D.L. NICKRENT. 2011-onwards. Co's Digital Flora of the Philippines. www.philippineplants.org. Accessed on January 13, 2016.
- PHILIPPINES GIS DATA CLEARINGHOUSE (PhilGIS). 2016. <http://www.philgis.org>.
- SEAMEO REGIONAL CENTER FOR GRADUATE STUDY AND RESEARCH IN AGRICULTURE (SEARCA). 2004. Biological Resources Assessment. Samar Island Natural Park. Philippines. Final Report. United Nations Development Programme (UNDP).
- TRAVAGLIA, C., A.F. BAES AND L.M. TOMAS. 1978. Geology of Samar Island. Bureau of Soils, Department of Agriculture – United Nations Development Programme – Food and Agriculture Organization of the United Nations. Manila. Pp. 149.
- UNITED NATIONS DEVELOPMENT PROGRAMME – GLOBAL ENVIRONMENT FACILITY (UNDP-GEF). 2014. <http://www.undp.org>.
- VALLEJO, B. 2011. The Philippines in Wallacea. Pp. 27–42. *In* Telnov, D. (ed.). Biodiversity, Biogeography and Nature Conservation in Wallacea and New Guinea, Vol. I. The Entomological Society of Latvia.
- YAP, S.L. 2010. Phylogeography and Demography of Common Plant Species from the Philippine Islands. PhD Dissertation. The University of Michigan.

Appendix 1. Reconnaissance surveys and fieldwork conducted in various localities across Samar Island (2013–2016).

Appendix 1. Reconnaissance surveys and fieldwork conducted in various localities across Samar Island (2013–2016).										
LOCALITY	TOWN	PROVINCE	Dec. 2013	DATES SURVEYED						
				Feb. 2014	Mar. 2014	Jul. 2015	Oct. 2015	Dec. 2015	Feb. 2016	
Lowland evergreen rainforests										
1. Mt. Huraw	San Jose de Buan	Samar			X				X	
2. Ulot River, Tula	Paranas	Samar		X			X		X	
3. Arizona, Cansulabao	Hinabangan	Samar		X	X		X		X	
4. Km 16, Tinabanan	Marabut	Samar						X		
5. Luisita	Victoria	N. Samar	X							
Forests over limestone										
6. Tenani	Paranas	Samar	X	X	X	X	X		X	
7. San Isidro	Paranas	Samar							X	
8. Sohoton Cave	Basey	Samar				O				
9. San Rafael	Taft	E. Samar				X		X	X	
Forests over ultramafic rocks										
10. Guinmayuhan	Balangiga	E. Samar							X	
11. Carapdapan	Salcedo	E. Samar					O		X	
12. Vigan	General MacArthur	E. Samar							X	
Other forest types (mangrove, beach, and freshwater swamp forests)										
13. Coastal areas	Borongan	E. Samar	X	X					O	
14. Roadsides	Quinapondan-Giporlos	E. Samar	O					O	O	
15. Dolongan	Basey	Samar					O		O	
16. Buri Airport, Cabugawan	Catbalogan	Samar	O						X	
17. Biri Rock Formation	Allen	N. Samar					O			
18. Coastal areas	Catarman-Lavezares	N. Samar	X					O		
19. Coastal areas	Motiong	Samar	X							
20. Coastal areas	Daram & Zumarraga	Samar	X							

Legend: X = actual fieldwork was conducted, O = visual surveys only.

Legend: X = actual fieldwork was conducted, O = visual surveys only.

Appendix 2. Palm species surveyed on Samar Island and observed distribution in different forest types.

Species	Local name	Forest Types (as per Fernando et al. 2008) ¹			
		A	B	C	D
Climbing palms (rattans)					
1. <i>Calamus aidae</i> Fernando	<i>ilihian, ulasi, ulisi</i>	X	(X)	X	-
2. <i>Calamus daemonoropoides</i> Fernando*	<i>sambunotan, uai balau</i>	X	(X)	-	-
3. <i>Calamus discolor</i> Mart.	<i>uban-uban, sungo-ng-orang</i>	X	X	-	-
4. <i>Calamus elmerianus</i> Becc.*	<i>pudlos</i>	X	-	X	-
5. <i>Calamus flispadix</i> Becc.	<i>tawin, pudlos</i>	X	-	X	-
6. <i>Calamus megaphyllus</i> Becc.*	<i>abangan, malubagacay</i>	X	-	X	-
7. <i>Calamus merrillii</i> Becc.	<i>parasan, pagiti</i>	X	(X)	X	-
8. <i>Calamus mollis</i> Blanco	<i>nukot, sungo</i>	X	X	X	-
9. <i>Calamus moseleyanus</i> Becc.	<i>malubagacay</i>	X	-	X	-
10. <i>Calamus multinervis</i> Becc.*	<i>lokuan</i>	(X)	X	-	-
11. <i>Calamus ochrolepis</i> (Becc.) W.J.Baker*	<i>balala, likbong, nukot</i>	X	X	X	-
12. <i>Calamus ornatus</i> Blume	<i>kalape</i>	X	X	X	-
13. <i>Calamus pedicellaris</i> (Becc.) W.J.Baker* <i>hiyod</i>		X	-	-	-
14. <i>Calamus siphonosphathus</i> Mart. var. <i>dransfieldii</i> Baja-Lapis*	<i>pi-jai</i>	X	-	-	-
15. <i>Calamus symphysipus</i> Mart.	<i>uai-bangkau, palanog</i>	-	X	-	-
16. <i>Calamus urdanetanus</i> (Becc.)W.J.Baker*-		X	-	-	-

Appendix 2 (continued).

Species	Local name	Forest Types (as per Fernando et al. 2008) ¹			
		A	B	C	D
17. <i>Calamus usitatus</i> Blanco	<i>pudlos</i>	X	(X)	-	-
18. <i>Calamus vidalianus</i> Becc.*	<i>sambunotan, tipunan, gachan</i>	X	-	X	-
19. <i>Calamus</i> sp. " <i>tagda</i> "	<i>tagda</i>	-	-	X	-
20. <i>Calamus</i> sp. " <i>pudlos-uban</i> "	<i>pudlos-uban</i>	-	X	-	-
21. <i>Calamus</i> sp. " <i>pudlos-liitan</i> "	<i>pudlos-liitan</i>	-	X	-	-
22. <i>Calamus</i> sp. " <i>tagsa-on</i> "	<i>tagsa-on</i>	X	-	-	-
23. <i>Calamus</i> sp. " <i>malubagacay-bilog</i> "	<i>malubagacay-bilog</i>	X	-	X	-
24. <i>Korthalsia laciniosa</i> (Griff.) Mart.	<i>talubag-angan, pahan, danan</i>	X	X	-	-
25. <i>Plectocomia elongata</i> Mart. ex Blume var. <i>philippinensis</i> Madulid*	<i>uai babae</i>	X	X	X	-
Erect palms (non-rattans)					
26. <i>Areca caliso</i> Becc.	<i>bugiking</i>	X	X	X	-
27. <i>Areca catechu</i> L.	<i>bunga, mama</i>	Cultivated			
28. <i>Areca costulata</i> Becc.	<i>ajibi, muren</i>	X	X	X	-
29. <i>Areca</i> sp. " <i>mudulan</i> "	<i>mudulan</i>	X	(X)	X	-
30. <i>Arenga pinnata</i> (Wurmb) Merr.	<i>kaong</i>	Cultivated			
31. <i>Caryota cumingii</i> Lodd. ex Mart.	<i>pugahan, bagsang</i>	X	X	X	-
32. <i>Caryota rumphiana</i> Mart.	<i>pugahan, bagsang</i>	X	X	X	-

Appendix 2 (continued).					
Species	Local name	Forest Types (as per Fernando et al. 2008) ¹			
		A	B	C	D
		Cultivated^			
33. <i>Cocos nucifera</i> L.	<i>lubi</i>				
34. <i>Corypha utan</i> Lam.	<i>buri</i>	(X)	-	-	X
35. <i>Heterospathe elata</i> Scheff.	<i>saguisi</i>	-	-	-	X
36. <i>Heterospathe intermedia</i> (Becc.) Fernando	<i>marighoi</i>	X	X	-	-
37. <i>Heterospathe</i> sp. “ <i>saguisi-mina</i> ”	<i>saguisi-mina</i>	-	-	X	-
38. <i>Livistona saribus</i> (Lour.) Merr. ex A. Chev.*	-	(X)	-	-	X
39. <i>Metroxylon sagu</i> Rottb.	<i>lumbia, lubi-lubi</i>	-	-	-	X
40. <i>Nypa fruticans</i> Wurmb	<i>nipa</i>	-	-	-	X
41. <i>Oncosperma</i> sp. “ <i>anibong-kinis</i> ”	<i>anibong-kinis</i>	X	(X)	X	-
42. <i>Orania decipiens</i> Becc.	<i>banga</i>	X	-	X	-
43. <i>Orania</i> sp. “ <i>banga-igang</i> ”	<i>banga-igang</i>	-	X	-	-
44. <i>Pinanga copelandii</i> Becc.	<i>muren, muring</i>	X	X	X	-
45. <i>Pinanga insignis</i> Becc.	<i>buyungaw</i>	X	-	X	-
46. <i>Pinanga samarana</i> Becc.	<i>tagibunga, salangisog</i>	X	X	X	-
47. <i>Pinanga</i> sp. “ <i>tagibungang-liitan</i> ”	<i>tagibungang-liitan</i>	X	X	X	-
48. <i>Ptychospermatinae</i> gen. & sp. indet.	<i>amuring</i>	-	X	-	-

Appendix 2 (continued).				
Species	Local name	Forest Types (as per Fernando et al. 2008) ¹		
49. <i>Saribus</i> cf. <i>merrillii</i> (Becc.) Bacon & W.J. Baker ^{^^}	<i>anahaw, bigo-bukid</i>	A	B	D
50. <i>Saribus rotundifolius</i> (Lam) Blume	<i>anahaw</i>	X	-	-
		X	X	X
Legend:				
1 Habitats: A = lowland evergreen. B = over limestone, C = over ultramafic rocks, D = swampy (fresh or brackish)				
Taxa in bold are Philippine endemic species (which is also likely to apply to possibly new species). Those with asterisk (*) are new records for the island.				
The symbol '(X)' indicates that the species was encountered very rarely in such forest type.				
[^] The reported population of native <i>C. nucifera</i> (Gruezo & Harries 1984) in the Guianan peninsula was not visited.				
^{^^} This palm superficially resembles <i>S. merrillii</i> based on gross vegetative morphology (Dowe, pers. com.), but fertile collections are needed to verify its identity.				

INTERNATIONAL PALM SOCIETY
Comparative Statement of Activities
Year End 2015 and 2016
(unaudited)

<u>EARNED & CONTRIBUTED REVENUES</u>	TOTALS	TOTALS
	<u>12/31/2015</u>	<u>12/31/2016</u>
Earned Revenue and Gains:		
Memberships	35,089	31,217
Publications, Books, Royalties & Merchandise	7,395	3,722
Advertising	900	1,386
Prepaid Airmail Shipping	728	476
** Singapore/Sarawak Biennial Net after expenses**	182,304	(145,245)
Investment Interest, Dividend & Gains	1,263	45,681
Total Earned Revenue and Gains:	227,679	(62,763)
Contributions and Donations		
Donations > 1,000	72	1,459
Donations < 1,000	0	21,000
EARNED & CONTRIBUTED REVENUES	227,751	(40,304)
 <u>EXPENSES</u>		
Program Costs		
Printing - Palms Journal	28,236	25,559
Book Warehousing, Storage & Insurance - Austin	36,973	38,464
Website	9,256	12,469
Accounting and Office Expense	8,781	16,245
Bank and Brokerage Fees	12,433	10,721
Commissions	180	0
Paypal & Credit Card Fees	2,111	2,022
Officers and Directors Liability Insurance	3,030	3,045
Travel - Board Meeting	1,674	0
Grants	22,897	17,000
Miscellaneous/Suspended Items	61	454
TOTAL EXPENSE	125,632	125,979
 NET INCOME	 <u>102,119</u>	 <u>(166,283)</u>

An Unusual Form of *Reinhardtia latisecta* in Cultivation

JOHN DRANSFIELD

*Herbarium,
Royal Botanic Gardens, Kew,
Richmond Surrey, TW9 3AE, UK
j.dransfield@kew.org*

AND

JEFF MARCUS

*Floribunda Palms and Exotics,
Mountain View, Hawai'i, USA
info@floribundapalms.com*

1. *Reinhardtia latisecta* "compact form" at Floribunda Palms; note the short-stemmed habit and erect inflorescences emerging above the leaves, one white inflorescence at anthesis is just visible in the background.



This beautiful palm, growing in the garden of Floribunda Palms and Exotics in Mountain View, the island of Hawai'i, has perplexed visitors (Fig. 1).

The palm was received in the 1990s as a seedling from Dr. Arden Dearden of Cairns, Australia, carrying the name *Hydriastele montana*, a species from New Guinea. Dr. Dearden does not know with certainty where his original plant came from; he thinks, perhaps, seed was sold by the late De Armand Hull, possibly from Panama, and grown on by the late Queensland palm grower, Maria Boggs. Dr Dearden has lost any such plants in his own garden.

In contrast, the mystery palm has flourished in JM's collection and has been commented on by many visitors. Now that the palm is performing well, regularly flowering and fruiting, the true affinities of the plant with the New World genus *Reinhardtia* are obvious. It is a very striking, beautiful dwarf palm, with its gleaming white inflorescences at anthesis, followed by crimson infructescences, held above the leaves, with an overall aspect unlike other species of *Reinhardtia*. Yet, the stems are

2. *Reinhardtia latisecta* "compact form": details of leaves, showing the rather inconspicuous windows, and infructescence.





3. Seedlings of three different *Reinhardtia* compared: A. *R. latisepta*. B. *R. latisepta* "compact form." C. *R. gracilis*.

covered with a network of fibers formed by the disintegration of the leaf sheaths and there is no crownshaft; the leaves are irregularly divided into leaflets with unequal, almost toothed tips, and the base of the lamina displays the windows that are so much a feature of most species of *Reinhardtia* (Fig.2). The inflorescence is interfoliar and bears both staminate and pistillate flowers. Staminate flowers have approximately 29 stamens, while the pistillate flower has a tricarpellate, triovulate gynoecium and many staminodes. The fruit is one-seeded, and the endosperm is ruminant. All these characters are congruent with its being a species of *Reinhardtia*.

Currently, six species of the genus are accepted, viz: *R. gracilis*, *R. paiewonskiana*, *R. latisepta*, *R. elegans*, *R. simplex* and *R. koschnyana* (see Henderson 2002). Henderson (2002) carried out a phenetic and phylogenetic analysis of the genus, confirming the species delimitation of Moore (1957) with the addition of *R. paiewonskiana* described from the Dominican Republic (Read et al. 1987). Henderson did not provide descriptions or keys. In the treatment of Moore, the only detailed monograph, the Floribunda plant keys to *R. latisepta*, based on the indumentum on the petioles, stamen number and endosperm ruminant. However, the appearance of the Floribunda plant is at

odds with *R. latisecta*. *Reinhardtia latisecta* is usually a relatively tall clustering palm with stems to 8 m tall and about 7 cm in diameter. The leaves are divided into two to three broad leaflets on each side of the rachis, occasionally with a few single-fold leaflets in between, and usually display striking “windows” (short basal splits) at the proximal end of the folds in the broad segments. In the Floribunda palm, the stems are very short, scarcely exceeding 50 cm tall, and the leaves are much smaller and are irregularly divided, into many leaflets, the basal pair being multifold and displaying windows, while most other leaflets are single-fold, some even displayed in different planes (Fig. 2). Finally, the inflorescences are erect and exserted from the leaves (Fig. 1), whereas in *R. latisecta* they are usually held below the crown. However, we can see no differences between the flowers and fruit of the Floribunda palm and *R. latisecta*.

At first we thought that the Floribunda palm might be an undescribed taxon, but with the lack of differences in flowers and fruit and with the limited material available to us we have decided to determine it as an unusual

compact form of *R. latisecta*. This compact form appears to come true from seed and, indeed, there are some slight differences in seedling morphology between *Reinhardtia latisecta* as grown at Floribunda and *R. latisecta* “compact form” (Fig. 3).

This is a beautiful palm and we believe it has considerable potential as an ornamental. By featuring it in this article, we draw attention to it and with the informal moniker “compact form” provide a reference point for it in horticulture while we have insufficient evidence to describe it formally.

LITERATURE CITED

- HENDERSON, A. 2002. Phenetic and phylogenetic analysis of *Reinhardtia* (Palmae). *Amer. J. Bot.* 89: 1491–1502.
- MOORE, H.E. 1957. Synopses of various genera of Arecoideae. 23. *Reinhardtia*. *Gentes Herb.* 8: 541–576.
- READ, R.W., T.A. ZANONI AND M. MEJIA. 1987. *Reinhardtia paiewonskiana* (Palmae), a new species for the West Indies. *Brittonia* 39: 20–25.

Photo Feature



Basselinia cf. *deplanchei* growing on Kouakoue Massif, New Caledonia. Several superficially similar species in this genus are all highly variable, making field identification difficult. Among these species, those at higher elevations in more exposed conditions, like this specimen, tend to have stiff, narrow, undivided leaves, while those in lower or more protected sites can be pinnately dissected. These species are uniquely adapted to grow on the island's distinctive ultramafic soils, which tend to be composed of high concentrations of magnesium, iron, nickel and other metals at levels that would be toxic to most plants.

ADAM BLACK
Peckerwood Garden

Index to Volume 61

- A new species of *Pritchardia* from the Marquesas Islands with notes on the genus in French Polynesia 139
- Adorador, J.T. & E.S. Fernando: Palms of Samar Island, Philippines 161
- Aiphanes* 9
- Aiphanes acaulis* 9, 12, 14, 15
- Aiphanes buenaventurae* 9, 12, 14–16
- Aiphanes hirsuta* 12
- An unusual form of *Reinhardtia latisecta* in cultivation 197
- Archontophoenix* 35
- Archontophoenix alexandrae* 24–26, 35
- Archontophoenix cunninghamiana* 23, 26
- Areca* 28, 30, 34, 161, 171, 174, 176, 177
- Areca aliciae* 26, 34
- Areca caliso* 175–177, 193
- Areca catechu* 34, 187, 193
- Areca costulata* 168, 175, 176, 193
- Areca jokowi* 4
- Areca monostachya* 26, 28
- Areca rheophytica* 68
- Areca sapida* 23, 30
- Areca* sp. “mudulan” 174, 176, 193
- Areca triandra* 26, 34
- Arenga pinnata* 193
- Asterogyne martiana* 12
- Astrocaryum malybo* 12, 14, 15
- Astrocaryum standleyanum* 12
- Attalea allenii* 12
- Attalea iguadummat* 10–12, 15
- Bactris* 14
- Bactris barronis* 12
- Bactris brongniartii* 12
- Bactris charnleyae* 11, 12, 15
- Bactris chocoensis* 12, 14
- Bactris coloradonis* 12
- Bactris gasipaes* 12
- Bactris glandulosa* 12
- Bactris gracilior* 12
- Bactris major* 12
- Bactris manriquei* 12, 14
- Bactris maraja* 12
- Bactris obovata* 12
- Bactris pilosa* 12
- Bactris setulosa* 12, 14
- Bacularia* 22, 26, 28, 33
- Baron Ferdinand von Mueller, the “Princeps of Australian Botany,” and a historical account of his Australasian palms 21
- Basselinia* cf. *deplanchei* 201
- Bernal, R., as co-author 5
- Bismarckia* 79
- Bismarckia nobilis* 137
- Black, A.: Photo Feature 201
- Borassus* 79
- Butaud, J.-F. and D.R. Hodel: A new species of *Pritchardia* from the Marquesas Islands with notes on the genus in French Polynesia 139
- Butaud, J.-F., as co-author 139
- Calamus* 166, 171, 174, 175, 176
- Calamus aidae* 175, 177, 180, 183, 186, 188, 192
- Calamus australis* 26
- Calamus daemonoropoides* 165, 171, 175, 186, 192
- Calamus discolor* 177, 179, 183, 186, 192
- Calamus elmerianus* 176, 180, 192
- Calamus filispadix* 175–177, 186, 192
- Calamus loherianus* 180
- Calamus megaphyllus* 176, 180, 183, 186, 192
- Calamus merrillii* 175, 183, 187, 188, 192
- Calamus microsphaerion* 179
- Calamus mindorensis* 176, 177
- Calamus mollis* 175, 186, 192
- Calamus moseleyanus* 171, 176, 183, 192
- Calamus muelleri* 32
- Calamus multinervis* 171, 176, 177, 179, 183, 192
- Calamus obstruens* 26
- Calamus ochrolepis* 166, 171, 175, 177, 180, 192
- Calamus ornatus* 175, 183, 186, 192
- Calamus pedicellaris* 176, 192
- Calamus reyesianus* 176
- Calamus siphonopathus* var. *dransfieldii* 171, 176, 192
- Calamus* sp. “malubagacay-bilog” 176, 180, 193
- Calamus* sp. “pudlos-liitan” 179, 180, 186, 188, 193
- Calamus* sp. “pudlos-uban” 179, 180, 193
- Calamus* sp. “tagda” 180, 184, 186, 188, 193
- Calamus* sp. “tagsa-on” 159, 176, 193
- Calamus symphysipus* 192
- Calamus urdanetanus* 165, 171, 192
- Calamus usitatus* 183, 193
- Calamus vidalianus* 176, 180, 183, 193
- Calypstrogyne costatifrons* 3, 12, 14
- Cano, Á., as co-author 5
- Cano, Á., H.F. Manrique, S.E. Hoyas-Gómez, N. Echavarría, A. Upegui, M.F. González, G. Galeano & R. Bernal: Palms of the Darien Gap (Colombia-Panama) 5
- Carpenteria acuminata* 33
- Caryota albertii* 26, 32
- Caryota cumingii* 175, 183, 193
- Caryota mitis* 160
- Caryota rumphiana* 175, 187, 193
- Caryota urens* 26
- Castañeda-Noa, I, as co-author 123

- Castillo Mont, J.J., D.R. Hodel, F.W. Stauffer & J.W. Tregear: The Pacaya palm, *Chamaedorea tepejilote*: observations of its cultivated and wild forms in Guatemala 109
- Castillo Mont, J.J., as co-author 109
- Chamaedorea* 11, 14, 109, 113, 117
- Chamaedorea allenii* 12
- Chamaedorea cataractarum* 65–68
- Chamaedorea costaricana* 12
- Chamaedorea deneversiana* 12
- Chamaedorea lucidifrons* 12
- Chamaedorea macrospadix* 12
- Chamaedorea matae* 12]
- Chamaedorea pinnatifrons* 12
- Chamaedorea ponderosa* 11, 12, 14, 15
- Chamaedorea pygmaea* 12, 16
- Chamaedorea seifrizii* 67
- Chamaedorea subjectifolia* 12, 15
- Chamaedorea tepejilote* 12, 105, 109, 110, 112, 114, 115, 117
- Chamaedorea warszewiczii* 12, 14
- Chamaedorea woodsoniana* 8, 12
- Chamaerops humilis* 56, 140
- Clinostigma* 33
- Coccothrinax* × *angelae* 45–49, 52
- Coccothrinax* × *angelae*, the first natural hybrid of the genus 45
- Coccothrinax* 45, 46, 83, 84
- Coccothrinax clarensis* 89
- Coccothrinax clarensis* subsp. *clarensis* 84, 89
- Coccothrinax crinita* subsp. *brevicrinis* 46, 47, 49
- Coccothrinax macroglossa* 84
- Coccothrinax miraguama* subsp. *roseocarpa* 46, 47
- Coccothrinax* sp. “azul” 84
- Coccothrinax* sp. “blue leaves” 84
- Coccothrinax spirituana* 55, 83–89
- Cocos* 34
- Cocos normanbyi* 34
- Cocos nucifera* 12, 26, 143, 145, 186, 194, 195
- Copernicia* 14
- Corypha* 69
- Corypha taliera* 33
- Corypha umbraculifera* 140, 160
- Corypha utan* 186, 194
- Cryosophila kalbreyeri* 13
- Dahme, M.: Observations on reproduction in *Kerriodoxa elegans* 135
- Desmoncus cirrifer* 13
- Desmoncus myriacanthos* 13
- Desmoncus obovoideus* 13
- Dictyocaryum lamarckianum* 8, 13
- Dowe, J., Baron Ferdinand von Mueller, the “Princeps of Australian Botany,” and a historical account of his Australasian palms 21
- Dransfield, J. & J. Marcus: An unusual form of *Reinhardtia latisepta* in cultivation 197
- Dransfield, J., Book review 102
- Dransfield, J.: Natalie Uhl (1919–2017) 119
- Dransfield, J.: Photo Feature 138
- Dypsis ovobontsira* 64
- Echavarria, N., as co-author 5
- Elaeis guineensis* 108
- Elaeis oleifera* 10, 13
- Euterpe longevaginata* 13
- Euterpe oleracea* 10, 13, 67, 68
- Euterpe precatoria* 13
- Exploring Robinson Crusoe or in search of *Juania australis* 56
- Fernando, E.S., as co-author 161
- Financial statement 196
- Galeano, G., as co-author 5
- García-Lahera, J.P., as co-author 83
- Gardiner, L.M., as co-author 69
- Gardiner, L.M., D. Rabehevitra, R. Letsara & A. Shapcott: *Tahina spectabilis*: an exciting new discovery in Madagascar ten years on 69
- Geonoma* 14, 117
- Geonoma calyptrogynoides* 13, 14
- Geonoma cuneata* 13
- Geonoma cuneata* subsp. *linearis* 67
- Geonoma deversa* 13
- Geonoma frontinensis* 13
- Geonoma interrupta* 13
- Geonoma lehmanii* 13
- Geonoma linearis* 67
- Geonoma maxima* subsp. *dispersa* 13
- Geonoma procumbens* 13
- Geonoma triandra* 13, 14, 17
- Geonoma undata* 13
- González, M.F., as co-author 5
- Griffith, M.P., as co-author 41
- Guihaia* 4
- Guihaia lancifolia* 4
- Hamann, G.: Exploring Robinson Crusoe or in search of *Juania australis* 56
- Hedyscepe canterburyana* 26, 28, 29
- Hemithrinax* 123, 124, 126–132
- Hemithrinax ekmaniana* 105, 107, 123–126
- Hemithrinax ekmaniana*, jewel of the Cuban palms 123
- Heterospathe* 161, 171, 174, 188
- Heterospathe elata* 183, 194
- Heterospathe intermedia* 164, 177, 178, 194
- Heterospathe philippinensis* 183
- Heterospathe* sp. “saguisi-mina” 180, 185, 194
- Hodel, D.R., as co-author 45, 109, 123, 139
- Howea belmoreana* 26–28
- Howea forsteriana* 26–28
- Hoyas-Gómez, S.E., as co-author 5
- Hurtado, R., as co-author 91
- Hydriastele montana* 26, 35, 198
- Hydriastele rheophytica* 68
- Hydriastele wendlandiana* 26, 28
- Hyospathe elegans* 13

- Hyospathe pittieri* 13, 14
Hyphaene 160
Iriartea deltoidea 13
Juania australis 53, 57, 59, 60, 62, 63
Kentia 28, 30, 33
Kentia acuminata 33
Kentia beccarii 26, 35
Kentia belmoreana 26–28
Kentia canterburyana 26, 28, 29
Kentia forsteriana 26–28
Kentia monostachya 28
Kentia mooreana 26, 28, 33
Kentia wendlandiana 26, 28
Kentiopsis 30
Kerriodoxa elegans 107, 135, 156
Korthalsia laciniosa 193
Lepidorrhachis mooreana 26, 33
 Letsara, R., as co-author 69
Licuala 30, 33
Licuala muelleri 32–34
Licuala peltata 33
Licuala ramsayi 26, 30–34
 Lima, M.R., as co-author 123
 Lima, M.R., as co-author 45
Linospadix 28
Linospadix monostachyos 26, 28
Livistona 23, 26, 28, 30, 32–35
Livistona alfredii 1, 26, 36
Livistona australis 23, 36
Livistona decora 32
Livistona drudei 26, 32, 33
Livistona humilis 26, 34, 35
Livistona inermis 32–35
Livistona leichhardtii 26, 28, 34, 35
Livistona mariae 26, 30, 32, 33, 35, 36
Livistona ramsayi 30–34
Livistona saribus 173, 183, 186, 194
 Lopez, R. et al. 2015: Palmeiras Nativas do Brasil, reviewed 102
 Magellan, T.M. & M.P. Griffith: Status of *Pseudophoenix sargentii* on Elliott Key, Florida 41
Manicaria saccifera 10, 13
 Manrique, H.F., as co-author 5
 Marcus, J., as co-author 197
 Martínez-Pentón, L.R., as co-author 83
Mauritia flexuosa 99, 108
Metroxylon sagu 183, 186, 194
 Metz, X., notice of death 4
 Moraes R., M., as co-author 91
 Moraes R., M., J. Roncal & R. Hurtado: Uses of *Parajubaea torallyi*, a vulnerable palm of Bolivia 91
 Moya López C.E., as co-author 123
 Moya López, C.E., R.M. Verdecia Perez, J.P. García-Lahera & L.R. Martínez-Pentón: The *Coccothrinax* “azul” from Sancti Spiritus, Cuba 83
 Natalie Uhl (1919–2017) 119
Nengella montana 35
Nephrosperma van-houtteanum 56
Normanbya 22, 26, 33
Normanbya normanbyi 34
Nypa fruticans 108, 183, 186, 194
 Observations on reproduction in *Kerriodoxa elegans* 135
Oenocarpus bataua 13, 99
Oenocarpus minor 13
Oncosperma 161, 174, 175, 177
Oncosperma gracilipes 175
Oncosperma horridum 176
Oncosperma sp. “anibong-kinis” 170, 175, 194
Orania 161, 174, 179
Orania decipiens 173, 176, 179, 180, 187, 194
Orania littoralis 179
Orania sp. “banga-igang” 181, 188, 194
 Palm Literature 102
 Palmeiras Nativas do Brasil, reviewed 102
 Palms of Samar Island, Philippines 161
 Palms of the Darién Gap (Colombia-Panama) 5
Parajubaea sunkha 99
Parajubaea torallyi 91–101
Phoenicophorium borsignianum 56
Phoenix roebelenii 68
Pholidostachys dactyloides 13
Pholidostachys panamensis 11, 13, 15, 17
 Photo Feature 64, 138, 201
Phytelephas macrocarpa 13
Pinanga 161, 166, 171, 174, 175, 180, 187
Pinanga copelandii 159, 175, 177, 194
Pinanga insignis 172, 176, 187
Pinanga rivularis 67
Pinanga samarana 171, 175–177, 180, 187
Pinanga sp. “tagibungang-liitan” 169, 175, 187, 194
Plectocomia elongata var. *philippinensis* 166, 171, 175, 180, 187, 193
 President’s message 102, 196
Prestoea acuminata 13
Prestoea decurrens 13
Pritchardia 139, 140, 145, 149–151
Pritchardia glabrata 149
Pritchardia mitiaraana 139, 143, 148–153
Pritchardia pacifica 139, 140, 149, 151, 152
Pritchardia pacifica var. *marquisensis* 143, 151, 152
Pritchardia pericularum 143, 149
Pritchardia tahuatana 140–152
Pritchardia thurstonii 139, 149, 151, 152
Pritchardia thurstonii 35
Pritchardia vuylstekeana 143, 149
Pseudophoenix sargentii 41–43
Ptychosperma 23, 35
Ptychosperma alexandrae 24–26
Ptychosperma beatriceae 26, 35

- Ptychosperma normanbyi* 34
 Rabehevitra, D., as co-author 69
 Rakotoarinivo, M.: Photo Feature 64
Raphia taedigera 10, 13
Ravenea madagascariensis 138
Reinhardtia 117, 198, 199
Reinhardtia elegans 199
Reinhardtia gracilis 8, 13, 18, 199
Reinhardtia koschnyana 13, 19, 199
Reinhardtia latisecta 199, 200
Reinhardtia latisecta "compact form" 159, 197–200
Reinhardtia paiewonskiana 199
Reinhardtia simplex 13
Rhopalostylis sapida 23, 30
 Rodríguez Lima, M., I. Castañeda-Noa, C.E. Moya López, D. Suárez Oropesa & D.R. Hodel: *Hemithrinax ekmaniana*, jewel of the Cuban palms 123
 Roncal, J., as co-author 91
Sabal maritima 49
Sabal mauritiiformis 13
Sabinaria 7, 8, 15
Sabinaria magnifica 11, 13, 15, 18
Saribus cf. *merrillii* 173, 176, 195
Saribus rotundifolius 167, 175, 186, 195
Seaforthia 23
Seaforthia elegans 26
 Seed dispersal in *Chamaedorea cataractarum* 65
 Shapcott, A., as co-author 69
Socratea exorrhiza 13
 Status of *Pseudophoenix sargentii* on Elliott Key, Florida 41
 Stauffer, F.W., as co-author 109
 Suárez Oropesa, D., as co-author 123
 Suárez Oropesa, D., M. Rodríguez Lima & D.R. Hodel: *Coccothrinax* × *angelae*, the first natural hybrid of the genus 45
Syagrus smithii 108
Synechanthus 117
Synechanthus dasystachys 4
Synechanthus warszewiczianus 13
Tahina 69, 70, 72, 74, 76–81
Tahina spectabilis 69–75, 79–82, 104
Tahina spectabilis: an exciting new discovery in Madagascar ten years on 69
 The *Coccothrinax* "azul" from Sancti Spiritus, Cuba 83
 The Pacaya palm, *Chamaedorea tepejilote*: observations of its cultivated and wild forms in Guatemala 109
Thrinax 45, 124
Thrinax radiata 42
 Tregear, J.W., as co-author 109
Trithrinax compacta 123
 Uhl, N.W., notice of death 56, 119
 Upegui, A., as co-author 5
 Uses of *Parajubaea torallyi*, a vulnerable palm of Bolivia 91
 Verdecia Perez, R.M., as co-author 83
Welfia regia 13
Wettinia 108
Wettinia aequalis 8, 13, 14
Wettinia donosoensis 108
Wettinia kalbreyeri 108
Wettinia quinaria 13
Wettinia radiata 13
 Zona, S.: Seed dispersal in *Chamaedorea cataractarum* 65

UNITED STATES POSTAL SERVICE® (All Periodicals Publications Except Requester Publications)

1. Publication Title Palms Quarterly	2. Publication Number 7 8 2 - 7 2 0	3. Filing Date 19 Sep 2017
4. Issue Frequency Quarterly	5. Number of Issues Published Annually 4	6. Annual Subscription Price \$45.00
7. Complete Mailing Address of Known Office of Publication (Not printer) (Street, city, county, state, and ZIP+4®) International Palm Society, 9300 Sandstone St., Austin, Travis Co., TX 78737-1135		
8. Complete Mailing Address of Headquarters or General Business Office of Publisher (Not printer) International Palm Society, 9300 Sandstone St., Austin, TX 78737-1135		
9. Full Names and Complete Mailing Addresses of Publisher, Editor, and Managing Editor (Do not leave blank) John Dransfield, Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, United Kingdom Scott Zona, 6111 SW 64 Ave., South Miami, FL 33143		

10. Owner (Do not leave blank. If the publication is owned by a corporation, give the name and address of the corporation immediately followed by the names and addresses of all stockholders owning or holding 1 percent or more of the total amount of stock. If not owned by a corporation, give the names and addresses of the individual owners. If owned by a partnership or other unincorporated firm, give its name and address as well as those of each individual owner. If the publication is published by a nonprofit organization, give its name and address.)

11. Known Bondholders, Mortgagees, and Other Security Holders Owning or Holding 1 Percent or More of Total Amount of Bonds, Mortgages, or Other Securities. If none, check box ☒ None

12. Tax Status (For completion by nonprofit organizations authorized to mail at nonprofit rates) (Check one)
☒ Has Not Changed During Preceding 12 Months
☐ Has Changed During Preceding 12 Months (Publisher must submit explanation of change with this statement)

13. Publication Title
Palms

14. Issue Date for Circulation Data Below
September - 2017

15. Extent and Nature of Circulation	Average No. Copies Each Issue During Preceding 12 Months	No. Copies of Single Issue Published Nearest to Filing Date
a. Total Number of Copies (Net press run)	993	1020
b. Paid Circulation (By Mail and Outside the Mail)	606	627
c. Total Paid Distribution (Sum of 15b (1), (2), (3), and (4))	0	0
d. Free or Nominal Rate Distribution (By Mail and Outside the Mail)	104	121
e. Total Free or Nominal Rate Distribution (Sum of 15d (1), (2), (3), and (4))	238	223
f. Total Distribution (Sum of 15c and 15e)	947	971
g. Copies not Distributed (See Instructions to Publishers #4 (page #3))	1	1
h. Total (Sum of 15f and g)	0	0
i. Percent Paid (15c divided by 15f times 100)	2	2
	3	3
	950	974
	43	46
	993	1020
	99.7	99.7

* If you are claiming electronic copies, go to line 16 on page 3. If you are not claiming electronic copies, skip to line 17 on page 3.

16. Electronic Copy Circulation

16. Electronic Copy Circulation	Average No. Copies Each Issue During Preceding 12 Months	No. Copies of Single Issue Published Nearest to Filing Date
a. Paid Electronic Copies		
b. Total Paid Print Copies (Line 15c) + Paid Electronic Copies (Line 16a)		
c. Total Print Distribution (Line 15f) + Paid Electronic Copies (Line 16a)		
d. Percent Paid (Both Print & Electronic Copies) (16b divided by 16c x 100)		

☒ I certify that 50% of all my distributed copies (electronic and print) are paid above a nominal price.

17. Publication of Statement of Ownership
☒ If the publication is a general publication, publication of this statement is required. Will be printed in the **December 2017** issue of this publication.
☐ Publication not required.

18. Signature and Title of Editor, Publisher, Business Manager, or Owner
Managing Editor
Date
19 Sep. 2017

I certify that all information furnished on this form is true and complete. I understand that anyone who furnishes false or misleading information on this form or who omits material or information requested on the form may be subject to criminal sanctions (including fines and imprisonment) and/or civil sanctions (including civil penalties).

PS Form 3526, July 2014 (Page 1 of 4) (see instructions page 4) PSN: 7530-01-000-9931 PRIVACY NOTICE: See our privacy policy on www.usps.com

UNITED STATES POSTAL SERVICE® (All Periodicals Publications Except Requester Publications)

16. Electronic Copy Circulation	Average No. Copies Each Issue During Preceding 12 Months	No. Copies of Single Issue Published Nearest to Filing Date
a. Paid Electronic Copies		
b. Total Paid Print Copies (Line 15c) + Paid Electronic Copies (Line 16a)		
c. Total Print Distribution (Line 15f) + Paid Electronic Copies (Line 16a)		
d. Percent Paid (Both Print & Electronic Copies) (16b divided by 16c x 100)		

☒ I certify that 50% of all my distributed copies (electronic and print) are paid above a nominal price.

17. Publication of Statement of Ownership
☒ If the publication is a general publication, publication of this statement is required. Will be printed in the **December 2017** issue of this publication.
☐ Publication not required.

18. Signature and Title of Editor, Publisher, Business Manager, or Owner
Managing Editor
Date
19 Sep. 2017

I certify that all information furnished on this form is true and complete. I understand that anyone who furnishes false or misleading information on this form or who omits material or information requested on the form may be subject to criminal sanctions (including fines and imprisonment) and/or civil sanctions (including civil penalties).

PS Form 3526, July 2014 (Page 3 of 4) PRIVACY NOTICE: See our privacy policy on www.usps.com

The IPS is grateful for the support of the following patrons

2016

Platinum (\$5000 and up)

Dr. Tom Jackson, California, USA, underwriting PALMS 61(1).
Ms. Jill Menzel, Florida, USA, in support of the Endowment Fund.

Gold (\$1000 to \$4999)

Ms. Sally Sneed Betts, Florida, USA,
in memory of her parents, Melvyn & Phyllis Sneed.
Mr. Leland Lai, California, USA.

Silver (\$500 to \$999)

Mr. Anthony Gray, Vancouver, Canada.

Bronze

Mr. Andrew Grant, California, USA.
Mr. Keith Thibodeaux, Louisiana, USA.
Dr. Lazaro Priegues, Florida, USA.

2017

Platinum (\$5000 and up)

Ms. Jill Menzel, Florida, USA, in support of
the Larry Noblick Fund for Research.

Gold (\$1000 to \$4999)

Ms. Sally Sneed Betts & Ms. Sarah Morlang, USA,
in memory of their parents, Melvyn & Phyllis Sneed.

Silver (\$500 to \$999)

Palm Society of South Texas, USA

Bronze

If you would like to make a gift, however large or small, please contact IPS
President Ray Hernandez, subtropicoofcancer@hotmail.com or any IPS Board
member. To give on-line, visit www.palms.org.

