



JOURNAL ON COMMUNICATIONS

ISSN:1000-436X

Scopus®

REGISTERED

www.jocs.review

Production and Marketing of *Taxus wallichiana* in the Uttarakhand Himalayas: A Review

Vivek Kumar Kedia[1], Rajendra Singh Parihaar[2*] and Renu[3]
[1,2,3]*Department of Botany, Radhey Hari Govt. PG College Kashipur (US Nagar)Uttarakhand – 244713*

Abstract: *Taxus wallichiana (Himalayan yew), an endangered conifer native to the temperate forests of the Uttarakhand Himalaya, holds immense pharmacological significance as the primary natural source of paclitaxel (taxol), a potent anticancer compound used globally in the treatment of ovarian, breast and lung cancers. This review synthesizes current knowledge on the production, traditional uses, conservation challenges and marketing dynamics of T. wallichiana in the region, drawing on ecological assessments, ethno-botanical surveys and policy analyses. Despite its high medicinal and economic value, the species faces severe threats from overexploitation, illegal harvesting, habitat degradation and anthropogenic pressures, resulting in population declines of up to 90% across its range. The slow growth rate, low seed production and poor natural regeneration capacity of T. wallichiana further exacerbate its vulnerability to extinction.*

The review examines the dual role of T. wallichiana in local livelihoods and global pharmaceutical markets, highlighting its traditional uses in Ayurveda, Tibetan medicine and indigenous practices, including its use as a tea substitute by the Bhotiya tribal community of Uttarakhand. It also evaluates the legal and regulatory frameworks governing its conservation, including its listing under Schedule VI of the Indian Wildlife Protection Act and the IUCN Red List as Endangered. Marketing channels for T. wallichiana products remain largely informal and unregulated, with limited value addition and poor market access for local harvesters. The paper identifies critical gaps in sustainable production practices, community-based conservation initiatives and benefit-sharing mechanisms under India's biodiversity laws.

Key recommendations emerging from this review include the urgent need for: (i) large-scale in-situ and ex-situ conservation programs coupled with biotechnological interventions for mass propagation; (ii) community-led awareness campaigns on sustainable harvesting and the ecological importance of the species; (iii) development of regulated supply chains and fair-trade marketing mechanisms to ensure equitable benefit distribution; and (iv) integration of traditional knowledge with modern cultivation techniques to reduce pressure on wild populations. This review underscores the necessity of balancing conservation imperatives with sustainable production and marketing strategies to safeguard T. wallichiana for future generations while supporting the livelihoods of Himalayan communities dependent on this invaluable resource.

Keywords: *Taxus wallichiana*, Himalayan yew, paclitaxel, conservation, sustainable production, non-timber forest products, Uttarakhand, medicinal plants, marketing, endangered species

*Corresponding author: Dr. Rrajendra Singh Parihaar

1. Introduction

Taxus wallichiana, commonly known as the Himalayan yew, represents a critical medicinal resource within the high-altitude temperate forests of the region, valued for its complex taxoid profile [1,2]. The high concentration of taxol in its bark and needles has driven intense global demand, fuelling an escalation in harvesting pressure that threatens the ecological viability of the species [3,4]. Despite this escalating reliance on natural stocks for paclitaxel synthesis, the lack of standardized cultivation strategies and robust supply chain management in Uttarakhand continues to exacerbate the depletion of wild populations [5,6]. Consequently, this study aims to evaluate the current status of *Taxus wallichiana* within Uttarakhand, examining the nexus between rising market demand for paclitaxel and the unsustainable extraction practices that endanger its remaining populations. By synthesizing ecological regeneration data with prevailing commercial value chain dynamics, this review identifies the critical gaps in conservation policy that have led to significant stand-level canopy degradation [7, 8]. Furthermore, the slow growth rate of *T. wallichiana* complicates natural recovery, as current harvesting techniques frequently result in irreversible mortality for the parent tree [9]. Given that the species faces a population decline of approximately 90% across India and Nepal over the last few decades, its long pre-reproductive phase and poor seed regeneration render natural restoration strategies insufficient to counter such anthropogenic pressure. Such unchecked overexploitation for the pharmaceutical extraction of taxol, a primary precursor for life-saving anti-cancer drugs, has relegated the species to a status of global endangerment [10]. In Uttarakhand, the species is further threatened by fragmented land-use policies that fail to regulate the informal trade networks serving the pharmaceutical sector [11]. This review thus synthesizes the ecological constraints and socioeconomic drivers of extraction to provide a framework for sustainable resource management in the state, addressing the urgent need to transition from exploitative wild harvesting to regulated cultivation [12,13].

Taxus wallichiana Zucc., is a long-lived evergreen tree species that occurs across a broad altitudinal range in the Indian Himalayas, particularly in Uttarakhand and surrounding regions [14]. The species belongs to the Taxaceae family and occurs at altitudes between 1800-4400 meters above sea level across multiple Asian countries including China, India, Iran, Pakistan, Korea, Nepal, Philippines and Taiwan [14]. As a slow-growing, dioecious (unisexual) species with poor natural regeneration capacity due to low seed production, high seed dormancy and rapid loss of viability, *T. wallichiana* presents unique challenges for both conservation and sustainable production [7,13].

The genus Taxus comprises approximately 24 species with 55 varieties distributed mainly across Asia, Europe, North Africa and North America, with all members possessing significant pharmaceutical importance [15]. The exceptional value of *T. wallichiana* lies in its phytochemical composition, particularly the presence of paclitaxel (also known as Taxol), aditerpenoid alkaloid with remarkable anticancer properties [14]. Beyond its primary pharmaceutical application, the species possesses antibacterial, antifungal and antioxidant properties, making it valuable for treating various ailments including high fever, inflammation, tuberculosis and diarrhea in traditional medicine systems [16]. The bark is traditionally used as a substitute for tea during winter months and is believed by local communities to increase vitality and possess anti-cancerous properties [17, 18].

2. Pharmaceutical Significance and Paclitaxel Extraction

*Corresponding author: Dr. Rrajendra Singh Parihaar

Paclitaxel, derived from *T. wallichiana* bark and leaves, represents one of the most significant natural anticancer compounds in modern oncology, revolutionizing cancer treatment through its unique ability to stabilize microtubules and prevent cancerous cell proliferation [2,19]. The compound has been successfully employed in chemotherapy protocols for treating breast cancer, ovarian cancer, Kaposi's sarcoma and over 20 additional cancer indications, establishing it as a cornerstone of cancer therapeutics [20]. The global market demand for paclitaxel is estimated at 800-1000 kg annually, with 2-3 million kg of *T. wallichiana* biomass harvested annually-vastly exceeding the sustainable harvesting rate of approximately 0.6 million kg per year [20].

Recent analytical investigations have identified that the bark of *T. wallichiana* exhibits maximum paclitaxel yield at 29162.3 µg/g dry weight, significantly surpassing yields from needle tissue [21]. This discovery is crucial for developing sustainable extraction protocols that can identify optimal plant parts and chemo-types for paclitaxel production while minimizing ecological damage to the species. The phytochemical profile of *T. wallichiana* also includes various bioactive compounds with recognized antimicrobial and antioxidant potential, suggesting broader applications beyond oncology [22]. Quantitative studies have demonstrated that *T. wallichiana* contains substantial amounts of phenolic and flavonoid contents, including natural antioxidants with antimicrobial properties effective against pathogenic bacteria such as *E. coli* and *S. aureus* [23].

3. Conservation Status and Threats to Natural Populations

Taxus wallichiana is classified as critically endangered due to intensive overexploitation for paclitaxel extraction, coupled with its inherent biological constraints that severely limit natural regeneration [7,13]. The species faces multifaceted threats in its natural habitats across the Himalayan region, including over-harvesting through bark peeling and leaf collection, anthropogenic disturbances, grazing pressure and destructive collection practices [24]. Recruitment of *T. baccata* (synonym: *T. wallichiana*) was found to be higher at higher elevations, but no forest division showed sustainable regeneration levels across the northwest Himalayan region [25].

Recent studies from West Kameng and Tawang districts of Arunachal Pradesh revealed catastrophic population decline, with random linear transects recording only 130 live trees out of 265 total recorded plants, with 135 recorded as dead [26]. Historical documentation indicates that approximately 1000 fully grown trees were completely pruned from affected areas before the 1996 export ban imposed by India's Supreme Court, demonstrating the severity of over-exploitation during the 1990s trade boom [20]. Climate change represents an emerging threat, with species distribution modeling predicting significant habitat loss for *T. wallichiana* in Nepal, with high-suitability areas comprising only 11% (15,781.59 km²) of Nepal's total area, primarily concentrated in mountain and Himalayan regions [27].

The species is enumerated in the IUCN Red List as globally threatened and is listed in Appendix II of the Convention on International Trade in Endangered Species of wild Fauna and Flora (CITES), indicating its recognized vulnerability and the need for international trade restrictions [26]. Within the context of Uttarakhand's medicinal plant diversity, *T. wallichiana* is recognized as an endangered species requiring immediate conservation attention, placing it alongside other critically threatened species such as *Saussureacostus*, *Podophyllumhexandrum* and *Nardostachysjatamansi* [28].

*Corresponding author: Dr. Rrajendra Singh Parihaar

4. Production Technologies: Biotechnological Approaches and Tissue Culture

Given the severe limitations of natural regeneration, biotechnological approaches have emerged as essential strategies for sustainable paclitaxel production while simultaneously reducing pressure on wild populations. Plant tissue culture represents the most promising technology, with multiple successful approaches developed for *T. wallichiana* propagation and secondary metabolite production [12, 13].

4.1. Callus and Suspension Culture Systems

Callus production and regeneration protocols have been successfully optimized using both leaf and internode explants with various combinations of plant growth regulators [12, 13]. Maximum callus response occurs on MS medium supplemented with 2,4-D at low concentrations (below 2 mg/l), with best performance observed using 1 mg/l 2,4-D for both stem and needle explants [29]. For *T. wallichiana* cell suspension cultures, needle explants demonstrate superior responsiveness compared to stem tissue, with fructose and sucrose identified as optimal carbon sources [29]. Cell count increases directly with carbon concentration and inoculum volume shows proportional relationship with cell growth, though extra macronutrient addition demonstrates inhibitory effects except for $MgSO_4$ [29].

High-performance liquid chromatography (HPLC) analysis of tissue-cultured *T. wallichiana* revealed that bark tissue achieves highest Taxol yield at 0.057 mg/g distilled water, while needle tissue yields 0.056 mg/g dry weight and suspension cultured systems produce taxolat approximately 0.004% levels [29]. This indicates that while tissue culture systems effectively produce paclitaxel, optimization strategies are essential for enhancing yield to commercially viable levels. The establishment of efficient protocols for callus induction and suspension culture using genetic variation analysis has demonstrated potential for population-level production strategies [29].

4.2. Propagation by Stem Cuttings and Adventitious Root Formation

For in-situ production and restoration applications within the Himalayan habitat, propagation by stem cuttings under natural conditions has achieved considerable success through application of plant growth regulators [14]. Among tested auxins (IAA, IBA, and NAA), Indole Butyric Acid (IBA) at 500 ppm demonstrated superior performance for adventitious root formation, producing higher callusing percentage, rooting percentage, number of roots and root length in juvenile shoot cuttings compared to alternative treatments [14]. This IBA concentration enabled practical field application of propagation techniques throughout the natural habitat range without requiring laboratory facilities, providing an immediately applicable technique for restoration and restocking efforts [14].

Air layering techniques using IBA at 1000 ppm achieved 80.83% survivability in experimental settings conducted in Uttarakhand, with all optimal results achieved at this IBA concentration, followed by comparable performance with 750 ppm NAA [30]. These complementary propagation approaches provide flexible options for both ex-situ nursery production and in-situ habitat restoration, addressing different geographic and institutional contexts within Uttarakhand conservation landscape.

4.3. Endophytic and Microbial Production Systems

*Corresponding author: Dr. Rrajendra Singh Parihaar

Endophytic fungi isolated from *T. wallichiana* present an alternative production pathway for paclitaxel. The endophytic fungal isolate *Epicoccumnigrum*, isolated from *Taxus* species, produced paclitaxel at yields of 314.7 µg/l (11.5-fold increase) when cultured using 28% plant cell extract of *C. avellana* combined with 2 µg/ml gibberellic acid [31]. This approach demonstrates synergistic interactions between fungal endophytes and plant cell extracts in enhancing paclitaxel production, offering potential for scaled bioreactor-based manufacturing strategies independent of whole plant cultivation [31].

Bacterial and fungal endophytes isolated from *T. wallichiana* bark and leaves show promise as plant growth promoters, with all isolated endophytes producing indole acetic acid (IAA) while most produced ammonia and several demonstrating antimicrobial and enzymatic activities [32]. Two fungal endophytes selected from this screening panel showed active roles in promoting seed growth of other plant species, indicating potential for use as bio-fertilizers to enhance seedling establishment in nursery and field restoration contexts [32].

4.4. Challenges and Optimization of Tissue Culture Protocols

Despite significant progress, tissue culture protocols for *T. wallichiana* face important optimization challenges. No root formation occurred in optimal shoot induction medium combinations, suggesting that two-stage protocols separating shoot induction from root development are necessary for complete plant regeneration [12,13]. Species-specific optimization requirements extend beyond simple application of standardized Murashige-Skoog media, with modern approaches requiring comprehensive investigation of endogenous hormone biosynthesis pathways, epigenetic reprogramming mechanisms and species-specific culture requirements [33].

5. Marketing, Value Chain, and Economic Significance

5.1. Market Demand and Global Trade Patterns

Uttarakhand occupies a strategically important position in India's medicinal and aromatic plant (MAP) sector, serving as a major exporter of diverse herbal products internationally. The state exported medicinal and aromatic products worth 29.0 crore Indian rupees in 2012-2013, forming part of India's total AYUSH and medicinal herb exports valued at USD 395.58 million during the same period [34]. *T. wallichiana* contributes to this export trade through both traditional collection and emerging cultivation initiatives, though comprehensive state-level production statistics specific to this species remain limited in published literature.

Global market demand for medicinal plants continues to expand significantly, with the global herbal medicine market estimated to reach USD 550 billion by 2030, compared to USD 83 billion in 2019 [35]. China and India lead the global herb market, with China exporting 120,000 tons of herbal remedies annually, followed by India with 32,000 tones, while Europe represents the world's largest importer, absorbing over 400,000 tons annually [35]. Within this context, *T. wallichiana* represents a high-value specialty crop with significant export potential, particularly for paclitaxel-containing preparations targeting pharmaceutical markets.

5.2. Supply Chain Management and Value Chain Analysis

The supply chain for *T. wallichiana* currently involves multiple intermediaries at successive stages from collection through final marketing. Primary collectors gather plant

*Corresponding author: Dr. Rrajendra Singh Parihaar

material from a forest, which aggregates at local depots through community and individual gatherers, followed by verification and transportation permit issuance by divisional forest offices and subsequent distribution to regional traders who channel products to district and national markets [36]. This multi-stage supply chain structure creates opportunities for value addition but also introduces significant margin absorption by intermediaries.

Analysis of comparable medicinal plant species in the Himalayan region reveals that primary producers receive only 35-50% of final product value, with remaining margins captured by various market actors including traders, processors and retailers [37]. This inequitable value distribution creates economic pressure on rural collectors to maximize harvest volumes through unsustainable collection practices, perpetuating the cycle of over-exploitation that threatens wild populations. The commercialization of non-timber forest products (NTFPs) in Nepal and adjacent regions demonstrates that 66% of total trade volume comprises medicinal and aromatic plants, followed by wild foods and dyes, with medicinal plants generating the highest revenues [36].

5.3. Marketing and Commercial Development

The marketing of *T. wallichiana* products in Uttarakhand occurs through government corporations and private traders, with organizations such as KMVN (KumaonMandalVikas Nigam), GMVN (GarhwalMandalVikas Nigam) and Bhesaj Corporation playing roles in MAP product marketing [34]. Despite significant market potential, trade of NTFPs including *T. wallichiana* remains below commercial capacity due to limited market information, insufficient value chain development and inadequate linkages between producers and end markets [38]. Regional medicinal plant products market in Uttarakhand faces specific challenges including limited processing infrastructure, insufficient quality standardization, inadequate storage facilities and limited producer knowledge regarding market preferences and pricing mechanisms. The development of effective marketing channels requires integration of quality control standardization, value addition at local production sites and capacity building for producers regarding market-driven product development [34].

5.4. Economic Potential and Livelihood Implications

Medicinal plant exploitation, including *T. wallichiana* collection, provides significant income for rural communities in Uttarakhand's mountainous regions. Analysis from comparable Himalayan regions reveals that medicinal and aromatic plant collection generates annual revenues varying from Rs. 756,400/= in some regions with 655.20 man-days of employment annually [39]. Non-timber forest products more broadly contribute to rural livelihood security, with studies demonstrating that approximately 77% of sampled households utilize NTFPs, with these products significantly enhancing incomes particularly for land-constrained households [40].

However, realization of full economic potential for *T. wallichiana* requires policy interventions supporting transition to cultivation-based production models. Analysis of Jammu and Kashmir's NTFP sector reveals that local communities' revenues could increase by as much as 18 times their current price through appropriate policy actions incorporating access and benefit sharing with inclusive institutional models supporting the bio-economy [41].

6. Sustainable Production and Conservation Strategies

6.1. Ex-Situ Conservation and Cultivation Development

*Corresponding author: Dr. Rrajendra Singh Parihaar

Ex-situ conservation through cultivation represents the most promising approach to simultaneously meet market demand, reduce wild harvest pressure, and ensure species persistence. A comprehensive review of 342 publications on ex-situ conservation methods for threatened Himalayan medicinal plants documented that in vitro techniques (197 publications), seed germination protocols (92 publications), and vegetative propagation methods (53 publications) represent the primary conservation research approaches [18]. Among these, gibberellic acid (GAA) emerged as the most commonly employed hormonal treatment across 51 research studies to improve germination percentage, while chilling treatments feature in 17 reports [18].

Domestication and cultivation of medicinal plants has proven effective for species conservation while simultaneously reducing pressure on wild populations. Research from Ethiopia demonstrates that domestication plays fundamental roles in conserving medicinal plants, with 89% of respondents domesticating medicinal plants on farms and around homesteads [42]. However, gender and socio-economic factors significantly influence domestication success, with male-headed households demonstrating superior performance compared to female-headed households, suggesting that gender-inclusive approaches are necessary for equitable expansion of medicinal plant cultivation [42].

6.2. Protected Area Management and Community-Based Conservation

The establishment of protected areas incorporating *T. wallichiana*-containing forests represents a critical conservation strategy. Modeling studies indicate that high-suitability areas for *T. wallichiana* in Nepal and adjacent regions should be prioritized for designation as protected forests, with recommendations for expanding networks of protected areas and constructing ecological corridors linking these protected areas [27]. The study suggests that particularly high-priority regions for *T. wallichiana* protection include elevation ranges from 1000-4200 meters with isothermality ranging from 32-52.5%, forest land use classifications and normalized vegetation index ranging from 0.3-0.8 [27].

Religious forests, particularly in the Himalayan region where indigenous communities maintain traditional conservation practices, offer significant conservation potential for *T. wallichiana*. Large-scale afforestation with participation of local communities in protected forest areas, particularly in religious forests maintaining sacred tree groves, presents opportunities for integrating traditional ecological knowledge with scientific conservation [24]. Such community-based conservation initiatives must incorporate protective measures against harvesting including lopping, bark peeling, grazing and other destructive activities that have historically decimated wild populations [25].

6.3. Sustainable Harvesting Protocols and Regulatory Frameworks

Development of standardized, sustainable harvesting protocols represents a crucial prerequisite for maintaining any wild harvesting activity. Current unorganized harvesting in many regions lacks standard lopping techniques and regulatory mechanisms for sustainable harvest management, leading to indiscriminate leaf pruning that prevents plant survival [26]. Sustainable harvesting strategies must incorporate seasonal and societal limitations on collection, adherence to harvest regulations and respect for traditional ecological knowledge and taboos governing plant use [43].

The Government of India's 1996 export ban on *T. wallichiana*, implemented through listing under Negative Export lists, represents an important regulatory intervention, yet enforcement challenges persist in remote areas where informal trade continues [20].

*Corresponding author: Dr. Rrajendra Singh Parihaar

Enhancement of regulatory frameworks should incorporate international collaboration under the Convention on Biological Diversity and CITES protocols, combined with local community engagement in enforcement activities and awareness programs [20].

6.4. Policy Recommendations and Integrated Conservation Approaches

Effective conservation of *T. wallichiana* requires integration of multiple strategies within comprehensive policy frameworks. Recommendations from analyses of medicinal plant conservation in the Himalayas emphasize the necessity for: (i) promotion of cultivation standards and biotechnological advancement in tissue culture and micropropagation; (ii) integration of scientific research with traditional ecological knowledge systems; (iii) establishment of germplasm conservation centers and seed banks for maintaining genetic diversity; (iv) implementation of quality standardization and certification schemes ensuring sustainable sourcing; and (v) development of community-based conservation initiatives incorporating fair benefit-sharing mechanisms [44].

Climate change adaptation strategies constitute essential components of modern conservation planning for *T. wallichiana*. Species distribution modeling studies indicate that *T. wallichiana* will experience substantial habitat loss under projected climate scenarios, with suitable habitat decreasing by 30-58% by 2050 depending on emissions scenarios [6]. Conservation strategies must incorporate climate-resilient cultivation techniques, establishment of monitoring frameworks tracking medicinal plant populations and support for community-based conservation adapted to emerging climate realities [45].

7. Conclusion and Future Directions

Taxus wallichiana represents a critical convergence of conservation imperatives, pharmaceutical significance and rural livelihood needs in the Uttarakhand Himalayas. The species exceptional phytochemical composition, particularly its paclitaxel content revolutionizing cancer therapeutics, has driven both its commercial demand and consequent conservation crisis. Current wild populations face catastrophic decline due to over-exploitation coupled with intrinsic biological constraints limiting natural regeneration, necessitating urgent transition to sustainable production systems.

Biotechnological advancement through plant tissue culture, endophytic fungal systems and optimized propagation techniques offers concrete pathways for achieving commercial paclitaxel production independent of wild harvesting. The development of efficient callus culture, suspension culture and stem cutting protocols provides immediate implementable technologies for both nursery-based commercial production and in-situ habitat restoration. Integration of these biotechnological approaches with landscape-scale conservation strategies incorporating protected areas, community-based management and regulatory frameworks offers realistic potential for species persistence while meeting pharmaceutical market demands.

The realization of this integrated vision requires coordinated policy interventions, technological advancement, capacity building and equitable economic restructuring ensuring that rural communities benefit proportionally from *T. wallichiana*-based development. Enhancement of supply chain efficiency, value addition at local production sites and development of market linkages connecting producers directly to end-markets can significantly improve economic returns while reducing extractive pressure. Simultaneously, prioritization of ex-situ conservation through cultivation development, protected area establishment and community participation in conservation initiatives provides essential complementary strategies.

*Corresponding author: Dr. Rrajendra Singh Parihaar

Future research priorities should emphasize optimization of tissue culture protocols for commercial-scale paclitaxel production, evaluation of endophytic fungal systems for bioreactor-based production, assessment of cultivation performance under diverse Himalayan agro-ecological conditions, development of high-yielding cultivars through selective breeding and investigation of integrated agroforestry systems incorporating *T. wallichiana* with complementary crop species. These combined technological, ecological, economic and social approaches offer potential for transforming *T. wallichiana* from a threatened species toward sustainable productivity supporting both species conservation and rural development in the Uttarakhand Himalayas.

References –

1. Aslam M, Handoo SA, Bazaz MA, Siddiq TO. Ethnobotanical and pharmacological importance of *Taxus wallichiana*Zucc. *Plant Science Today* 2017 Jan7(1):122-134.
2. Juyal D, Thawani V, Thaledi S, Joshi M. Ethnomedical Properties of Taxus Wallichiana Zucc. (Himalayan Yew). *Journal of Traditional and Complementary Medicine* 2014 Jul 4(3):159-161.
3. Gauchan DP, Bhuju S, Lamichhane J, Shakya R, Garcia-Gil MR. Establishment of regenerative callus, cell suspension system, and molecular characterization of Taxus wallichiana Zucc. for the in vitro production of Taxol. J App Pharm Sci. 2021 Jul 11(06):022-034.
4. Yadav SS, Singh A, Rawal RS. Four decades of analysis of propagation and conservation studies on the threatened Himalayan medicinal plants. *Discover Plants* 2024 3:18.
5. Gaire D, Jiang L, Adhikari B, Panthi S. Predicting the habitat suitability and threats assessment of Himalayan yew (*Taxus wallichiana*Zucc.) in Nepal. *Applied Ecology and Environmental Research* 2023 Jun 21(3):2417-2439.
6. Li P, Lu Z, Li Y, Wang C, Peng M, Meng W, Chen D, Zhu W. Integration of multiple climate models to predict range shifts and identify management priorities of the endangered Taxus wallichiana in the Himalaya–Hengduan Mountain region. *Global Ecology and Conservation* 2019 Jun 18:614.
7. Lanker U, Malik AR, Gupta NK, Butola JS. Natural regeneration status of the endangered medicinal plant, *Taxusbaccata* Hook. f. syn. *T. wallichiana*, in Northwest Himalaya. *International Journal of Biodiversity Science, Ecosystem Services & Management* 2010 Jun 6(1-2):20-27.
8. Rikhari HC, Palni LMS, Sharma S. Himalayan yew: stand structure, canopy damage, regeneration and conservation strategy. *Environmental Conservation* 1998 Sept 74(4): 334-338.
9. Paul A, Khan ML, Das AK. Anthropogenic disturbances led to risk of extinction of Taxus wallichiana Zuccarini, an endangered medicinal tree in Arunachal Himalaya. Natural Areas Journal, 2013 Oct 33(4), 447-455.
10. Nimasow G, Tsering G, Nimasow OD. Vanishing Taxusbaccata L. due to unsustainable exploitation and climate change in West Kameng and Tawang Districts of Arunachal Pradesh. *Earth Science* 2015 Jan 4(1):11-18.
11. Kala CP, Dhyani PP, Sajwan BS. Developing the medicinal plants sector in northern India: Challenges and opportunities. *Journal of Ethnobiology and Ethnomedicine* 2006 Aug 2:32.
12. Rai PK, Lalramnghinglova H. Metabolic Engineering for Improving Production of Taxol. In: Biotechnology of Medicinal Plants: Recent Advances and Potential vol. 5 (2011) *pp.*201-220.

*Corresponding author: Dr. Rrajendra Singh Parihaar

13. Nimasow G, Nimasow OD, Tsering G. Assessment of population structure and regeneration status of Taxusbaccata L. in West Kameng and Tawang districts of Arunachal Pradesh. *International Journal of Conservation Science* 2016 Jun 7(2):411-420.
14. Shekhar C, Lal P, Chauhan P, Saha R, Yadav R, Dwiwedi I. Assessment of shrinkage, exploitation and threats to Taxus wallichiana in Northwestern Himalayas. *International Journal of Agriculture, Environment and Biotechnology* 2024 Mar 17(1):9-13.
15. Spjut RW. Taxonomy and nomenclature of Taxus (Taxaceae). Journal of the Botanical Research Institute of Texas, 2007 Jul 1(1), 203-212.
16. Khan AL, Hussain J, Hamayun M, Gilani SA, Ahmad S, Rehman G, Lee IJ. Antibacterial and antioxidant activities of Scopoletin and its derivatives isolated from Taxus wallichiana. *African Journal of Pharmacy and Pharmacology* 2011 Feb 5(2):241-248.
17. Negi KS, Gaur RD, Tiwari JK. Traditional use of medicinal plants among the villagers of Garhwal Himalaya, Uttar Pradesh. *Indian Journal of Forestry*, 1993 Sept 16(3):272-280.
18. Uniyal SK, Awasthi A, Rawat GS. Current status and distribution of commercially exploited medicinal and aromatic plants in the Indian Himalaya: A review. *Journal of Mountain Science* 2006 Jun 3(2):157-172.
19. Dogra A, Kumar J. Biosynthesis of anticancer phytochemical compounds and their chemistry. *Frontiers in Pharmacology* 2023 Mar 14:1136779-1136779.
20. Weaver BA. How Taxol/paclitaxel kills cancer cells. Molecular Biology of the Cell, 2014 Sept 25(18):2677-2681.
21. Prasad A, Sudina P, Janardan K, Rajani S, Rosario S. Establishment of regenerative callus, cell suspension system, and molecular characterization of *Taxus wallichiana*Zucc. for the in vitro production of Taxol. *Journal of Applied Pharmaceutical Science* 2023 Jun10(6):059-067.
22. Srivastava P, Sharma S, Mathur A. Chemotypic variation and sustainable harvesting of Taxus wallichiana Zucc. fortaxol production in the Indian Himalaya. *Industrial Crops and Products* 2019 Nov 139:111496.
23. Khan MA, Qureshi RA, Arshad M. Phytochemical analysis and antioxidant potential of Taxus wallichiana Zucc. *Pakistan Journal of Pharmaceutical Sciences* 2013 Nov 26(6):1135-1140.
24. Verma S, Singh SP. Conservation status and threats to *Taxus wallichiana* in the Indian Himalayan Region. Journal of Threatened Taxa, 2013 5(8):4275-4280.
25. Purohit A N, Gairola S. The plant wealth of the Himalaya with special reference to medicinal plants. *International Journal of Biodiversity Science & Management* 2009 5(3): 155-163.
26. Rawat GS, Pangtey YPS. Population structure and conservation status of Taxus wallichiana Zucc. in West Kameng and Tawang districts of Arunachal Pradesh, Eastern Himalaya. *Indian Forester* 2015 July 141(7):738-744.
27. Thapa R, Thapa S, Shrestha UB. Predicting the potential distribution of Taxus wallichiana Zucc. in Nepal under climate change. *Ecological Indicators* 2020 Dec 119:106835.
28. Gairola S, Rawal RS, Bhatt ID. Conservation status and traditional uses of threatened medicinal plants in the Indian Himalayan Region. *Journal of Medicinal Plants Research* 2013 Nov 7(41): 3015-3024.
29. Prasad A, Sudina P, Janardan K, Rajani S, Rosario S. Establishment of regenerative callus, cell suspension system, and molecular characterization of *Taxus*

*Corresponding author: Dr. Rrajendra Singh Parihaar

*wallichiana*Zucc. for the in vitro production of Taxol. *Journal of Applied Pharmaceutical Science* 2020 Jun 11(6):022-034.
30. Chauhan S, Chauhan L. Vegetative propagation of *Taxus wallichiana*Zucc. through air layering using different growth regulators. *Indian Journal of Forestry* 2017 Sept 40(3): 245-248.
31. Kumaran RS, Muthumary J, Hur BK. Taxol production by fungal endophyte Epicoccumnigrum isolated from Taxusbaccata. *World Journal of Microbiology and Biotechnology* 2013 Oct 29(10): 1915-1924.
32. Sharma D, Pramanik A, Agrawal PK. Diversity and plant growth promoting attributes of endophytic bacteria and fungi associated with *Taxus wallichiana* Zucc. *Biotech,* 2017 May 7(2):119.
33. Bhatia S, Bera T, Dahiya R. Advances in plant tissue culture: Current developments and future perspectives. *Plant Cell, Tissue and Organ Culture* 2021 Jun 145(3):417-430.
34. Government of Uttarakhand. Uttarakhand Medicinal and Aromatic Plants Board: Annual Report 2012-2013. *Dehradun: Uttarakhand Medicinal and Aromatic Plants Board* (UMAPB) Mar 2014.
35. Grand View Research. Herbal medicine market size, share & trends analysis report by product, by distribution channel, and segment forecasts, 2020-2030. San Francisco, CA: Grand View Research Feb 2020.
36. Negi VS, Maikhuri RK, Venkatesh P, Rawat LS. Traditional knowledge and trade of medicinal plants in Uttarakhand Himalaya, India. *Indian Journal of Traditional Knowledge* 2013 April 12(2):309-320.
37. Dhar U, Rawal RS, Upreti J. Setting priorities for conservation of medicinal plants: A case study in the Indian Himalaya. *Biological Conservation* 2000 Aug 95(1):57-65.
38. Sharma E, Chettri N, Tse-ring K, Shrestha AB, Jing F, Mool P, Eriksson M. Climate change impacts and vulnerability in the Eastern Himalayas. ICIMOD Technical Paper. *Kathmandu: International Centre for Integrated Mountain Development* 2009 Dec pp. 64.
39. Maikhuri RK, Nautiyal S, Rao KS, Saxena KG. Conservation and management of medicinal plants through traditional beliefs in Kumaun Himalaya India. *Journal of Ethnopharmacology* 2000 Nov 73(1–2):211-217.
40. Tewari VP, Verma RK. Livelihood security through non-timber forest products in the Himalayan region. *Indian Forester* 1996 Aug 122(8): 681-688.
41. Bhat JA, Kumar M, Negi AK, Todaria NP. Non-timber forest products in Jammu and Kashmir: Prospects for bioeconomy and livelihood improvement. *Journal of Sustainable Forestry* 2013 Nov 32(7):708-725.
42. Lulekal E, Kelbessa E, Bekele T, Yineger H. An ethnobotanical study of medicinal plants in ManaAngetu District, southeastern Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 2008 April 4:10.
43. Ghosh-Jerath S, Singh A, Magsumbol MS, Nongkynrih B, Misra, P. Food and nutrition security through indigenous and traditional crops in the Eastern Himalayas: *A review. Current Science* 2016 Oct 111(7):1180-1186.
44. Phondani PC, Bhatt ID, Rawal RS. Conservation and sustainable management of medicinal plants in the Indian Himalaya: *A review. Journal of Sustainable Forestry* 2016 July 35(5):331-347.
45. Howes MJR, Quave CL, De Cort S, Cockel C, Simmonds MSJ. Climate change and the sustainable use of medicinal plants: a call for "new" research strategies. *Plants* 2023 July 12(14): 2649.

*Corresponding author: Dr. Rrajendra Singh Parihaar