

The complete chloroplast genome of *Microcycas calocoma* (Miq.) A. DC. (Zamiaceae, Cycadales) and evolution in Cycadales

Aimee Caye G. Chang^{1,2,3}, Qiang Lai¹, Tao Chen², Tiejiao Tu¹, Yunhua Wang², Esperanza Maribel G. Agoo⁴, Jun Duan¹ and Nan Li²

¹ South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China

² Shenzhen Fairy Lake Botanical Garden, Chinese Academy of Sciences, Shenzhen, China

³ University of Chinese Academy of Sciences, Beijing, China

⁴ Department of Biology, De La Salle University, Manila, Philippines

ABSTRACT

Cycadales is an extant group of seed plants occurring in subtropical and tropical regions comprising putatively three families and 10 genera. At least one complete plastid genome sequence has been reported for all of the 10 genera except *Microcycas*, making it an ideal plant group to conduct comprehensive plastome comparisons at the genus level. This article reports for the first time the plastid genome of *Microcycas calocoma*. The plastid genome has a length of 165,688 bp with 134 annotated genes including 86 protein-coding genes, 47 non-coding RNA genes (39 tRNA and eight rRNA) and one pseudogene. Using global sequence variation analysis, the results showed that all cycad genomes share highly similar genomic profiles indicating significant slow evolution and little variation. However, identity matrices coinciding with the inverted repeat regions showed fewer similarities indicating that higher polymorphic events occur at those sites. Conserved non-coding regions also appear to be more divergent whereas variations in the exons were less discernible indicating that the latter comprises more conserved sequences. Phylogenetic analysis using 81 concatenated protein-coding genes of chloroplast (cp) genomes, obtained using maximum likelihood and Bayesian inference with high support values (>70% ML and = 1.0 BPP), confirms that *Microcycas* is closest to *Zamia* and forms a monophyletic clade with *Ceratozamia* and *Stangeria*. While *Stangeria* joined the Neotropical cycads *Ceratozamia*, *Zamia* and *Microcycas*, *Bowenia* grouped with the Southern Hemisphere cycads *Encephalartos*, *Lepidozamia* and *Macrozamia*. All *Cycas* species formed a distinct clade separated from the other genera. *Dioon*, on the other hand, was outlying from the rest of Zamiaceae encompassing two major clades—the Southern Hemisphere cycads and the Neotropical cycads. Analysis of the whole cp genomes in phylogeny also supports that the previously recognized family—Stangeriaceae—which contained *Bowenia* and *Stangeria*, is not monophyletic. Thus, the cp genome topology obtained in our study is congruent with other molecular phylogenies recognizing only a two-family classification (Cycadaceae and Zamiaceae) within extant Cycadales.

Submitted 5 September 2019

Accepted 27 November 2019

Published 13 January 2020

Corresponding authors

Jun Duan, duanj@scib.ac.cn

Nan Li, andreali1997@126.com

Academic editor

Victoria Sosa

Additional Information and
Declarations can be found on
page 16

DOI 10.7717/peerj.8305

© Copyright

2020 Chang et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Molecular Biology, Plant Science

Keywords Microcycas, cpDNA, Chloroplast genomes, Cycadales, Phylogeny, Zamiaceae, Cycads, Evolution, Plastids

INTRODUCTION

Chloroplasts are essential organelles for photosynthesis in plant tissues. Their primary function is to fix carbon by harnessing solar energy to synthesize carbohydrates, pigments and amino acids (Hong *et al.*, 2017; Liu *et al.*, 2017). These plant organelles, alongside mitochondria, contain their own DNA and a unique set of genes and are also referred to as plastomes (Wu & Chaw, 2015) although the bulk of gene content is in the nucleus of most eukaryotic organisms (Lodish *et al.*, 2000). Generally, the chloroplast (cp) consists of a large single copy (LSC) region and a small single copy (SSC) region located directly across from each other. Between these two regions are two inverted repeat (IR) sequences (IR_A and IR_B). These four regions make up a single quadripartite circular structure about 120–160 kb in length (Liu *et al.*, 2017; Lee *et al.*, 2006). However, some reports and more recent data from over 1,500 complete genome sequences of chloroplasts from land plants showed that their size ranged up to 170 kb (Smith, 2017; Shaw *et al.*, 2007).

Chloroplast genomes prove to be advantageous over nuclear genomes in ease of handling and analyzing due to their reduced complexity in structure and dimensions. Despite the relatively smaller genome size, a large amount of data can be obtained using cp genomes such as nucleotide substitutions, pseudogenes, intron losses and variation in genome size. Compared to nuclear and mitochondrial DNA, cp genomic regions are more stable and conserved due to their maternal lineage (Zhong *et al.*, 2011) and lack of recombination events (Martin *et al.*, 2005). Information gathered from cp genome sequences are also extensively used in addressing uncertainties in evolutionary relationships in various taxa, reconstructing phylogenetic relationships, and gaining a deeper understanding of biogeographical, structural and functional diversity across all organisms (Jiang, Hinsinger & Strijk, 2016). These features of chloroplast or plastid DNA (cpDNA or ptDNA) make them ideal models for genome-wide evolutionary studies (Asaf *et al.*, 2016; Choi, Chung & Park, 2016; Jiang, Hinsinger & Strijk, 2016).

Cycads (Cycadophyta) are one particular plant group where studies of molecular phylogeny have been broadly applied (Liu *et al.*, 2018; Gorelick *et al.*, 2014; Lu *et al.*, 2014; Salas-Leiva *et al.*, 2013; Chaw *et al.*, 2005; Hill *et al.*, 2003). Because of their ancestry and close morphological resemblance to some pteridophytes and angiosperms (Xiao & Moller, 2015), cycads belong to the earliest lineage among the five major groups of seed plants (Cycadales, Coniferales, Ginkgoales, Gnetales and angiosperms) that date back as far as the Carboniferous Period about 300 Ma (Jiang, Hinsinger & Strijk, 2016; Wu & Chaw, 2015; Wu *et al.*, 2007). Evidences were plant fossils discovered from late Permian determined to be morphologically-related to extant cycads (Gao & Thomas, 1989). However, Nagalingum *et al.* (2011) considered that species diversification in cycads occurred recently by analyzing PHYTOCHROME P (PHYP) nuclear gene, in which it showed that the age of extant cycads dates back to only 12 million years. But regardless of

various hypotheses, cycads as a key node in the evolutionary phylogeny of green living things remain true.

Cycads are the second largest group of gymnosperms. They comprise putatively three families, 10 genera and 356 extant species (Calonje, Stevenson & Osborne, 2019; Feng et al., 2017; Jiang, Hinsinger & Strijk, 2016). Currently, assessment on the status of cycads based on IUCN Red List reports that 19.8% are near threatened, 23.1% are vulnerable, 20.4% are endangered, 17.3% are critically endangered and 1.2% are already extinct in the wild (IUCN, 2019). Owing to their esthetic nature, cycads are significant in horticulture (Jiang, Hinsinger & Strijk, 2016; Wu & Chaw, 2015). Unfortunately, exploitation for personal and ornamental use, their slow reproductive rate and habitat destruction have resulted in significant declines in cycad populations and poses a threat to the longevity of the plants and to seed production (Pinares et al., 2009; Stevenson, Vovides & Chemnick, 2003). One species currently on the verge of extinction is *Microcycas calocoma* (Miq.) A. DC., a species endemic to western Cuba (north to central of Pinar del Rio Province). It grows in montane forests and on limestone in lowlands and grasslands between 50 and 250 m and is currently listed as critically endangered (IUCN, 2019; Bösenberg, 2010). At present, the monotypic *M. calocoma* is represented by less than 1,000 individuals worldwide. It was previously reported that the female cones are becoming non-receptive to fertilization causing failure in producing viable seeds (Whitelock, 2002). However, this claim still requires research for evidences and validation. Most probable factor to unsuccessful fertilization might be due to the limited cycad-pollinator interactions. *Pharaxonotha* is a taxon of beetles known to pollinate New World Zamiaceae such as *Zamia*, *Dioon* and *Ceratozamia*, wherein the species *Papilio esperanzae* was first described in Cuba as potential pollinator of *M. calocoma* found breeding in male cones and feeds on pollen (Chaves & Genaro, 2005). According to Chaves & Genaro (2005), this discovery paves a way to investigate factors that influence natural reproduction in *Microcycas* that could also address why some populations fail to reproduce. Other threats commonly encountered by cycads species could be factors induced by humans (e.g., intentional private use, over-collection, habitat destruction, grazing, fire, etc.), by climate (e.g., drought, floods, etc.) and by pressures linked to the biology of cycads some of which were already mentioned above such as reproductive failure, dispersal ability and limited number of identified pollinators to facilitate conservation, among others (IUCN, 2019; Mankga & Yessoufou, 2017; Chaves & Genaro, 2005).

Of the 10 accepted genera in Cycadales, only *Microcycas* lacks a completely sequenced chloroplast genome. This study therefore aimed to determine for the first time the whole chloroplast genome of *M. calocoma* to elucidate the variation and evolution in chloroplast genomes in Cycadales.

MATERIALS AND METHODS

DNA extraction, library preparation and sequencing

Five grams of fresh leaves of *M. calocoma* was collected from a plant cultivated in the China National Cycad Conservation Center at Shenzhen Fairy Lake Botanical Garden (Chen T. 2017011701, SZG) for cpDNA isolation using a modified extraction method of

McPherson et al. (2013). After DNA isolation, one µg of purified DNA was fragmented to build short-insert libraries with an insert size of 350 bp following the manufacturer's instructions (Illumina). The short fragments were then sequenced using an Illumina HiSeq 4000 sequencing system at Total Genomics Solution (TGS) in Shenzhen, China (*Borgstrom, Lundin & Lundeberg, 2011*).

Genome assembly

Approximately 1,538 Mb of raw reads with an average read length of 140 bp was generated from paired-end sequencing. Raw sequence reads were then filtered prior to genome assembly to remove adaptor sequences and 1,244 Mb of cleaned data was used to obtain 165,688 bp assembled chloroplast genome. The cp genome of *Microcycas* was reconstructed by combining de novo and reference-guided assemblies (*Cronn et al., 2008*). The following steps were performed: (1) clean sequence reads were assembled into contigs using SOAPdenovo v2.04 (*Li et al., 2010*) (<https://sourceforge.net/projects/soapdenovo2/>) short sequence assembly software, then the assembly results were constructed and optimized according to the paired-end and overlapping reads by filling the gaps and removing redundant sequence segments using GapCloser v1.12 (<http://soap.genomics.org.cn/soapdenovo.html>); (2) for additional guidance in the assembly, BLAST was used to align the contigs to the reference genome *Bowenia serrulata* (GenBank ID: NC_026036.1), which was selected for having the closest sequence length (165,695 bp) with *Microcycas* among available non-Cycadaceae complete chloroplast genome, then aligned contigs (>80% similarity and query coverage) were organized according to the reference genome; and (3) obtained reads were mapped to the assembled draft cp genomes that were visualized by OrganellarGenomeDRAW v1.2 (*Lohse, Drechsel & Bock, 2007*).

Genome annotation and sequence analysis

Using default parameters to predict protein-coding genes, transfer RNA (tRNA) genes, and ribosome RNA (rRNA) genes, annotation of the chloroplast genes was performed using online processing tools, DOGMA (*Wyman, Jansen & Boore, 2004*) and CpGAVAS software (<http://www.herbalgenomics.org/cpgavas/>). A whole chloroplast genome BLAST (*Altschul et al., 1990*) search ($E\text{-value} \leq 1e-5$, minimal alignment length percentage $\geq 40\%$) was performed against 5 databases: KEGG (Kyoto Encyclopedia of Genes and Genomes) (*Kanehisa et al., 2004, 2006; Kanehisa, 1997*), COG (Clusters of Orthologous Groups), NR (Non-Redundant Protein Database databases) (*Tatusov et al., 2003; Tatusov, Koonin & Lipman, 1997*), Swiss-Prot (*Magrane, 2011*), and GO (Gene Ontology) (*Ashburner et al., 2000*) to check and combine annotated genes with their amino acid sequences and corresponding functional annotation information. Finally, the circular *M. calocoma* chloroplast genome map was drawn using OGDRAW v1.2 (*Lohse, Drechsel & Bock, 2007*). The sequence was then deposited and can be viewed in National Center for Biotechnology Information (NCBI) database (<http://www.ncbi.nlm.nih.gov/nuccore>) with accession number MN577566.

Sequence alignment and phylogenetic analyses


Complete chloroplast sequences available in the NCBI database were retrieved and included in the dataset for phylogenetic analyses. All currently available sequenced whole cp genomes under Cycadales and Ginkgoales were obtained. A total of 15 sequences were included in the data set (Table S1).

Eighty-one protein-coding genes encompassing LSC, SSC and IR regions and excluding intergenic spacers were extracted from the plastid genomes followed by alignment of each gene using MAFFT multiple aligner with default parameters in Geneious Prime v2019.0.3 (<https://www.geneious.com>, *Kearse et al., 2012*). Ambiguous regions that have low nucleotide similarities causing large alignment gaps were then manually removed from some genes, after which, all aligned genes were concatenated resulting in total length of 55,971 bp prior to phylogenetic analyses. The data matrix was subjected to maximum likelihood (ML) phylogenetic reconstruction using RAxML v.8 (*Stamatakis, 2014*) with GTR + I + G substitution model selected by jmodeltest2 and 1,000 bootstrap replicates. Bayesian inference (BI) was also calculated using MrBayes 3.2.6 using a GTR model of nucleotide substitution and a gamma distribution rate variation across sites (*Ronquist & Huelsenbeck, 2003*) setting four MCMC running for one million generations with sampling every 1,000 generations and the first 25% discarded as burn-in. Branches with bootstrap values >75 for ML and Bayesian posterior probabilities >0.95 for BI were considered as highly supported. Both ML and BI analyses were conducted in CIPRES Science Gateway (www.phylo.org) (*Miller, Pfeiffer & Schwartz, 2010*). Alignment of sequence identities were plotted using mVISTA online platform (*Frazer et al., 2004; Mayor et al., 2000*). A total of 10 genera of Cycadales were included to analyze similarity profiles on coding and noncoding regions across the genomes. Shuffle-LAGAN alignment program was selected as a parameter prior to run the analysis (*Brudno et al., 2003*) as this setting performs global alignment that detects rearrangements and inversions.

RESULTS

Plastome size and features of *M. calocoma*

The genome size of the complete chloroplast DNA of *M. calocoma* is 165,688 bp in length with 86 protein-coding genes (88,778 bp), 39 tRNA genes (9,050 bp) and eight ribosomal RNA genes (2,906 bp). In *M. calocoma*, the *tufA* gene contains five premature stop codons within the sequence therefore making it a pseudogene. One hundred and thirty three genes were organized in a circular quadripartite structure consisting of large and SSC regions separated by two inverted repeat regions (Fig. 1) composed of 90,651 bp, 22,741 bp and a pair of 26,184 bp regions, respectively. In the entire cp genome of *M. calocoma*, four protein-coding (*ndhB*, *rps7*, *rps12*, *ycf2*), four rRNA (*rrn16*, *rrn23*, *rrn4.5*, *rrn5*) and seven tRNA genes (*trnA*-UGC, *trnH*-GUG, *trnI*-GAU, *trnL*-CAA, *trnN*-GUU, *trnR*-ACG, *trnV*-GAC) were duplicated in the IR regions (Table 1). The protein-coding gene *rps12* was also trans-spliced in the LSC aside from being duplicated in the IRs. The presence of two split genes (*trnK*-UUU/*matK* and *atpB/atpE*) in

Full-size  DOI: [10.7717/peerj.8305/fig-1](https://doi.org/10.7717/peerj.8305/fig-1)

the LSC were also found. The overall GC content of *M. calocoma* (39.6) falls within the range of sequenced chloroplast genomes of cycads which range from 39.4% to 40.1%. LSC, SSC and IR regions have 38.8%, 37.2% and 41.8% GC content, respectively.

Table 1 Genes annotated in *Microcycas calocoma*. List of genes encoded by *Microcycas calocoma*.

Gene category	Gene names	Database accessions
ATP Synthase	<i>atpA, atpB, atpE, atpF, atpH, atpI</i>	GO:0005524 GO:0009507 GO:0015986 GO:0045263 GO:0009507 GO:0016020
NADH dehydrogenase	<i>ndhA, ndhB^(x2), ndhC, ndhD, ndhE, ndhF, ndhG, ndhH, ndhI, ndhJ, ndhK</i>	GO:0003954
Cytochrome b/f complex	<i>petA, petB, petD, petG, petL, petN</i>	GO:0009512
Photosystem I	<i>psaA, psaB, psaC, psaI, psaJ</i>	GO:0009522
Photosystem II	<i>psbA, psbB, psbC, psbD, psbE, psbF, psbH, psbI, psbJ, psbK, psbL, psbM, psbN, psbT, psbZ</i>	GO:0009523
RubisCO large subunit	<i>rbcL</i>	GO:0009507
Ribosomal protein genes (large subunits)	<i>rpl14, rpl16, rpl2, rpl20, rpl22, rpl23, rpl32, rpl33, rpl36</i>	GO:0015934
Ribosomal protein genes (small subunits)	<i>rps11, rps12^(x3), rps14, rps15, rps16, rps18, rps19, rps2, rps3, rps4, rps7^(x2), rps7, rps8</i>	GO:0015935
RNA Polymerase	<i>rpoA, rpoB, rpoC1, rpoC2</i>	GO:0009507 GO:0003677 GO:0009507 GO:0009536
Ribosomal RNA genes	<i>rrn16^(x2), rrn23^(x2), rrn4.5^(x2), rrn5^(x2)</i>	GO:0003735 GO:0003743
Transfer RNA genes	<i>trnA-UGC^(x2), trnC-GCA, trnD-GUC, trnE-UUC, trnF-GAA, trnM-CAU, trnG-GCC, trnG-UCC, trnH-GUG^(x2), trnI-GAU^(x2), trnK-UUU, trnL-CAA^(x2), trnL-UAG, trnM-CAU, trnN-GUU^(x2), trnP-GGG, trnP-UGG, trnQ-UUG, trnR-ACG^(x2), trnR-CCG, trnR-UCU, trnS-GCU, trnS-GGA, trnS-UGA, trnT-GGU, trnT-UGU, trnV-GAC^(x2), trnV-UAC, trnW-CCA, trnY-GUA</i>	GO:0030533 GO:0006412
ATP-dependent protease	<i>clpP</i>	GO:0008462
Maturase	<i>matK</i>	GO:0009536
Hypothetical chloroplast reading frames	<i>ycf1, ycf12, ycf2^(x2), ycf3, ycf4</i>	GO:0016021 GO:0016021 GO:0009507 GO:0009507 GO:0015979
Acetyl-CoA carboxylase	<i>accD</i>	GO:0009536
C-type cytochrome synthesis gene	<i>ccsA</i>	GO:0009507
Envelope membrane protein	<i>cemA</i>	GO:0016020
Photochlorophyllide reductase	<i>chlB, chlL, chlN</i>	GO:0016630
Translational initiation factor	<i>infA</i>	GO:0009536

Plastome features in Cycadales

Thirteen completely-sequenced chloroplast genomes, representing nine of the ten genera of cycads, were in GenBank during our study. The addition of the plastid genome of *M. calocoma* makes known the sequences of all genera of Cycadales. Summary of the genome features in all sequenced cp genomes of cycads was shown in (Table 2).

The smallest genome with 161,815 bp is in *Dioon spinolosum* while *Macrozamia mountperriensis* consisting of 166,341 bp has the largest. Despite significant difference in

Table 2 Gene features of *Microcycas calocoma*. Characteristics of complete chloroplast genomes of Cycadales showing genome lengths, gene numbers and GC content. Data are retrieved from GenBank and accession numbers are indicated.

Species (Accession No.)	Genome size (bp)	LSC (bp)	SSC (bp)	IRs (bp)	Number of genes					GC content (%)
					Total	Coding	tRNA	rRNA	Pseudo-genes	
<i>Microcycas calocoma</i> (MN577566)	165,688	90,651	22,741	26,148	134	86	39	8	1	39.6
<i>Cycas revoluta</i> (NC_020319.1)	162,489	88,977	23,376	25,068	157	109	39	8	1	39.4
<i>Cycas debaoensis</i> (KM459003)	162,094	88,852	23,088	25,076	133	87	37	8	1	39.4
<i>Cycas debaoensis</i> (KU743927)	162,092	88,854	23,088	25,076	133	87	37	8	1	39.4
<i>Cycas taitungensis</i> (NC_009618.1)	163,403	90,216	23,039	25,074	169	122	38	8	1	39.5
<i>Cycas panzhihuaensis</i> (NC_031413.1)	162,470	88,932	23,488	25,045	157	109	39	8	1	39.4
<i>Stangeria eriopus</i> (NC_026041.1)	163,548	89,850	23,006	25,346	134	81	39	8	6	39.5
<i>Bowenia serrulata</i> (NC_026036.1)	165,695	90,733	23,156	25,903	134	87	38	8	1	39.9
<i>Dioon spinolosum</i> (NC_027512.1)	161,815	88,756	23,355	24,852	135	87	39	8	1	40.1
<i>Encephalartos lehmannii</i> (NC_027514.1)	165,822	90,724	23,302	25,898	135	87	39	8	1	39.9
<i>Macrozamia mountperriensis</i> (NC_027511.1)	166,341	91,171	23,334	25,918	135	87	39	8	1	39.8
<i>Lepidozamia peroffskyana</i> (NC_027513.1)	165,939	90,804	23,299	25,918	135	87	39	8	1	39.9
<i>Ceratozamia hildae</i> (NC_026037.1)	165,733	90,487	22,973	26,137	135	87	39	8	1	39.7
<i>Zamia furfuracea</i> (NC_026040.1)	164,953	90,441	23,228	25,642	135	87	39	8	1	39.7

genome length, both species contain 135 genes (87 protein-coding, 39 tRNA, eight rRNA, one pseudogene) indicating that an increase in plastome size does not denote addition of genes or functions. The number of genes in cycads typically ranges from 133 to 135 except in *C. revoluta* and *C. panzhihuaensis*, both of which have 157 genes, and *C. taitungensis* which has 169 genes, the highest number in cycads. These three species of *Cycas* also have the highest number of protein-coding genes among cycads with 109 genes in *C. revoluta* and *C. panzhihuaensis* and 122 genes in *C. taitungensis*. Variation in the number of tRNAs was also observed. *Cycas debaoensis* has 37, *C. taitungensis* and *B. serrulata* both have 38, while the remaining cycads have 39 tRNAs. In comparison, the number of ribosomal RNA (rRNA), eight, is the same in all cycads.

The majority of cycads have 87 protein-coding genes and one pseudo-*tufA* gene. Interestingly, *Stangeria eriopus* has 81 coding genes and six reported pseudogenes-*chlL*, *chlN*, *rpl23*, *tufA*, *chlB* and *matK* genes (Wu & Chaw, 2015). *M. calocoma* contains one




Figure 2 Sequence identity plots of various genera in Cycadales using mVISTA. Sequence variations among 10 genera in Cycadales using *C. debaoensis* as base reference. Complete chloroplast genomes generated by mVISTA. Regions in pink indicate conserved non-coding sequences, purple are conserved exons, while white-colored regions identify more variable sites. The Y-axis represents percent identity ranging from 50% to 100%. IR junctions are indicated in parentheses to show LSC, SSC and IR regions.

Full-size [DOI: 10.7717/peerj.8305/fig-2](https://doi.org/10.7717/peerj.8305/fig-2)

tufA pseudogene ($\Psi tufA$) which is also in all cycad genomes. Jiang, Hinsinger & Strijk (2016) reported that the $\Psi tufA$ gene of *C. debaoensis* was one base pair longer than in *C. taitungensis* (723 bp). In *M. calocoma*, this pseudogene is 1,233 bp in length.




Figure 3 cpDNA tree obtained using ML and BI. Maximum likelihood (ML) tree and Bayesian inference (BI) tree of concatenated data matrix of 81 protein-coding genes from complete chloroplast genome sequences of 10 genera of Cycadales available in GenBank. *Ginkgo biloba* was used as outgroup. Bootstrap support values are given at the nodes annotated as ML/BI. Branches with bootstrap values >75 for ML and Bayesian posterior probabilities >0.95 for BI were considered as highly supported.

Full-size [DOI: 10.7717/peerj.8305/fig-3](https://doi.org/10.7717/peerj.8305/fig-3)

Sequence variation in cycads

Using one representative per genus, mVISTA produced a plot showing peaks and valleys where pink regions denote conserved non-coding sequences (CNS). White regions indicate sites within the genome that are indistinctly aligned, suggesting more variable sites. These regions happen to coincide with the inverted repeat regions of all chloroplast genomes suggesting that IRs are prone to polymorphisms in comparison with the LSC and SSC regions. High degree of synteny in the coding regions was observed across all genomes of Cycadales when *C. debaoensis* was used as a base reference, as indicated by the purple-colored blocks (conserved exons) for all genomes (Fig. 2). Due to the lack of other representative complete chloroplast genomes under Zamiaceae, a *Cycas* species was used as reference. Since a *Cycas* species was used as base reference, only *C. panzhihuaensis* showed an obviously unique pattern indicating that it shares more similar regions with the reference as they are more closely related, setting it apart from the identity plots of the rest of the species which all belong to Zamiaceae. Divergence in CNS regions was higher whereas variations in coding regions were less discernible. Most divergent regions were apparent in the intergenic spacers among all cp genomes. As shown in our alignment, these highly varied sites include regions between *psbA-matK*, *petA-psbJ*, *petL-psbE*, *ndhF-ccsA*, among others. Nonetheless, these plots suggest high similarities in the nucleotide sequences within Cycadales. It also denotes that slow evolution occurs in Cycadales where gene order is highly conserved and a reduced amount of gene rearrangements occur.

Phylogenetic analyses

The phylogeny obtained from the complete chloroplast sequences of cycads using ML and BI revealed several consistencies from previous studies employing various gene segments

from nuclear and/or chloroplast DNA (Fig. 3). Both trees obtained showed significant support (>70 for ML and >0.95 Bayesian Posterior Probability for BI) and consistency in tree topologies. These include the monophyly of the three major clades: (1) all *Cycas* spp., (2) the Encephalartoid clade or “Southern Hemisphere clade” consisting of *Encephalartos*-*Lepidozamia*-*Macrozamia* genera, and (3) the “Neotropical clade” comprising of *Ceratozamia*-*Microcycas*-*Zamia* group. Moreover, *Dioon* was confirmed as sister to the rest of Zamiineae (Liu et al., 2018; Lu et al., 2014; Gorelick et al., 2014; Salas-Leiva et al., 2013).

Phylogenetic analyses also showed that *Microcycas* is closer to *Zamia* than to other genera of cycads which is congruent with previous studies using chloroplast *matK* and *rbcL* genes, *trnK* intron, nuclear rDNA ITS gene and concatenated nuclear and plastid DNA sequences (Lu et al., 2014; Salas-Leiva et al., 2013; Chaw et al., 2005; Hill et al., 2003). Furthermore, the monogeneric Cycadaceae forms the basal-most clade wherein *C. panzhihuaensis* and the sisters *C. taitungensis* and *C. revoluta* diverged from *C. debaoensis*. The phylogenetic placements of *Bowenia* and *Stangeria* were controversial, where older references recognize that they belong to the same family Stangeriaceae (Stevenson, 1992; Stevenson, 1981). In our study, these two genera separated into different clades. Even though showing weaker ML support, *Bowenia* (72 ML and 1.0 BPP) grouped with the Encephalartoid clade while *Stangeria*, with significant support (90 ML and 1.0 BPP), was sister to *Ceratozamia* clustered in the Neotropical clade. As the results suggest that the African cycad *Stangeria* is closest to Neotropical cycads instead of its supposed closest relatives in terms of morphology (*Bowenia*) and biogeographic location (*Encephalartos*), this interesting classification will be further discussed below, as well as the placement of *Bowenia* with the Southern Hemisphere cycads.

DISCUSSION

Plastomes of cycads

Based on previously-completed sequences, 100–130 unique genes are contained within chloroplast genomes of cycads (Jiang, Hinsinger & Strijk, 2016; Wu & Chaw, 2015; Wu et al., 2007). Approximately 79 of these genes code for proteins involved in metabolic pathways for photosynthesis whereas the remaining 30 are responsible for coding transfer RNAs (tRNAs) and four are responsible for ribosomal RNAs (rRNAs) (Hong et al., 2017; Dong et al., 2013; Yang et al., 2013). In this study, the whole chloroplast genome of *Microcycas calcoma* was presented and 13 complete chloroplast genomes of cycads representing nine genera within Cycadales were used for comparison. Hence, all the ten genera within Cycadales now have at least one species with complete chloroplast genome.

The gene organization, content and order in cycad genomes appear to be highly conserved in all ten genera. This agrees with previous reports (Rai et al., 2003; Wu et al., 2007) and supports the slow evolution of cycads in which the changes that occur in the genomes are very minimal since they diverged in the early Jurassic (Nagalingum et al., 2011; Wu & Chaw, 2015). Our study showed that even the universal barcode loci *matK* and *rbcL* are not highly variable in cycads. On the contrary, we observed that *ycf1*, *ycf2* and *accD* genes are highly polymorphic sites across all cycad plastomes.

The nonfunctional gene, *tufA*, was inherited from the charophyte lineage of green algae that once coded for a functional protein elongation factor (Wu et al., 2007). Functional *tufA* genes are present in the cpDNA of algae and bacteria; and are also known to be transferred from the cpDNA to the nuclear ribosomal DNA in rice, legumes, tobacco and *Arabidopsis* (Wu et al., 2007; Sugita, Murayama & Sugiura, 1994; Baldauf & Palmer, 1990). Pseudo-*tufA* genes are also in the genomes of *Anthoceros* (hornwort) and *Ginkgo*. In cycads, the highest frequency of pseudonized genes (6) were present in *S. eriopus* (Wu et al., 2007).

Sequence variations and evolution in Cycadales

Sequence identity plots of cycad genomes identified highly conserved coding (purple) regions. Non-coding (pink) regions appeared more divergent whereas white areas mostly coinciding with IR regions also showed less sequence similarities. According to Wu & Chaw (2015), gene conversions frequently occur in IRs, which favors a GC-biased (gBGC) substitution mechanism. gBGC is believed to be the mechanism employed by cycads in amending plastome-wide mutations, therefore playing a role in evolutionary stasis and genome stability of chloroplast genomes (Wu & Chaw, 2015). The results of the global sequence alignment agreed with the phylogeny employing 81 protein-coding genes, obtained using ML and BI with high support values (>70% and = 1.0 BPP). It showed that the three major clades that grouped together had common identity profiles, thus denoting close relationships. These three are the *Cycas* group, the Australian genus *Bowenia* joining the Southern Hemisphere group (*Encephalartos-Lepidozamia-Macrozamia*), and South African genus *Stangeria* joining the Neotropical group (*Ceratozamia-Microcycas-Zamia*). Geographical patterns and distribution of cycads coincide with our phylogenetic analyses using cp genomes but with few exceptions discussed in the succeeding sections such as the case of African genera *Stangeria* and *Encephalartos* which were separated into different clades.

The use of whole chloroplast genomes shows a clear delineation separating Cycadaceae from the rest of the cycads. But for Zamiaceae, numerous studies have shown that the taxonomic placement of *Bowenia*, *Stangeria* and *Dioon* was ambiguous and their phylogeny remains unresolved (Gorelick et al., 2014; Bogler & Francisco-Ortega, 2004). In this study, *Dioon* was basal among the three genera, *Bowenia* formed a monophyletic group with *Encephalartos-Lepidozamia-Macrozamia* and *Stangeria* was sister to *Ceratozamia* forming a monophyletic clade with *Microcycas* and *Zamia*. This is congruent with other phylogenies wherein *Dioon* separated from other genera and contained two major clades (Salas-Leiva et al., 2013; Bogler & Francisco-Ortega, 2004; Hill et al., 2003). Morphological synapomorphies that can be correlated in our cpDNA topology include the presence of stomata on the sporangia of all Zamiaceae cycad species (Dehgan, Schutzman & Almira, 1993); *Encephalartos*, *Lepidozamia* and *Macrozamia* all have lateral lobes in the megasporophylls (Stevenson, 1990); *Microcycas* and *Zamia* have asymmetrically-branched trichomes (Stevenson, 1990); *Encephalartos* and *Lepidozamia* have a common mucilage chemistry not found in other cycads (De Luca et al., 1982); and the Australian *Macrozamia* and African *Encephalartos* both having pith structures intersected by gum canals and irregular cauline vascular bundles, in contrast to the

peduncular vascular bundles found in *Dioon*, *Stangeria*, *Ceratozamia* and *Zamia* (Berry, 1918). Moreover, *Stangeria* and *Bowenia* share few similarities with some members of their respective clades such as the formation of tracheids in *Stangeria* and *Zamia* are scalariform, and the orientation of vascular bundles in the tertiary cambium of *Bowenia* and *Macrozamia* are both inverted (Govil, 2014; Berry, 1918).

Stevenson (1981) previously described a family Boweniaceae for the genus *Bowenia* but striking morphological similarities eventually unite *Bowenia* and *Stangeria* in Stangeriaceae (Hill et al., 2003; Stevenson, 1992). These characters similar to both genera are vascularized stipules, amphivasal cotyledon bundles, leaflet traces coming from more than one rachis bundle, uneven production of cataphylls, ovules connected under the megasporophyll stalk, and purple sarcotesta. Although morphological trees group these two genera together, molecular data always show weak affinity of the two suggesting that these characters evolved independently from the two genera (Hill et al., 2003; Bogler & Francisco-Ortega, 2004). A study by Griffith et al. (2012) using combined DNA sequences, morphology and phenology data also did not show evidences of strong connection between these two genera. *Bowenia* formed a distinct clade second to *Dioon* within Zamiaceae whereas *Stangeria* was embedded in a clade with *Ceratozamia*, *Zamia* and *Microcycas* (Griffith et al., 2012). The results of our study suggest that *Bowenia* and *Stangeria* are indeed taxonomically distant from each other despite having similar morphological features and being placed previously in the same family Stangeriaceae. The topology obtained in our study therefore supports the formal classification of only two families, Cycadaceae and Zamiaceae, within extant Cycadophyta (Christenhusz et al., 2011). The previous placement of *Bowenia* and *Stangeria* under the same family was in fact inaccurate (Gorelick et al., 2014; Salas-Leiva et al., 2013; Chaw et al., 2005; Bogler & Francisco-Ortega, 2004; Hill et al., 2003) based on molecular data.

Differences in chromosome number and karyotype were also linked to the phylogenetic separation of *Bowenia* ($2n = 18$) and *Stangeria* ($2n = 16$) (Rastogi & Ohri, 2019; Moretti, 1990). This also provides an explanation to the phylogenetic affinity of *Stangeria* to *Ceratozamia* as they had the most similar karyotypes in Cycadales both having 12 metacentric, two submetacentric and two telocentric chromosomes (Kokubugata et al., 2001). Mapping the distribution of 18S and 26S rDNA sites in the somatic chromosomes of both genera also revealed similar dispersed patterns (Kokubugata, Hill & Kondo, 2002). In addition, mapping of 5S rDNA site is similarly situated at the interstitial region of two metacentric chromosomes. However in *Ceratozamia*, it was positioned proximal to the center, whereas in *Stangeria*, it was located near the distal end. This indicates that at least one paracentric inversion might have occurred in the past evolution of these two genera (Kokubugata, Vovides & Kondo, 2004).

Salas-Leiva et al. (2013) were not able to specify morphological synapomorphies that support the relation of *Stangeria* with Neotropical cycads, as well as *Bowenia* with Southern Hemisphere cycads. This led us to search for shared morphological characters that would unite *Stangeria* and *Bowenia* with their respective clades in our whole cpDNA tree. One similarity observed is the obscurity of veins in the leaflets of *Bowenia* and Southern Hemisphere cycads, whereas in the clade comprising *Stangeria* and Neotropical

cycads, the veins tend to be more embossed. However, this needs further validation especially in some large genera such as *Encephalartos* and *Zamia* as they seem to exhibit more diversity in articulation of veins. Another shared characteristic is the type of stem wherein cycads can be divided into two categories. In *Dioon*, *Stangeria*, *Ceratozamia*, *Zamia* and *Microcycas*, the stem is monoxyle characterized by narrow cambiums with only one ring of secondary xylem and no apparent successive growth. Whereas in *Bowenia*, *Encephalartos*, *Lepidozamia* and *Macrozamia*, also including *Cycas*, the stem is polyxyle characterized by multiple layers of cortical cambiums and secondary thickening (Govil, 2014; Berry, 1918; Chamberlain, 1912). Moreover, Dehgan & Dehgan (1988) studied the pollen morphology within Cycadales and determined that *Bowenia* resembled a psilate exine surface and have relatively thin walls and similar alveolar arrangement with the pollen of all Southern Hemisphere cycads. On the other hand, *Stangeria* shares a foveolate exine surface and relatively thick walls with all Neotropical cycads. These characteristics can also be attributed to the pollination biology of Cycadales. Norstog (1987) hypothesized three pollination modes in cycads: anemophily (pollination by wind), entomophily (pollination by insects), and amphiphily (anemophily succeeded by entomophily). Although all genera are likely to be amphiphilous, taxa whose pollen surface lack ornamentation and walls are relatively thin (*Macrozamia*, *Lepidozamia*, *Encephalartos*, *Bowenia*) are presumably anemophilous as the smooth, non-sticky and thin surface makes the pollen easier to be dispersed by wind (Dehgan & Dehgan, 1988; Norstog, 1987; Thanikaimoni, 1986). On the contrary, cycads whose pollen surface consist of coarse pits and walls are relatively thick (*Ceratozamia*, *Dioon*, *Microcycas*, *Stangeria*, and *Zamia*) are probably entomophilous as the rough and thick surface seem to be more attuned to pollination by insects (Dehgan & Dehgan, 1988; Norstog, 1987; Tang, 1987; Punt, 1986). In addition, pollen grain morphology of *Dioon* and *Stangeria* were found to be highly similar having subcircular outline and foveolate exine surface. Aside from having a single vascular ring, both genera also has a common leaflike megasporophyll which was presumed to be a shared primitive feature that was conserved independently (Dehgan & Dehgan, 1988).

The African genera *Encephalartos* and *Stangeria* were separated into different clades suggesting that their evolutionary origin may have occurred before South American, African and Australian continents split (Hill et al., 2003) that accounts for their shared morphological and molecular features with other genera in Zamiaceae. The morphological similarities discussed in the previous section somehow provide biogeographical linkage between the African *Stangeria* and the cycads endemic to the Americas (*Dioon*, *Ceratozamia*, *Microcycas* and *Zamia*). In our hypothesis, the lack of diverse morphological synapomorphies to resolve taxonomic placements of *Dioon*, *Stangeria* and *Bowenia* might be due to the fact that only a number of extant cycads remain and the lack of these presumed extinct species are causing discrepancies in consolidating biogeographical, morphological and molecular phylogenies. Hence we place an emphasis on the hypothesis that around 200-135 MYA in Gondwana (Bogler & Francisco-Ortega, 2004; Jokat et al., 2003), the southern part of Pangaea where Africa, the Americas, Antarctica, India and Australia were once connected, there might exist other ancestral cycads species common

between (1) *Dioon-Stangeria*, (2) *Stangeria-Ceratozamia*, (3) *Stangeria-Zamia-Microcycas*, (4) *Stangeria-Bowenia*, and (5) *Encephalartos-Lepidozamia-Macrozamia*. To support this theory, the latter two would therefore require common ancestor/s situated in Antarctica wherein the shift in climate and environment after fragmentation of the land masses caused eventual extinction of the presumed cycad ancestors. This theory is very likely as cycad fossils from the genus *Eostangeria* was already discovered in North America and Europe, believed to have existed before the final division of Pangaea supercontinent (Uzunova, Palamarev & Kvacek, 2001; Kvacek & Manchester, 1999). A fossil cycad taxa, *Restrepophyllum*, was also discovered in Patagonia, Argentina and found to be morphologically similar to *Zamia* as well as few shared characters with *Stangeria*. This suggests a possible northward migration of *Zamia* from South America and that members of *Zamiaceae* and putative *Stangeriaceae* were widespread in Patagonia during the Early Cretaceous (Passalia, Del Fueyo & Archangelsky, 2010). Other fossil cycad taxa include genera *Dioonopsis* found in Northeast Japan (Horiuchi & Kimura, 1987) and western North America (Erdei, Manchester & Kvaček, 2012), *Encephalartites* (Takimoto & Ohana, 2016) in Northeast Japan, *Crossozamia* in Taiyuan, China (Gao & Thomas, 1989), *Antarcticycas* in Antarctica (Smoot, Taylor & Delevoryas, 1985) and *Zamia* fossil species discovered in Panama (Erdei et al., 2018). Most of these fossil discoveries were found to be morphologically linked to extant cycads under *Zamiaceae*, providing plausible explanations to the diverse morphological characters observed in *Zamiaceae* species compared to the more conserved characters in *Cycadaceae*. Phylogenetic studies were mostly restricted on extant cycads and these lost ancestors might be the missing links that could bridge the gap on the taxonomic issues of *Dioon* and the affinity of *Stangeria* and *Bowenia* with Neotropical cycads and Southern Hemisphere cycads, respectively, despite their differences in distribution and morphology. Only upon integration of extinct and extant cycads will we gain profound understanding on this intriguing group of plants.

CONCLUSION

The use of complete chloroplast genome proved to be advantageous in the phylogenetic reconstruction of *Cycadales* due to its conserved genomic structure and arrangement. Likewise, it holds sufficient genetic information to discriminate variation between species. Here, we present the 165,688 bp sequence of the whole chloroplast genome of *M. calocoma* which is within the range of all previously sequenced cycad genomes. The phylogenetic analysis of *Cycadales* confirmed that *Microcycas* is closely related to *Zamia* and sister to *Ceratozamia*, and that *Stangeria* and *Bowenia* are in different groups; thus stressing that their previous placement in a single family was definitely inappropriate. Hence, our resulting taxonomic classification using chloroplast genomes are consistent with all other molecular phylogenies of cycads (Salas-Leiva et al., 2013; Christenhusz et al., 2011; Bogler & Francisco-Ortega, 2004; Hill et al., 2003).

Furthermore, sequence variation analyses showed that all cycad genomes share highly similar genomic profiles confirming their slow evolution and reduced mutations in comparison with other gymnosperms and seed plants. No rearrangements in the genome structure were detected. This was expected as cycads are known to have a static

evolutionary history, undergoing few changes in morphology since the Jurassic Period (Wu & Chaw, 2015; Nagalingum et al., 2011). However, the common identity profiles were shared less in the inverted repeat regions as well as non-coding sequence regions, suggesting their potential use as regions of interest in studying cycad evolution and phylogeny. In Zamiaceae, only a single species representing each genus has a completely-sequenced chloroplast genome. Thus, adding more sequences to large genera, such as *Zamia* and *Encephalartos*, is crucial to validate existing phylogenetic data and to bridge knowledge gaps regarding cycads species divergence in land plant evolution. Furthermore, the need to expand our knowledge on the field of paleobotany, pollination biology and anatomical and morphological characterization of cycads opens up research opportunities to address incongruence between morphological and molecular phylogenies in Cycadales.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research was supported by the special financial project from the National Forestry and Grassland Administration for Rescues and Breeding of Wild Rare and Endangered Species (Li Nan 2017, 2018, 2019), the Shenzhen Urban Management Bureau (Li Nan 201411) and the Shenzhen Key Laboratory of Tropical and Subtropical Plant Diversity, Fairy Lake Botanical Garden, Chinese Academy of Sciences (Chen Tao 2017–2019), and South China Botanical Garden (Duan Jun 2017–2019), Chinese Academy of Sciences. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Forestry and Grassland Administration for Rescues and Breeding of Wild Rare and Endangered Species: 2017, 2018, 2019.

Shenzhen Urban Management Bureau: 201411.

Shenzhen Key Laboratory of Tropical and Subtropical Plant Diversity, Fairy Lake Botanical Garden, Chinese Academy of Sciences: 2017–2019.

South China Botanical Garden: 2017–2019.

Chinese Academy of Sciences.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Aimee Caye G. Chang conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Qiang Lai performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

- Tao Chen conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Tieyao Tu performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Yunhua Wang conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Esperanza Maribel G. Agoo analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Jun Duan conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Nan Li conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The complete chloroplast genome of *Microcycas calocoma* is available at GenBank: [MN577566](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8305#supplemental-information>.

REFERENCES

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215(3):403–410 DOI 10.1016/S0022-2836(05)80360-2.
- Asaf S, Khan AL, Khan AR, Waqas M, Kang SM, Khan MA, Lee SM, Lee IJ. 2016. Complete chloroplast genome of *Nicotiana otophora* and its comparison with related species. *Frontiers in Plant Science* 7(843):1–12 DOI 10.3389/fpls.2016.00843.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene ontology: tool for the unification of biology. *Nature Genetics* 25(1):25–29 DOI 10.1038/75556.
- Baldauf SL, Palmer JD. 1990. Evolutionary transfer of the chloroplast tufA gene to the nucleus. *Nature* 15:262–265.
- Berry EW. 1918. Paleobotany: a sketch of the origin and evolution of floras. In: Walcott CD, ed. *Report of the Secretary of the Smithsonian Institution*. Washington, D.C.: Government Printing Office, 289–407.
- Bogler DJ, Francisco-Ortega J. 2004. Molecular systematic studies in cycads: evidence from trnL intron and ITS2 rDNA sequences. *Botanical Review* 70(2):260–273 DOI 10.1663/0006-8101(2004)070[0260:MSSICE]2.0.CO;2.
- Borgstrom E, Lundin S, Lundeberg J. 2011. Large scale library generation for high throughput sequencing. *PLOS ONE* 6(4):e19119 DOI 10.1371/journal.pone.0019119.
- Brudno M, Malde S, Poliakov A, Do CB, Couronne O, Dubchak I, Batzoglou S. 2003. Global alignment: finding rearrangements during alignment. *Bioinformatics* 19(Suppl. 1):i54–i62 DOI 10.1093/bioinformatics/btg1005.

- Bösenberg JD. 2010. *Microcycas calocoma*. IUCN Red List of Threatened Species e.T42107A10647674 DOI 10.2305/IUCN.UK.2010-3.RLTS.T42107A10647674.enRLTS.T42107A10647674.en.
- Calonje M, Stevenson DW, Osborne R. 2019. The world list of cycads. Available at <http://www.cycadlist.org>.
- Chamberlain CJ. 1912. Morphology of *Ceratozamia*. *Botanical Gazette* 53(1):1–19 DOI 10.1086/330707.
- Chaves R, Genaro JA. 2005. A new species of *Pharaxonotha* (Coleoptera: Erotylidae), probable pollinator of the endangered Cuban cycad, *Microcycas calocoma* (Zamiaceae). *Insecta Mundi* 19(3):143–150.
- Chaw SM, Walter TW, Chang CC, Hu SH, Chen SH. 2005. A phylogeny of cycads (Cycadales) inferred from chloroplast *matK* gene, *trnK* intron, and nuclear rDNA ITS region. *Molecular Phylogenetics and Evolution* 37(1):214–234 DOI 10.1016/j.ympev.2005.01.006.
- Choi KS, Chung MG, Park SJ. 2016. The complete chloroplast genome sequences of three Veroniceae species (Plantaginaceae): comparative analysis and highly divergent regions. *Frontiers in Plant Science* 7(355):1–8.
- Christenhusz MJM, Reveal JL, Farjon A, Gardner MF, Mill RR, Chase MW. 2011. A new classification and linear sequence of extant gymnosperms. *Phytotaxa* 19(1):55 DOI 10.11646/phytotaxa.19.1.3.
- Cronn R, Liston A, Parks M, Gernandt DS, Shen RK. 2008. Multiplex sequencing of plant chloroplast genomes using Solexa sequencing-by-synthesis technology. *Nucleic Acids Research* 36(19):e122 DOI 10.1093/nar/gkn502.
- De Luca P, Moretti A, Sabato S, Gigliano GS. 1982. A comparative study of cycad mucilages. *Phytochemistry* 21(7):1609–1611 DOI 10.1016/S0031-9422(82)85026-7.
- Dehgan B, Dehgan NB. 1988. Comparative pollen morphology and taxonomic affinities in Cycadales. *American Journal of Botany* 75(10):1501–1516 DOI 10.1002/j.1537-2197.1988.tb11224.x.
- Dehgan B, Schutzman BA, Almira F. 1993. Utilization of scanning electron microscopy on the study of surface features in Cycadales. In: Stevenson DW, Norstog KJ, eds. *The Biology, Structure and Systematics of the Cycadales. Proceedings of the Second International Conference on Cycad Biology*. Milton: Palm & Cycad Societies of Australia, 228–235.
- Dong W, Xu C, Cheng T, Zhou S. 2013. Complete chloroplast genome of *Sedum sarmentosum* and chloroplast genome evolution in Saxifragales. *PLOS ONE* 8(10):e77965 DOI 10.1371/journal.pone.0077965.
- Erdei B, Calonje M, Hendy A, Espinosa N. 2018. A review of the Cenozoic fossil record of the genus *Zamia* L. (Zamiaceae, Cycadales) with recognition of a new species from the late Eocene of Panama-evolution and biogeographic inferences. *Bulletin of Geosciences* 93(2):185–204 DOI 10.3140/bull.geosci.1671.
- Erdei B, Manchester SR, Kvaček Z. 2012. *Dioonopsis* Horiuchi et kimura leaves from the Eocene of western North America: a cycad shared with the Paleogene of Japan. *International Journal of Plant Sciences* 173(1):81–95 DOI 10.1086/662654.
- Feng Z, Lv Y, Guo Y, Wei HB, Kerp H. 2017. Leaf anatomy of a late Palaeozoic cycad. *Biology Letters* 13(11):20170456 DOI 10.1098/rsbl.2017.0456.
- Frazer KA, Pachter L, Poliakov A, Rubin EM, Dubchak I. 2004. VISTA: computational tools for comparative genomics. *Nucleic Acids Research* 32:W273–W279 DOI 10.1093/nar/gkh458.
- Gao ZF, Thomas BA. 1989. A review of fossil cycad megasporophylls, with new evidence of *Crossozamia* Pomel and its associated leaves from the lower Permian of Taiyuan. *China Review of Palaeobotany and Palynology* 60(3–4):205–223 DOI 10.1016/0034-6667(89)90044-4.

- Gorelick R, Fraser D, Zonneveld BJM, Little DP. 2014. Cycad (Cycadales) chromosome numbers are not correlated with genome size. *International Journal of Plant Sciences* 175(9):986–997 DOI 10.1086/678085.
- Govil CM. 2014. *Gymnosperms: extinct and extant*. Meerut: Krishan Prakashan.
- Griffith MP, Calonje MA, Stevenson DW, Husby CE, Little DP. 2012. Time, place, and relationships: cycad phenology in a phylogenetic and biogeographic context. *Memoirs of the New York Botanical Garden* 106:59–81.
- Hill KD, Chase MW, Stevenson DW, Hills HG, Schutzman B. 2003. The families and genera of cycads: a molecular phylogenetic analysis of Cycadophyta based on nuclear and plastid DNA sequences. *International Journal of Plant Sciences* 164(6):944–948 DOI 10.1086/378538.
- Hong SY, Cheon KS, Yoo KO, Lee HO, Cho KS, Suh JT, Kim SJ, Nam JH, Sohn HB, Kim YH. 2017. Complete chloroplast genome sequences and comparative analysis of *Chenopodium quinoa* and *C. album*. *Frontiers in Plant Science* 8(1696):1–12.
- Horiuchi J, Kimura T. 1987. *Dioonopsis nipponica* gen. et. sp. nov., a new cycad from the palaeogene of Japan. *Review of Palaeobotany and Palynology* 51(1–3):213–225 DOI 10.1016/0034-6667(87)90031-5.
- IUCN. 2019. *The IUCN Red List of Threatened Species*, Version 2019-2. Available at <https://www.iucnredlist.org>.
- Jiang GF, Hinsinger DD, Strijk JS. 2016. Comparison of intraspecific, interspecific and intergeneric chloroplast diversity in Cycads. *Scientific Reports* 6(1):31473 DOI 10.1038/srep31473.
- Jokat W, Boebel T, König M, Meyer U. 2003. Timing and geometry of early Gondwana breakup. *Journal of Geophysical Research: Solid Earth* 108(B9):2428.
- Kanehisa M. 1997. A database for post-genome analysis. *Trends in Genetics* 13(9):375–376 DOI 10.1016/S0168-9525(97)01223-7.
- Kanehisa M, Goto S, Hattori M, Aoki-Kinoshita KF, Itoh M, Kawashima S, Katayama T, Araki M, Hirakawa M. 2006. From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Research* 34:D354–D357 DOI 10.1093/nar/gkj102.
- Kanehisa M, Goto S, Kawashima S, Okuno Y, Hattori M. 2004. The KEGG resource for deciphering the genome. *Nucleic Acids Research* 32:D277–D280 DOI 10.1093/nar/gkh063.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12):1647–1649.
- Kokubugata G, Hill KD, Kondo K. 2002. Ribosomal DNA distribution in somatic chromosomes of *Stangeria eriopus* (Stangeriaceae, Cycadales) and molecular-cytotaxonomical relationships to the other cycad genera. *Brittonia* 54(1):1–5 DOI 10.1663/0007-196X(2002)054[0001:RDDISC]2.0.CO;2.
- Kokubugata G, Hill KD, Wilson GW, Kondo K, Randall LM. 2001. A comparison of chromosome number and karyotype in somatic chromosomes of *Bowenia* and *Stangeria* (Stangeriaceae, Cycadales). *Edinburgh Journal of Botany* 58(3):475–481 DOI 10.1017/S0960428601000786.
- Kokubugata G, Vovides AP, Kondo K. 2004. Mapping 5S ribosomal DNA on somatic chromosomes of four species of *Ceratozamia* and *Stangeria eriopus* (Cycadales). *Botanical Journal of the Linnean Society* 145(4):499–504 DOI 10.1111/j.1095-8339.2004.00299.x.

- Kvacek Z, Manchester SR. 1999. *Eostangeria* Barthel (extinct Cycadales) from the Paleogene of western North America and Europe. *International Journal of Plant Sciences* 160(3):621–629 DOI 10.1086/314152.
- Lee SB, Kaittani C, Jansen RK, Hostetler JB, Tallon LJ, Town CD, Daniell H. 2006. The complete chloroplast genome sequence of *Gossypium hirsutum*: organization and phylogenetic relationships to other angiosperms. *BMC Genomics* 7(61):1–12 DOI 10.1186/1471-2164-7-1.
- Li R, Zhu H, Ruan J, Qian W, Fang X, Shi Z, Li Y, Li S, Shan G, Kristiansen K, Li S, Yang H, Wang J, Wang J. 2010. De novo assembly of human genomes with massively parallel short read sequencing. *Genome Research* 20(2):265–272 DOI 10.1101/gr.097261.109.
- Liu LX, Li R, Worth JRP, Li X, Li P, Cameron KM, Fu CX. 2017. The complete chloroplast genome of chinese bayberry (*Morella rubra*, Myricaceae): implications for understanding the evolution of Fagales. *Frontiers in Plant Science* 8(968):1–15.
- Liu J, Zhang S, Nagalingum NS, Chiang YC, Lindstrom A, Gong X. 2018. Phylogeny of the gymnosperm genus *Cycas* L. (Cycadaceae) as inferred from plastid and nuclear loci based on large-scale sampling: evolutionary relationships and taxonomical implications. *Molecular Phylogenetics and Evolution* 127:87–97 DOI 10.1016/j.ympev.2018.05.019.
- Lodish H, Berk A, Zipursky SL, Matsudaira P, Baltimore D, Darnell J. 2000. *Molecular cell biology*. Fourth Edition. New York: W. H. Freeman.
- Lohse M, Drechsel O, Bock R. 2007. OrganellarGenomeDRAW (OGDRAW): a tool for the easy generation of high-quality custom graphical maps of plastid and mitochondrial genomes. *Current Genetics* 52(5–6):267–274 DOI 10.1007/s00294-007-0161-y.
- Lu Y, Ran JH, Guo DM, Yang ZY, Wang XQ. 2014. Phylogeny and divergence times of gymnosperms inferred from single-copy nuclear genes. *PLOS ONE* 9(9):e107679 DOI 10.1371/journal.pone.0107679.
- Magrane M. 2011. UniProt knowledgebase: a hub of integrated protein data. *Database: The Journal of Biological Databases & Curation* 2011:bar009 DOI 10.1093/database/bar009.
- Mankga LT, Yessoufou K. 2017. Factors driving the global decline of cycad diversity. *AoB Plants* 9(4):plx022 DOI 10.1093/aobpla/plx022.
- Martin W, Deusch O, Stawski N, Grunheit N, Goremykin V. 2005. Chloroplast genome phylogenetics: why we need independent approaches to plant molecular evolution. *Trends in Plant Science* 10(5):1360–1385.
- Mayor C, Brudno M, Schwartz JR, Poliakov A, Rubin EM, Frazer KA, Pachter LS, Dubchak I. 2000. VISTA: visualizing global DNA sequence alignments of arbitrary length. *Bioinformatics* 16(11):1046–1047 DOI 10.1093/bioinformatics/16.11.1046.
- McPherson H, Van Der Merwe M, Delaney SK, Edwards MA, Henry RJ, McIntosh E, Rymer RD, Milner ML, Siow J, Rossetto M. 2013. Capturing chloroplast variation for molecular ecology studies: a simple next generation sequencing approach applied to a rainforest tree. *BMC Ecology* 13(1):8 DOI 10.1186/1472-6785-13-8.
- Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES science gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop (GCE)*, 14 November 2010, New Orleans, LA, USA. 1–8.
- Moretti A. 1990. Karyotype data on North and Central American Zamiaceae (Cycadales) and their phylogenetic implications. *American Journal of Botany* 77(8):1016–1029 DOI 10.1002/j.1537-2197.1990.tb13597.x.
- Nagalingum NS, Marshall CR, Quental TB, Rai HS, Little DP, Mathews S. 2011. Recent synchronous radiation of a living fossil. *Science* 334(6057):796–799 DOI 10.1126/science.1209926.

- Norstog K. 1987. Cycads and the origin of insect pollination. *American Journal of Science* 75:270–279.
- Passalia MG, Del Fueyo G, Archangelsky S. 2010. An early Cretaceous zamiaceous cycad of South West Gondwana: *Restrepophyllum* nov. gen. from Patagonia, Argentina. *Review of Palaeobotany and Palynology* 161(3–4):137–150 DOI 10.1016/j.revpalbo.2010.04.001.
- Pinares A, Gonzalez-Astorga J, Vovides AP, Lazcano J, Vendrame WA. 2009. Genetic diversity of the endangered endemic *Microcycas calocoma* (Miq.) A. DC (Zamiaceae, Cycadales): implications for conservation. *Biochemical Systematics and Ecology* 37(4):385–394 DOI 10.1016/j.bse.2009.07.006.
- Punt W. 1986. Functional factors influencing pollen form. In: Blackmore S, Ferguson IK, eds. *Pollen and Spores: Form and Function*. Orlando: Academic Press.
- Rai HS, Obrien HE, Reeves PA, Olmstead RG, Graham SW. 2003. Inference of higher-order relationships in the cycads from a large chloroplast data set. *Molecular Phylogenetics and Evolution* 29:350–359.
- Rastogi S, Ohri D. 2019. Karyotype evolution in cycads. Epub ahead of print 5 October 2019. *Nucleus*.
- Ronquist F, Huelsenbeck JP. 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(12):1572–1574 DOI 10.1093/bioinformatics/btg180.
- Salas-Leiva DE, Meerow AW, Calonje M, Griffith MP, Francisco-Ortega J, Nakamura K, Stevenson DW, Lewis CE, Namoff S. 2013. Phylogeny of cycads based on multiple single-copy nuclear genes: congruence of concatenated parsimony, likelihood and species tree inference methods. *Annals of Botany* 112(7):1263–1278 DOI 10.1093/aob/mct192.
- Shaw J, Lickey EB, Schilling EE, Small RL. 2007. Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: the tortoise and the hare III. *American Journal of Botany* 94(3):275–288 DOI 10.3732/ajb.94.3.275.
- Smith DR. 2017. Does cell size impact chloroplast genome size? *Frontiers in Plant Science* 8:1–6.
- Smoot EL, Taylor TN, Delevoryas T. 1985. Structurally preserved fossil plants from Antarctica. I. *Antarcticycas*, gen. nov., a Triassic cycad stem from the Beardmore Glacier Area. *American Journal of Botany* 72(9):1410–1423 DOI 10.1002/j.1537-2197.1985.tb08399.x.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9):1312–1313 DOI 10.1093/bioinformatics/btu033.
- Stevenson DW. 1981. Observations on ptyxis, phenology, and trichomes in the Cycadales and their systematic implications. *American Journal of Botany* 68(8):1104–1114 DOI 10.1002/j.1537-2197.1981.tb06394.x.
- Stevenson DW. 1990. Morphology and systematics of the Cycadales. *Memoirs of the New York Botanical Garden* 57:8–55.
- Stevenson DW. 1992. A formal classification of the extant cycads. *Brittonia* 44(2):220–223 DOI 10.2307/2806837.
- Stevenson DW, Vovides A, Chemnick J. 2003. Regional overview: new world. In: Donaldson JS, ed. *Cycads. Status Survey and Conservation Action Plan*. Gland and Cambridge: IUCN/SSC Cycad Specialist Group, IUCN, 31–38.
- Sugita M, Murayama Y, Sugiura M. 1994. Structure and differential expression of two distinct genes encoding the chloroplast elongation factor Tu in tobacco. *Current Genetics* 25:164–168.
- Takimoto H, Ohana T. 2016. *Encephalartites nipponensis* sp. nov., from the Jurassic Tochikubo Formation (Oxfordian), Somanakamura Group, Northeast Japan. *Paleontological Research* 20(3):261–267 DOI 10.2517/2015PR035.

- Tang W. 1987.** Insect Pollination in the Cycad *Zamia pumila* (Zamiaceae). *American Journal of Botany* **74**(1):90–99.
- Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA. 2003.** The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* **4**(1):41 DOI [10.1186/1471-2105-4-41](https://doi.org/10.1186/1471-2105-4-41).
- Tatusov RL, Koonin EV, Lipman DJ. 1997.** A genomic perspective on protein families. *Science* **278**(5338):631–637 DOI [10.1126/science.278.5338.631](https://doi.org/10.1126/science.278.5338.631).
- Thanikaimoni G. 1986.** Pollen apertures: form and function. In: Blackmore S, Ferguson IK, eds. *Pollen and Spores: Form and Function*. Orlando: Academic Press, 119–136.
- Uzunova K, Palamarev E, Kvacek Z. 2001.** *Eostangeria ruzinciniana* (Zamiaceae) from the middle Miocene of Bulgaria and its relationship to similar taxa of fossil Eostangeria, and extant *Chigua* and *Stangeria* (Cycadales). *Acta Palaeobot* **41**(2):177–193.
- Whitelock LM. 2002.** *The Cycads*. Portland: Timber Press.
- Wu CS, Chaw SM. 2015.** Evolutionary stasis in Cycad plastomes and the first case of plastome GC-biased gene conversion. *Genome Biology and Evolution* **7**(7):2000–2009 DOI [10.1093/gbe/evv125](https://doi.org/10.1093/gbe/evv125).
- Wu CS, Wang YN, Liu SM, Chaw SM. 2007.** Chloroplast genome (cpDNA) of *Cycas taitungensis* and 56 cp protein-coding genes of *Gnetum parvifolium*: insights into cpDNA evolution and phylogeny of extant seed plants. *Molecular Biology and Evolution* **24**(6):1366–1379 DOI [10.1093/molbev/msm059](https://doi.org/10.1093/molbev/msm059).
- Wyman SK, Jansen RK, Boore JL. 2004.** Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* **20**(17):3252–3255 DOI [10.1093/bioinformatics/bth352](https://doi.org/10.1093/bioinformatics/bth352).
- Xiao LQ, Moller M. 2015.** Nuclear ribosomal ITS functional paralogs resolve the phylogenetic relationships of a late-miocene radiation Cycad *Cycas* (Cycadaceae). *PLOS ONE* **10**(1):e0117971 DOI [10.1371/journal.pone.0117971](https://doi.org/10.1371/journal.pone.0117971).
- Yang JB, Tang M, Li HT, Zhang ZR, Li DZ. 2013.** Complete chloroplast genome of the genus *Cymbidium*: lights into the species identification, phylogenetic implications and population genetic analyses. *BMC Evolutionary Biology* **13**(84):1–12 DOI [10.1186/1471-2148-13-1](https://doi.org/10.1186/1471-2148-13-1).
- Zhong Z-R, Li N, Qian D, Jin J-H, Chen T. 2011.** Maternal inheritance of plastids and mitochondria in *Cycas* L. (Cycadaceae). *Molecular Genetics and Genomics* **286**(5–6):411–416 DOI [10.1007/s00438-011-0653-9](https://doi.org/10.1007/s00438-011-0653-9).