

Multichromosomal Mitochondrial Genome of *Punica granatum*: Comparative Evolutionary Analysis and Gene Transformation from chloroplast genomes

Lijuan Feng

Shandong Institute of Pomology

Chuanzeng Wang

Shandong Academy of Agricultural Sciences

Xuemei Yang

Shandong Institute of Pomology

Mengmeng An

Zibo Academy of Agricultural Sciences

Yanlei Yin (✉ yylei66@sina.com)

Shandong Institute of Pomology

Research Article

Keywords: *Punica granatum*, mitogenome, multi-chromosomal structure, evolutionary relationship, intracellular gene transfer

Posted Date: July 10th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3140766/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at BMC Plant Biology on October 25th, 2023. See the published version at <https://doi.org/10.1186/s12870-023-04538-8>.

Abstract

Background *Punica granatum*

is an essentially important fruit tree that has important economic, medicinal and ornamental properties. At present, there are few reports on the mitochondrial genome of pomegranate. To further understand the organization, variation, and evolution of pomegranate mitogenomes, the *P. granatum* mitogenome was sequenced and assembled in the study.

Results

The genome structure was multi-chromosomes with seven circular chromosomes, measuring 382,774 bp in length with a 45.91% GC content. It contained 74 genes, including 46 protein-coding genes, 25 tRNA genes, and three rRNA genes. There were 188 pairs of dispersed repeats with lengths greater than or equal to 30, which were mainly reverse complementary repeats. The mitogenome analysis identified 114SSRs and 466 RNA editing sites. Analyses of codon usage, nucleotide diversity and gene migration from chloroplast to mitochondrial were also conducted. The collinear and comparative analysis of mitochondrial structures between *P. granatum* and its proximal species indicated that *P. granatum* was closely related to *Lagerstroemia indica*. Phylogenetic examination based on the mitogenome also confirmed the evolutionary relationship.

Conclusion

The results provided the predominant information for pomegranate evolutionary biology and to promote the species germplasm utilization.

Introduction

Pomegranate (*Punica granatum* L.) belongs to the genus *Punica* of the Lythraceae family in scientific research, and is an essentially important fruit tree in the world [1–3]. It is native to Central Asia region and grows mainly in tropical and subtropical countries such as India, Iran, China, and the USA [4–5]. The consumption of pomegranates has shown a remarkable increase due to their exceptional economic, nutritional, medicinal and ornamental value. The different parts (fruit peel, juice, seeds, flowers, leaves, bark) are rich in bioactive constituents, and have various medicinal advantages to provide high antioxidant, antineoplastic, anticancer and phytonutrient capacity [6–9]. At present, pomegranate related industries are expanding rapidly, and plays an important role in the economic, cultural and ecological fields.

Mitochondria are semi-autonomous organelles with a mitochondrial genome found in virtually all eukaryotic cells, which synthesize Adenosine Triphosphate (ATP) through the tricarboxylic acid cycle and

oxidative phosphorylation to provide energy for cells [10–11]. Mitochondria are involved in numerous metabolic processes and perform crucial roles in cell differentiation, apoptosis, cell development, and cell division by converting biomass energy into chemical energy for daily activities [12–14]. Plant chloroplast genomes have revealed conserved genome structure, but the mitochondrial genome has very low collinearity due to extensive rearrangement and repeat sequences, whereas its protein-coding genes are relatively conserved [12, 15]. Plant mitogenomes are species specific and possess many unique features, which vary considerably in length, gene order, and gene content [16–17]. The structures of mitogenomes are normally depicted as a complex and dynamic circular structure with a large amount of unconserved DNA of unknown function [18]. In some species, the mitochondrial genomes have linear conformations, branched and multichromosomal structures [14, 19]. The above characteristics of the mitochondrial genome provide helpful information for evolutionary and phylogenetic studies.

The mitochondrial genomes of 545 plant species have been assembled and reported [14], such as *malus domestica* [20], *Satsuma mandarin* [21], and kiwifruit [22], etc. To date, the chloroplast genomes of approximately 114 species of *P. granatum* have been released to the NCBI database (accessed on 28 June 2023), but the mitochondrial genome of only one pomegranate variety has just been identified and published [23]. Therefore, more in-depth researches of the mitochondrial genome of *P. granatum* are of great significance for its effective utilization and genetic improvement.

In this study, we sequenced, assembled and annotated the complete mitochondrial genome of *P. granatum* ‘Taishanhong’. The mitogenome characteristics, repetitive sequences, SSR identification, and RNA editing prediction were investigated. Further analyses regarding species synteny and phylogeny were carried out for the determination of phylogenetic positions and molecular diversity of the myrtales. The gene transformation of chloroplast genome to mitochondrion DNA was analyzed. It was anticipated that current study would serve as a foundation for a deeper comprehension and investigation of evolution variations and genomic breeding of pomegranate mitochondrial genomes.

Results

Genomic features of *P. granatum* mitochondrial genome

The *P. granatum* mitochondrial genome was sequenced, assembled, and annotated, which is now archived in GenBank (accession number OQ973289, OQ973290, OQ973291, OQ973292, OQ973293, OQ973294 and OQ973295). The mitochondrial genome of *P. granatum* was assembled into 7 circular chromosomes with lengths ranging from 3,728 bp to 126,124 bp, with a total length of 382,774 bp (Fig. 1 and Fig. S1). The total GC content was 45.91%, ranging between 45.42% and 52.60% among chromosomes (Table S1). It encoded 74 genes, including 46 mitochondrial protein-coding genes, 25 tRNA genes, and three rRNA genes (*rrn5*, *rrn18*, and *rrn26*). The total GC contents of CDS, tRNA and rRNA genes were 42.35%, 50.85% and 52.14%, respectively (Table S1).

The protein-coding genes included 34 core genes and nine variable genes (Table 1). The core genes included five ATP synthase genes (*atp1*, *atp4*, *atp6*, *atp8* and *atp9*), nine NADH dehydrogenase genes

(*nad1*, *nad2*, *nad3*, *nad4*, *nad4L*, *nad5*, *nad6*, *nad7* and *nad9*), four cytochrome c biogenesis genes (*ccmB*, *ccmC*, *ccmFC* and *ccmFN*), three cytochrome c oxidase genes (*cox1*, *cox2* and *cox3*), one Maturases gene (*matR*), one membrane transport protein gene (*mttB*) and one ubiquinol cytochrome c reductase gene (*cob*). The variable genes included four large subunits of ribosomal protein (*rpl10*, *rpl16*, *rpl2* and *rpl5*), seven small subunits of ribosomal protein (*rps1*, *rps3*, *rps4*, *rps7*, *rps12*, *rps14* and *rps19*), and one succinate dehydrogenase (*sdh4*). Among these, the genes *atp9*, *mttB*, *nad2* and *nad5* had two copies. The *nad1*, *nad2*, *nad5* and *nad7* genes contained four introns and *nad4* gene had three introns, while *ccmFC*, *cox2*, *rpl2* and *rps3* had one intron. The *trnC-GCA*, *trnM-CAT*, *trnP-TGG* and *trnS-TGA* genes were multi-copy genes. The *trnA-TGC*, *trnI-GAT* and *trnI-TAT* genes contained one intron.

Table 1
Gene composition in the mitogenome of *P. granatum*

Group of Genes	Gene name	Length	Start codon	Stop codon	Amino acid
ATP synthase	atp1	1536	ATG	TAA	512
	atp4	597	ATG	TAG	199
	atp6	747	ATG	TGA	249
	atp8	480	ATG	TAA	160
	atp9(2)	225	ATG	CGA(TGA)	75
Cytochrome c biogenesis	ccmB	621	ATG	TGA	207
	ccmC	753	ATG	TGA	251
	ccmFc	1356	ATG	TAA	452
	ccmFn*	1737	ATG	TAA	579
Ubichinol cytochrome c reductase	cob	1890	ATG	TAA	630
Cytochrome c oxidase	cox1	1065	ATG	TAA	355
	cox2*	783	ATG	TAA	261
	cox3	798	ATG	TGA	266
Maturases	matR	1926	ATG	TAA	642
Transport membrane protein	mttB(2)	342	ATG	TGA	114
NADH dehydrogenase	nad1****	978	ACG(ATG)	TAA	326
	nad2(2)****	1467	ATG	TAA	489
	nad3	357	ATG	TAA	119
	nad4***	1488	ATG	TGA	496
	nad4L	303	ATG	TAA	101
	nad5(2)****	2013	ATG	TAA	671
	nad6	618	ATG	TAA	206
	nad7****	1185	ATG	TAG	395
	nad9	573	ATG	TAA	191

Note: Numbers after gene names are the number of copies. The superscripts * represent the number of contained introns.

Group of Genes	Gene name	Length	Start codon	Stop codon	Amino acid
Ribosomal proteins (LSU)	rpl10	489	ATG	TAA	163
	rpl16	249	ATG	TAA	83
	rpl2	1014	ATG	TAA	338
	rpl5	564	ATG	TAA	188
Ribosomal proteins (SSU)	rps1	624	ATG	TAA	208
	rps12*	378	ATG	TGA	126
	rps14	303	ATG	TAG	101
	rps19	285	ATG	TAA	95
	rps3	1710	ATG	TAA	570
	rps4	1047	ATG	TAA	349
	rps7	447	ATG	TAA	149
	sdh4	387	ATG	CGA(TGA)	129
Ribosomal RNAs	rrn18	1929	-	-	-
	rrn26	3403	-	-	-
	rrn5	121	-	-	-
Transfer RNAs	trnA-TGC*	65	-	-	-
	trnC-GCA(2)	71/73	-	-	-
	trnD-GTC	74	-	-	-
	trnE-TTC	72	-	-	-
	trnF-GAA	74	-	-	-
	trnG-GCC	72	-	-	-
	trnH-GTG	74	-	-	-
	trnI-GAT*	72	-	-	-
	trnI-TAT*	76	-	-	-
	trnK-TTT	73	-	-	-

Note: Numbers after gene names are the number of copies. The superscripts * represent the number of contained introns.

Group of Genes	Gene name	Length	Start codon	Stop codon	Amino acid
	trnM-CAT(3)	74/73/74	-	-	-
	trnN-GTT	72	-	-	-
	trnP-TGG(3)	74/75/75	-	-	-
	trnQ-TTG	72	-	-	-
	trnS-GCT	88	-	-	-
	trnS-TGA(2)	93/87	-	-	-
	trnV-GAC	72	-	-	-
	trnW-CCA	74	-	-	-
	trnY-GTA	83	-	-	-
Note: Numbers after gene names are the number of copies. The superscripts * represent the number of contained introns.					

Condon usage and RSCU analysis

The ATG was the most frequent start codon for protein-coding genes, while the *nad1* gene was exception with the initiating codon ACG (ATG). The stop codons TAA, TAG, TGA, and CGA (TGA) were identified (Table 1). These results indicated that the C to U RNA editing phenomenon was found in the start or stop codons.

The relative synonymous codon usage (RSCU) analysis of *P. granatum* mitochondrial genome was shown in Fig. 2. It contained 10, 445 codons excluding termination codons in protein-coding genes regions. The most frequent codons used were UUU (Phe), AUU (Ile) and GAA (Glu) and were used 300 times, while CUG (Met), UUG (Met) and UAG (Ter) were rarely found (Table S2). We found that most RSCU values of the codons ending with A or T were higher than 1.0, while most of those ending with C or G had RSCU values of less than 1. Codon usage was generally strongly biased toward A or T (U) at the third codon position in the *P. granatum* mitogenome. Similar results had been found in the mitogenome of other plants [17, 24].

Repeat sequence analysis

A total of 188 pairs of repeats with a length greater than or equal to 30 bp were found, including 97 pairs of reverse complementary repeats, 91 pairs of forward repeats. The longest reverse complementary repeat was 355 bp, while the longest forward repeat was 24,984 bp (Table S3).

The simple repeated sequences (SSRs) with motifs of one to six bases are abundant in higher plants. In the study, a total of 141SSRs were detected in the *P. granatum* mitogenome, including 58 (41.13%) mono-, 19 (13.48%) di-, 18 (12.77%) tri-, 35(24.82%) tetra-, 9 (6.38%) penta-, and 2 (1.42%) hexanucleotide repeats (Fig. 3A). Among them, monomeric and dimeric SSRs accounted for 54.61% of the total SSRs. Mononucleotide repeats of A/T (34.75%) were more prevalent than the other repeat types, and dinucleotide repeats of AG/CT (7.80%) were the second most numerous.

Prediction of RNA editing sites

RNA editing events were identified for 36 unique PCGs based on online website predictions. There were 466 potential RNA editing sites distributed among all PCGs, and all of which were C-to-U base editing (Fig. 3B and Table S4). The *nad4* gene was the most edited and possessed 39 potential RNA editing sites of the mitochondrial genes. This was followed by the *ccmB* gene with 36 RNA editing events. The edited number of *rps1* gene was the lowest and had only one potential RNA editing events among all mitochondrial genes.

The total number of hydrophilic-hydrophobic type induced by RNA editing was 224 sites, which had the highest proportion at 48.07% (Table 2). The hydrophobic-hydrophobic and hydrophilic-hydrophilic type was 31.33% (146 sites) and 12.23% (57 sites), respectively. The hydrophilic-stop number was the lowest and the proportion was 1.07% (5 sites). Among them, there were 10 site conversions of CCT to TTT and 8 conversions of CCC to TTC.

Table 2
Prediction of RNA editing sites in *P. granatum* mitogenome

Type	RNA-editing	Number	Percentage
hydrophilic-hydrophilic	CAC (H) = > TAC (Y)	6	
	CAT (H) = > TAT (Y)	16	
	CGC (R) = > TGC (C)	6	
	CGT (R) = > TGT (C)	29	
	total	57	12.23%
hydrophilic-hydrophobic	ACA (T) = > ATA (I)	5	
	ACC (T) = > ATC (I)	1	
	ACG (T) = > ATG (M)	5	
	ACT (T) = > ATT (I)	4	
	CGG (R) = > TGG (W)	32	
	TCA (S) = > TTA (L)	71	
	TCC (S) = > TTC (F)	30	
	TCG (S) = > TTG (L)	38	
	TCT (S) = > TTT (F)	38	
	total	224	48.07%
hydrophilic-stop	CAA (Q) = > TAA (X)	3	
	CGA (R) = > TGA (X)	2	
	total	5	1.07%
hydrophobic-hydrophilic	CCA (P) = > TCA (S)	6	
	CCC (P) = > TCC (S)	8	
	CCG (P) = > TCG (S)	3	
	CCT (P) = > TCT (S)	17	
	total	34	7.30%
hydrophobic-hydrophobic	CCA (P) = > CTA (L)	36	
	CCC (P) = > CTC (L)	12	
	CCC (P) = > TTC (F)	8	
	CCG (P) = > CTG (L)	29	

Type	RNA-editing	Number	Percentage
	CCT (P) = > CTT (L)	23	
	CCT (P) = > TTT (F)	10	
	CTC (L) = > TTC (F)	8	
	CTT (L) = > TTT (F)	12	
	GCA (A) = > GTA (V)	1	
	GCG (A) = > GTG (V)	5	
	GCT (A) = > GTT (V)	2	
	total	146	31.33%
	All	466	100%

Nucleotide diversity and comparative analysis of mitochondrial structure

The nucleotide diversity (π) values of 38 regions were calculated and ranged from 0 to 0.10582, with an average of 0.03132 (Fig. 4A and Table S5). The π value of *gene10.nad9* region was highest in these regions, which was 0.07259 and 0.0698 in *gene37.atp9* and *gene33.rnr18* regions, respectively. The lower π values suggested that the mitogenome sequences of *P. granatum* were highly conserved.

The mitogenome structures of *P. granatum* and its proximal species belonged to the Myrtales were comparative analyzed using CGVIEW software (Fig. 4B). There were high similarities in the mitochondrial structure of *P. granatum* and *Eucalyptus grandis* (NCBI Number: MG925370.1), *Lagerstroemia indica* (NCBI Number: KX641464.1) and *Medinilla magnifica* (NCBI Number: MT043351.1).

Phylogenetic and collinearity analysis

The mitochondrial genome of proximate species of pomegranate is rarely reported. To further explore the evolutionary relationships of *P. granatum*, the phylogenetic tree was constructed by maximum likelihood method (Fig. 5). Phylogenetic analysis was performed on 30 conserved mitochondrial PCGs (*atp1*, *atp4*, *atp6*, *atp8*, *atp9*, *ccmC*, *ccmFc*, *ccmFn*, *cob*, *cox1*, *cox2*, *cox3*, *matR*, *nad1*, *nad2*, *nad3*, *nad4*, *nad4L*, *nad5*, *nad6*, *nad7*, *nad9*, *rpl5*, *rps1*, *rps3*, *rps4*, *rps12*, *rps14* and *sdh4*) from 11 species. The mitochondrial genomes of *Malus domestica*, *Vitis vinifera* and *Fragaria orientalis* were set as outgroups. The phylogenetic tree strongly supports (100% bootstrap support) the close phylogenetic relationship between *P. granatum* and *Lagerstroemia indica*.

The collinearity analysis of *Lagerstroemia indica*, *Eucalyptus grandis*, *Medinilla magnifica* and *P. granatum* were performed with two methods (Fig. 6). The mitogenome of *P. granatum* and *Lagerstroemia indica* had more collinear forward alignment and reverse complementary alignment

sequences, which was followed by *P. granatum* and *Eucalyptus grandis* (Fig. 6A). There were many homologous collinear blocks between the *P. granatum* mitochondrial genome and other three proximate species (Fig. 6B). Some gaps illustrates that these sequences are unique to the species and have no homology with the rest of the species. The results suggest that the *P. granatum* mitochondrial genome has undergone a lot of genomic rearrangement with close species.

Intracellular gene transfer of *P. granatum* organelle genomes

The *P. granatum* mitogenome sequence was approximately 2.41 times longer than its cp genome (NCBI Number: MK603511, 1158,638 bp). According to sequence similarity analysis, there were 22 homologous fragments between the mitogenome and chloroplast genome, with a total length of 19,889 bp, accounting for 5.20% of the total mitogenome (Fig. 7 and Table S6). By annotating these homologous sequences, 11 complete genes were also found on 22 homologous fragments, including nine tRNA genes (*trnD-GUC*, *trnH-GUG*, *trnI-GAU*, *trnM-CAU*, *trnP-UGG*, *trnS-UGA*, *trnV-GAC*, *trnW-CCA*, *trnN-GUU*) and two rRNA genes (*rrn5* and *rrn16*). There were 35 gene sequence of chloroplast transfer to mitochondria in *P. granatum* (Table S7). The length of *atpA_len507* gene was the longest with a length of 471bp. The *ndhA_len363* gene had the shortest length (35bp).

Discussion

Structure and characteristic of *P. granatum* mitogenome

The genome structure of plant mitochondrial genomes has always been mapped as a single master circle and a collection of subgenomic circles because of the large, direct, repeat recombination [25–26]. For example, the mitochondrial genomes of bitter melon [11] and *Vitex rotundifolia* [27] were mapped as a molecular circle. The mitochondrial genome of *Chaenomeles speciosa* was multi-chromosomes with two circular chromosomes [28] and *Paphiopedilum micranthum* with 26 circular chromosomes [29], while the *Q. acutissima* mitogenome was a branched structure of two circular molecules and one linear molecule [30]. In the present study, the *P. granatum* ‘Taishanhong’ mitogenome was represented by seven circular chromosomes structure (Fig. 1) with a total length of 382,774bp. There was significant difference in the structure (a single linear molecule) and size (404,807 bp) of *P. granatum* ‘Qingpi Sweet’ mitogenome [23]. The difference of mitochondrial genome between two varieties of pomegranate provided new insights into the function and structure of the mitochondrial genomes in pomegranate species.

GC content is an important factor for assessing species. The total GC content of *P. granatum* mitogenome was 45.91%, which was highest in the rRNA genes. The result was comparable to that of *Phaseolus vulgaris* (45.11%) [16], Bitter Melon (45.60%) [11], and *Bupleurum chinense* (45.68) [31], which was higher than that in *Mangifera sylvatica* (44.66%) [32]. The mitogenome of *P. granatum* ‘Taishanhong’ encoded 74 genes, including 46 protein-coding genes, 25 tRNA genes, and three rRNA genes, which was higher than that in *P. granatum* ‘Qingpi Sweet’(60 genes) [23].

The intron contents of the plant mitochondrial genes are conserved in each lineage but are variable among different lineages [33]. In the present study, the trans-spliced genes *nad1*, *nad2*, *nad5*, *nad7* contained four introns, and *nad4* gene had three introns. These trans-spliced introns were postulated to have evolved from one common ancestor by fragmentation of a cis-spliced arrangement [34].

Identification of codon usage and repeat sequences

Most PCGs were the typical ATG start codon, while *nad1* gene uses the ACG as initiation codons in the *P. granatum* mitogenome. The phenomenon also has been found in other studies, which were considered to be altered by RNA editing modificationp [35–36].

Codon usage preference is usually considered to be the result of the tendency to develop relative equilibrium within cells during species evolution, and codons with a RSCU greater than 1 are considered to be used by amino acid preference [17, 37–39]. The result indicated that Phe, Ile and Glu were the most common amino acids, and Met and Ter were much less common in the *P. granatum* mitogenome. This is different from other studies in *Camellia sinensis*[40] and *Clematis acerifolia* [41].

Repeated sequences are widely present in the mitochondrial genome, which are generally vital for intermolecular recombination, structural variations and extreme mitogenome sizes [42–44]. In this study, a total of 188 pairs of repeats were observed, including 97 pairs of reverse complementary repeats and 91pairs of forward repeats. These repetitive sequences play a crucial role in the assembly of the *P. granatum* mitogenome [45]. There were a total of 141 SSRs were detected and mononucleotide repeats of A/T were the most frequent, which was consistent with other research findings [11, 25, 46–47]. It provided a large number of reference markers for the evaluation of pomegranate germplasm diversity and species identification.

RNA editing analysis

RNA editing is a very common phenomenon in the mitochondrial genomes of higher plants. Almost all of the transcription products of mitochondrial PCGs are subjected to varying degrees of RNA editing but rarely occur in rRNA, tRNA, and introns [48–50]. RNA editing can affect the composition of the final mitochondrial proteome, which can create start and stop codons in the mitochondrial mRNA molecules [51–52]. Previous researches had uncovered approximately 421RNA editing sites within 26 genes in *Acer truncatum*[17] and 486 RNA editing sites within 31 genes in *Phaseolus vulgaris* [16]. There were 466 potential RNA editing sites distributed among 36 PCGs, and all of which were C-to-U base editing in *P. granatum* mitogenome. The result was similar to the study in *Salvia officinalis* [53], *Leucaena trichandra* [54] and *Pereskia aculeata* [55]. The *nad4* gene possessed 39 potential RNA editing sites and was the most edited of pomegranate mitogenome, which was supported by other studies [14, 39].

Phylogenetic relationships and syntenic analysis

Changes in the size and structure of plant mitochondrial genomes are evident, but functional genes remain conserved [56]. The nucleotide diversity (π) reveals the variation of nucleotide sequence in

different species. Our data showed that the pi value of *gene10.nad9* region was highest in these regions. The lower pi values suggested that the mitogenome sequences of *P. granatum* were highly conserved.

There were high similarities in the mitochondrial structure of *P. granatum* and *Eucalyptus grandis*, *Lagerstroemia indica* and *Medinilla magnifica* (Fig. 4B). The phylogenetic tree strongly supports the close phylogenetic relationship between *P. granatum* and *Lagerstroemia indica*. It was further confirmed from the molecular level that the pomegranate belonged to the Lythraceae family [57]. The co-linearity between *P. granatum* and *Lagerstroemia indica* was also high with large homozygous co-linear blocks, suggesting that the two species have undergone extensive rearrangement phenomena in the long evolutionary process [58]. The result provided an effective way to understand species evolution of pomegranate, and the specific mechanism and process need to be further studied.

Intracellular gene transfer of *P. granatum* organelle genomes

Gene transfer from chloroplasts to mitochondrial genomes is common during long-term plant evolution. The discovery of intracellular gene transfers can contribute to the accurate assembly of chloroplast and mitochondrial genomes and to the understanding of organelle genome evolution [59–60]. We found 22 homologous fragments between the pomegranate mitogenome and chloroplast genome, with a total length of 19,889 bp, accounting for 5.20% of the total mitogenome. There were 35 gene sequence of chloroplast transfer to mitochondria in *P. granatum*, while the length of *atpA*_len507 gene was the longest with a length of 471bp. Previous studies have found that tRNA genes occasionally migrated from the chloroplast to mitochondria [16]. The nine complete tRNA genes of homologous fragments were identified from the chloroplast of *P. granatum*, which suggested that they may play a role in normal functions [61]. Therefore, larger-scale sampling is required better to understand the evolution of phylogenetic analyses and molecular markers of pomegranate. These studies will provide a substantial foundation for understanding pomegranate mitogenome evolution and facilitate mitochondrial-based breeding.

Conclusions

In this study, we assembled and annotated the complete mitogenome of *P. granatum* cultivar ‘Taishanhong’. The genome structure was multi-chromosomes with seven circular chromosomes, and total length was 382,774 bp and GC content was 45.91%. In addition, the codon usage, sequence repeats, RNA editing and chloroplast to mitochondrion DNA transformation were also analyzed. There were high similarities in the mitochondrial structure between *P. granatum* and its proximal species. Evolutionary and collinear analysis showed that *P. granatum* was closely related to *Lagerstroemia indica*. This study provided information on the genetic characteristics, phylogenetic relationships, species identification and biological evolution of *P. granatum*.

Methods and materials

Plant materials, DNA extraction and sequencing

The tissue culture seedlings of *P. granatum* cultivar 'Taishanhong' were planted in the tissue culture room at Shandong Institute of Pomology (Taian, Shandong, China). Plants were kept in the dark for 14 d to obtain etiolated seedlings. Then, the fresh leaves were obtained and cleaned with DEPC water and kept in a freezer at -80°C.

The total genomic DNA was extracted by a DNA plant extraction kit (Tiangen, Beijing, China). After the genomic DNA of the sample was qualified, the DNA was fragmented using ultrasound. DNA fragments were purified, terminal repair, A was added to the 3' end, and sequencing splices were connected. The fragment size was selected by agarose gel electrophoresis and PCR amplification was performed to form sequencing library. The constructed libraries were inspected first, and qualified libraries were sequenced using Illumina Novaseq6000 platform by Nanjing Genepioneer Biotechnologies (Nanjing, China).

The Fastp (version 0.20.0, <https://github.com/OpenGene/fastp>) was used to filter the raw data, removing primer sequences and sequencing junctions, discarding reads with quality scores below Q5, and removing N values greater than 5. Finally, we obtained a total of 21,083,687 clean reads. The same DNA sample was also used for Oxford Nanopore sequencing based on the promethION platform. The triple sequenced data were filtered using Filtlong v0.2.1 software, and counted using Perl scripts.

Assembly and annotation of the mitochondrial genome

The minimap2 v2.1 software was used to compare the reference gene sequence (plant mitochondrial core gene) with the original three-generation data [62]. The sequences with alignment length greater than 50bp were selected as candidate sequences, and the sequences with more alignment genes and higher alignment quality were selected as seed sequences. The seed sequences were compared with original third-generation sequencing data. The sequences with overlap greater than 1kb and similarity greater than 70% were screened and obtained all third-generation sequencing data of mitochondrial genome. The third-generation datas were corrected by canu software and bowtie2 v2.3.5.1 was used to compare the second-generation data to the corrected sequence [63]. The Unicycler (v0.4.8) default parameter comparison was used to concatenate the above second-generation data with the corrected third-generation data. Finally, a master circular genome was generated by merging them manually.

The encoded proteins and rRNAs were compared to published plant mitochondrial sequences using BLAST, and further manual adjustments were made based on closely related species. The tRNA was annotated using tRNAscanSE (<http://lowelab.ucsc.edu/tRNAscan-SE/>) software [64]. ORFs were annotated using Open Reading Frame Finder (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>). The minimum length was set as 102bp, and redundant sequences and sequences with overlap with known genes were excluded. Sequences with length greater than 300 were compared with the nr library and annotated. The mitochondrial genome map was constructed using OGDRAW (<https://chlorobox.mpimp->

golm.mpg.de/OGDraw.html). RNA editing sites were predicted using the Plant Predictive RNA Editor (PREP) suite (<http://prep.unl.edu/>).

Analysis of RSCU and repeat sequences

Relative Synonymous Codon Usage (RSCU) is the ratio of the actual frequency of the use codon/theoretical frequency of the use codon. The RSCU of the mitochondrial genome of *P. granatum* was analyzed using a self-encoded Perl script, to screen for a unique CDS and determine the number of codons per gene. The blastn v2.10.1 software was used to identify the repeat sequences.

Phylogenetic tree construction

The CDS was used to make the maximum likelihood evolutionary tree. MAFFT software (v7.427) was used for multi-sequence alignment of inter-species sequences. The matched sequences were connected from head to tail, and trimAl (v1.4.rev15) was used to trim. The jmodeltest-2.1.10 software was used to predict the model after trimming, and the model was determined as GTR type. The maximum likelihood evolutionary tree was constructed using RAxML v8.2.10 (<https://cme.h-its.org/exelixis/software.html>) software. The GTRGAMMA model and bootstrap = 1000 were selected as the parameter.

Genomic comparison analysis with related species

The Mafft software was used for global alignment of homologous gene sequences of different species, and dnasp5 was used to calculate the pi value of each gene. The comparative analysis of mitochondrial genome structure between *P. granatum* and proximal species was carried out using the CGVIEW (http://stothard.afns.ualberta.ca/cgview_server/) software. The nucmer (4.0.0beta2) software with maxmatch parameter was used for genome alignment between other sequences and assembly sequences, and Dot-plot was generated. The blastn (2.10.1+) software was used to compare the assembled species and the selected species in turn, and Multiple Synteny plots were drawn. The parameters were set as follows: -word_size was 7 and E-value was 1e-5. The blast software was used to find homologous sequences between chloroplasts and mitochondria genome, and the similarity was set to 70% with an E-value of 1E-5. The homologous fragments were visualized using circos v0.69-5 software.

Declarations

Acknowledgements

Not applicable.

Author contributions

Y.Y.: Conceptualization. F.L. and W.C.: analyzed the data, F.L. and Y.X.: wrote original manuscript. A.M. and Y.Y.: reviewed and edited the manuscript. F.L., W.C. and Y.Y.: methodology. F.L. and Y.Y.: project administration. Y.Y.: funding acquisition. All authors reviewed the manuscript and agreed to the submission.

Funding

This work was supported by the Shandong Key Research and Development Projects (2022TZXD009, 2021LZGC007), Agricultural Science and Technology Innovation Engineering Discipline Team of Shandong Academy of Agricultural Sciences (CXGC2022D02, CXGC2023A12).

Data Availability

The raw sequencing data from the Illumina and Nanopore platforms generated during this study are available in GenBank. The associated BioProject and BioSample and SRA numbers are PRJNA971780 and SAMN35054861. SRA numbers of second and third generation data for the Illumina sequencing reads are SRR24519284 and SRR24554674, respectively. The mitogenome sequences were released into GenBank soon with the following accession numbers: OQ973289, OQ973290, OQ973291, OQ973292, OQ973293, OQ973294 and OQ973295. The plant samples are stored at the Shandong Institute of Pomology, Shandong, China (voucher numbers: SD-SL-026).

Ethics approval and consent to participate

Not applicable.

Ethics approval and consent to participate

All plant materials used in this study were grown and collected according to institutional and national guidelines. *P. granatum* ‘Taishanhong’ was a common cultivar and was obtained commercially. The study, including plant samples, complies with relevant institutional, national, and international guidelines and legislation. No specific permits were required for plant collection.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Chen L, Ren Y, Zhao J, Wang Y, Liu X, Zhao X, Yuan, Z. Phylogenetic analysis of wild pomegranate (*Punica granatum* L.) based on its complete chloroplast genome from Tibet, China. *Agronomy*. 2023; 13:126.
2. Shi J, Gao H, Wang S, Wu W, Tong R, Wang S, Li M, Jian Z, Wan R, Hu Q, et al. Exogenous arginine treatment maintains the appearance and nutraceutical properties of Hardand soft-seed pomegranates in cold storage. *Front. Nutr.* 2022; 9: 828946.
3. Trainin T, Harel-Beja R, Bar-Ya'akov I, Ben-Simhon Z, Yahalomi R, Borochoy-Neori H, Ophir R, Sherman M, Doron-Faigenboima A, Holland D. Fine mapping of the “black” peel color in pomegranate (*Punica granatum* L.) strongly suggests that a mutation in the anthocyanidin reductase (ANR) gene is responsible for the trait. *Front. Plant Sci.* 2021; 12: 642019.
4. Gao L, Zhang L, Liu J, Zhang X, Lu Y. Analysis of the volatile flavor compounds of pomegranate seeds at different processing temperatures by GC-IMS. *Molecules*. 2023; 28: 2717.
5. Hadree J, Shahidi F, Mohebbi M, Abbaspour M. Evaluation of effects of spray drying conditions on physicochemical properties of pomegranate juice powder enriched with pomegranate peel phenolic compounds: modeling and optimization by RSM. *Foods*. 2023; 12: 2066.
6. Anean HA, Mallasiy LO, Bader DMD, Shaat HA. Nano edible coatings and films combined with zinc oxide and pomegranate peel active phenol compounds has been to extend the shelf life of minimally processed pomegranates. *Materials*. 2023; 16:1569.
7. Ökmen G, Giannetto D, Fazio F, Arslan K. Investigation of pomegranate (*Punica granatum* L.) flowers' antioxidant properties and antibacterial activities against different staphylococcus species associated with bovine mastitis. *Vet. Sci.* 2023; 10: 394.
8. Magangana TP, Makunga NP, Fawole OA, Stander MA, Opara UL. Antioxidant, antimicrobial, and metabolomic characterization of blanched pomegranate peel extracts: effect of cultivar. *Molecules*. 2022; 27: 2979.
9. Wang Q, Yuan T, Zhu X, Song G, Wang D, Li L, Huang M, Gong J. The phenolics, antioxidant activity and in vitro digestion of pomegranate (*Punica granatum* L.) peels: an investigation of steam explosion pre-treatment. *Front. Nutr.* 2023; 10: 1161970.
10. Li X, Zhe M, Huang Y, Fan W, Yang J, Zhu, A. The evolution of mitochondrial genomes between two *cymbidium* sister species: dozens of circular chromosomes and the maintenance and deterioration of genome synten. *Genes*. 2023; 14: 864.
11. Niu Y, Zhang T, Chen M, Chen G, Liu Z, Yu R, Han X, Chen K, Huang A, Chen C, et al. Analysis of the complete mitochondrial genome of the bitter melon (*Momordica charantia*). *Plants*. 2023; 12:1686.
12. Lu C, Gao LZ, Zhang QJ. A high-quality genome assembly of the mitochondrial genome of the oil-tea tree *Camellia gigantocarpa* (Theaceae). *Diversity*. 2022; 14: 850.
13. Zhang N, Li Y, Halanych KM, Kong L, Li QA comparative analysis of mitochondrial ORFs provides new insights on expansion of mitochondrial genome size in Arcidae. *BMC Genomics*. 2022; 23:809.
14. You C, Cui T, Zhang C, Zang S, Su Y, Que Y. Assembly of the complete mitochondrial genome of *Gelsemium elegans* revealed the existence of homologous conformations generated by a repeat

- mediated recombination. *Int. J. Mol. Sci.* 2023; 24:527.
15. Li J, Li J, Ma Y, Kou L, Wei J, Wang W. The complete mitochondrial genome of okra (*Abelmoschus esculentus*): using nanopore long reads to investigate gene transfer from chloroplast genomes and rearrangements of mitochondrial DNA molecules. *BMC Genomics.* 2022; 23: 481.
 16. Bi C, Lu N, Xu Y, He C. Characterization and analysis of the mitochondrial genome of common bean (*Phaseolus vulgaris*) by comparative genomic approaches. *Int. J. Mol. Sci.* 2020; 21:3778.
 17. Ma Q, Wang Y, Li S, Wen J, Zhu L, Yan K, Du Y, Ren J, Li S, Chen Z, et al. Assembly and comparative analysis of the first complete mitochondrial genome of *Acer truncatum* Bunge: a woody oil-tree species producing nervonic acid. *BMC Plant Biol.* 2022; 22: 29.
 18. Wu ZQ, Liao XZ, Zhang XN, Tembrock LR, Broz A. Genomic architectural variation of plant mitochondria-A review of multichromosomal structuring. *J. Syst. Evol.* 2022; 60:160–168.
 19. Tsujimura M, Kaneko T, Sakamoto T, Kimura S, Shigyo M, Yamagishi H, Terachi T. Multichromosomal structure of the onion mitochondrial genome and a transcript analysis. *Mitochondrion.* 2019; 46:179–186.
 20. Goremykin VV, Lockhart PJ, Viola R, Velasco R. The mitochondrial genome of *malus domestica* and the import-driven hypothesis of mitochondrial genome expansion in seed plants. *Plant J.* 2012; 71: 615-626.
 21. Zhang S, Yin Z, Wu X, Li C, Xie K, Deng X, Grosser JW, Guo W. Assembly of Satsuma mandarin mitochondrial genome and identification of cytoplasmic male sterility–specific ORFs in a somatic cybrid of pummelo. *Tree.Genet. Genomes.* 2020; 16: 84.
 22. Wang S, Li D, Yao X, Song Q, Wang Z, Zhang Q, Zhong C, Liu Y, Huang H. Evolution and diversification of kiwifruit mitogenomes through extensive whole-genome rearrangement and mosaic loss of intergenic sequences in a highly variable region. *Genome Biol. Evol.* 2019; 11(4):1192–1206.
 23. Lu G, Zhang K, Que Y, Li Y. Assembly and analysis of the first complete mitochondrial genome of *Punica granatum* and the gene transfer from chloroplast genome. *Front. Plant Sci.* 2023; 14: 1132551.
 24. Ke SJ, Liu DK, Tu XD, He X, Zhang MM, Zhu MJ, Zhang DY, Zhang CL, Lan SR, Liu ZJ. Apostasia mitochondrial genome analysis and monocot mitochondria phylogenomics. *Int. J. Mol. Sci.* 2023; 24:7837.
 25. Guo Y, Li Z, Jin S, Chen S, Li F, Wu H. Assembly and comparative analysis of the complete mitochondrial genome of two species of *Calla Lilies* (*Zantedeschia*, Araceae). *Int. J. Mol. Sci.* 2023; 24: 9566.
 26. Putintseva YA, Bondar EI, Simonov EP, Sharov VV, Oreshkova NV, Kuzmin DA, Konstantinov YM, Shmakov VN, Belkov VI, Sadovsky MG, et al. Siberian larch (*Larix sibirica* Ledeb.) mitochondrial genome assembled using both short and long nucleotide sequence reads is currently the largest known mitogenome. *BMC Genom.* 2020; 21:654.
 27. Yu X, Duan Z, Wang Y, Zhang Q, Li W. Sequence analysis of the complete mitochondrial genome of a medicinal plant, *Vitex rotundifolia* Linnaeus f. (Lamiales: Lamiaceae). *Genes.* 2022; 13: 839.

28. Cao P, Huang Y, Zong M, Xu Z. De Novo Assembly and comparative analysis of the complete mitochondrial genome of *Chaenomeles speciosa* (Sweet) Nakai revealed the existence of two structural isomers. *Genes*. 2023; 14: 526.
29. Yang JX, Dierckxsens N, Bai MZ, Guo YY. Multichromosomal mitochondrial genome of *Paphiopedilum micranthum*: compact and fragmented genome, and rampant intracellular gene transfer. *Int. J. Mol. Sci.* 2023; 24:3976.
30. Liu D, Guo H, Zhu J, Qu K, Chen Y, Guo Y, Ding P, Yang H, Xu T, Jing Q, et al. Complex physical structure of complete mitochondrial genome of *Quercus acutissima* (Fagaceae): a significant energy plant. *Genes*. 2022; 13: 1321.
31. Qiao Y, Zhang X, Li Z, Song Y, Sun Z. Assembly and comparative analysis of the complete mitochondrial genome of *Bupleurum chinense* DC. *BMC Genomics*. 2022; 23:664.
32. Niu Y, Gao C, Liu J. Complete mitochondrial genomes of three *Mangifera* species, their genomic structure and gene transfer from chloroplast genomes. *BMC Genomics* 2022; 23:147.
33. Terasawa K, Odahara M, Kabeya Y, Kikugawa T, Sekine Y, Fujiwara M, Sato N. The mitochondrial genome of the moss *Physcomitrella patens* sheds new light on mitochondrial evolution in land plants. *Mol. Biol. Evol.* **2007**; 24: 699–709.
34. Guo W, Zhu A, Fan W, Adams RP, Mower JP. Extensive shifts from cis- to trans-splicing of Gymnosperm mitochondrial introns. *Mol. Biol. Evol.* 2020; 37: 1615–1620.
35. Yang J, Wariss H, Tao L, Zhang R, Yun Q, Hollingsworth P, Dao Z, Luo G, Guo H, Ma Y, Sun W. De novo genome assembly of the endangered *Acer yangbiense*, a plant species with extremely small populations endemic to Yunnan Province, China. *Gigascience*. 2019; 8(7): giz085.
36. Ye N, Wang X, Li J, Bi C. Assembly and comparative analysis of complete mitochondrial genome sequence of an economic plant *Salix suchowensis*. *PeerJ*. 2017; 5(1): e3148.
37. Shi S, Liu Y, Xia R, Qin L. Comprehensive analysis of codon usage in *Quercus* chloroplast genome and focus on *psbA* gene. *Genes*. 2022; 13(11):2156.
38. Zeng Y, Shen L, Chen S, Qu S, Hou N. Codon usage profiling of chloroplast genome in Juglandaceae. *Forests*. 2023;14(2):378.
39. Wu P, Xiao W, Luo Y, Xiong Z, Chen X, He J, Sha A, Gui M, Li Q. Comprehensive analysis of codon bias in 13 *Ganoderma* mitochondrial genomes. *Front. Microbiol.* 2023; 14: 1170790.
40. Li J, Tang H, Luo H, Tang J, Zhong N, Xiao L. Complete mitochondrial genome assembly and comparison of *Camellia sinensis* var. *Assamica* cv. *Duntsa*. *Front. Plant Sci.* 2023; 14: 1117002.
41. Liu D, Qu K, Yuan Y, Zhao Z, Chen Y, Han B, Li W, El-Kassaby Y. A, Yin Y, Xie X, Tong B, Liu H. Complete sequence and comparative analysis of the mitochondrial genome of the rare and endangered *Clematis acerifolia*, the first clematis mitogenome to provide new insights into the phylogenetic evolutionary status of the genus. *Front. Genet.* 2023; 13:1050040.
42. Guo W, Felix G, Fan W, Young G. Ginkgo and Welwitschia Mitogenomes reveal extreme contrasts in gymnosperm mitochondrial evolution. *Mol Biol Evol.* 2016, 33(6): 1448–1460.

43. Xiong Y, Lei X, Bai S, Xiong Y, Liu W, Wu W, Yu Q, Dong Z, Yang J, Ma X. Genomic survey sequencing, development and characterization of single- and multi-locus genomic SSR markers of *Elymus sibiricus* L. BMC Plant Biol. 2021; 21: 3.
44. Li J, Xu Y, Shan Y, Pei X, Yong S, Liu C, Yu J. Assembly of the complete mitochondrial genome of an endemic plant, *Scutellaria tsinyunensis*, revealed the existence of two conformations generated by a repeat-mediated recombination. Planta. 2021; 254: 36.
45. Dong, S, Zhao, C, Chen, F, Liu, Y, Zhang, S, Wu, H, Zhang L, Liu Y. The complete mitochondrial genome of the early flowering plant *nymphaea colorata* is highly repetitive with low recombination. BMC Genomics. 2018; 19: 614.
46. Shan Y, Li J, Zhang X, Yu J. The complete mitochondrial genome of *Amorphophallus albus* and development of molecular markers for five *Amorphophallus* species based on mitochondrial DNA. Front. Plant Sci. 2023; 14: 1180417.
47. Sheng W, Deng J, Wang C, Kuang Q. The garden asparagus (*Asparagus officinalis* L.) mitochondrial genome revealed rich sequence variation throughout whole sequencing data. Front. Plant Sci. 2023; 14:1140043.
48. Habib S, Dong S, Liu Y, Liao W, Zhang S. The complete mitochondrial genome of *Cycas debaoensis* revealed unexpected static evolution in gymnosperm species. PLoS ONE. 2021; 16(7): e0255091.
49. He P, Xiao GH, Liu H, Zhang LH, Zhao L, Tang MJ, Huang S, An YJ, Yu JN. Two pivotal RNA editing sites in the mitochondrial *atp1* mRNA are required for ATP synthase to produce sufficient ATP for cotton fiber cell elongation. New Phytol. 2018; 218:167–182.
50. Shtratnikova V Y, Schelkunov M I, Penin A A, Logacheva M D. Mitochondrial genome of the nonphotosynthetic mycoheterotrophic plant *Hypopitys monotropa*, its structure, gene expression and RNA editing. PeerJ. 2020; 8: e9309.
51. Gerke P, Szovenyi P, Neubauer A, Lenz H, Gutmann B, McDowell R, Small I, Schallenberg-Rudinger M, Knoop V. Towards a plant model for enigmatic U-to-C RNA editing: The organelle genomes, transcriptomes, editomes and candidate RNA editing factors in the hornwort *Anthoceros agrestis*. New Phytol. 2020; 225: 1974–1992.
52. Mower JP, Palmer JD. Patterns of partial RNA editing in mitochondrial genes of *Beta vulgaris*. Mol. Genet. Genom. 2006; 276: 285–293.
53. Yang H, Chen H, Ni Y, Li J, Cai Y, Wang J, Liu C. Mitochondrial genome sequence of *Salvia officinalis* (Lamiales: Lamiaceae) suggests diverse genome structures in Cogenetic species and finds the stop gain of genes through RNA editing events. Int. J. Mol. Sci. 2023; 24: 5372.
54. Kovar L, Nageswara-Rao M, Ortega-Rodriguez S, Dugas DV, Straub S, Cronn R, Strickler SR, Hughes CE, Hanley KA, Rodriguez DN, et al. PacBio-based mitochondrial genome assembly of *Leucaena trichandra* (Leguminosae) and an intrageneric assessment of mitochondrial RNA editing. Genome Biol. Evol. 2018; 10(9): 2501–2517.
55. Zhang X, Shan Y, Li J, Qin Q, Yu J, Deng H. Assembly of the complete mitochondrial genome of *Pereskia aculeata* revealed that two pairs of repetitive elements mediated the recombination of the

- genome. *Int. J. Mol. Sci.* 2023; 24: 8366.
56. Hong Z, Liao XZ, Ye YJ, Zhang NN, Yang ZJ, Zhu WD, Gao W, Sharbrough J, Tembrock LR, Xu DP, et al. A complete mitochondrial genome for fragrant Chinese rosewood (*Dalbergia odorifera*, Fabaceae) with comparative analyses of genome structure and intergenomic sequence transfers. *BMC Genom.* 2021; 22: 672.
57. Usha T, Middha SK, Babu D, Goyal AK, Das AJ, Saini D, Sarangi A, Krishnamurthy V, Prasannakumar MK, Saini DK, Sidhalinghamurthy KR. Hybrid assembly and annotation of the genome of the Indian *Punica granatum*, a superfood. *Front. Genet.* 2022; 13: 786825.
58. Cole L. W, Guo W, Mower JP, Palmer J. D. High and variable rates of repeat-mediated mitochondrial genome rearrangement in a genus of plants. *Mol. Biol. Evol.* 2018; 35(11): 2773-2785.
59. Gui ST, Wu ZH, Zhang HY, Zheng YZ, Zhu ZX, Liang DQ, Ding Y. The mitochondrial genome map of *Nelumbo nucifera* reveals ancient evolutionary features. *Sci Rep.* 2016; 6:11.
60. Nguyen VB, Giang VNL, Waminal NE, Park HS, Kim NH, Jang W, Lee J, Yang T. Comprehensive comparative analysis of chloroplast genomes from seven *Panax* species and development of an authentication system based on species-unique single nucleotide polymorphism markers. *J Ginseng Res.* 2020; 44(1):135–44.
61. Kitazaki K, Kubo T, Kagami H, Matsumoto T, Fujita A, Matsuhira H, Matsunaga M, Mikami T. A horizontally transferred Trna (Cys) gene in the sugar beet mitochondrial genome: Evidence that the gene is present in diverse angiosperms and its transcript is aminoacylated. *Plant J. Cell Mol. Biol.* 2011; 68: 262–272.
62. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics.* 2018; 34: 3094–3100.
63. Koren S, Walenz BP, Berlin K, Miller JR, Phillippy AM. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res.* 2017; 27: 722–736.
64. Chan PP, Lowe T M. tRNAscan-SE: Searching for tRNA genes in genomic sequences. *Methods Mol Biol.* 2019; 1962:1–14.

Figures



Figure 1

Circular Map of the *P. granatum* mitochondrial genome. The genome consisted of 7 circular chromosomes. The genes that code in the forward direction are on the outside of the circle, and the genes that code in the opposite direction are on the inside of the circle. The inside gray circle represents GC content.




Figure 2

Relative synonymous codon usage (RSCU) in the *P. granatum* mitochondrial genome. The different amino acids are shown on the x-axis. RSCU values are the number of times a particular codon is observed relative to the number of times that codon would be expected for a uniform synonymous codon usage. The bottom square represents all the codons that code for each amino acid, and the height of the top column represents the sum of all the codon RSCU values.




Figure 3

The number of SSR motifs and RNA editing sites predicted in *P. granatum* mitogenome. **(A)** The number of identified SSR motifs. **(B)** The number of RNA editing sites identified in each PCGs.




Figure 4

Nucleotide diversity and comparative analysis of *P. granatum* mitochondrial structure. (A) The gene nucleotide variability (pi) values. (B) Comparative analysis of mitochondrial structure of *P. granatum* and proximal species. The outer two circles describe the length and direction of the genome. The circles inside represent similar results compared to other reference genomes. The black circles represent the GC content.




Figure 5

The evolutionary relationship between *P. granatum* and homologous species.




Figure 6

Collinear analysis of mitochondrial sequence with two methods. (A) The horizontal coordinate in each box indicates the assembled sequence, and the vertical coordinate indicates the other sequences. The red line in the box indicates the forward alignment, and the blue line indicates the reverse complementary alignment. **(B)** The boxes in each row represent a genome, and the lines in the middle represent homologous regions.




Figure 7

Homologous fragments of chloroplast and mitochondrial sequence in *P. granatum*. Chloroplast is chloroplast sequence and chr is mitochondrial sequence. Genes from the same complex are labeled with the same color, with homologous sequences represented at the center line junctions.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.docx](#)