



OPEN ACCESS

EDITED BY

Zi-Piao Ye,
Jinggangshan University, China

REVIEWED BY

Changfang Zhou,
Nanjing University, China
Xiaolong Yang,
Nantong University, China

*CORRESPONDENCE

Ole Pedersen

✉ opedersen@bio.ku.dk

Jianfeng Hua

✉ jfhua@cncb.net

RECEIVED 14 December 2023

ACCEPTED 04 March 2024

PUBLISHED 19 March 2024

CITATION

Guo J, Xue J, Yin Y, Pedersen O and Hua J (2024) Response of underwater photosynthesis to light, CO₂, temperature, and submergence time of *Taxodium distichum*, a flood-tolerant tree. *Front. Plant Sci.* 15:1355729. doi: 10.3389/fpls.2024.1355729

COPYRIGHT

© 2024 Guo, Xue, Yin, Pedersen and Hua. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Response of underwater photosynthesis to light, CO₂, temperature, and submergence time of *Taxodium distichum*, a flood-tolerant tree

Jinbo Guo^{1,2,3}, Jianhui Xue^{1,2,3}, Yunlong Yin^{1,3}, Ole Pedersen^{4,5*} and Jianfeng Hua^{1,3*}

¹Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Nanjing, China, ²College of Biology and the Environment, Nanjing Forestry University, Nanjing, China, ³Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Jiangsu Key Laboratory for the Research and Utilization of Plant Resources, Nanjing, China, ⁴Department of Biology, University of Copenhagen, Copenhagen, Denmark, ⁵School of Biological Sciences, The University of Western Australia, Crawley, WA, Australia

Introduction: Partial or complete submergence of trees can occur in natural wetlands during times of high waters, but the submergence events have increased in severity and frequency over the past decades. *Taxodium distichum* is well-known for its waterlogging tolerance, but there are also numerous observations of this species becoming partially or complete submerged for longer periods of time. Consequently, the aims of the present study were to characterize underwater net photosynthesis (P_N) and leaf anatomy of *T. distichum* with time of submergence.

Methods: We completely submerged 6 months old seedling of *T. distichum* and diagnosed underwater (P_N), hydrophobicity, gas film thickness, Chlorophyll concentration and needles anatomy at discrete time points during a 30-day submergence event. We also constructed response curves of underwater P_N to CO₂, light and temperature.

Results: During the 30-day submergence period, no growth or formation new leaves were observed, and therefore *T. distichum* shows a quiescence response to submergence. The hydrophobicity of the needles declined during the submergence event resulting in complete loss of gas films. However, the Chlorophyll concentration of the needles also declined significantly, and it was there not possible to identify the main cause of the corresponding significant decline in underwater P_N . Nevertheless, even after 30 days of complete submergence, the needles still retained some capacity for underwater photosynthesis under optimal light and CO₂ conditions.

Discussion: However, to fully understand the stunning submergence tolerance of *T. distichum*, we propose that future research concentrate on unravelling the finer details in needle anatomy and biochemistry as these changes occur during submergence.

KEYWORDS

bald cypress, contact angle, flood tolerance, gas films, hydrophobicity, swamp cypress, low oxygen quiescence syndrome

1 Introduction

Partial or complete submergence of trees can occur in natural wetlands during times of high waters. Accordingly, more than 1,000 species of trees and bushes in Pantanal (one of the world's largest tropical wetlands) become submerged every year in the wet season when the River Negro rises up to 10 m above its water level in the dry season (Parolin, 2009). However, trees can also face complete submergence in man-made wetlands such as at the banks of the Three Gorges Reservoir, where *Taxodium distichum* has been introduced in an attempt to stabilize the steep banks (Wang et al., 2019; Ding et al., 2021; He et al., 2021). Since the Three Gorges dam has first stored water in June 2003, plants growing at lower elevations, including the cultivated *T. distichum*, have experienced periodic complete submergence every year. In this case, *T. distichum* not only survived but grew from seedling to tree (Li et al., 2006). We propose that the outstanding flood tolerance of *T. distichum* is partly a result of its remarkable ability to photosynthesize under water, which slows down carbohydrate depletion and protects the tissue from anoxia via O_2 production during extended periods of submergence.

In air, gases diffuse 10,000-fold faster than in water, and therefore, CO_2 and O_2 generally restrict photosynthesis and respiration of submerged terrestrial plants. Consequently, submerged aquatic plants have evolved a number of key shoot and root traits involved in facilitating CO_2 or O_2 exchange with the floodwater including, but not limited to, thin leaf lamina composed of only two cell layers, thin or completely absent leaf cuticle, chloroplasts in the leaf epidermis, and aerenchyma to facilitate internal aeration (Sculthorpe, 1967). However, even in the presence of these extreme adaptations, CO_2 availability can still limit underwater photosynthesis (Madsen and Sand-Jensen, 1991; Maberly and Madsen, 2002), and about half of the world's aquatic plant species have thus evolved the ability to use bicarbonate (HCO_3^-) as an alternative inorganic carbon source in photosynthesis (Prins and Elzenga, 1989; Iversen et al., 2019). Lacking most of these key leaf traits, the photosynthetic rates of submerged terrestrial plants are significantly lower than those of aquatic plants regardless of whether underwater photosynthesis is measured at ambient or elevated CO_2 levels (Colmer et al., 2011). Similarly, the availability of molecular O_2 can restrict underwater respiration of submerged terrestrial plants, and an O_2 pressure of almost twice that of atmospheric equilibrium is needed to saturate respiration (Colmer and Pedersen, 2008). A recent meta-analysis encompassing 112 species of both aquatic and terrestrial plants have clearly demonstrated that partial or complete submergence lead to significant declines in tissue O_2 status particularly during darkness when the only source of O_2 for underwater respiration is O_2 dissolved in the floodwater (Herzog et al., 2023). However, some species of wetland plants form numerous adventitious roots emerging from the stem and hanging into the floodwater as response to partial or complete submergence (Rich et al., 2013; Zhang et al., 2017; Lin et al., 2023). Such roots are referred to as aquatic adventitious roots and have been shown to act as "physical gills" by facilitating uptake of O_2 from the floodwater (Ayi et al., 2016).

Some terrestrial plants, including the focal species of the present study, possess superhydrophobic leaves, and these have been shown to enhance gas exchange with the floodwater. Upon submergence, superhydrophobic leaves retain a thin gas film visible as a silvery sheen from the leaf surface (Pedersen and Colmer, 2012). Gas film formation on submerged leaves was first reported for deepwater rice, wheat, barley, and oats, where the beneficial effects on carbon fixation was also first reported (Raskin and Kende, 1983). Later, a series of studies reported leaf gas film formation during submergence in several species of wild wetland plants, where gas films were retained on partially or completely submerged leaves (Colmer and Pedersen, 2008; Pedersen et al., 2009; Winkel et al., 2016). The increased CO_2 exchange caused by leaf gas films results in enhanced underwater net photosynthesis rate (P_N), and generally underwater P_N is 6- to 10-fold higher in the presence of leaf gas films compared with leaves without superhydrophobic leaves or with leaves where the gas films have been experimentally removed (Colmer and Pedersen, 2008; Pedersen et al., 2009; Winkel et al., 2011; Verboven et al., 2014; Konnerup et al., 2017; Winkel et al., 2017). Although the beneficial effects of leaf gas films have emphasized CO_2 exchange for underwater photosynthesis, leaf gas films have also been shown to significantly enhance internal aeration. Accordingly, removal of hydrophobicity (and thereby also the leaf gas films) by brushing with a dilute detergent resulted in steep declines in O_2 status of belowground tissues both in rice (Winkel et al., 2013) and in a wild wetland plant (Winkel et al., 2011), clearly demonstrating the crucial importance of leaf gas films for internal aeration of submerged terrestrial plants.

Low light availability under water may also restrict photosynthesis and submergence can invoke shade acclimation of submerged terrestrial leaves. In natural water bodies, the light intensity under water is lower than that in the air above not only because light is being reflected at the surface but also because light is being absorbed by water itself, by suspended particles such as algae, and by colored dissolved organic matter (Kirk, 1994). Murky floodwaters with algal blooms and/or high amounts of colored dissolved organic matter offer even less light for submerged plants with only 0.5% of the surface insolation left at deep floods (Vervuren et al., 2003). Consequently, many terrestrial plants respond to submergence by shade acclimations in their leaves, and these acclimations involve a reduction in leaf thickness, a thinner cuticle, thinner cell walls, and therefore a lower leaf mass area all resulting in better tissue O_2 status (Mommer et al., 2007) and enhanced underwater P_N due to better CO_2 exchange (Mommer et al., 2004). Interestingly, the strong beneficial effect of leaf gas films on CO_2 uptake is to a certain extent counteracted by the reflection of light at low light intensities; i.e., at low light, the silvery sheen of gas films reflects light and results in lower underwater P_N (Winkel et al., 2017), showing that leaf gas films can also be disadvantageous during submergence in a low-light environment.

Trees and bushes forming the riparian vegetation often become partial or completely submerged when the river rises. However, poor flood tolerance of terrestrial plants leads to decreases in species richness as flooding intensity increases, leaving only the most flood-tolerant species to form the riparian vegetation (Garssen et al., 2017). Consequently, there are only two genera of Central European

Trees showing very high flood tolerance (i.e., species of *Alnus* and *Salix*), and these are characterized with the formation of adventitious roots, lenticels, and aerenchyma in response to flooding (Glenz et al., 2006). However, the model species of the present study, *T. distichum* (L.) Rich, also shows extraordinary flood tolerance. It is a deciduous tree of the *Taxodium* genus, native to North America and Mexico where it forms large natural stands mostly in coastal plains affected by tide, in marshes with poor drainage, and in lowlands with periodic flooding (Wang et al., 2022; Guo et al., 2023). Owing to its outstanding flood tolerance, *T. distichum* has broad application prospects and is promoted for use in ecosystem restoration and construction of wetlands (Ding et al., 2021; He et al., 2021). More recently, it was found to have excellent tolerance to long-term periodic submergence in the water-level-fluctuating zone of the Three Gorges Reservoir (Wang et al., 2016, 2019). Interestingly, we found that *T. distichum* had a higher survival rate when submerged in winter than in summer, which may be because the activity of enzymes involved in underwater P_N is affected by water temperature like other enzymes; thus, it is necessary to explore the response of underwater P_N to temperature.

Consequently, the aims of the present study were to characterize underwater P_N and leaf anatomy of *T. distichum* with time of submergence and to establish light, CO_2 , and temperature response curves of the underwater P_N . Aerial photosynthesis of *T. distichum* has been thoroughly investigated (Wang et al., 2016; Taylor and Smith, 2017), but its capacity for underwater photosynthesis has not yet been evaluated. Interestingly, it was previously reported that *T. distichum* possesses a superhydrophobic leaf cuticle (Neinhuis and Barthlott, 1997), which should result in gas film formation during submergence. We therefore hypothesized that (i) some photosynthesis takes place when submerged, but the rate is strongly limited by light and CO_2 showing a characteristic relationship with temperature; (ii) the underwater P_N of *T. distichum* declines with time of submergence; (iii) and the decline in P_N is linked to loss of leaf hydrophobicity and thereby the beneficial role of leaf gas films. Our study therefore fills an important gap related to the complete lack of knowledge related to the photosynthetic capacity of woody plants under water and the associated mechanistic understanding of flood tolerance of trees.

2 Materials and methods

2.1 Plant materials and growth conditions

Leaf material for characterization of key photosynthetic parameters was sampled from a 4-year old *T. distichum* (L.) Rich at the Institute of Botany, Jiangsu Province and Chinese Academy of Sciences (32°05' N, 118°83' E) during midmorning from 9 to 10 a.m. The plant height was 2.6 m high and the diameter at breast height was 2.32 cm, which was in a rapid growth phase. Young but fully expanded and healthy branchlets with green needle-like leaves of the linear-lanceolate type were chosen for these experiments in June–July, whereas scale-like and appressed needles were disregarded.

For the long-term submergence experiment, seeds of *T. distichum* were sown in a seedling tray and after germination transferred to pots

(23 cm upper diameter, 15 cm basal diameter, and 22 cm high) filled with a mixture of potting soil and sandy clay. The seedlings were grown in a greenhouse (temperature: $23 \pm 2^\circ\text{C}$; humidity: 60%–70%) for 2 months and then moved outdoors for another 4 months, and the pots were watered daily with tap water. 30 healthy seedlings with an average height of 45 cm were selected for the experiment.

2.2 CO_2 versus underwater P_N

Underwater P_N was measured following the approach of Pedersen et al. (2013). In brief, artificial floodwater was prepared according to Smart and Barko (1985) with a final alkalinity of $2.0 \text{ mol H}^+ \text{ equivalent m}^{-3}$. Considering that the underwater photosynthetic CO_2 saturation concentration of submerged leaves of terrestrial wetland plants is approximately 20–75 times or even higher than the atmospheric equilibrium concentration ($\sim 18 \text{ mmol m}^{-3}$ free CO_2) (Pedersen et al., 2009), a CO_2 concentration range of 10–2,000 mmol m^{-3} was set. In addition, compared with high CO_2 concentration, underwater P_N changes more significantly under low CO_2 concentration; thus, six concentration gradients (10, 25, 50, 100, 200, and 500 mmol m^{-3}) were set for low concentration and three concentration gradients (1,000, 1,500, and 2,000 mmol m^{-3}) were set for high concentration. Prior to the pH adjustment, the solution was purged with N_2 to reduce the O_2 concentration approximately 30%–50% of air equilibrium to prevent photorespiration during incubation (Setter et al., 1989). The artificial floodwater was then siphoned into 44-mL glass vials and two pieces of 3-mm glass beads were added to each vial to ensure the mixing during incubation. One branchlet with needles (approximately 1 cm^2 or approximately 15.8 mg fresh mass) before the vial was sealed with a glass lid (no headspace or gas bubbles present). The vials were mounted on a vertically rotating disk (10 rpm) and inundated in a constant temperature bath at 25°C and illuminated with a photon flux of $1,000 \mu\text{mol m}^{-2} \text{ s}^{-1}$ (see below). Vials without tissue served as blanks.

After 60 min, the vials were retrieved and the O_2 concentration was measured using an O_2 optode (OPTO-MR, Unisense, Denmark) inserted into the vial. The needles in the vial were then neatly placed on a clean white background board while a ruler is placed to take the photo. Make sure the leaves do not overlap each other when taking the photo; ImageJ software was used to measure the exact leaf area (Schneider et al., 2012). The photosynthetic rate was calculated using the following equation:

$$P_N (\mu\text{mol O}_2 \text{ m}^{-2} \text{ s}^{-1}) = \frac{\Delta\text{O}_2 (\mu\text{mol O}_2 \text{ L}^{-1}) V_{\text{vial}} (\text{L})}{t (\text{sec}) A (\text{m}^2)}$$

where ΔO_2 is the difference in O_2 concentration in vial with tissue and blanks, V_{vial} is the volume of the vials, i.e., 0.044 L, t is the incubation time, and A is the area of the needles.

2.3 Light versus underwater P_N

Floodwater and tissue were prepared as above but with a fixed CO_2 concentration of 500 mmol m^{-3} . Compared with high light

intensity, underwater P_N changes more significantly under low light intensity; thus, six gradients (0, 50, 100, 200, 300, and 500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) were set for low light intensity and three gradients (1,000, 1,500, and 2,000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) were set for high light intensity. Different light intensities were established by using a high-pressure Na lamp at various distances and by regulating the voltage. For zero light, the vials were wrapped in aluminum foil. The light intensities were measured using a spherical PAR (photosynthetically active radiation) sensor (QSL2101, Biospherical Instruments Inc., USA). As for the CO_2 response (see above), samples were incubated for 60 min.

2.4 Temperature versus underwater P_N

Floodwater and tissue were prepared as above but with a fixed CO_2 concentration of 500 mmol m^{-3} and PAR at 1,000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. The temperature was set to 10, 15, 20, 25, 30, or 35°C using a combination of an immersion heater (300W, SUNSUN, Zhejiang, China) and a water cooler (TECO, Taiwan, China) to achieve a stable temperature, which was monitored in real time by a temperature electrode (Temp-UniAmp thermosensor, Unisense, Denmark). As for the CO_2 response (see above), samples were incubated for 60 min.

2.5 Long-term submergence

A 30-day submergence experiment with two treatments (drained controls or completely submerged) was conducted using 6-month-old *T. distichum* seedlings. Fifteen pots with one plant in each were transferred to three plastic tanks (depth, 69 cm; volume, 122 L) filled with tap water, with five seedlings (technical replicates) in each tank (true replicates). The water was changed every 2 days, and the hose was placed at the bottom of the bucket to ensure that the water was completely replaced. Another 15 plants served as controls, and these were unsubmerged and kept under conditions (photoperiod and temperature) similar to those for submerged plants and irrigated every 2 days. Underwater P_N was measured on healthy needles on days 7, 14, 21, and 31 using a PAR of 1,000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and a CO_2 concentration of 500 mmol m^{-3} (see experimental procedure above) and leaves were transported in water back to the laboratory to minimize damage and exposure to air. It should be noted that the underwater P_N of the submerged leaves of terrestrial wetland plants will be severely limited by the CO_2 availability at atmospheric equilibrium CO_2 concentrations. In order to more accurately evaluate the underwater photosynthetic capacity of submerged leaves, a CO_2 concentration higher than atmospheric equilibrium was required, and 500 mmol m^{-3} is set for this experiment.

During the first 8 days of the submergence experiment, submerged leaves were harvested every other day. These were observed and photographed with scanning electron microscopy (SU8100, Hitachi Scientific Instruments, Japan). Subsequently, stomatal density (number of stomata per mm^{-2}) and stomatal

index (ratio of number of stomata to the total number of epidermal cells including stomata) were calculated from these images (Hegde and Krishnaswamy, 2021; Li et al., 2022).

At the end, healthy leaves were sampled and then transported into water to minimize damage and re-exposure to air. Chlorophyll measurements were conducted on submerged leaves as well as controls using ethanol extractions and absorbance of the extract was measured on a spectrophotometer (UV-1800, Shanghai Mepuda Instrument Co., LTD, China). Chlorophyll was calculated using the equations in Zhang et al. (2020).

Finally, cross-sections for microscopy were prepared from paraffin-embedded needles and later studied using visible light microscopy (BX53F, Olympus, Tokyo, Japan).

2.6 Influence leaf gas films on underwater P_N

To investigate the effects of leaf gas films on underwater P_N , four healthy 6-month-old *T. distichum* seedlings were selected. Two fully unfolded branchlets were sampled from each plant and then they were divided into two groups. To remove hydrophobicity and thereby prevent formation of leaf gas films, one group was brushed five times, on both sides, with a fine paintbrush dipped into 0.01% (v/v) Triton X. After that, they were washed for 5 s, three times, in artificial floodwater without Triton X (Teakle et al., 2014; Winkel et al., 2014). The other group was untreated and served as control with each group having four replicates. Underwater P_N was measured as described above with 500 $\text{mmol CO}_2 \text{ m}^{-3}$ under a photon flux of 1,000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ at 25°C.

2.7 Assessment of gas film thickness and needle hydrophobicity

Leaf gas film thickness was measured following the approach of Raskin and Kende (1983). In brief, the buoyancy of a branchlet was measured on five replicates using a four-digit balance with a hook underneath before and after removal of hydrophobicity using 0.01% Triton X; see above. Next, the area of the needles was determined (see above), and the gas film thickness (m) was calculated as gas film volume (m^3) divided by needle area (m^2).

Surface hydrophobicity was assessed by measuring the contact angle of a 1- mm^3 droplet of water on the needle surfaces following Sikorska et al. (2017). Branchlets with needles were held horizontal using a glue stick. Water droplets were applied to the lamina of 10 replicate needles (5 on the adaxial side and 5 on the abaxial side), and photographed at $\times 35$ magnification using a horizontally positioned dissecting microscope (MZ62, Mshot, China) and a digital camera. The droplet contact angles were measured using ImageJ (ImageJ v.1.43U, National Institutes of Health, Bethesda, MD, USA).

Finally, specific leaf area (SLA) was measured by determining the area (see above) and dry mass of needles, where the needles were

dried for 48 h at 60°C. SLA was calculated as area (m^2) divided by dry mass (kg).

2.8 Data analysis

We used non-linear regression to fit models derived from FvCB (CO_2 response) (Liang and Liu, 2017) and light response was fitted according to the Ye model (Ye et al., 2013). The temperature optimum was modeled using a Gaussian function and a standard exponential function was used to predict temperature coefficient Q_{10} (Pedersen et al., 2016). The temperature coefficient Q_{10} represents the relative change of underwater P_N with every 10°C change in temperature. The data were processed using Excel 2016 and graphed with Origin software (2021 64Bit, Electronic Arts Games, USA). Results were expressed as means $\pm$ standard deviation. All statistical tests were conducted using SPSS 16.0s (SPSS Inc., USA) including that of the Duncan's multiple range test. Unless otherwise stated, a probability level of 0.05 was used.

3 Results

3.1 Submergence of *Taxodium* during times of high water level

Waterlogging, partial submergence, and even complete submergence of *Taxodium* is a recurrent phenomenon during

times of high water levels. When the water first starts rising, the soil becomes flooded, resulting in waterlogging of vast areas of trees (Figure 1A), and as the water continues to rise, the lower branches become submerged (Figures 1B, C). Ultimately, the entire canopy is under water (Figure 1D), and gas exchange with the atmosphere is no longer possible. The floodwater in the Yangtze River is murky as a result of suspended materials (Figure 1D), and therefore the light environment also changes upon submergence, resulting in reduced photosynthesis due to the combination of low light and restricted CO_2 availability. The habitat photos in Figure 1 clearly demonstrate the relevance of our study, where we are aiming to characterize the ability of *T. distichum* to continue photosynthesizing under water with emphasis on response to CO_2 and light availabilities, and temperature.

3.2 Response of underwater net photosynthesis to CO_2 , light, and temperature by *Taxodium distichum*

We characterized the response of underwater net photosynthesis (P_N) to dissolved CO_2 , light availability, and temperature, which are relevant environmental parameters during submergence of *T. distichum*. Submerged branchlets with needles produced O_2 when incubated in the artificial floodwater in the light visible as bubble formation on the needles (Figure 2A). The response of P_N (i.e., net O_2 consumption) to dissolved CO_2 at 25°C and a photon flux of 1,000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ showed a




FIGURE 1

Habitat photos of *Taxodium* growing along the banks of the Three Gorges Reservoir, which is part of the Yangtze River. The species depicted is hybrid between *Taxodium distichum* and *Taxodium mucronatum*, which has been planted on the banks in an attempt to reduce erosion as the annual water level fluctuations are up to 175 m (Wang et al., 2019). (A) The initial phase of flooding resulting in waterlogging, but as the water continues to rise, the low branches become submerged (B, C). Finally, the entire canopy is under water (D) and may remain so for up to 120 days and still survive (Yang et al., 2023).




FIGURE 2 Response of underwater photosynthesis to CO_2 , light, and temperature of submerged *Taxodium distichum* branchlets. In (A), a section of a branchlet is incubated in artificial floodwater in a glass vial, and the gas bubbles forming on the leaves show that O_2 is being produced in underwater photosynthesis. (B) Underwater net photosynthesis (P_N) at $\text{PAR} = 1,000 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ as a response to CO_2 dissolved in the floodwater followed a saturation curve and is fitted to the means using the FvCB model ($R^2 = 0.99$). (C) Similarly, underwater P_N was fitted to a general light response curve (Ye model, $R^2 = 0.997$) measured with $500 \mu\text{mol L}^{-1}$ dissolved CO_2 in the floodwater to enable estimation of $P_{N\text{max}}$ ($10.97 \mu\text{mol O}_2 \text{ m}^{-2} \text{ s}^{-1}$). In (D), the response of P_N to temperature is shown with $500 \mu\text{mol L}^{-1}$ dissolved CO_2 in the floodwater and $\text{PAR} = 1,000 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, and using a Gaussian fit revealed an optimum for P_N at 25°C . Data points in B–D show the mean $\pm$ SD ($n = 4$).

typical saturation response, increasing as CO_2 was raised to $2,000 \text{ mmol m}^{-3}$ (Figure 2B). The FvCB model estimated the maximum carboxylation rate (V_{cmax}), maximum electron transfer rate (J_{max}), and day respiratory rate (R_{day}) to be $19.73 \mu\text{mol m}^{-2} \text{ s}^{-1}$, $39.34 \mu\text{mol m}^{-2} \text{ s}^{-1}$, and $0.3 \mu\text{mol m}^{-2} \text{ s}^{-1}$, respectively. However, the CO_2 compensation point (Γ^*) and CO_2 saturation point ($C_{i,\text{TUP}}$) are $37.43 \text{ mmol m}^{-3}$ and $2,334 \text{ mmol m}^{-3}$, which are approximately 2-fold and 130-fold atmospheric equilibrium ($\sim 18 \text{ mmol m}^{-3}$ free CO_2), respectively (Table 1).

We diagnosed P_N at contrasting light availabilities with $500 \mu\text{mol CO}_2 \text{ L}^{-1}$ in the floodwater. As expected, underwater P_N also followed a saturation response with increasing light availability. In darkness, the dark respiration (R_{dark}) of branchlets with needles was $1.31 \mu\text{mol O}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Figure 2C). Using the equation from Ye et al. (2013), the light compensation point (I_c) and light saturation point (I_m) were $65.38 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ and $1,666.67 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, respectively. $P_{N\text{max}}$ was estimated to $10.97 \mu\text{mol O}_2 \text{ m}^{-2} \text{ s}^{-1}$ at the given environmental conditions, i.e., dissolved CO_2 at $500 \mu\text{mol L}^{-1}$ at 25°C (Table 1).

Underwater P_N followed an exponential increase with increasing temperature in the tested interval from 10 to 25°C whereafter it steeply decreased with increasing temperature. Using a Gaussian model, we estimated the temperature optimum for underwater P_N in *T. distichum* to 25°C with $500 \mu\text{mol L}^{-1}$ dissolved CO_2 in the floodwater at a photon flux of $1,000 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Figure 2D). Using the same dataset, but without considering temperatures exceeding the optimum for underwater

TABLE 1 *Taxodium distichum* response curve parameter estimation and goodness of fit.

Parameters	Estimated value
P_N–CO_2 response curve (FvCB model)	
V_{cmax}	$19.73 \mu\text{mol m}^{-2} \text{ s}^{-1}$
J_{max}	$39.34 \mu\text{mol m}^{-2} \text{ s}^{-1}$
R_{day}	$0.30 \mu\text{mol m}^{-2} \text{ s}^{-1}$
$C_{i,\text{TUP}}$	2334 mmol m^{-3}
Γ^*	$37.43 \text{ mmol m}^{-3}$
R^2	0.99025
P_N–light response curve (Ye model)	
α	0.0221
$P_{N\text{max}}$	$10.97 \mu\text{mol m}^{-2} \text{ s}^{-1}$
R_{dark}	$1.31 \mu\text{mol m}^{-2} \text{ s}^{-1}$
I_m	$1666.67 \mu\text{mol m}^{-2} \text{ s}^{-1}$
I_c	$65.38 \mu\text{mol m}^{-2} \text{ s}^{-1}$
R^2	0.997

V_{cmax} , maximum carboxylation rate; J_{max} , maximum electron transfer rate; R_{day} , daily respiration rate; $C_{i,\text{TUP}}$, CO_2 saturation point; Γ^* , CO_2 compensation point; R^2 , coefficient of determination; α , initial quantum efficiency; $P_{N\text{max}}$, maximum net photosynthetic rate at light saturation; R_{dark} , dark respiration rate; I_m , light saturation point; I_c , light compensation point.

P_N , we estimated the Q_{10} of P_N to 1.84 demonstrating the strong dependence of underwater P_N on environmental temperature.


3.3 Hydrophobicity and gas film retention by needles of *Taxodium distichum*

At the onset of submergence, *T. distichum* forms a thin gas film on its needles and therefore we aimed at characterizing hydrophobicity, gas film thickness, and other key features known to influence underwater P_N . Macroscopically, the needles are very similar on their adaxial and abaxial sides, but stomatal density differs with more than fourfold higher density of stomata on the abaxial side (Figures 3A–D, G). However, the water-repellent traits were similar, showing a contact angle of 146° on both sides, and although these angles only render the needles hydrophobic (and not superhydrophobic) (Koch and Barthlott, 2009), the hydrophobicity was nevertheless sufficient to initially retain a 35-μm-thick gas layer upon submergence (Figures 3E–G). The needles of *T. distichum* in air have been observed to repel water, and the gas film formed underwater is directly visible as a silvery sheen (Figures 3H, I), and its well-described facilitation of underwater photosynthesis is evident from the bubble formation when the branchlets are submerged in CO₂-rich water in the light (Figure 3J). We also manipulated needle hydrophobicity to enable a direct comparison of underwater P_N of needles with or without a gas film, and we found that the gas film increased underwater P_N 2.1-fold as

compared with needles where gas film formation was prevented (Figure 3K).

3.4 Response of *Taxodium distichum* to long-term submergence

The fact that *Taxodium* can become completely submerged for several months prompted us to conduct a controlled laboratory experiment where we submerged 6-month-old plants for 30 days. During the first 24 days of submergence, the hydrophobicity was lost and the leaf cuticle gradually became colonized with bacteria with the first bacterial cells appearing already after 2 days of submergence (Figure 4A). The colonization of bacteria was accompanied by a decline in hydrophobicity, and leaf gas films dramatically decreased during the first few days of submergence (Figure 4B). This loss of gas films resulted in a significant decline in underwater P_N as indicated by the significant positive correlation between gas film thickness and under P_N (Figure 4C). Key stomatal features (stomatal density and stomatal index) did not change during the first 8 days of submergence (Figure 4D). However, over the entire submergence period of 30 days, significant changes took place at the needle level. The needles started yellowing (Figure 5A), and the yellow was also reflected in a significant decline in chlorophylls (Figure 5B). The combined effect of loss of hydrophobicity and the decline in chlorophylls resulted in a steep decline in underwater photosynthesis already



within the first 8 days of submergence with a predicted $T_{1/2}$ in P_N of 1.85 days (Figure 5C). In addition to the biochemical changes in chlorophyll concentration, the needles also underwent anatomical changes during the 30 days of submergence as all of the palisade cells degraded (Figures 5D, E). Importantly, even with the loss of palisade tissues and the significant declines in chlorophylls, the needles maintained some capacity for underwater P_N during the entire submergence period as underwater P_N never declined below $1\ \mu\text{mol O}_2\ \text{m}^{-2}\ \text{s}^{-1}$.

4 Discussion

Partial or complete submergence of trees is a common phenomenon in several natural or man-made wetlands (Parolin, 2009) and yet the ability of the leaves to photosynthesize under water had not yet previously been studied. In the present study, we found that the needles of *T. distichum* are hydrophobic and retain a thin gas film during submergence, and the gas films greatly enhanced underwater P_N through their beneficial effect on gas exchange between needles and floodwater. We also found that the

needle hydrophobicity was lost with time of submergence, but even after a month of complete submergence, the needles still maintained some capacity for underwater P_N . Nevertheless, the needles had undergone structural and biochemical changes with loss of mesophyll cells and significant declines in chlorophyll concentrations. Below, we are discussing these findings in the context of existing knowledge on flood tolerance of terrestrial plants with emphasis on beneficial leaf traits such as leaf hydrophobicity, gas film formation, and SLA, and we also identify areas of exploration to fill in the many knowledge gaps that still remain.

4.1 Hydrophobicity of *Taxodium distichum* needles and gas film formation

Terrestrial leaves generally perform poorly under water due to the restricted gas exchange in water compared with air resulting in restricted O_2 uptake for respiration or CO_2 uptake for photosynthesis. The poor performance has been clearly demonstrated using the model plant, *Rumex palustris*, showing




FIGURE 4 Changes in needle surface structure, gas film thickness, and underwater P_N with time of submergence of *Taxodium distichum*. Scanning electron micrographs (A) show the changes in surface structure immediately before submergence (day 0, control) and days 2, 4, 6, and 8. Decline in gas film thickness (B) during the 30 days of submergence and the relationship between gas film thickness and underwater P_N (C). A Pearson correlation analysis showed a correlation coefficient of 0.83 and $p < 0.01$. The table (D) shows stomatal aperture, density, opening rate and index, and the same time points. Data are means $\pm$ SD, $n = 3-5$. Different letters within the same column of data indicate $p < 0.05$ (Duncan's multiple comparison).

that underwater P_N of aerial leaves was only 0.5% of the rate in air (Mommer et al., 2006). In stark contrast, leaves of *R. palustris* formed under water could attain rates of underwater P_N at 35% of that in air showing the great benefit of leaf acclimation to underwater gas exchange. However, leaf acclimation is only a feasible strategy for long-term submergence, as production of new aquatic leaves requires reallocation of carbohydrates to fuel leaf. Instead, superhydrophobic leaves that retain a gas film under water have been shown to be a very competitive solution to enhance gas

exchange without further investment in leaf acclimation (Colmer and Pedersen, 2008).

Hydrophobicity can be characterized using the contact angle of a microscopic water droplet. Accordingly, leaf cuticles with contact angles exceeding 150° are classified as superhydrophobic (with rice being a typical example) (Kwon et al., 2014), whereas those with contact angles less than 150° —but larger than 90° (Koch and Barthlott, 2009)—are hydrophobic. In the case of *T. distichum*, the contact angles were just below the 150° cut (146° , Figure 3G),




FIGURE 5

Response to long-term submergence by *Taxodium distichum*. (A) shows habitus photos of an unsubmerged control plant and a plant that has been completely submerged for 30 days. In (B), chlorophylls (Chl_a , Chl_b , and Chl_{a+b}) are shown for unsubmerged control needles and needles that have been submerged for 30 days. (C) shows underwater net photosynthesis (P_N) with time of submergence along with control measurements on unsubmerged branchlets at each sampling point. Data are means $\pm$ SD ($n = 4$), and * and ** indicate $p < 0.05$ and $p < 0.01$, respectively (one-tailed Student's t -test), and the half-life of P_N was calculated using an exponential decay function. Below, (D, E) show cross-sections of an unsubmerged needle and a needle that has been submerged for 30 days. To the left, arrowheads point at palisade tissues, and to the right, * indicates missing palisade tissues.

but the needles nevertheless retained a gas film upon submergence (Figure 3I). The gas films forming on the needles of *T. distichum* were of similar thickness (35 μm , Figure 3G) to those formed by rice leaves (30–60 μm) (Winkel et al., 2014; Kurokawa et al., 2018; Mori et al., 2019) or by wheat leaves (20–40 μm) (Konnerup et al., 2017). Consequently, it is not surprising that the beneficial effects on gas exchange and underwater P_N were significant.

In water, gas diffusion is slow and therefore physiological processes relaying on gas exchange can become restricted by slow substrate supply such as CO_2 for photosynthesis. However, leaf gas films greatly enhance gas exchange (Verboven et al., 2014), and we found that needles with gas films achieved twofold higher photosynthetic rates compared with needles that had the hydrophobicity removed and where gas films therefore did not form (Figure 3K). It has previously been found that leaf gas films can increase underwater P_N up to three- to sevenfold (Colmer and Pedersen, 2008), but the photosynthesis in these experiments was assessed at lower (200 μM) external CO_2 concentrations, where the beneficial effect of gas films on gas exchange is more pronounced (Winkel et al., 2017). Interestingly, the underwater P_N obtained at saturating light and CO_2 levels in the present study matched those of P_N in air; i.e., in both environments, the rates were approximately 10–12 $\mu\text{mol O}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Figures 2B, C) (Neufeld, 1983; Mommer et al., 2006), underlining the significant effect of gas films on the needles of *T. distichum*. Consequently, we propose that a key reason for the stunning flood tolerance of *T. distichum* is its ability to maintain a substantial photosynthetic activity during submergence resulting in both carbohydrate and O_2 production. However, the realized photosynthetic rates greatly depend on both light and CO_2 availability under water and, to a large extent, temperature as well.

4.2 Influence of CO_2 , light, and temperature on underwater P_N of *Taxodium distichum*

CO_2 uptake by *T. distichum* followed a classical FvCB response curve. However, underwater P_N remained negative until the CO_2 compensation point at 37.43 mmol m^{-3} was reached; below, the needles of *T. distichum* consumed more O_2 than they produced. The CO_2 compensation point is equivalent to approximately twofold that of atmospheric equilibrium ($\sim 18 \text{ mmol m}^{-3}$ free CO_2), demonstrating the importance of net CO_2 production in the floodwater in order for the underwater P_N to become positive and therefore result in significant carbohydrate production. The $C_{i,\text{TUP}}$ (CO_2 saturation point) was estimated to 2,334 mmol m^{-3} , or 130-fold atmospheric equilibrium, which is higher than that of submerged rice (Winkel et al., 2013) and submerged wheat (Winkel et al., 2017), and even higher than that of submerged *Hordeum marinum* (Pedersen et al., 2010). It has been demonstrated that the underwater P_N capacity of submerged plants was severely limited at atmospheric equilibrium CO_2 concentrations (Pedersen et al., 2009). Although some studies found that the CO_2 concentration recorded in flooded rice fields was 20–180 times the atmospheric equilibrium concentration (360–

3,240 mmol m^{-3}) (Setter et al., 1987), the diffusion rate of the gas in water is 10,000 times lower than in air, resulting in underwater P_N being still limited (Pedersen et al., 2013). CO_2 availability limitations are a long-standing challenge for submerged plants, but interestingly, rice, wheat, and *H. marinum* retain leaf gas films when submerged, which has been shown to significantly enhance gas exchange. The overall similarity of the CO_2 response in *T. distichum* to the other three terrestrial species with leaf gas films is likely due to the physical effect of the gas films facilitating the exchange between needles and floodwater rather than physiological similarities among these distantly related species.

Light utilization by *T. distichum* under water also showed a saturating response to light when assessed with 500 $\mu\text{mol CO}_2 \text{ L}^{-1}$ in the floodwater. As for CO_2 , underwater P_N was initially negative at low light levels and only reached positive values (I_c) at PAR > 65.38 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Figure 2C). When PAR is higher than the saturation point (I_m) of 1,666.67 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, underwater P_N reaches a maximum value ($P_{N\text{max}}$) of 10.97 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, which is very difficult to achieve due to light loss; thus, underwater P_N is generally limited by PAR. The initial quantum efficiency (α) of 0.0221 absorbed was in the same order of magnitude as that of submerged wheat (Winkel et al., 2017) and *Phalaris arundinacea* (a terrestrial wetland species also forming leaf gas films upon submergence) (Vervuren et al., 1999; Winkel et al., 2017), whereas two other species without gas films utilized light much less efficiently (*Rumex crispus* and *Arrhenatherum elatius*) (Vervuren et al., 1999). These findings emphasize the importance of gas films also for light use efficiency as the light use relies not only on incident light reaching the leaf surfaces but also on entry of CO_2 .

In addition to CO_2 and light, underwater P_N of *T. distichum* was also strongly affected by the environmental temperature. In the temperature range tested, i.e., 10 to 35°C, there was a strong positive relationship between temperature and underwater P_N until the temperature optimum was reached at 25°C after which P_N declined with increasing temperature (Figure 2D). This is a common response of underwater P_N to rising temperature as also demonstrated for two tidal seagrass species, *Thalassia hemprichii* and *Enhalus acoroides* (Pedersen et al., 2016). Being tropical species, these seagrasses showed a temperature optimum for underwater P_N at 33°C, i.e., 8°C above that of *T. distichum*. The Q_{10} of underwater P_N in *T. distichum* was somewhat lower (1.8) than that of *T. hemprichii* (2.0) and *E. acoroides* (2.8) (Pedersen et al., 2016), but it nevertheless shows the strong dependency of temperature for underwater P_N also in *T. distichum*. This is an important point to consider when extrapolating the current laboratory findings to the field situation since the water temperature in the Yangtze River can fluctuate from 11 to 22°C during the time of the year when the trees on the banks become submerged (Yu et al., 2021).

4.3 Responses of *Taxodium distichum* to long-term submergence

The trees growing on the banks of the Yangtze River can become partially or completely submerged for up to 120 days and

still survive (Yang et al., 2023), and we therefore tested the response of *T. distichum* to long-term submergence. Six-month old seedlings were completely submerged for a period of 30 days with sampling of leaf tissue during the period at discrete time points. Leaf gas films quickly diminished, and the decline in gas film thickness was accompanied by a decline in underwater P_N (Figure 4C). The strong relationship between gas film thickness and underwater P_N during long-term submergence has previously been observed in rice (Winkel et al., 2014), but our study represents the first to demonstrate this relationship for a tree species.

The gas films forming on the surfaces of the submerged needles of *T. distichum* persist longer than in other species tested with hydrophobic cuticles. In the present study, the gas films were detectable up to 24 days of submerged after which they had totally vanished (Figure 4B). This is longer than observed for rice, which represents the only other species with superhydrophobic leaves, where gas film thickness have been followed during a submergence event. Here, it was found that gas film thickness was below the detection limit already after 7 days of submergence (Winkel et al., 2014). In *T. distichum*, the needles maintained their ability to photosynthesize also after the gas films were lost at a rate of approximately $1 \mu\text{mol O}_2 \text{ m}^{-2} \text{ s}^{-1}$, which is similar to the photosynthetic rate of rice and wheat once the leaf gas films of these species have also disappeared due to long-term submergence (Winkel et al., 2014; Konnerup et al., 2017). It is still not fully understood why the loss in hydrophobicity occurs during submergence. However, the present study as well as one on wheat (Konnerup et al., 2017) clearly demonstrated that a biofilm was established during submergence (Figure 3I), but if this biofilm is the result or the cause of loss of hydrophobicity remains unknown.

Distinct anatomical and biochemical changes occurred during the long-term submergence event. The decline in total chlorophylls was significant with initial values at $2.1 \text{ mg g}^{-1} \text{ DM}$ and only $1.0 \text{ mg g}^{-1} \text{ DM}$ after 30 days of submergence (Figure 5B). While this decline will have significant consequences for the light capturing capabilities, the decline in submerged rice was even more pronounced where the initial levels are at approximately $18 \text{ mg g}^{-1} \text{ DM}$ to less than $5 \text{ mg g}^{-1} \text{ DM}$ in only 2 weeks. The decrease in chlorophyll concentration observed in the present study may be one of the important reasons for the decrease in underwater P_N , as it has been shown that underwater P_N of rice is positively correlated with leaf chlorophyll concentration (Winkel et al., 2014). The significant decrease in chlorophyll concentration is likely a result of palisade tissue loss in needles (Figures 5D, E). To our knowledge, similar observations are missing in the literature as previous studies have focused on anatomy of leaves formed during the submergence event (Mommer et al., 2007) and not on acclimation of already existing leaves. The lysis of palisade tissues observed towards the end of the 30-day submergence event might be accompanied by water infiltration in the newly formed cavities, and such water-filled cavities would slow down intra-tissue diffusion of O_2 and CO_2 (Armstrong, 1980). Interestingly, the parallel decline in chlorophyll concentration and leaf gas film thickness makes it difficult to identify the primary causal effect of the observed decline in underwater P_N .

5 Conclusions and perspectives

During the 30-day submergence period, no growth or formation new leaves were observed, and therefore, *T. distichum* shows a quiescence response to submergence (cf. Bailey-Serres and Voisenek, 2008). The hydrophobicity of the needles declined during the submergence event, resulting in loss of gas films. However, the chlorophyll concentration of the needles also declined significantly, and it was therefore not possible to identify the main cause of the corresponding significant decline in underwater P_N .

Several questions still remain unresolved in order to fully understand the striking ability of *T. distichum* to withstand partial or complete submergence for months. We propose that future research concentrate on unraveling the finer details in needle anatomy and biochemistry as these changes occur during submergence. For example, the lysis of palisade tissues should be further studied in order to understand if the lysis is merely a consequence of senescence processes or if the lysis is actively controlled via programmed cell death with the aim of acclimating the leaves to a low-light environment and the slow diffusion of gases in water. We also suggest to investigate if changes in cuticle structure take place beyond those involved in surface hydrophobicity. A thinning of the cuticle would greatly enhance diffusion of O_2 and CO_2 from the floodwater to the needles' tissues and thereby enhance the supply of O_2 for dark respiration or CO_2 for underwater photosynthesis. In addition, whether the bacteria colonized on leaf surface will cause negative impacts on the leaf cells and hence affect hydrophobicity and photosynthesis, and the difference in underwater P_N capacity and detailed submergence tolerance mechanisms of *T. distichum* seedlings and big trees are also interesting questions that are worthy of further investigation.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding authors.

Author contributions

JG: Writing – original draft, Visualization, Software, Data curation. JX: Methodology, Writing – review & editing, Validation, Supervision. YY: Writing – review & editing, Supervision, Resources. OP: Writing – original draft, Methodology, Funding acquisition, Data curation. JH: Writing – review & editing, Methodology, Funding acquisition, Conceptualization.

Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. This work

was supported by grants from the Jiangsu Special Fund on Technology Innovation of Carbon Dioxide Peaking and Carbon Neutrality, grant no. BE2022420, and the Jiangsu Long-term Scientific Research Base for *Taxodium* Rich Breeding and Cultivation, grant no. LYKJ(2021)05. OP was supported by the Carlsberg Foundation, grant no. CF23-0039.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Armstrong, W. (1980). "Aeration in higher plants," in *Advances in botanical research* (Elsevier) 7, 225–332. doi: 10.1016/S0065-2296(08)60089-0
- Ayi, Q., Zeng, B., Liu, J., Li, S., Van Bodegom, P. M., and Cornelissen, J. H. C. (2016). Oxygen absorption by adventitious roots promotes the survival of completely submerged terrestrial plants. *Ann. Bot.* 118, 675–683. doi: 10.1093/aob/mcw051
- Bailey-Serres, J., and Voesenek, L. (2008). Flooding stress: acclimations and genetic diversity. *Annu. Rev. Plant Biol.* 59, 313–339. doi: 10.1146/annurev.arplant.59.032607.092752
- Colmer, T., and Pedersen, O. (2008). Oxygen dynamics in submerged rice (*Oryza sativa*). *New Phytol.* 178, 326–334. doi: 10.1111/j.1469-8137.2007.02364.x
- Colmer, T. D., Winkler, A., and Pedersen, O. (2011). A perspective on underwater photosynthesis in submerged terrestrial wetland plants. *AoB Plants* 2011, plr030. doi: 10.1093/aobpla/plr030
- Ding, D., Liu, M., Arif, M., Yuan, Z., Li, J., Hu, X., et al. (2021). Responses of ecological stoichiometric characteristics of carbon, nitrogen, and phosphorus to periodic submergence in mega-reservoir: growth of *Taxodium distichum* and *Taxodium ascendens*. *Plants* 10, 2040. doi: 10.3390/plants10102040
- Garssen, A. G., Baatrup-Pedersen, A., Riis, T., Raven, B. M., Hoffman, C. C., Verhoeven, J. T., et al. (2017). Effects of increased flooding on riparian vegetation: Field experiments simulating climate change along five European lowland streams. *Global Change Biol.* 23, 3052–3063. doi: 10.1111/gcb.13687
- Glenz, C., Schlaepfer, R., Iorgulescu, I., and Kienast, F. (2006). Flooding tolerance of Central European tree and shrub species. *For. Ecol. Manage.* 235, 1–13. doi: 10.1016/j.foreco.2006.05.065
- Guo, J., Xue, J., Hua, J., Yin, Y., Creech, D. L., and Han, J. (2023). Research status and trends of *taxodium distichum*. *HortScience* 58, 317–326. doi: 10.21273/HORTSCI17045-22
- He, X., Wang, T., Wu, K., Wang, P., Qi, Y., Arif, M., et al. (2021). Responses of swamp cypress (*Taxodium distichum*) and Chinese willow (*Salix matsudana*) roots to periodic submergence in mega-reservoir: Changes in organic acid concentration. *Forests* 12, 203. doi: 10.3390/f12020203
- Hegde, S. M., and Krishnaswamy, K. (2021). Measurement of stomatal density stomatal index in some species of terrestrial orchids. *Ann. Agri Bio Res.* 26, 173–177.
- Herzog, M., Pellegrini, E., and Pedersen, O. (2023). A meta-analysis of plant tissue O₂ dynamics. *Funct. Plant Biol.* 10, 1071/FP22294. doi: 10.1071/FP22294
- Iversen, L. L., Winkler, A., Båstrup-Spohr, L., Hinke, A. B., Alahuhta, J., Baatrup-Pedersen, A., et al. (2019). Catchment properties and the photosynthetic trait composition of freshwater plant communities. *Science* 366, 878–881. doi: 10.1126/science.aay5945
- Kirk, J. T. O. (1994). *Light and photosynthesis in aquatic ecosystems* (New York: Cambridge Univ Press). doi: 10.1017/CBO9780511623370
- Koch, K., and Barthlott, W. (2009). Superhydrophobic and superhydrophilic plant surfaces: an inspiration for biomimetic materials. *Philos. Trans. R. Soc. A: Mathematical Phys. Eng. Sci.* 367, 1487–1509. doi: 10.1098/rsta.2009.0022
- Konnerup, D., Winkler, A., Herzog, M., and Pedersen, O. (2017). Leaf gas film retention during submergence of 14 cultivars of wheat (*Triticum aestivum*). *Funct. Plant Biol.* 44, 877–887. doi: 10.1071/FP16401
- Kurokawa, Y., Nagai, K., Huan, P. D., Shimazaki, K., Qu, H., Mori, Y., et al. (2018). Rice leaf hydrophobicity and gas films are conferred by a wax synthesis gene (LGF 1) and contribute to flood tolerance. *New Phytol.* 218, 1558–1569. doi: 10.1111/nph.15070
- Kwon, D. H., Huh, H. K., and Lee, S. J. (2014). Wettability and impact dynamics of water droplets on rice (*Oryza sativa* L.) leaves. *Experiments fluids* 55, 1–9. doi: 10.1007/s00348-014-1691-y
- Li, C., Zhong, Z., and Liu, Y. (2006). Effect of soil water changes on photosynthetic characteristics of *Taxodium distichum* seedlings in the hydro-fluctuation belt of the Three Gorges Reservoir area. *Front. Forestry China* 1, 163–169. doi: 10.1007/s11461-006-0013-9
- Li, Q., Zhou, L., Chen, Y., Xiao, N., Zhang, D., Zhang, M., et al. (2022). Phytochrome interacting factor regulates stomatal aperture by coordinating red light and abscisic acid. *Plant Cell* 34, 4293–4312. doi: 10.1093/plcell/koac244
- Liang, X.-Y., and Liu, S.-R. (2017). A review on the FvCB biochemical model of photosynthesis and the measurement of A-Ci curves. *Chin. J. Plant Ecol.* 41, 693–706. doi: 10.17521/cjpe.2016.0283
- Lin, C., Peralta Ogorek, L. L., Liu, D., Pedersen, O., and Sauter, M. (2023). A quantitative trait locus conferring flood tolerance to deepwater rice regulates the formation of two distinct types of aquatic adventitious roots. *New Phytol.* 238, 1403–1419. doi: 10.1111/nph.18678
- Maberly, S. C., and Madsen, T. V. (2002). Freshwater angiosperm carbon concentrating mechanisms: processes and patterns. *Funct. Plant Biol.* 29, 393–405. doi: 10.1071/PP01187
- Madsen, T. V., and Sand-Jensen, K. (1991). Photosynthetic carbon assimilation in aquatic macrophytes. *Aquat. Bot.* 41, 5–40. doi: 10.1016/0304-3770(91)90037-6
- Mommer, L., Pedersen, O., and Visser, E. J. W. (2004). Acclimation of a terrestrial plant to submergence facilitates gas exchange under water. *Plant Cell Environ.* 27, 1281–1287. doi: 10.1111/j.1365-3040.2004.01235.x
- Mommer, L., Pons, T. L., and Visser, E. J. (2006). Photosynthetic consequences of phenotypic plasticity in response to submergence: *Rumex palustris* as a case study. *J. Exp. Bot.* 57, 283–290. doi: 10.1093/jxb/erj015
- Mommer, L., Wolters-Arts, M., Andersen, C., Visser, E. J., and Pedersen, O. (2007). Submergence-induced leaf acclimation in terrestrial species varying in flooding tolerance. *New Phytol.* 176, 337–345. doi: 10.1111/j.1469-8137.2007.02166.x
- Mori, Y., Kurokawa, Y., Koike, M., Malik, A. I., Colmer, T. D., Ashikari, M., et al. (2019). Diel O₂ dynamics in partially and completely submerged deepwater rice: leaf gas films enhance internodal O₂ status, influence gene expression and accelerate stem elongation for 'snorkelling' during submergence. *Plant Cell Physiol.* 60, 973–985. doi: 10.1093/pcp/pcz009
- Neinhuis, C., and Barthlott, W. (1997). Characterization and distribution of water-repellent, self-cleaning plant surfaces. *Ann. Bot.* 79, 667–677. doi: 10.1006/anbo.1997.0400
- Neufeld, H. S. (1983). Effects of light on growth, morphology, and photosynthesis in baldcypress (*Taxodium distichum* (L.) Rich.) and pondcypress (*T. ascendens* Brongn.) seedlings. *Bull. Torrey Botanical Club* 110, 43–54. doi: 10.2307/2996516
- Parolin, P. (2009). Submerged in darkness: adaptations to prolonged submergence by woody species of the Amazonian floodplains. *Ann. Bot.* 103, 359–376. doi: 10.1093/aob/mcn216
- Pedersen, O., and Colmer, T. D. (2012). Physical gills prevent drowning of many wetland insects, spiders and plants. *J. Exp. Biol.* 215, 705–709. doi: 10.1242/jeb.065128
- Pedersen, O., Colmer, T. D., Borum, J., Zavala-Perez, A., and Kendrick, G. A. (2016). Heat stress of two tropical seagrass species during low tides—impact on underwater net photosynthesis, dark respiration and diel *in situ* internal aeration. *New Phytol.* 210, 1207–1218. doi: 10.1111/nph.13900
- Pedersen, O., Colmer, T. D., and Sand-Jensen, K. (2013). Underwater photosynthesis of submerged plants – recent advances and methods. *Front. Plant Sci.* 4. doi: 10.3389/fpls.2013.00140

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2024.1355729/full#supplementary-material>

- Pedersen, O., Malik, A. I., and Colmer, T. D. (2010). Submergence tolerance in *Hordeum marinum*: dissolved CO₂ determines underwater photosynthesis and growth. *Funct. Plant Biol.* 37, 524–531. doi: 10.1071/FP09298
- Pedersen, O., Rich, S. M., and Colmer, T. D. (2009). Surviving floods: leaf gas films improve O₂ and CO₂ exchange, root aeration, and growth of completely submerged rice. *Plant J.* 58, 147–156. doi: 10.1111/j.1365-3113X.2008.03769.x
- Prins, H. B. A., and Elzenga, J. T. M. (1989). Bicarbonate utilization: function and mechanism. *Aquat. Bot.* 34, 59–83. doi: 10.1016/0304-3770(89)90050-8
- Raskin, I., and Kende, H. (1983). How does deep water rice solve its aeration problem? *Plant Physiol.* 72, 447–454. doi: 10.1104/pp.72.2.447
- Rich, S. M., Pedersen, O., Ludwig, M., and Colmer, T. D. (2013). Shoot atmospheric contact is of little importance to aeration of deeper portions of the wetland plant *Meionectes brownii*; submerged organs mainly acquire O₂ from the water column or produce it endogenously in underwater photosynthesis. *Plant Cell Environ.* 36, 213–223. doi: 10.1111/j.1365-3040.2012.02568.x
- Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675. doi: 10.1038/nmeth.2089
- Sculthorpe, C. D. (1967). *The biology of aquatic vascular plants* (London: Edward Arnold Ltd).
- Setter, T., Kupkanchanakul, T., Kupkanchanakul, K., Bhekasut, P., Wiengweera, A., and Greenway, H. (1987). Concentrations of CO₂ and O₂ in floodwater and in internodal lacunae of floating rice growing at 1–2 metre water depths. *Plant Cell Environ.* 10, 767–776. doi: 10.1111/1365-3040.ep11604764
- Setter, T. L., Waters, I., Wallace, I., Bekhasut, P., and Greenway, H. (1989). Submergence of rice I. Growth and photosynthetic response to CO₂ enrichment of floodwater. *Aust. J. Plant Physiol.* 16, 251–263. doi: 10.1071/PP9890251
- Sikorska, D., Papierowska, E., Szatylowicz, J., Sikorski, P., Suprun, K., and Hopkins, R. J. (2017). Variation in leaf surface hydrophobicity of wetland plants: the role of plant traits in water retention. *Wetlands* 37, 997–1002. doi: 10.1007/s13157-017-0924-2
- Smart, R., and Barko, J. (1985). Laboratory culture of submersed freshwater macrophytes on natural sediments. *Aquat. Bot.* 21, 251–263. doi: 10.1016/0304-3770(85)90053-1
- Taylor, S. A., and Smith, W. K. (2017). “Effect of increased salinity exposure on photosynthetic indicators and water status of bald cypress (*Taxodium distichum*) seedlings”. In: *RURALS: Review of Undergraduate Research in Agricultural and Life Sciences* 11 (1), 1. Available at: <http://digitalcommons.unl.edu/rurals/vol11/iss1/1>.
- Teakle, N. L., Colmer, T. D., and Pedersen, O. (2014). Leaf gas films delay salt entry and enhance underwater photosynthesis and internal aeration of *M. elilottii* sicutus submerged in saline water. *Plant Cell Environ.* 37, 2339–2349. doi: 10.1111/pce.12269
- Verboven, P., Pedersen, O., Ho, Q. T., Nicolai, B. M., and Colmer, T. D. (2014). The mechanism of improved aeration due to gas films on leaves of submerged rice. *Plant Cell Environ.* 37, 2433–2452. doi: 10.1111/pce.12300
- Vervuren, P., Beurskens, S., and Blom, C. (1999). Light acclimation, CO₂ response and long-term capacity of underwater photosynthesis in three terrestrial plant species. *Plant Cell Environ.* 22, 959–968. doi: 10.1046/j.1365-3040.1999.00461.x
- Vervuren, P. J. A., Blom, C. W. P. M., and De Kroon, H. (2003). Extreme flooding events on the Rhine and the survival and distribution of riparian plant species. *J. Ecol.* 91, 135–146. doi: 10.1046/j.1365-2745.2003.00749.x
- Wang, C., Li, C., Wei, H., Xie, Y., and Han, W. (2016). Effects of long-term periodic submergence on photosynthesis and growth of *Taxodium distichum* and *Taxodium ascendens* saplings in the hydro-fluctuation zone of the Three Gorges Reservoir of China. *PloS One* 11, e0162867. doi: 10.1371/journal.pone.0162867
- Wang, T., Wei, H., Ma, W., Zhou, C., Chen, H., Li, R., et al. (2019). Response of *Taxodium distichum* to winter submergence in the water-level-fluctuating zone of the Three Gorges Reservoir region. *J. Freshw. Ecol.* 34, 1–17. doi: 10.1080/02705060.2018.1470041
- Wang, Z., Yin, M., Creech, D. L., and Yu, C. (2022). Microsporogenesis, pollen ornamentation, viability of stored *Taxodium distichum* var. *distichum* pollen and its feasibility for cross breeding. *Forests* 13, 694. doi: 10.3390/f13050694
- Winkel, A., Colmer, T. D., Ismail, A. M., and Pedersen, O. (2013). Internal aeration of paddy field rice (*O. ryza sativa*) during complete submergence—importance of light and floodwater O₂. *New Phytol.* 197, 1193–1203. doi: 10.1111/nph.12048
- Winkel, A., Colmer, T. D., and Pedersen, O. (2011). Leaf gas films of *Spartina anglica* enhance rhizome and root oxygen during tidal submergence. *Plant Cell Environ.* 34, 2083–2092. doi: 10.1111/j.1365-3040.2011.02405.x
- Winkel, A., Herzog, M., Konnerup, D., Floytrup, A. H., and Pedersen, O. (2017). Flood tolerance of wheat—the importance of leaf gas films during complete submergence. *Funct. Plant Biol.* 44, 888–898. doi: 10.1071/FP16395
- Winkel, A., Pedersen, O., Ella, E., Ismail, A. M., and Colmer, T. D. (2014). Gas film retention and underwater photosynthesis during field submergence of four contrasting rice genotypes. *J. Exp. Bot.* 65, 3225–3233. doi: 10.1093/jxb/eru166
- Winkel, A., Visser, E. J., Colmer, T. D., Brodersen, K. P., Voesenek, L. A., Sand-Jensen, K., et al. (2016). Leaf gas films, underwater photosynthesis and plant species distributions in a flood gradient. *Plant Cell Environ.* 39, 1537–1548. doi: 10.1111/pce.12717
- Yang, Q., Bejarano, M. D., Ma, W., Salam, M., Pu, B., Wei, H., et al. (2023). Effects of long-term submergence on non-structural carbohydrates and N and P concentrations of *Salix matSudana* along the Three Gorges Reservoir. *Hydrobiologia* 850, 2035–2047. doi: 10.1007/s10750-023-05215-5
- Ye, Z. P., Suggett, D. J., Robakowski, P., and Kang, H. J. (2013). A mechanistic model for the photosynthesis–light response based on the photosynthetic electron transport of photosystem II in C₃ and C₄ species. *New Phytol.* 199, 110–120. doi: 10.1111/nph.12242
- Yu, S., Sheng, L., Zhan, L., and Niluo, Z. (2021). Analysis of temperature variation characteristics in the three gorges reservoir area after impoundment of the three gorges dam. *Acta Ecologica Sin.* 41, 384–389. doi: 10.1016/j.chnaes.2020.06.003
- Zhang, J., Sun, H., Gao, D., Qiao, L., Liu, N., Li, M., et al. (2020). Detection of canopy chlorophyll content of corn based on continuous wavelet transform analysis. *Remote Sens.* 12, 2741. doi: 10.3390/rs12172741
- Zhang, Q., Huber, H., Beljaars, S. J. M., Birnbaum, D., De Best, S., De Kroon, H., et al. (2017). Benefits of flooding-induced aquatic adventitious roots depend on the duration of submergence: linking plant performance to root functioning. *Ann. Bot.* 120, 171–180. doi: 10.1093/aob/mcx049