

Can parasitism by *Tamarixia radiata* (Hymenoptera: Eulophidae) affect the movement and oviposition behavior of *Diaphorina citri* (Hemiptera: Psyllidae)?

Juliana S. Oliveira (✉ julianabioso@gmail.com)

University of São Paulo (USP)

Carolina Reigada

Federal University of São Carlos (UFSCar)

Alexandre J.F. Diniz

University of São Paulo (USP)

Idemauro A.R. Lara

University of São Paulo (USP)

Rafael A. Moral

Maynooth University

José R.P. Parra

University of São Paulo (USP)

Research Article

Keywords: Aggregation, intraspecific competition, choice of oviposition and feeding site, displacement, Asian citrus psyllid.

Posted Date: August 12th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1908974/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Journal of Insect Behavior on December 29th, 2022. See the published version at <https://doi.org/10.1007/s10905-022-09811-6>.

Abstract

Diaphorina citri is the vector of the bacteria that cause HLB. The psyllid acquires the pathogen after feeding on infected plants and transmits it to other trees as it moves through the grove. The psyllid's movement is directly affected by abiotic and biotic factors, such as the presence of conspecific insects and natural enemies. This study evaluated the effects of the presence of conspecific nymphs either not parasitized or parasitized by *Tamarixia radiata* on the movement and oviposition of adult female *D. citri*. *D. citri* females were released into cages containing a row of four equidistant *Murraya paniculata* plants. The first plant, the release point, had one of three conditions: no nymphs; non-parasitized nymphs (high and low densities); or previously parasitized nymphs. At 24, 48, and 72 h following release, the females on each plant in the cage were counted. At the end of the observations (72 h), the females were removed, and the eggs laid on the plants were counted. Discrete data analysis showed that in the presence of a high density of non-parasitized nymphs, adult psyllids were more homogeneously distributed among the plants. However, the highest oviposition rates were observed on plants without nymphs. On the other hand, in the presence of nymphs, the largest number of females remained at the release site and more eggs were deposited on plants with parasitized nymphs. This information will help to improve psyllid management techniques and releases of their natural enemy, according to their distribution in the grove.

Introduction

Biotic and abiotic factors directly affect the ability of living beings to move, as they are decisive in determining the ability of species to disperse and survive (Dingle 1972; Stinner et al. 1983; Minter et al. 2018; Naranjo 2019). Understanding the effects of biotic and abiotic factors on the behavior of insects during their movement and search for oviposition and feeding sites can significantly contribute to management programs for insect agricultural pests (Bullock et al. 2002; Mazzi and Dorn 2012).

The Asian citrus psyllid *Diaphorina citri* Kuwayama, 1908 (Hemiptera: Psyllidae), which transmits the bacteria that cause “huanglongbing” (HLB), is native to Asia. Due to its dispersal capacity, which increases from the immature to the adult stage (Sétamou et al. 2020), the species is now present on most continents (Halbert 1998; French et al. 2001; Bové 2006; Shimwela et al. 2016; Oke et al. 2020).

Although *D. citri* adults tend to remain in the same grove after their arrival, they continually move over short or long distances within the orchard (Boina et al. 2009; Kobori et al. 2011; Lewis-Rosenblum et al. 2015; Tomaseto et al. 2016, 2019). During these movements, adults and nymphs can acquire the bacteria (HLB) by feeding on an infected plant for 30 minutes or more (Capoor et al. 1974; Hung et al. 2004; Pelz-Stelinski et al. 2010). The adult becomes an HLB transmitter 3 days after infection (Xu et al. 1988) and the nymphs transmit HLB after they become adults (Inoue et al. 2009; Pelz-Stelinski et al. 2010). Infected plants become unviable after 4–5 years, and their fruits drop before ripening and have lower sugar content (Grafton-Cardwell et al. 2013; Dala-Paula et al. 2019).

Considered the most destructive citrus disease, in 2014 HLB reduced orange production in Florida by approximately 43% (Singerman and Useche 2016) and has decimated more than 50 million trees in Brazil (CDA 2019). The disease significantly reduces citrus production in Africa (Garnier and Bové 1996; FAO 2002), Asia (Martinez and Wallace 1968; Bhavakul et al. 1982; Bové 1986; Sharma 2008; Shivankar et al. 2000), and Oceania (Fraser 1978). Infestation reduction is achieved by eliminating infected plants, using healthy seedlings for planting, and managing the vector insect through application of chemical products and/or release of natural enemies, mainly *Tamarixia radiata* Waterston, 1992 (Hymenoptera: Eulophidae), an ectoparasitoid that attacks the psyllid nymphs (Diniz et al. 2020; Garcia et al. 2021).

The movement of psyllids in orchards is influenced by the presence and density of shoots available on the host plants, temperature, relative humidity, and wind (Yamamoto et al. 2001; Lewis-Rosenblum et al. 2015; Tomaseto et al. 2017; Garcia et al. 2019; Antolinez et al. 2021). In addition to these factors, the frequency and pattern of movement of *D. citri* adults in the search for host plants and the selection of microhabitats can also be influenced by competitive interactions of conspecifics and by encounters with natural enemies (Fritz 1982; Nufio and Papaj 2001). Natural control agents can, in addition to directly reducing populations, indirectly affect the behavior, physiology, morphology, and development of their prey/hosts. This non-consumptive effect was evidenced in studies with parasites (Holmes and Bethel 1972; Price 1980) and predators (Nelson et al. 2004; Hermann and Thaler 2014; Seo et al. 2018) that found reductions in the population growth of aphids, *Acyrtosiphon pisum* (Homoptera: Aphididae), *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae), and *Diaphorina citri* caused only by the presence of predatory insects, such as damsel bugs (Hemiptera: Nabidae), *Podisus maculiventris* (Heteroptera: Pentatomidae), and *Hippodamia convergens* (Coleoptera: Coccinellidae), respectively. Little information is available on this effect caused by parasitoids.

Evaluation of the effects of intraspecific competition (i.e., density of conspecifics) and the presence of natural enemies on the pattern of movement and oviposition of *D. citri* adults and the consequent potential for HLB infection/transmission in citrus orchards can contribute significantly to the improvement of techniques used in integrated management (IPM) of this insect pest. This study evaluated the movement of psyllid females and the pattern of egg distribution in experimental cages containing host plants in different situations: (i) healthy plants (i.e., without psyllid nymphs), (ii) plants with *D. citri* nymphs, and (iii) plants with nymphs parasitized by *T. radiata*.

Materials And Methods

Colony of insects and host plants

For this study, nymphs and adults (females) of *D. citri* and adults of *T. radiata* were obtained from the rearing colonies at the Laboratory of Insect Biology of the University of São Paulo –ESALQ/USP, Piracicaba, maintained following the method of Parra et al. (2016). The host plants used were orange jessamine, *Murraya paniculata* (L.) Jack, kept in a greenhouse under natural conditions and with automatic ventilation that maintains the temperature between 25 and 28°C.

Experimental design

In order to evaluate the effects of the presence of conspecifics and the presence of natural enemies on the movement and oviposition pattern of *D. citri* during the search for available plants for food and development of its offspring, a group of adult psyllid females ($n = 90$), 9 days old (average period of onset of oviposition) (Nava et al. 2007) was used. Prior to the beginning of the experiment, these females were kept for 9 days in cages containing males, to ensure mating. The female psyllids were transferred to fabric tunnel cages ($200 \cdot 60 \cdot 70$ cm), each containing 4 tubes (250 ml) with 4 orange jessamine shoots per tube and spaced equidistantly from each other (Fig. 1). The shoots had an average length of 1 cm, as this is the most attractive size for oviposition (Diniz 2013). After the insects were released, the number of females on each plant (A, B, C, and D) was counted to assess the movement of insects on the available plants over 24, 48 and 72 h. At the end of 72 h, the total number of eggs laid on each plant was also counted. The cages were kept in the greenhouse under the same conditions described above.

In the first set of experiments, the effect of intraspecific competition on the movement and distribution of *D. citri* eggs was evaluated. Inside the experimental cages, 4th and 5th instar psyllid nymphs were placed on the first plant (plant A, release site). Two nymph densities (*i.e.*, treatments) were established: (i) low density, with 15 nymphs present; and (ii) high density, with 70 nymphs present. A control consisted of plants with no nymphs. Ten replicates were performed for each treatment and control.

In the second set of experiments, the effect of the presence of natural enemies on the movement and distribution behavior of *D. citri* eggs was evaluated, using two treatments: (i) presence of 25 non-parasitized *D. citri* nymphs on plant A and (ii) presence of 25 nymphs previously parasitized by *T. radiata* on plant A. Ten replicates were performed for each treatment.

Data analysis

For the analysis of the movement of insects between plants over time (24, 48, and 72 h) an extension of the discrete-time transition models for clustered nominal data was used, according to the procedure proposed by Lara et al. (2020). In this procedure, a multinomial distribution was assumed for the response variable, with five categories: Plant A, Plant B, Plant C, Plant D, and none, each representing the preference of insect groups, and the generalized logit models were fitted (Agresti, 2013). The first-order Markov stochastic assumption was used to fit the model, *i.e.*, the number of insects on each of the plants was incorporated into the linear predictor of the model as additional covariates. To decide between a stationary process or not, the likelihood ratio test proposed by Lara et al. (2017) was used. If the stationary model is assumed, in practice, insects move through the vases, following a homogeneous pattern over time.

For the analysis of the variation of the total number of eggs deposited on the four plants (A, B, C, and D) after 72 h, considering the possible effect of treatments (experiment 1: high nymph densities, low nymph density, and absence of nymphs; experiment 2: parasitized nymphs and non-parasitized nymphs), the Generalized Linear Model was fitted assuming a Poisson distribution, considering the Quasi-Poisson

model when overdispersion is found, which is a common phenomenon with count data with a variance much greater than the mean (McCullagh and Nelder, 1989). The best statistical model was chosen through comparisons between the adjustments of the complete models and models with the reduction of explanatory variables by the Akaike criterion (AIC). The quality of the adjustments obtained by the models was assessed using the “half-normal-plot” graph with simulated envelopes at the 95% level (Moral et al. 2017). All analyses were performed using the R software (R Core Team 2019).

Results

Experiment 1: Movement rate of females and egg distribution pattern of Diaphorina citri in the presence of non-parasitized nymphs

Time variation (24, 48, and 72 h), covariates for psyllid occupation options (plants A, B, C, D, and none), and the presence of nymphs (low or high density) significantly affected the movement of the adult females of *D. citri* (LRV = 55.3394, df = 28, $p < 0.01$; LRV = 746.67, df = 16, $p < 0.01$ and LRV = 13,640.51, df = 8, $p < 0.01$, respectively). The results showed that, in the control group, in which nymphs were absent, the average proportion of insects remained higher on plant A (insect release point) during the three evaluation periods (24, 48, and 72 h). The proportion of insects on plant A when nymphs were present, however, was lower, and over time was reduced in both treatments, at low and high nymph densities (Fig. 2). With the passage of time, the females tended to avoid remaining on the plants with nymphs, moving toward the other plants in the cages. The proportion of psyllids elsewhere than on the plants was also higher and increased with time when nymphs were present, indicating a greater tendency to move (Table 1).

Table 1 Numbers (mean $\pm$ standard error) of insects found on the four plants (A, B, C, D) after release of *Diaphorina citri* on plant A (with the presence or absence of nymphs in low or high density), over 3 days.

Treatment	Period (h)	Mean no. of insects			
		Plant A	Plant B	Plant C	Plant D
Absence of nymphs (Control)	24	19.5 ± (2.5) Aa	8.4 ± (1.9) Aab	3.9 ± (1.2) Ab	5.1 ± (1.1) Ab
	48	18.9 ± (2.3) Aa	9.0 ± (2.3) Aab	4.2 ± (0.9) Ab	6.0 ± (1.3) Ab
	72	15.6 ± (2.3) Aa	10.1 ± (2.1) Aab	4.3 ± (1.1) Ab	7.6 ± (1.7) Ab
Low nymph density	24	12.1 ± (2.8) Aa	6.0 ± (1.1) Aab	3.9 ± (1.1) Ab	6.9 ± (2.0) Aab
	48	6.0 ± (2.1) ABa	6.7 ± (0.9) Aa	4.5 ± (1.3) Aa	7.5 ± (2.1) Aa
	72	3.8 ± (1.6) Ba	5.0 ± (1.1) Aa	5.3 ± (1.5) Aa	5.7 ± (1.9) Aa
High nymph density	24	8.0 ± (1.6) Aa	8.1 ± (1.8) Aa	3.6 ± (0.6) Ab	9.1 ± (1.6) Aa
	48	3.4 ± (0.9) Bb	7.1 ± (2.0) Aab	2.7 ± (0.5) Ab	10.1 ± (2.0) Aa
	72	2.4 ± (1.1) Bb	8.0 ± (2.1) Aa	4.5 ± (0.7) Aab	8.5 ± (2.1) Aa

Means followed by different letters indicate a significant difference in the number of insects present on each plant, within each treatment at significance level $\alpha = 5\%$ (likelihood ratio test).

The presence of nymphs (low or high density) and the distance of the plants from the release point (plant A) significantly affected the distribution of eggs on the different plants ($F_{6,108} = 3.83$, $p < 0.01$). In the control group, the highest number of eggs was found on plant A. In plants B, C, and D, fewer eggs were laid and the numbers of eggs deposited on these plants did not differ significantly (Table 2). The presence of nymphs at the release point (plant A) resulted in changes in the distribution of eggs by females. When the density of nymphs on plant A was low, the distribution of eggs was more uniform among the plants in the cage, with no significant difference among plants. However, when the density of nymphs on plant A was high, more eggs were observed on plants D, B, and C, in that order, 72 h after the females were released into the cages (Table 2).

Table 2 Numbers (mean ± SD) of eggs of *Diaphorina citri* found on the four plants (A, B, C, D), in the presence (low or high density) or absence of nymphs on plant A (insect release point) 72 h after adults were released.

Treatment	Mean no. of eggs				Total
	Plant A	Plant B	Plant C	Plant D	
Absence of nymphs (Control)	205.7 ($\pm$ 63.5) Aa	74.7 ($\pm$ 22.7) Ba	42.1 ($\pm$ 13.7) Ba	58.2 ($\pm$ 31.3) Ba	3807 A
Low density of nymphs	26.6 ($\pm$ 10.3) Ab	49.6 ($\pm$ 16.4) Aa	27.4 ($\pm$ 10.9) Aa	39.8 ($\pm$ 20.5) Aa	1434 B
High density of nymphs	8.9 ($\pm$ 3.9) Ab	87.0 ($\pm$ 41.7) Ba	27.1 ($\pm$ 12.4) Aa	93.9 ($\pm$ 30.8) Ba	2169 B
Total	2412	2113	966	1919	

Means followed by different letters indicate a significant difference in the number of eggs present on each plant, within each treatment at significance level $\alpha = 5\%$ (likelihood ratio test).

*Experiment 2: Movement rate of females and pattern of egg distribution of *Diaphorina citri* in the presence of non-parasitized nymphs and nymphs parasitized by *Tamarixia radiata**

The variation in time (24, 48, and 72 h), the covariates for occupation options by the psyllid (plants A, B, C, D, and none), and the presence of nymphs (parasitized or not parasitized) significantly affected the movement on the plants by adult females of *D. citri* (LRV = 60.44, df = 24, $p < 0.01$; LRV = 405.92, df = 16, $p < 0.01$ and LRV = 3258.7, df = 4, $p < 0.01$, respectively). The results showed that plant A, the site of adult release and the presence of nymphs, had more adult psyllids over the observation period, compared to the other plants (B, C, and D) (Fig. 3 and Table 3). Over the observation period, in both cases, the proportion of psyllids on plant A was similar. The proportion of psyllids on the other plants increased with time, decreasing the tendency to move (Fig. 3).

Table 3 Numbers (mean $\pm$ standard error) of insects, *Diaphorina citri*, found on the four plants (A, B, C, D), in the presence of non-parasitized or parasitized nymphs on plant A, over 3 days.

Treatment	Period (h)	Mean no. of insects			
		Plant A	Plant B	Plant C	Plant D
Non-parasitized nymphs	24	14.1 ± 2.7 Aa	6.6 ± 1.1 Ab	8.0 ± 1.6 Aab	6.9 ± 1.0 Ab
	48	12.1 ± 2.2 Aa	6.0 ± 1.2 Ab	7.7 ± 1.4 Aab	6.6 ± 0.7 Ab
	72	13.5 ± 2.5 Aa	7.8 ± 1.7 Aa	8.6 ± 1.8 Aa	8.0 ± 1.3 Aa
Parasitized nymphs	24	12.2 ± 1.5 Aa	7.3 ± 1.5 Ab	3.5 ± 0.7 Ac	4.0 ± 0.5 Ac
	48	11.7 ± 1.5 Aa	7.0 ± 1.4 Ab	3.3 ± 0.7 Ac	4.4 ± 0.5 Ac
	72	13.1 ± 1.9 Aa	7.7 ± 1.6 Aab	4.3 ± 1.1 Ab	6.3 ± 1.2 Ab

Means followed by different letters indicate a significant difference in the number of insects present on each plant, within each treatment at significance level $\alpha = 5\%$ (likelihood ratio test).

Regarding the distribution of eggs on the four plants, there was no effect of plant and parasitism interaction ($F_{3,72} = 0.3017$, $p = 0.8241$). The plant effect was also not significant ($F_{3,75} = 0.0349$, $p = 0.9912$). However, the presence of non-parasitized or parasitized nymphs had a significant but marginal effect ($F_{1,78} = 3.74$, $p = 0.0571$) on the distribution of eggs on the four plants. In both treatments, there was no significant difference between the number of eggs deposited among plants (Table 4). Eggs were present under parasitized nymphs, an unexpected occurrence (Fig. 4).

Table 4 Numbers (mean ± SD) of eggs of *Diaphorina citri* found on the four plants (A, B, C, D), in the presence of non-parasitized or parasitized nymphs on plant A, 72 h after the release of adults.

Treatment	Mean	Mean no. of eggs			
		Plant A	Plant B	Plant C	Plant D
Non-parasitized nymphs	49.9 (± 7.7) a	40.5	41.9	37.6	40.0
Parasitized nymphs	30.0 (± 6.5) b	(± 11.2) a	(± 12.4) a	(± 10.7) a	(± 6.9) a
Total		810	839	747	800

Means followed by different letters indicate a significant difference in the mean number of eggs in each treatment and the number of eggs on each plant, within each treatment at the significance level $\alpha = 5\%$ (likelihood ratio test).

Discussion

In this study, the presence of *D. citri* nymphs affected the movement behavior of adults. In the absence of nymphs, adult females tended to remain at the release site (plant A). When nymphs (parasitic or non-parasitized) were present at the release site (plant A), the adult females tended to disperse to other plants, reducing aggregation. The presence of nymphs also altered the females' oviposition behavior. In the absence of nymphs, oviposition rates were higher on the plant with the highest aggregation of adults (plant A). However, when non-parasitized or parasitized nymphs were present on plant A, the number of eggs deposited on plant A was reduced.

Psyllid females showed aggregate behavior at the release site (plant A) in situations where nymphs were absent or present in low/medium densities. The tendency to remain at the place of release or arrival in an orchard was also observed in the field, by Arakawa and Miyamoto (2007) in their wind-tunnel experiment, and by Boina et al. (2009), who used protein markers to observe the bidirectional movement of insects between managed and unmanaged areas. These results support the idea that psyllids have low active mobility, even though they have a flight capacity of up to 2.4 km (Martini et al. 2014a).

The aggregate behavior observed may also be due to the attraction of *D. citri* to volatiles emitted by plants damaged by feeding of conspecifics present (Groot et al. 2003; Blackmer et al. 2004). In addition to the psyllid's attraction to citrus plants and orange jasmine (Patt and Sétamou 2010), its attraction to psyllid-damaged plants and its aggregation behavior were observed by Hijaz (2013) and Patt et al. (2018), who linked this result to increased release of salicylic acid (SA). Mann et al. (2012) previously made this observation in a study with citrus plants. Although the study was carried out with plants infected with HLB, they observed an increase in methyl salicylate (MeSA) emission in plants that attracted more insects. Martini et al. (2014b), in their study with curry plants, also observed aggregated behavior of females of *D. citri* on damaged plants when fewer than 10 individuals (females) were present. In the absence of damaged plants, the presence of more than 10 females had a repellent effect on other females, which were also not attracted by the presence of males of the species, although males were attracted by females.

In the absence of nymphs at the release site, the present results indicate that plants with more psyllids also had more eggs deposited. In this species, as in others such as *Cassida rubiginosa* Müller (Coleoptera: Chrysomelidae) (Fujiyama et al., 2008) and *Thaumastocoris peregrinus* Carpintero & Dellapé (Hemiptera: Thaumastocoridae) (Martínez and Bianchi 2010; Wilcken et al. 2010), the feeding and oviposition sites overlap, since *D. citri* adults and nymphs both prefer young growing shoots (Hall et al. 2013; Cifuentes-Arenas et al. 2018). This aggregation behavior of the adult and nymphal stages benefits the species, since it indicates better local abiotic conditions for development and reproduction (Sétamou et al. 2016; Cifuentes-Arenas et al. 2018). On the other hand, this aggregation also entails competition for space and food, in addition to increasing the risk of infections and the action of natural enemies (Martínez et al. 2014b), which may explain the change in the adults' behavior as the process continued, with an increase in the density of nymphs on plant A.

Knowing that the presence of conspecifics at a location indicates quality and the degree of competition, a decrease in numbers of both adult and immature insects is expected at these locations (Davis et al. 2011), as was observed here in the treatments with low and high density of nymphs. Bawin et al. (2014), Guo et al. (2014), and Kohandani et al. (2015) found similar results and determined that the presence of conspecifics above a certain threshold is required to trigger a “displacer” effect on insects. Although this “displacer” effect was observed in this study in both experimental scenarios (presence of non-parasitized nymphs and presence of parasitized nymphs; Tables 1 and 2), eggs were still deposited at these “occupied” sites, even if other plants of the same quality were available for feeding and oviposition in the same cage.

As observed in other studies on the impact of the presence of natural enemies on the reproductive rate of hosts/prey (Holmes and Bethel 1972; Price 1980; Nelson et al. 2004; Ninkovic et al. 2013; Hermann and Thaler 2014), *D. citri* reduced the number of offspring due to the presence of a control agent; however, it laid eggs in the presence of the parasitoid, although the parasitoid was still developing within the psyllid nymphs (Fauvergue and Quilici 1991; Hoy and Nguyen 2000). *Diaphorina citri* also showed this behavior in another study (Seo et al. 2018), where it perceived and was repelled by the chemical traces of a natural enemy, the predator *Hippodamia convergens* Guérin-Ménéville (Coleoptera: Coccinellidae), reducing its oviposition rate and excretions.

Detection of the presence of natural enemies has important consequences for the survival rate of an insect and can manifest at the level of predator/prey and host/parasitoid interactions as well as in intraguild competition (predator/parasitoid) interactions. For example, Milosavljević et al. (2021) found that the presence of *Linepithema humile* (Mayr) (Hymenoptera: Formicidae) strongly reduced the control of *D. citri* by *T. radiata*, since the ants defended the psyllids from the parasitoids in exchange for food. Thus, the development of behaviors by insects that minimize their risks in the presence of conspecifics and natural enemies is necessary from a biological point of view. For example, *T. radiata* marks the nymphs of *D. citri* with a pheromone after parasitizing them, to prevent superparasitism (Mann et al, 2010) and can detect volatiles to avoid predating of their nymphs (Milosavljević et al. 2021).

The results obtained here reinforce the recommendation that *D. citri* control tactics should be focused on the nymphal stages, since in addition to contributing more effectively to the spread of HLB (Inoue et al. 2009, Diniz et al. 2021), the presence of nymphs can stimulate adults to disperse, also contributing to the spread of the disease.

Conclusion

The results showed that *D. citri* has an aggregated behavior, with a tendency to low mobility in the absence of nymphs. However, this behavior changes when nymphs are present, in low or high densities and whether or not the nymphs are parasitized. *D. citri* distinguishes unparasitized nymphs from nymphs parasitized by *T. radiata*, and in this case, lays eggs close to the parasitized nymphs.

Declarations

Acknowledgments

We thank the Fund for Citrus Protection (FUNDECITRUS) (project Tamarixia 878) for financial support (scholarship); the São Paulo Research Foundation (FAPESP) (process 2018/02317-5) as part of the São Paulo Advanced Research Center for Biological Control (SPARCBIO) for material resources; and Dr. Janet W. Reid (JWR Associates) for English language editing. Carolina Reigada is grateful to the Coordination for the Improvement of Higher Education Personnel (CAPES) (PNDP: 2014.1.93.11.0) for financial support.

Author contributions

Juliana S. Oliveira, Carolina Reigada, Alexandre J.F. Diniz and José R.P. Parra contributed to the conception and design of the study. Material preparation and data collection were performed by Juliana S. Oliveira, data analysis was performed by Juliana S. Oliveira, Carolina Reigada, Idemauro A.R. Lara and Rafael A. Moral. The first version of the manuscript was written by Juliana S. Oliveira and Carolina Reigada and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Disclosure

The authors have no conflict of interest to declare.

Funding Declaration

This study was funding by Citrus Protection Fund (FUNDECITRUS) (Tamarixia 878 project), São Paulo Research Foundation (FAPESP) (process 2018/02317-5) and Coordination for the Improvement of Personnel from Higher Education (CAPES) (PNDP: 2014.1.93.11.0).

References

1. Agresti A (2013) Categorical Data Analysis, 3rd ed. Wiley, New York.
2. Antolinez CA, Moyneur T, Martini X, Rivera MJ (2021) High temperatures decrease the flight capacity of *Diaphorina citri* Kuwayama (Hemiptera: Liviidae). *Insects* 12: 394.
3. Arakawa K, Miyamoto K (2007) Flight ability of Asian citrus psyllid, *Diaphorina citri* Kawayama (Homoptera: Psyllidae) measured by a flight mill. *Res Bull PI Prot Jpn* 43: 23–26.
4. Bawin T, De Backer L, Dujeu D et al. (2014) Infestation level influences oviposition site selection in the tomato leafminer *Tuta absoluta* (Lepidoptera: Gelechiidae). *Insects* 5: 877–884.
5. Bhavakul K, Intavimolsri S, Vichitrananda S, Kratureuk C, Prommintara M (1982) The current citrus disease situation in Thailand with emphasis on citrus greening. *Proc Intl Soc Citricult* 4: 464–466.

6. Blackmer JL, Rodriguez-Saona C, Byers JA, Shope KL, Smith JP (2004) Behavioral response of *Lugus Hesperus* to conspecifics and headspace volatiles of alfalfa in a Y-tube olfactometer. *J Chem Ecol* 30:1547–1564.
7. Boina DR, Meyer WL, Onagbola EO, Stelinski LL (2009) Quantifying dispersal of *Diaphorina citri* (Hemiptera: Psyllidae) by immunomarking and potential impact of unmanaged groves on commercial citrus management. *Environ Entomol* 38:1250–1258.
8. Bové JM (1986) Greening in the Arabian Peninsula: towards new techniques for its detection and control. *FAO Plant Protect Bull* 34: 7–14.
9. Bové JM (2006) Huanglongbing: a destructive, newly emerging, century-old disease of citrus. *J Plant Pathol* 88:7–37.
10. Bullock JM, Kenward RE, Hails RS (2002) *Dispersal Ecology: the 42nd Symposium of the British Ecological Society*. Blackwell, Oxford, pp 458.
11. Capoor SP, Rao DG, Viswanath SM (1974) Greening disease of citrus in the Deccan Trap Country and its relationship with the vector, *Diaphorina citri* Kuwayama pp 43–49. In LG Weathers, M Cohen [eds.], *Proceedings of the 6th Conference of the International Organization of Citrus Virologist*. University of California, Riverside, California, USA.
12. CDA – Coordenadoria de Defesa Agropecuária (2019) Dados da Citricultura Paulista. In: Dados da Citricultura Paulista. <http://www.defesaagropecuaria.sp.gov.br/www/gdsv/index?action=dadosCitriculturaPaulista>. Accessed 23 February 2022.
13. Cifuentes-Arenas JC, de Goes A, de Miranda MP, Beattie GAC, Lopes SA (2018) Citrus flush shoot ontogeny modulates biotic potential of *Diaphorina citri*. *PLoS One* 13: e0190563.
14. Dala-Paula BM, Plotto A, Bai J et al. (2019) Effect of Huanglongbing or Greening disease on orange juice quality, a review. *Front Plant Sci* 9:1976.
15. Davis JM, Nufio CR, Papaj DR (2011) Resource quality or competition: why increase resource acceptance in the presence of conspecifics? *Beha Ecol* 22:730–737.
16. Dingle H (1972) Migration strategies of insects. *Science* 175:1327–1335.
17. Diniz AJF (2013) Otimização de criação de *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) e *Tamarixia radiata* (Waterson, 1922) (Hymenoptera:Eulophidae), visando a produção em larga escala do parasitoide e avaliação de seu estabelecimento em campo. Thesis., University of São Paulo (USP).
18. Diniz AJF, Garcia AG, Alves GR, Reigada C, Vieira JM, Parra JRP (2020) The enemy is outside: Releasing the parasitoid *Tamarixia radiata* (Hymenoptera: Eulophidae) in external sources of HLB inocula to control the Asian Citrus Psyllid *Diaphorina citri* (Hemiptera: Liviidae). *Neotrop Entomol* 49:250–257.
19. Diniz AJF, Alves GR, Vieira JM, Parra JRP (2021) Manejo do psílideo dos citros com uso do parasitoide *Tamarixia radiata*: uma nova abordagem em controle biológico. In: Parra JRP, Pinto AS, Nava DE, Oliveira CR, Diniz AJF (eds) *Controle Biológico com Parasitoides e Predadores na Agricultura Brasileira*, Fealq, Piracicaba, pp 127–158.

20. FAO (2002) Saudi citrus industry thrives in desert. Food and Agriculture Organization, United Nations, Rome.
21. Fauvergue X, Quilici S (1991) Étude de certains paramètres de la biologie de *Tamarixia radiata* (Waterston, 1992) (Hymenoptera: Eulophidae), ectoparasitoïde primaire de *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae), vecteur asiatique du greening des agrumes. *Fruits* 46:179–185.
22. Fraser LR (1978) Recognition and control of citrus virus diseases in Australia. *Proc Intl Soc Citricult* 3:178-181.
23. French JV, Kahlke CJ, da Graça JV (2001) First Record of the Asian Citrus Psylla, *Diaphorina citri* Kuwayama (Homoptera: Psyllidae), in Texas. *Subtrop Plant Sci* 53:14–15.
24. Fritz RS (1982) Selection for host modification by insect parasitoids. *Evolution* 36:283-288.
25. Fujiyama N, Torii C, Akabane M, Katakura H (2008) Oviposition site selection by herbivorous beetles: a comparison of two thistle feeders, *Cassida rubiginosa* and *Henosepilachna niponica*. *Entomol Exp Appl* 128:41–48.
26. Garcia AG, Diniz AJF, Parra JRP (2019) A fuzzy-based index to identify suitable areas for host-parasitoid interactions: Case study of the Asian citrus psyllid *Diaphorina citri* and its natural enemy *Tamarixia radiata*. *Bio Control* 135:135–140.
27. Garcia AG, Jamielniak JA, Diniz AJF, Parra JRP (2021) The importance of Integrated Pest Management to flatten the Huanglongbing (HLB) curve and limit its vector, the Asian citrus psyllid. *Entomol Gen.* <https://doi.org/10.1127/entomologia/2021/1247>
28. Garnier M, Bové JM (1996) Distribution of the Huanglongbing (greening) *Liberobacter* species in fifteen African and Asian countries. In: *Proceedings of 13th conference IOCV, Riverside*, pp 388–391.
29. Grafton-Cardwell EE, Stelinski LL, Stansly PA (2013) Biology and management of Asian Citrus Psyllid, vector of the Huanglongbing pathogens. *Annu Rev Entomol* 58:413–432.
30. Groot AT, Hejiboer A, Visser JH, Dicke M (2003) Oviposition preference of *Lygocoris pabulinus* (Het., Miridae) in relation to plants and conspecifics. *J Appl Entomol* 127:65–71.
31. Guo H, Meng L, Wang Y, Zheng L, Li B (2014) Oviposition behavior of the predatory midge *Aphidoletes aphidimyza* in response to aphid patch quality. *J Insect Behav* 2:816–825.
32. Halbert SE (1998) Entomology Section. *Tri-Ology* 37: 5–9.
33. Hall DG, Richardson ML, Ammar ED, Halbert SE (2013) Asian citrus psyllid, *Diaphorina citri*, vector of citrus Huanglongbing disease. *Entomol Exp Appl* 146:207–223.
34. Hermann SL, Thaler JS (2014) Prey perception of predation risk: volatile chemical cues mediate non-consumptive effects of a predator on a herbivorous insect. *Oecologia* 176:669–676.
35. Hijaz F, El-Shesheny I, Killiny N (2013) Herbivory by the insect *Diaphorina citri* induces greater change in citrus plant volatile profile than does infection by the bacterium, *Candidatus Liberibacter asiaticus*. *Plant Signal Behav* 8:e25677.
36. Holmes JC, Bethel WM (1972) Modification of intermediate host behavior by parasites. *Zool J Linn Soc-Lond* 51:123–149.

37. Hoy MA, Nguyen R (2000) Classical biological control of Asian citrus psylla. *Citrus Indust* 81:48–50.
38. Hung TH, Hung SC, Chen CN, Hsu MH, Su HJ (2004) Detection by PCR of *Candidatus Liberibacter asiaticus*, the bacterium causing citrus huanglongbing in vector psyllids: application to the study of vector-pathogen relationships. *Plant Pathol* 53:96–102.
39. Inoue H, Ohnishi J, Ito T et al. (2009) Enhanced proliferation and efficient transmission of *Candidatus Liberibacter asiaticus* by adult *Diaphorina citri* after acquisition feeding in the nymphal stage. *Ann Appl Biol* 155:29–36.
40. Kobori Y, Nakata T, Ohto Y, Takasu F (2011) Dispersal of adult Asian citrus psyllid, *Diaphorina citri* Kuwayama (Homoptera: Psyllidae), the vector of citrus greening disease, in artificial release experiments. *Appl Entomol Zool* 46:27–30.
41. Kohandani F, Le Goff GJ, Hance T (2015) Does insect mother know under what conditions it will make their offspring live? *Insect Sci* 24:141–149.
42. Lara IAR, Hinde J, Taconeli CA (2017) An alternative method for evaluating stationarity in transition models. *J Stat Comput Simul* 87:1–19.
43. Lara IAR, Moral RA, Taconeli CA, Reigada C, Hinde J (2020) A generalized transition model for grouped longitudinal categorical data. *Biom J* 62:1837–1858.
44. Lewis-Rosenblum H, Martini X, Tiwari S, Stelinski LL (2015) Seasonal movement patterns and long-range dispersal of Asian citrus psyllid in Florida citrus. *J Econ Entomol* 108:3–10.
45. Mann RS, Qureshi JA, Stansly PA, Stelinski LL (2010) Behavioral response of *Tamarixia radiata* (Waterson) (Hymenoptera: Eulophidae) to volatiles emanating from *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) and citrus. *J. Insect Behav* 23:447–458.
46. Mann RS, Ali JG, Hermann SL et al. (2012) Induced Release of a Plant-Defense Volatile 'Deceptively' Attracts Insect Vectors to Plants Infected with a Bacterial Pathogen. *PLoS Pathog* 8:e1002610.
47. Martinez AL, Wallace JM (1968) *Citrus* leaf-mottle-yellows disease in the Philippines and transmission of the causal virus by a psyllid, *Diaphorina citri*. *Plant Dis Rep* 51:692-695.
48. Martínez G, Bianchi M (2010) Primer registro para Uruguay de la chinche del eucalipto, *Thaumastocoris peregrinus* Carpintero y Dellappé, 2006 (Heteroptera: Thaumastocoridae. *Agrociencia* 14:15–18.
49. Martini X, Hoyte A, Stelinski LL (2014a) Abdominal color of the Asian citrus psyllid (Hemiptera: Liviidae) is associated with flight capabilities. *Ann Entomol Soc Am* 107:842–847.
50. Martini X, Kuhns EH, Hoyte A, Stelinski LL (2014b) Plant volatiles and density-dependent conspecific female odors are used by Asian citrus psyllid to evaluate host suitability on a spatial scale. *Arth-Plant Int* 8:453–460.
51. Mazzi D, Dorn S (2012) Movement of insect pests in agricultural landscapes. *Ann Appl Biol* 160:97–113.
52. McCullagh P, Nelder JA (1989) "Binary data." *Generalized linear models*. Springer US 98: 148.

53. Milosavljević I, Morgan DJW, Massie RE, Hoddle MS (2021) Density dependent mortality, climate, and Argentine ants affect population dynamics of an invasive citrus pest, *Diaphorina citri*, and its specialist parasitoid, *Tamarixia radiata*, in Southern California, USA. *Biolog Control* 159:104627.
54. Minter M, Pearson A, Lim KS, Wilson K, Chapman JW, Jones CM (2018) The tethered flight technique as a tool for studying life-history strategies associated with migration in insects. *Ecol Entomol* 43:397–411.
55. Moral R, Hinde J, Demétrio C (2017) Half-normal and overdispersed models in R: The hnp package. *J Stat Softw* 81:1–23.
56. Naranjo SE (2019) Assessing insect flight behavior in the laboratory: A primer on flight mill methodology and what can be learned. *Ann Entomol Soc Am* 112:182–199.
57. Nava DE, Torres MLG, Rodrigues MDL, Bento JMS, Parra JRP (2007) Biology of *Diaphorina citri* (Hem., Psyllidae) on different hosts and at different temperatures. *J Appl Entomol* 131:709–715.
58. Nelson EH, Matthews CE, Rosenheim JA (2004) Predators reduce prey population growth by inducing changes in prey behavior. *Ecology* 85:1853–1858.
59. Ninkovic V, Feng Y, Olsson U, Pettersson J (2013) Ladybird footprints induce aphid avoidance behavior. *Biolog Control* 65:63–71. <https://doi.org/10.1016/j.biocontrol.2012.07.003>
60. Nufio CR, Papaj DR (2001) Host marking behavior in phytophagous insects and parasitoids. *Entomol Exp Appl* 99:273–293
61. Oke AO, Oladigbolu AA, Kunta M, Alabi OJ, Sétamou M (2020) First Report of the Occurrence of Asian Citrus Psyllid *Diaphorina citri* (Hemiptera: Liviidae), an Invasive Species in Nigeria, West Africa. *Sci Rep* 10:9418.
62. Parra JRP, Alves GR, Diniz AJF, Vieira JM (2016) *Tamarixia radiata* (Hymenoptera: Eulophidae) × *Diaphorina citri* (Hemiptera: Liviidae): Mass rearing and potential use of the parasitoid in Brazil. *J Integr Pest Manag* 7:1–11.
63. Patt JM, Sétamou M (2010) Responses of the Asian Citrus Psyllid to volatiles emitted by the flushing shoots of its Rutaceous host plants. *Environ Entomol* 39:618–624.
64. Patt JM, Robbins PS, Niedz R, McCollum G, Alessandro R. (2018) Exogenous application of the plant signalers methyl jasmonate and salicylic acid induces changes in volatile emissions from citrus foliage and influences the aggregation behavior of Asian citrus psyllid (*Diaphorina citri*), vector of Huanglongbing. *PLoS One* 13:e0193724.
65. Pelz-Stelinski KS, Brlansky RH, Ebert TA, Rogers ME (2010) Transmission parameters for *Candidatus Liberibacter asiaticus* by Asian citrus psyllid (Hemiptera: Psyllidae). *J Econ Entomol* 103:1531–1541.
66. Price PW (1980) *Evolutionary Biology of Parasites*. Princeton University Press, Princeton.
67. R Core Team (2019) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
68. Seo M, Rivera MJ, Stelinski LL (2018) Trial chemicals of the convergens Ladybird Beetle, *Hippodamia convergens*, reduce feeding and oviposition by *Diaphorina citri* (Hemiptera: Pysillidae)

- on citrus plants. *J Insect Behav* 31:298–308.
69. Sétamou M, Simpson CR, Alabi OJ, Nelson SD, Telagamsetty S, Jifon JL (2016) Quality matters: influences of citrus flush physicochemical characteristics on population dynamics of the Asian citrus psyllid (Hemiptera: Liviidae). *PLoS One* 11:e0168997.
70. Sétamou M, Flores D, French JV, Hall DG (2020) Dispersion patterns and sampling plants for *Diaphorina citri* (Hemiptera: Psyllidae) in citrus. *J Econ Entomol* 101:1478–1487.
71. Sharma DR (2008) Population dynamics in relation to abiotic factors and management of citrus psylla in Punjab. *Indian J Hortic* 65:417–422.
72. Shimwela MM, Narouei-Khandan HA, Halbert SE et al. (2016) First Occurrence of *Diaphorina citri* in East Africa, Characterization of the Ca. Liberibacter Species Causing Huanglongbing (HLB) in Tanzania, and Potential Further Spread of *D. Citri* and HLB in Africa and Europe. *Eur J Plant Pathol* 146:349–368.
73. Shivankar VJ, Rao CN, Singh S (2000) Studies on citrus psylla, *Diaphorina citri* Kuwayama: a review. *Agric Rev* 21:199–204.
74. Singerman A, Useche MP (2016) Impact of Citrus Greening on Citrus Operations in Florida. University of Florida/Institute of Food and Agricultural Sciences Extension. FE983.
75. Stinner RE, Barfield CS, Stimac JL, Dohse L (1983) Dispersal and movement of insect pests. *Ann Rev Entomol* 28:319–335.
76. Tomaseto AF, Krugner R, Lopes JRS (2016) Effect of plant barriers and citrus leaf age on dispersal of *Diaphorina citri* (Hemiptera: Liviidae). *J Appl Entomol* 140:91–102.
77. Tomaseto AF, Miranda MP, Moral RA, Lara IAR, Fereres A, Lopes JRS (2017) Environmental conditions for *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) take-off. *J Appl Entomol* 142:104–113.
78. Tomaseto AF, Marques RN, Fereres A et al. (2019) Orange jasmine as a trap control to control *Diaphorina citri*. *Sci Rep* 9:2070.
79. Wilcken CF, Soliman EP, de Sá LAN et al. (2010) Bronze bug *Thaumastocoris peregrinus* Carpintero and Dellapé (Hemiptera: Thaumastocoridae) on Eucalyptus in Brazil and its distribution. *J Plant Prot Res* 50:201–205.
80. Xu CF, Xia YH, Li KB & Ke C (1988) Further study of the transmission of citrus huanglongbing by a psyllid, *Diaphorina citri* Kuwayama. Proceedings of the 10th Conference of the International Organization of Citrus Virologists, 17–21 November 1986, Valencia, Spain (ed. by LW Timmer, SM Garnsey & L Navarro), pp 243– 248. International Organization of Citrus Virologists, University of California, Riverside, CA, USA.
81. Yamamoto PT, Paiva PEB, Gravena S (2001) Population dynamics of *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) in citrus orchards in the North of Sao Paulo State, Brazil. *Neotrop Entomol* 30:165–170.

Figures




Figure 1

Experimental cage to evaluate the movement and distribution pattern of *Diaphorina citri* eggs on *Murraya paniculata* plants.




Figure 2

Mean percentages of adult females of *Diaphorina citri* on *Murraya paniculata* plants in the absence and presence of nymphs in low and high densities over 24, 48, and 72 h.




Figure 3

Mean percentages of adult females of *Diaphorina citri* on *Murraya paniculata* plants in the presence of non-parasitized or parasitized nymphs over 24, 48, and 72 h.




Figure 4

A and **B** *Diaphorina citri* eggs under psyllid nymphs parasitized by *Tamarixia radiata*. Parasitism is evidenced by the presence of meconium and wax, on the back and around the body of the nymph, respectively.