

Antimalarial activity of the alkaloid, heptaphylline, and the furanocoumarin, imperatorin, from *Clausena anisata* against human *Plasmodium falciparum* malaria: Ex vivo trophozoitocidal, schizonticidal and gametocytocidal approach

Emmanuel Kofi Kumatia (✉ kofi2rhyme@yahoo.com)

Centre for Plant Medicine Research, Mampong-Akwapim

Felix Zoiku

University of Ghana

Alex Asase

Centre for Plant Medicine Research, Mampong-Akwapim

Nguyen Huu Tung

Phenikaa University

Research Article

Keywords: Malaria, Alkaloid, Coumarin, Plasmodium, Trophozoite, Schizont, Gametocyte, Cytotoxicity, Selectivity index

Posted Date: August 2nd, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2843793/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Malaria Journal on September 9th, 2023. See the published version at <https://doi.org/10.1186/s12936-023-04678-0>.

Abstract

Background: The erythrocytic stage life cycle of the malaria parasite, *Plasmodium falciparum*, consists of trophozoite, schizont and gametocyte stages in humans. Various antimalarial agents target different stages of the parasite to produce treatment outcomes. This study reports on the stage specific antimalarial activity of heptaphylline and imperatorin against human *P. falciparum* in addition to their cytotoxicity and selectivity indices (SI).

Methods: The compounds were isolated from *Clausena anisata* using column chromatography and their structures elucidated using NMR spectroscopy. The antimalarial activity was determined by measuring the trophozoitocidal, schizonticidal and gametocytocidal activities of the compounds using the SYBR green assay. Cytotoxicity was evaluated using the tetrazolium-based colorimetric assay.

Results: Heptaphylline and imperatorin produced trophozoitocidal, schizonticidal and gametocytocidal activities with IC₅₀s of 1.57 (0.2317) - 26.92 (0.3144) μ M with those of artesunate (the standard drug) being 0.00024 (0.0036) - 0.0070 (0.0013) μ M. In the cytotoxicity assay, the compounds produced CC₅₀s greater than 350 μ M and SI of 13.76 - 235.90. Also, the trophozoitocidal and schizonticidal activities of the compounds were more pronounced than their gametocytocidal activity. Imperatorin was 42.04% more trophozoitocidal than heptaphylline. But, heptaphylline has more schizonticidal and gametocytocidal properties than imperatorin.

Conclusion: Heptaphylline and imperatorin are therefore, promising antimalarial agents since, they possessed potent anti-malaria activity with weak cytotoxicity on RBCs. However, imperatorin is a better antimalaria prophylactic agent whereas heptaphylline is a better malaria treatment agent.

Background

Malaria is a major public health challenge which puts severe stress on health care systems, individuals and family incomes, especially in developing countries, as a result of hospitalizations and deaths [1, 2]. The *Plasmodium* parasite which causes malaria lives in the salivary glands of an infected *Anopheles* mosquito where it poses no problem to the vector. The parasite infects humans through mosquito bite and cause sickness. Over the years, efforts were made to treat, minimize and/or eradicate malaria infections by employing chemotherapeutic agents (E.g., artesunate or chloroquine) for treatment, prophylactic drugs (E.g., sulphadoxine pyrimidine) to prevent *P. falciparum* infections from developing into the disease, prevention of parasite transmission to humans. E.g., sleeping under insecticide treated nets to prevent mosquito bites, the use of insecticides to eradicate mosquitoes and clearing of bushes and drainages of stagnant water in and around human habitations to prevent the breeding of mosquitoes [3 – 5]. These programs have yielded great outcomes by reducing global malaria deaths from 1,817,000 in 2004 to 1,238,000 in 2010 and to 619,000 in 2022 [6, 7].

Despite, these successes, malaria infection is still on the rise as reported in the year 2021 [7]. Furthermore, the discovery of insecticide resistance mosquitoes and artesunate resistance *Plasmodium*

parasite in recent years, put the global efforts of malaria eradication at risk [8 - 10]. These challenges call for development of new drugs to fight malaria.

Medicinal plants are the greatest sources of diverse chemical entities which provides treasure troves of resources for novel drug discovery. *Clausena anisata* (Wild) Hook f. ex. Benth is a plant used in traditional medicine systems to treat a broad range of human ailments including infectious diseases such as schistosomiasis, leprosy, syphilis, gonorrhea, and malaria [11, 12]. Phytochemical investigations of the plant led to isolation and characterization of a host of compounds which belong to various classes of phytochemicals such as carbazole alkaloids, simple coumarins, furanocoumarins and phytosterols [13 - 15]. Although, some biological and pharmacological activities were reported for some of the individual compounds, their anti-malaria activity against the different stages of the human *P. falciparum* is not known. This study, therefore, seek to isolate, characterize and evaluate the anti-malarial activity of some of the constituents of *C. anisata* root bark against the trophozoite, schizont and gametocyte stages of human *P. falciparum* malaria and to also determine their cytotoxicity and selectivity indices.

Materials and methods

Chemicals and reagents

Petroleum ether (40 – 60 °C) and ethyl acetate were procured from Park Scientific Limited (Northampton, U.K.). Silica gel for column chromatography (40-60 µm particle size and 230 x 400 mesh size) and TLC plate (aluminum sheets coated silica gel 60F₂₅₄) were also obtained from Sorbent Technologies (Atlanta, GA, USA) and Merck Chemicals (Damstadt, Germany) respectively. Ethanol (99%) was acquired from Midland Ghana Limited, Tema.

Human *P. falciparum* malaria parasite

Human *P. falciparum* parasite was obtained from a clinically ill patient diagnose of *P. falciparum* malaria at Damfa Health Center, Accra, Ghana. The blood (1.0 mL) was aseptically collected into a sterile ACD tube with vacutainer needle and immediately transported on ice to the laboratory, after ethical clearance and consent of the patient were obtained.

Processing of plant material and isolation of compounds

C. anisata was identified and harvested in March, 2020, from Ayikumah, (in the Shai Osudoku District in the Greater Accra Region, Ghana). Identity of the plant was confirmed by comparism with an old voucher sample (CPMR 5101) in the herbarium of the Centre for Plant Medicine Research (CPMR), Mampong-Akuapem, Ghana. The root was washed with portable water, debarked and the bark sun-dried for six days before being pulverized. The extraction and chromatography were performed by modification of a previous method [13]. Briefly, about 500 g of the powder was macerated twice with 70% ethanol (4 L x 2) for 4 days each. The extracts were combined, filtered and the ethanol distilled off in vacuum at 45 °C. The aqueous extract was partitioned with petroleum ether (PE), 0.5 L x 4, and ethyl acetate (EtOAc), 0.5 L

x 4, to obtain petroleum ether, ethyl acetate and aqueous fractions. The dried ethyl acetate fraction (20.50g) was fractionated over 450 g of silica gel using PE-EtOAc-EtOH solvent systems. The process was started with PE-EtOAc system with addition of EtOAc at 10% incremental rate until 100% EtOAc was attained. EtOAc - EtOH system was then introduced at 10% incremental rate of EtOH until 100% EtOH was also attained. A total of 195 eluents were collected in an aliquot of about 200 mL. The eluents were then grouped into 8 major fractions (F1-F8) by comparison of their TLC chromatogram. F3 and F5 produced **CA1** and **CA2** respectively. **CA1** and **CA2** were then purified after repeated recrystallization in PE-EtOAc 5:1.

NMR spectroscopy analysis

NMR spectra (1D and 2D) of the compounds were obtained using NMR machine (Bruker FT-NMR Spectrometer Avance TM 500 MHz, Germany) equipped with Bruker\TopSpin3.6.3, JJ 10 NMR data analysis software and the chemical shifts resonance (δ) measured in ppm [13].

Evaluation of antimalarial activity of the compounds using SYBR green assay

Preparation of the assay solutions of the compounds

Approximately, 3 mg of each compound was added to 3 mL of 0.5% DMSO and vortexed to yield a stock solution of 1000 $\mu\text{g/mL}$. The solution was filtered with 0.2 μm pore size filter paper and diluted 10-fold with incomplete medium (ICM) to obtain a working solution of 100 $\mu\text{g/mL}$ which was then taken through a 9-fold serial dilution with ICM to obtain 100, 50, 25, 12.5, 6.25, 3.13, 1.56, 0.78 and 0.39 $\mu\text{g/mL}$ concentrations. The process was repeated for each compound to obtain the serially diluted solutions for the assay against the trophozoite, schizont and gametocyte stages respectively.

Cultivation of the human *P. falciparum* parasite

Efficacy of the compounds against the 3 stages of *P. falciparum* life cycle in humans was assessed ex vivo using a modified method of Baker et al. [16]. Continuous asexual culture of *P. falciparum* was maintained in an atmosphere of 90% N_2 , 5% CO_2 , and 5% O_2 at 37°C in complete medium (CM) consisting of (10.44 g/L, RPMI 1640, 5.94 g/L, HEPES, 5 g/L, AlbuMAX II, 50 mg/L hypoxanthine, 2.1 g/L sodium bicarbonate). The procedure was repeated for the schizont and the gametocyte stages.

Cultivation of trophozoites of human *P. falciparum* parasite

Parasite from the culture established above was then cultivated in O^+ RBCs and maintained in the incubator with daily media changes until a parasitemia of more than 5% trophozoites were obtained. The culture was then treated with 5% sorbitol to obtain a synchronized trophozoite stage. The parasite growth was monitored for 5 days by estimating percentage (%) parasitemia using Giemsa-stained slides and light microscope (x100 magnification) until parasitemia of more than 5% was recorded. Parasite mix of 2% hematocrit with 1% parasitemia were then prepared using uninfected blood to make a total of 10 mL in a CM for the plating [16].

Cultivation of schizonts of *P. falciparum* parasite

Firstly, the trophozoites were cultivated as described above and allowed to grow further to the schizonts at parasitemia of > 8%. The parasite was then synchronized with 60% Percoll (30 mL Percoll + 17 mL RPMI + 3 mL 10XPBS), transferred into a 15 mL falcon tube and then spined at 600 g for 5 minutes. The supernatant was discarded and the pellet washed twice in incomplete RPMI. Final pellet was suspended in CM to create 20% hematocrit in a 15 mL falcon tube. Approximately 7 mL of 60% Percoll was measured into a fresh falcon tube and the parasite suspension (2.5 mL) was slowly layered onto the Percoll. The tube was then spined at 700 g for 10 minutes to obtain the schizonts as a distinct reddish-brown band which was transferred into a clean falcon tube and adjusted to 2% hematocrit by addition of 20 mL uninfected RBCs in CM before plating with the various compounds. Percoll density gradient centrifugation described in this experiment was used to enhance the purification of the schizonts [17].

Cultivation of gametocytes of *P. falciparum* parasite

Gametocyte culture was initiated as described by Foley et al. [18], with some modifications at 4% hematocrit (200 µL of culture was measured into a new T-25 flask containing 5 mL CM). The culture was later transferred into a sterile T-75 culture flask after the parasitemia growth was above 8%. The parasite was allowed to grow further until 9-10% parasitemia was achieved by daily microscopy monitoring and media changes. The culture was then synchronized with 5% sorbitol to obtain only rings which were treated with freshly prepared CM and 50 mM N-acetyl glucosamine. This was noted as day 0. The flask was flushed with gentle flow of gas and then capped tightly and returned into the incubator. Spent media was removed from day 1 - 4 without taking the blood cells. 50 mM N-acetyl glucosamine in CM was added to the flask and then gassed before returning into the incubator. On day 5, the spent media was taken out and only CM was added to the flask and then gassed before returning into the incubator. The culture flask was maintained by repeated gassing procedure up to day 14. Percentage gametocyte was estimated by counting number of gametocyte or 1000-1500 RBCs. Parasite suspension was made and later plated with the compounds.

Compound plating

Approximately, 100 µL of each nine dilutions (100, 50, 25, 12.5, 6.25, 3.13, 1.56, 0.78 and 0.39 µg/mL) were plated in duplicates in a 96 well coastal plate. Artesunate (15 ng/mL) was serially diluted and plated alongside as the standard control. About 100 µL of the parasite suspension with 2% hematocrit and 1% parasitemia were added to each treated well starting from the 2nd to the 10th well. Furthermore, another 100 µL of the parasite suspension was added to each of the 11th well as a negative control. The procedure was repeated for the other compound. The plates were then, arranged in a modular chamber and gassed for 5 min with gas mixture of 90% N₂, 5% CO₂, and 5% O₂ at 37°C and then incubated at 37 °C for 72 hours.

SYBR green assay

The trophozoitocidal, schizonticidal, gametocytocidal assays of the compounds were performed according to a previous method [19], with some modifications. After the 72 hours, the plates were harvested and 100 μ L lysing buffer containing SYBR green was added to each well with thorough and gentle mixing prior to incubation in the dark for 30 minutes. The excitation and emission wavelengths of the samples were then measured at 470 and 520 nm respectively using fluorometer (FLUOstar OPTIMA with software version 2.20). The concentration that inhibits 50% of the growth of *P. falciparum* parasite (IC₅₀) was estimated from dose-response curves by non-linear regression analysis using Graph pad Prism version 7.0 Software (Graph Pad software, San Diego, CA, USA). The IC₅₀s were later converted to molar concentration.

Cytotoxicity assay

The toxicity of the compounds to RBCs was evaluated using the tetrazolium-based colorimetric assay as described [20], with some modifications. Briefly, 100 μ L of each compound with concentrations ranging from 6.25 - 100 μ g/mL was measured into separate wells of a 96-well microtiter plate in duplicates. Uninfected red blood cells (100 μ L) of 2% hematocrit were added to each well. The plates were then incubated at 37°C for 72 hours in a humidified incubator at 5% O₂ and CO₂. MTT solution (20 μ L of 7.5 mg/mL in phosphate buffered saline) was added to each well and the plate incubated again for 2 hours. Finally, Triton X-100 (150 μ L) in acidified isopropanol was added to each well to dissolve any formazan formed. The plates were then kept in the dark for 24 hours after which the optical densities were measured at 570 nm using a plate reader. The cytotoxicity concentration (CC₅₀) which is the concentration of the compound required to kill 50% of the cells was then determined by plotting concentration of the compounds on x-axis and percentage of cell viability on y-axis with dose-response curves.

$$\text{Cell viability (\%)} = \frac{A_0 - A_1}{A_0} \times 100$$

Where A₀ or A₁ is the mean absorbance of wells with untreated (vehicle) or test wells.

Selectivity indices (SI)

The SI of the compounds was calculated as:

$$SI = \frac{CC_{50}}{IC_{50}}$$

Statistical analysis

Each concentration of a respective compound was tested in duplicate and the IC₅₀ results presented as $\bar{X}$ (SD) i.e., mean (standard deviations).

Results

Characterization of the compounds

Characterization of CA1

CA1 was obtained as bright yellow crystals (6.56 g). R_f = 0.603 (PE- EtOAc, 20:1); 0.400 (PE-CF, 5:1). ^1H NMR (CDCl_3 , 500 MHz), δ 11.67 (1H, s, -NH), 9.95 (1H, s, -CHO), 8.22 (1H, s, H-4,), 8.00 (1H, d, H-5, J = 7.05 Hz), 7.29 (1H, ddd, H-6, J = 3.28, 3.16 and 7.40 Hz), 7.43 (1H, m, H-7, J = 3.35 Hz), 7.99 (1H, dd, J = 7.7, 1.0 Hz, H-8), 3.67 (2H, d, H-1', J = 6.90 Hz), 5.35 (2H, t, H-2', J = 5.93 Hz), 1.74 (3H, s, H-4'), 1.80 (3H, s, H-5'). ^{13}C NMR (CDCl_3 , 500 MHz), δ 195.44 (CHO), 157.86 (C-2), 140.16 (C-8a), 145.08 (C-9a), 115.49 (C-3), 125.90 (C-4), 119.82 (C-5), 125.36 (C-4b), 123.70 (C-6), 120.89 (C-7), 110.90 (C-8), 117.37 (C-4a), 109.08 (C-1), 25.75 (C-1'), 121.27 (C-2'), 134.21 (C-3'), 22.87 (C-4'), 18.15 (C-5'). These results correspond with the literature data for the carbazole alkaloid, heptaphylline [13]. Hence, CA1 was identified as heptaphylline.

Characterization of CA2

CA2 was obtained as white spongy crystals (10.56 g). R_f = 0.286 (PE-CF, 3:1); 0.706 (PE- EtOAc, 4:1). ^1H NMR (CDCl_3 , 500 MHz), δ 6.36 (1H, d, H-3; J = 9.6 Hz), 7.77 (1H, *d*, H-4, J = 9.4 Hz), 7.36 (1H, s, H-5), 7.69 (1H, d, H-2'', J = 8.00 Hz), 6.82 (1H, d, H-3'', J = 6.84 Hz), 5.00 (2H, d, H-1', J = 7.10 Hz), 5.61 (1H, t, H-2', J = 7.35 Hz), 1.72 (3H, s, H-4'), 1.74 (3H, s, H-5'). ^{13}C NMR (CDCl_3 , 500 MHz), δ 160.57 (C-2), 114.70 (C-3), 143.83 (C-4), 116.51 (C-4a), 113.60 (C-5), 125.88 (C-6), 148.62 (C-7), 131.68 (C-8), 144.25 (C-8a), 146.60 (C-2''), 106.73 (C-3''), 70.10 (C-1'), 119.70 (C-2'), 139.70 C-3'), 25.83 (C-4'), 18.13 (C-5'). These data are in agreement with those reported for imperatorin [13]. CA2 was therefore, identified as imperatorin.

The chemical structures of the isolated compounds are given below in Fig. 1.

Trophozoitocidal, schizonticidal and gametocytocidal activities of the compounds

The IC_{50} curves (Fig. 2) and the IC_{50} values (Table 1) below shows the inhibitory effects of imperatorin, heptaphylline and artesunate on the various stages of the life cycle of human *P. falciparum* parasite ex vivo.

Table 1 The IC_{50} (SD) of the compounds on the inhibition of the various stages of *P. falciparum*

Compound	Trophozoite	Schizont	Gametocyte
	IC_{50} / μM	IC_{50} / μM	IC_{50} / μM
Imperatorin	1.57 (0.2317)	8.97 (0.8935)	26.92 (0.3144)
Heptaphylline	2.23 (0.0801)	6.71 (1.0094)	20.87 (0.1762)
Artesunate	0.00024 (0.0036)	0.0070 (0.0013)	0.00024 (0.00004)

The compounds strongly inhibited all the three stages (the trophozoite, schizont and gametocyte) of the human *P. falciparum* malaria parasite. The minimum inhibitory concentration (IC₅₀) of imperatorin or heptaphylline against the trophozoite, schizont and gametocyte stages of the human *P. falciparum* were 1.57 (0.2317) – 26.92 (0.3144) µM and 2.23 (0.0801) - 20.87 (0.1762) µM respectively (Table 1). The furanocoumarin, imperatorin, was 42.04% more active against the trophozoites of the human *P. falciparum* malaria parasite than the carbazole alkaloid, heptaphylline. Apart from the trophozoite stage, heptaphylline produced the lowest IC₅₀ against the schizonts and the gametocytes compared to imperatorin. Hence, heptaphylline is a better schizonticidal (33.68% higher) and gametocytocidal (31.15% higher) agent than imperatorin.

Although the compounds produced very good inhibitory action on the various stages of the malaria disease, they were most active against the trophozoites followed by the schizonts. The lest activity of the compounds was observed on the gametocytes (Table 1).

3.3. Imperatorin and heptaphylline are weakly cytotoxic against human RBCs

The CC₅₀ and the SI of the compounds on the trophozoite, schizont and gametocyte stages of the parasite are reported below (Table 2).

Table 2 CC₅₀ and SI of the compounds against the various stages of human *P. falciparum*

Compound	CC ₅₀ / µM	<u>SI of the compounds on various stages of human <i>P. falciparum</i></u>		
		Trophozoite	Schizont	Gametocyte
Imperatorin	370.37	235.90	41.28	13.76
Heptaphylline	358.42	160.73	53.41	17.17
Artesunate	Nd	Nd	Nd	Nd

Nd = Not determined

Discussion

In humans, the malaria parasite undergoes development in three stages. Liver form schizonts which envelops thousands of merozoites developed from sporozoites inoculated into the human skin through a bite of an infected *Anopheles* mosquito, disintegrates and release the merozoites into the blood stream where they invade RBCs to commence the intraerythrocytic asexual life cycle phase of the parasite [21, 22]. The trophozoite stage is the first phase of this process where the merozoites invades the RBCs and occupy a small portion of its food vacuole in a form of a small round ring (ring or early trophozoite stage) and feed on the hemoglobin and protein content of the RBCs and grow bigger to occupy most of the RBCs’ cytoplasm (trophozoite or late trophozoite) stage. The second stage is called the schizont stage. At this stage, the trophozoite divides into several small daughter cells called merozoites. The mature exo-

erythrocytic schizonts erupts and release the merozoites along with RBCs and parasite remains (such as hemozoin and glycerophosphatidylinositol) into the blood stream [23]. The merozoites then invade healthy RBCs and the process continues [23]. In the gametocyte stage, the merozoites in some cells developed into male and female sexual forms called gametocytes, instead of undergoing asexual replication, which circulate in the blood where they are ingested by a vector mosquito during a blood meal, the male and the female gametocytes then meet in the mosquito and undergo various development to become sporozoites needed for human infection [23].

The trophozoite and the gametocyte stages of the parasite do not manifest the malaria ailment in the human host. The clinical display of the disease is produced by the schizonts stage of the parasite in unison with the eruption of the RBCs and the simultaneous release of parasite rubles and poisons which activates the host immune response to produce cytokines, free radicals and additional cellular constituents which are accountable for the symptoms like fever, perspirations, chills, fatigue, and other systemic injuries and deaths coupled with the disease [24 - 26]. Thus, in order to achieve the best treatment outcome and relief the symptoms of malaria the anti-malaria agent must be very active against the schizont stage of the parasite. All the tested compounds were very effective against the schizont stage of *P. falciparum*. Therefore, they are good treatment substances for malaria. The compound that will produce the best treatment outcome against clinical malaria among the tested compound is heptaphylline. Since, it was the most effective against the schizont stage of the parasite than imperatorin.

Furthermore, the gametocyte stage is responsible for perpetuating the survival/life cycle and reinfection of the malaria parasite from humans to mosquito and back to humans. The compounds also effectively inhibited the gametocyte stage of the *P. falciparum* malaria parasite. This shows that the compounds and artesunate can be used to effectively prevent the transmission of human *P. falciparum* malaria parasite to the mosquito vector, thereby disrupting the lifecycle of the parasite. The order of this effect was discussed in the results of the gametocyte stage.

Finally, the trophozoite stage of the *P. falciparum* malaria parasite is the initiation stage of malaria. Thus, if the trophozoites are destroyed, then the clinical manifestation of malaria and deaths which occur at the schizont stage, and the gametocyte/perpetuation stage of the malaria will be eradicated. Since, the compounds inhibited the trophozoite stage of *P. falciparum* parasite better than all other stages, indicates that the tested compounds can prevent malaria infection and transmission.

Artesunate is a semisynthetic analog of artemisinin (a plant-based antimalaria drug extracted from *Artemisia annua*). Artesunate is produced by chemically incorporating hydrophilic group into the structure of artemisinin hence making artesunate more efficacious drug than its parent drug, artemisinin and other artemisinin derivatives [8]. This suggest that modification of the structure of imperatorin and heptaphylline may also increase their efficacy as antimalarial agents. It was reported that imperatorin was inactive as antiplasmodial agent against *P. falciparum* D6 and W6 clones respectively [27]. Heptaphylline was also reported to demonstrate antiplasmodial activity against *P. falciparum* K1

multidrug resistance strain with IC_{50} of 3.2-6.4 $\mu\text{g/mL}$ [28]. However, this is the first report on the stage specific antimalaria activity of both imperatorin and heptaphylline against the clinical isolate of human malaria.

In the cytotoxic assay, both imperatorin and heptaphylline produced CC_{50} values which were greater than 350 μM against human RBCs with percent cell survival of greater than 50%. This indicates that the compounds were weakly toxic or safe to human RBCs even at the highest concentration of 350 μM . Additionally, the *Plasmodium* inhabits the human RBCs and produce ailment at the schizont stage. Since, heptaphylline and imperatorin are nontoxic to the RBCs indicates that the antimalaria activity of these compounds are intrinsic i.e., the compounds move across the membranes of the RBCs to kill the *Plasmodium* parasites in them without harming the RBCs themselves.

The selectivity index (SI) was determined as the ratio of the 50% cytotoxic concentration of the compounds against the RBCs (CC_{50}^{RBCs}) to the minimum inhibitory concentration (IC_{50}^{Pf}) of the compound against *P. falciparum*. De Souza et al. [29], reported that SI values > 10 indicates safety since, the concentration at which the test compound is effective against the parasite is smaller than the concentration at which the same compound is toxic to the human cell. In this study, the compounds produced SI > 13.736 - 211.864 (imperatorin) and > 17.152 - 160.771 (heptaphylline) respectively against the trophozoites, schizonts and gametocytes of *P. falciparum*. This indicates that both imperatorin and heptaphylline are nontoxic to human RBCs at all stages of malarial treatment. Imperatorin was also reported to be noncytotoxic on African monkey kidney fibroblast cell line (VERO) with an IC_{50} of 28.7 $\mu\text{g/mL}$ in the XTT assay [27].

Conclusion

This research has shown, for the first time, that the carbazole alkaloid, heptaphylline and the furanocoumarin, imperatorin possessed potent antimalarial activity as the result of their trophozoitocidal, schizonticidal and gametocytocidal actions on human *P. falciparum* parasite ex vivo. Furthermore, the results also indicated that both heptaphylline and imperatorin are weakly toxic to human RBCs with high selectivity indices against the trophozoites, schizonts and gametocytes of the human *P. falciparum* parasite. Heptaphylline and imperatorin are therefore promising antimalarial agents to be considered for development into the next generation of antimalaria drugs.

Declarations

Acknowledgments

Not applicable

Funding

This study did not receive any specific funding from any public or private institution.

Author credit statement

Emmanuel Kofi Kumatia conceived the research, isolated and purified the compounds, analyzed the NMR data and wrote the manuscript. Felix Zoiku performed the antimalaria and cytotoxicity assays and analyzed the results. Alex Asase performed identification, harvesting, processing and extraction. Nguyen Huu Tung elucidated the structures of the compounds. All authors read and revised the manuscript and approved its publication in this journal.

Availability of data and materials

The datasets produced and analyzed during this study in addition to the isolated compounds can be obtained from the corresponding author upon reasonable request.

Ethical statement

Ethical clearance with approval number NMIMR – IBR CPN082/17-18 was obtained from Noguchi Memorial Institute for Medical Research (NMIMR) Internal Review Board (IBR), Legon - Accra (Ghana).

Consent for publication

Not applicable.

Competing interest

The authors declare that they do not have any conflict of interest with regard to this study.

References


1. Tizifa TA, Kabaghe AN, McCann RS, Van den Berg H, Van Vugt M, Phiri KS. Prevention Efforts for Malaria. *Curr Trop Med Rep*. 2018; 5 (1): 41-50.
2. Chuma JM, Thiede M, Molyneux CS. Rethinking the economic costs of malaria at the household level: Evidence from applying a new analytical framework in rural Kenya. *Malar J*. 2006; 5: 76.
3. WHO. Global technical strategy for malaria 2016–2030. Geneva: World Health Organization; 2015. http://www.who.int/malaria/areas/global_technical_strategy/en/
4. WHO. Insecticide-treated mosquito net: a WHO position statement. Geneva: World Health Organization; 2007.
5. Diabate A, Tripet F. Targeting male mosquito mating behaviour for malaria control. *Parasit Vectors*. 2015; 8 (1):3 47. <https://doi.org/10.1186/s13071-015-0961-8>.
6. Murray CJL, Rosenfeld LC, Lim SS, Andrews KG, Foreman KJ, Haring D, et al. Global malaria mortality between 1980 and 2010: A systematic analysis. *Lancet*. 2012; 379 (9814): 413-31.
7. The World Health Organization (WHO). The World malaria report 2022. <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2022>. Accessed

on 06-02-2023.


8. Khana P. Antimalarial and anticancer properties of artesunate and other artemisinins: Current development. *Monatshefte für Chemie - Chemical Monthly*. 2021; 152: 387- 400.
9. Feachem RG, Phillips AA, Targett GA, Snow RW. Call to action: priorities for malaria elimination. *Lancet*. 2010; 376: 1517-21.
10. Cisse MBM, Keita C, Dicko A, Dengela D, Coleman J, Lucas B, et al. Characterizing the insecticide resistance of *Anopheles gambiae* in Mali. *Malar J*. 2015; 14: 327.
11. Hutchings A, Scott AH, Lewis G, Cunningham A. *Clausena anisata* (Wild.) Hook. F. ex Benth. Zulu medicinal plants: An inventory, Pietermaritzburg, South Africa, University of Natal Press. 1996; 153-4.
12. Uwaifo AO. The mutagenicities of seven coumarin derivatives and a furan derivative (Nimbolide) isolated from three medicinal plants. *J Toxicol Environ Health*. 1984; 13: 521-30.
13. Kumatia EK, Annan K, Dickson RA, Mensah AY, Amponsah IK, Appiah AA, et al. Antiinflammatory and analgesic effects in rodent models of ethanol extract of *Clausena anisata* roots and their chemical constituents *Nat Prod Commun*. 2017; 12 (1): 62-7.
14. Songue J, Kouman L, Dongo E, Mpondo TN, White RL. Chemical constituents from stem bark and roots of *Clausena anisata*. *Molecules*. 2012; 17: 13673-86.
15. Ito C, Katsuno S, Itoigawa M, Ruangrunsi N, Mukainaka T, Okuda M, et al. New carbazole alkaloids from *Clausena anisata* with antitumor promoting activity. *J Nat Prod*. 2000; 63 (1): 125-8.
16. Baker DA, Stewart LB, Large JM, Bowyer PW, Ansell KH, Jiménez-Díaz MB, et al., A potent series targeting the malarial cGMP-dependent protein kinase clears infection and blocks transmission. *Nat. Commun*. 2017; 8: 430.
17. Llorca-Batlle O, Michel-Todó L, Witmer K, Toda H, Fernandez-Becerra C, Baum J, et al. Conditional expression of PfAP2-G for controlled massive sexual conversion in *Plasmodium falciparum*. *Sci Adv*. 2020. <https://doi.org/10.1126/sciadv.aaz5057>.
18. Foley M, Deady LW, Ng K, Cowman AF, Tilley LJ. *Biol Chem*. 1994; 269: 6955 - 61.
19. Vossen MG, Pferschy S, Chiba P, Noed H. The SYBR Green I malaria drug sensitivity assay performance in low parasitemia samples. *Am J Trop Med Hyg*. 2010; 82 (3):398 - 401.
20. Denizot F, Lang R. Rapid colorimetric assay for cell growth and survival. Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *J Immunol Methods*. 1986; 89: 271–7.
21. Stiebler R, Timm BL, Oliveira PL, Hearne GR, Egan TJ, Oliveira MF. *Insect Biochem Mol*. 2010; 40: 284 - 292.
22. Khoshmanesh A, Dixon MWA, Kenny S, Tilley L, McNaughton D, Wood BR. Detection and quantification of early-stage malaria parasites in laboratory infected erythrocytes by attenuated total reflectance infrared spectroscopy and multivariate analysis. *Anal Chem*. 2014; 86: 4379 - 4386.
23. Perkins DJ, Were T, Davenport GC, Kempaiah P, Hittner JB, Ongecha JM. Severe malarial anemia: innate immunity and pathogenesis. *Int J Biol Sci*. 2011; 7: 1427-42.

24. Crutcher JM, Hoffman SL. Malaria. In: Baron S, editor. Medical Microbiology. 4th edition. Galveston (TX): University of Texas Medical Branch at Galveston; 1996. Chapter 83. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK8584/>
25. Schofield L, Hackett F. Signal transduction in host cells by a glycosylphosphatidylinositol toxin of malaria parasites. J Exp Med. 1993;177: 145 - 53.
26. Clark IA, Cowden WB. The pathophysiology of *falciparum* malaria. Pharmacol Ther 2003; 99: 221-60.
27. Rossa SA, Krishnavena K, Radwana MM, Takamatsua S, Burandt CL. Constituents of *Zanthoxylum flavum* and their antioxidant and antimalarial activities. Nat Prod Commun. 2008; 5: 791-4.
28. Yenjai C, Sripontan S, Sripajun P, Kittakoop P, Jintasirikul A, Tanticharoen M, et al. Coumarins and carbazoles with antiplasmodial activity from *Clausena harmandiana*. Planta Medica. 2000; 66: 277 - 9.
29. De Souza GE, Bueno RV, de Souza JO, Zanini CL, Cruz FC, Oliva G, et al. Antiplasmodial profile of selected compounds from malaria box: in vitro evaluation, speed of action and drug combination studies. Malar J. 2019; 18: 447.

Figures



Heptaphylline (CA1)



Imperatorin (CA2)

Figure 1

Chemical structures of the isolated compounds from *C. anisata* root bark.




Figure 2

Ex vivo inhibition of: trophozoite (A-C); schizont (D-F) and gametocyte (G-I) stages of human *P. falciparum* by imperatorin, heptaphylline and artesunate. The shape of Fig 2F reversed because the % parasitemia inhibition was calculated from the highest to the lowest concentration instead of the reverse. However, the IC₅₀ value was not affected by the shape of the graph.