

RESEARCH ARTICLE


The future of subalpine forests in the Southern Rocky Mountains: Trajectories for *Pinus aristata* genetic lineages

Sparkle L. Malone^{1,2*}, Anna W. Schoettle², Jonathan D. Coop³

1 Department of Biological Sciences, Florida International University, Miami, Florida, United States of America, **2** USDA Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, United States of America, **3** Biology, Western State Colorado University, Gunnison, Colorado, United States of America

☞ These authors contributed equally to this work.

* smalone@fiu.edu



OPEN ACCESS

Citation: Malone SL, Schoettle AW, Coop JD (2018) The future of subalpine forests in the Southern Rocky Mountains: Trajectories for *Pinus aristata* genetic lineages. PLoS ONE 13(3): e0193481. <https://doi.org/10.1371/journal.pone.0193481>

Editor: RunGuo Zang, Chinese Academy of Forestry, CHINA

Received: October 20, 2017

Accepted: February 12, 2018

Published: March 19, 2018

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: Data used in this analysis are available through the Forest Inventory and Analysis National Program (FIA) at <https://www.fia.fs.fed.us/tools-data/>. The climate and species distribution data is made available through the Forest Service at <http://charcoal.cnre.vt.edu/climate/species/>.

Funding: Funding for this work was provided by Rocky Mountain Research Station (SLM).

Competing interests: The authors have declared that no competing interests exist.

Abstract

Like many other high elevation alpine tree species, Rocky Mountain bristlecone pine (*Pinus aristata* Engelm.) may be particularly vulnerable to climate change. To evaluate its potential vulnerability to shifts in climate, we defined the suitable climate space for each of four genetic lineages of bristlecone pine and for other subalpine tree species in close proximity to bristlecone pine forests. Measuring changes in the suitable climate space for lineage groups is an important step beyond models that assume species are genetically homogeneous. The suitable climate space for bristlecone pine in the year 2090 is projected to decline by 74% and the proportional distribution of suitable climate space for genetic lineages shifts toward those associated with warmer and wetter conditions. The 2090 climate space for bristlecone pine exhibits a bimodal distribution along an elevation gradient, presumably due to the persistence of the climate space in the Southern Rocky Mountains and exclusion at mid-elevations by conditions that favor the climate space of other species. These shifts have implications for changes in fire regimes, vulnerability to pest and pathogens, and altered carbon dynamics across the southern Rockies, which may reduce the likelihood of bristlecone pine trees achieving exceptional longevity in the future. The persistence and expansion of climate space for southern bristlecone pine genetic lineage groups in 2090 suggests that these sources may be the least vulnerable in the future. While these lineages may be more likely to persist and therefore present opportunities for proactive management (e.g., assisted migration) to maintain subalpine forest ecosystem services in a warmer world, our findings also imply heightened conservation concern for vulnerable northern lineages facing range contractions.

Introduction

Tree species are increasingly threatened by anthropogenic environmental change including deforestation [1], native and introduced insects and pathogens [2], altered disturbance regimes

[3,4], and climate change [5–7]. Climate is a crucial factor controlling where tree species occur [8] and as the climate changes so then do species distributions. Studies have shown that many tree species respond to shifts in climate by migrating [9–12], which influences ecosystem structure and function with the potential to produce novel ecosystems [13].

Predicting species range shifts in response to climate change is a major challenge in conservation. A species is composed of a compilation of genetically differentiated populations with clinal variation along environmental gradients [14]. These populations may respond differently to climate change [15] and the species-wide genetic structure may change [16] as coverage by some populations contract and others expand and evolve [16,17]. Modeling the future distribution of suitable climate space for genetically divergent populations within a species can identify those that may be disproportionately represented or underrepresented in the future and therefore of current management or conservation interest.

Some of the most vulnerable ecosystems to climate change include subalpine forests [11,18] where growing space declines with elevation and species distributions are defined by distinct climatic gradients (altitudinal) and microclimate effects (i.e. topography, aspect, soil). Climate change is projected to be rapid and heightened in these habitats, where subalpine treelines are expected to move up in elevation with warming growing season temperatures [11]. To understand the potential vulnerability to climate change for long lived high-elevation species, we can evaluate changes in the suitable climate space for tree species. Suitable climate space is determined based on empirical relationships between observed species distributions and climate [19].

In the southern Rocky Mountains, Rocky Mountain bristlecone pine (*Pinus aristata*; henceforth, bristlecone pine) is a long-lived tree species (>500 years) [20] of high elevation forests (2750 to 3670 m elevation) where it occupies sites beyond the physiological limits of most other tree species. Co- and sub-dominant species of bristlecone pine forests include Engelmann spruce (*Picea engelmannii* Parry ex Engelm.), Douglas-fir (*Pseudotsuga menziesii* (Mirbel) Franco), subalpine or corkbark fir (*Abies lasiocarpa* (Hook.) Nutt.), limber pine, quaking aspen (*Populus tremuloides* Michx.), ponderosa pine (*Pinus ponderosa* Lawson), and, rarely, pinyon pine (*Pinus edulis* Engelm.). Although bristlecone pine has persisted through historical shifts in climate, a combination of genetic (i.e. low genetic diversity and limited gene flow), demographic (i.e., long generation times and low regeneration rates), and ecological (i.e., forest composition, disease, and insect outbreaks) characteristics suggest that it may be particularly vulnerable to contemporary and projected climate change [21].

Consistent with its narrow distribution, bristlecone pine has the lowest genetic diversity of the high-elevation five-needle pines [22] and is structured into populations with varying proportions of three genetic lineages that differ in diversity and associated climatic factors [22]. Lineage 1 is the most common and present throughout the range and some populations also contain significant proportions of lineage 2 or 3 [22]. Hereafter, populations dominated or solely composed of lineage 1 are referred to as lineage group 1, while those with either lineage 2 or 3 are referred to as lineage group 2 or lineage group 3, respectively. Lineage group 3 has the highest genetic diversity of the three groups and is most common on sites towards the northeastern and southeastern edges of the bristlecone pine core distribution. These sites are the warmest and wettest compared with sites occupied by the other two groups [22].

Although long-lived tree species have adapted repeatedly to past climate change [23], when tolerance limits are exceeded, survival is dependent on the species' phenotypic plasticity, ability to genetically adapt, or migrate to an area with a suitable climate space. Although alpine treelines have moved in the past in response to changing conditions [24,25], the rate of climate change in projections for the near future in many alpine areas is much greater than the rate of change in historical periods [26]. Climate induced range shifts are generally poleward or

upward, and the rate of loss from the trailing edge of a species range is likely to exceed the rate of expansion at the leading edge [12]. As growing space declines with elevation, high elevation ecosystems are likely to see changes in species composition as species migrate up slope.

The objective of this research is to use bristlecone pine population genetics, geographic distribution, and suitable climate space to quantify the species vulnerability to climate change in the near future (the year 2090). Specifically, we (1) evaluate changes in the suitable climate space for bristlecone pine, (2) determine the potential role of population genetic structure (i.e. genetic lineage group) in defining the suitable climate space, and (3) quantify the potential for range shifts in neighboring species and forest types. We hypothesize that, in the near future, (1) the climate space for bristlecone pine will decline; (2) due to its greater climate variability and associated genetic variation, the climate space for lineage group 3 will be more extensive than the climate space for groups 1 or 2; and (3) the climate space for lower elevation species will encroach on the area currently occupied by bristlecone pine.

Materials and methods

Study area

The core of the bristlecone pine range is in south-central Colorado and extends south into New Mexico along the Sangre de Cristo Mountains, with a disjunct population on the San Francisco Peaks in central Arizona [22, 27]. Common bristlecone pine forest types include: (1) sub-dominant spruce and fir species, (2) sub-dominant spruce and fir with aspen, (3) co-dominant limber pine, (4) dominant non-regenerating bristlecone pine, and (5) dominant spruce and fir species with a sub-dominant bristlecone pine. The northern edge of the bristlecone pine range coincides with the northern reach of the North American Monsoon, suggesting a dependence on mid- to late-summer precipitation [28]. This pioneer species primarily regenerates following fire and variation in forest structure is often driven by time since fire [29], although within-forest gaps can also support regeneration [30]. As time since fire increases, bristlecone pine can be replaced by other species over long-time periods on mesic sites.

Tree species location and genetic lineage data

Data from multiple sources were utilized. Forest Inventory and Analysis (FIA) plots were used to determine where all species of interest were present or absent. The USDA Forest Service maintains these permanent plots that systematically sample woody vegetation on forested and non-forested lands within the United States [32,33]. There are 162,409 FIA plots in the geographic area of interest (Arizona, Colorado, New Mexico, Utah, and Wyoming) with bristlecone pine occurring at 235 of these plots. The FIA datasets include a tabulation of the presence-absence of species and elevation. In accordance with FIA proprietary restrictions, true-plot locations are not available for publication. We used the publicly available FIA data supplemented with an additional 165 plot locations throughout the range of bristlecone pine [21, 34]. To understand how the true-plot vs publicly available plot locations might influence our results, we compared (1) the measured true FIA plot elevations to the publicly released plot elevations (i.e. public-plot location) and (2) the climate space defined by the true- and public-plot locations (S1 Fig). Differences in elevation ranged from 0.3 meters to 276 meters and 80% of the public-plot elevations were within 100 meters of the measured elevation of the true-plots. The climate space defined by Crookston et al. [35], who used the true FIA plots locations, was substantially smaller (72%) and exhibited 86% overlap with the climate space defined with the public FIA location data and supplemental plots. The climate space for the public FIA and supplemental plots captured the range and distribution of the Rocky Mountain Bristlecone pine trees. Studies of similar comparisons have also shown that the underlying

distributions of predictor variables sampled with true versus public coordinates are not significantly different at any extent [36].

The geographic location of the bristlecone pine lineage groups defined in Schoettle et al. [22] was used to assign a lineage group to all plots that contained bristlecone pine. Schoettle et al. [22] identified these dominant bristlecone pine genetic lineages. Genetic patterns were evaluated at sixteen sites across the four main mountain ranges (i.e., the Sawatch-Mosquito Range, the Front Range, the San Juan Mountains, and the Sangre de Cristo Mountains) that are in the core bristlecone pine distribution. Sites represented the elevational, latitudinal, and climatic gradients of the species. Genetic diversity was assessed with 21 isozyme loci and sampled trees were assigned to three genetic lineages that varied in diversity and admixture [22]. Geographic lineage group was determined by the proportion of trees in the three lineages. In lineage group 1, the majority of the trees were assigned to lineage 1. Lineage groups 2 and 3 included those sites with a higher proportion of individuals assigned to lineages 2 and 3, respectively. A single polygon was drawn for each lineage group to include all sites assigned to that group (Fig 1). The small disjunct population of bristlecone pine that occurs on the San Francisco Peaks was not evaluated in Schoettle et al. [22], and is included in this analysis as its own group. While Schoettle et al. [22] defined the genetic lineages and the geographic range of lineage groups, we build on this work by defining the suitable climate space for each lineage group. There were 176 plots (out of the 1866 total plots within the geographic range of lineage group 1) with bristlecone pine present in the lineage group 1, 76 plots (out of 747 total plots) in lineage group 2, 119 plots in lineage group 3 (out of 2642 total plots), and 29 plots (38 total plots) in the San Francisco Peaks.

Suitable climate space

Habitat suitability maps have been used with relatively good success to investigate a variety of scientific issues [37]. These tools statistically relate multiple abiotic habitat characteristics [38] with observed occurrences of a species, thus fitting the original definition of the Hutchinsonian environmental niche [38,39]. While these models are useful for understanding the current conditions associated with species locations, they only reflect a snapshot view of the expected relationship between environmental conditions and species presence. A major assumption is that the modeled species are in pseudo-equilibrium with their environment, an assumption that could be questioned for very long-lived tree species [40]. Although this approach doesn't explicitly require a mechanistic link between environmental gradients and population fitness [37], this approach can provide key insights into climatic thresholds that drive species success on the landscape.

The Random Forests classification and the regression tree package [41–43] was used to model species presence and absence based on 27 climate surfaces (Table 1) that were developed by Rehfeldt et al. [44]. This nonparametric approach is resistant to over fitting, multicollinearity, and spatial autocorrelation [41]. The regression tree analysis used 10 independent forests of 100 independent regression trees. Random Forests build each tree using a separate bootstrap sample of the data and withholds 36% of the data to compute the classification error. Indices of variable importance (mean decrease in accuracy) were used to drop the 12 least important variables after the first run. The regression procedure was then rerun nine more times with the remaining predictors, dropping the least important 1 to 3 predictors at each run until classification errors began to increase. The Random Forests run with the fewest variables selected prior to detecting an increase in classification error were considered the most parsimonious bioclimatic model for the genetic lineage or species. The estimated climate and projected climates of each location was run down the 100 regression trees in the final set. A single-




Fig 1. Subalpine and montane species range maps based on Little [31]. Subalpine species include lower subalpine (limber pine, lodgepole pine) and upper subalpine species (Engelmann spruce, subalpine fir, and Douglas fir). Ponderosa pine and quaking aspen make up the montane zone while pinyon-juniper represents lower elevation woodland species. The numbers (1, 2, and 3) in the delineated areas refer to the genetic lineage groups for Rocky Mountain bristlecone pine (red) [22].

<https://doi.org/10.1371/journal.pone.0193481.g001>

Table 1. Acronyms and definitions of climatic variables used in Random Forest procedures. Climate variable layers were developed by Rehfeldt et al. [44]. Variables significant in climate models are bolded.

Acronym	Description
Temperature	
D100	The sum of degree-days (Julian Date) $> 5^{\circ}\text{C} = 100$
DD0	The sum of days $> 0^{\circ}\text{C}$
DD5	The sum of days $> 5^{\circ}\text{C}$
FDAY	First freeze day (Julian Date)
FFP	Length of frost free period (days)
MAT	Mean Annual Temperature ($^{\circ}\text{C}$)
MMAX	Mean Maximum Temperature ($^{\circ}\text{C}$)
MTCM	Mean temperature in coldest month ($^{\circ}\text{C}$)
MTWM	Mean temperature in warmest month ($^{\circ}\text{C}$)
MMIN	Mean minimum temperature in coldest month ($^{\circ}\text{C}$)
MINDD0	Negative degree-days using minimum daily temperature for calculation (days)
SDAY	Date (Julian Date) of last freezing temperature in spring
FDAY	Date (Julian Date) of first freezing temperature in autumn
GSDD5	Mean degree-days $> 5^{\circ}\text{C}$ between SDAY and FDAY. MMAX Mean temperature in warmest month (days)
TDIFF	Summer–winter temperature differential: $\text{MTWM} - \text{MTCM}$ MAPDD5 $(\text{MAP} \times \text{DD5}) / 1000$ ($^{\circ}\text{C}$)
Precipitation	
MAP	Mean annual precipitation (mm)
GSP	April–September precipitation (mm)
SMRP	Summer Precipitation: July–August precipitation (mm)
SPRP	Spring Precipitation: April–May precipitation (mm)
WINP	Winter Precipitation: November–February precipitation (mm)
PRATIO	Growing season precipitation balance: GSP / MAP (mm)
SMRPB	Summer Precipitation Balance: $(\text{jul} + \text{aug} + \text{sep}) / (\text{apr} + \text{may} + \text{jun})$ (mm)
SMRSPRPB	Summer/Spring precipitation balance: $(\text{jul} + \text{aug}) / (\text{apr} + \text{may})$ (mm)
SDI	Summer dryness index: $(\text{GSDD5}) / 0.5 / \text{gsp}$
SDIMINDD0	$\text{SDI} \times \text{MINDD0}$
TDGSP	$\text{TDIFF} / \text{GSP}$ ($^{\circ}\text{C} / \text{mm}$)
MAPDD5	$(\text{MAP} \times \text{MINDD0}) / 1000$

<https://doi.org/10.1371/journal.pone.0193481.t001>

tree prediction is termed a vote. The votes were averaged and only mean votes greater than or equal to the 50% threshold were considered suitable climate for the species.

Once the suitable climate space was defined under current conditions, climate change projections for the year 2090 were used to estimate changes in suitable climate space. The year 2090 was selected to aid comparison to other western species and forest types that have been evaluated with similar methods [44]. We used a regional summary of the IS92a scenario (1% year⁻¹ increase in greenhouse gases after 1990) of the International Panel on Climate Change from General Circulation Models produced by the Canadian Centre for Climate Modeling and Analysis (CGCM2_ghga) [45]. The CGCM2_ghga is generally well respected, and provides an illustration of the potential effects of global warming on species realized climatic niche [45,46]. The regression tree approach has been used to define the realized climate niche for > 70 western tree species [44,46–52] and is used here to model the realized climate niche for bristlecone pine with all lineage groups combined and by individual lineage groups (i.e., lineage groups 1–3 and the San Francisco Peaks).

Shifts in species and forest types

To understand the potential for bristlecone pine forests to be replaced by other forest types, we merged the suitable climate space for the dominant tree species of important forest types or zones. Douglas fir, subalpine fir, and Englemann spruce constitute the upper subalpine forests. Limber pine and lodgepole pine were used to define the lower subalpine forests. Ponderosa pine and quaking aspen (i.e., lower elevation species) are grouped to represent species encroachment from the montane zone, and pinyon-juniper was evaluated to quantify southwestern lower elevation woodland species encroachment. The current and 2090 probability scores for suitable climate space were obtained from the USDA Forest Service—Moscow Forestry Sciences Laboratory (<http://charcoal.cnre.vt.edu/climate/species/>). The same random forests classification approach was used to develop these layers [44,47] and probabilities greater than or equal to 50% were considered suitable climate for forest types and zones.

We evaluated changes in the elevational range in climate space for each forest type or zone. Elevation data (ASTER Global DEM) was obtained from the USGS Earth Explorer (<https://earthexplorer.usgs.gov>). We used changes in the percent overlap with the bristlecone pine climate space to further evaluate the potential for different forest types to encroach on the space occupied by bristlecone pine. The inclusion of additional predictor variables representing the presence-absence of known competitors has been shown to increase the predictive power of models [44,53,54]. By incorporating a measure of the climate space of subalpine forest types, we include information about the physical conditions that are not accounted for by the environmental descriptors included in climate space models. Although this analysis does not conclusively demonstrate competitive exclusion or release, it can provide directional hypotheses that can be tested using observational or experimental field studies.

Results

Climatic differences between lineage groups

Average climate conditions varied across and between bristlecone pine lineage groups. Lineage group 1 had the lowest mean annual precipitation and growing season precipitation, while lineage group 2 had the lowest mean annual temperature, mean temperature in the coldest month, July average temperature, and frost-free period (days; Table 2). Lineage group 2 also had the highest growing season precipitation. Lineage group 3 occupied sites that generally had warmer mean annual and July temperatures, longer frost-free periods, and compared to lineage groups 1 and 2, the warmest mean temperature in the coldest month. Climate conditions in the San Francisco Peaks were most similar to lineage group 3 and exhibited the highest mean temperature in the coldest month and mean annual precipitation.

Suitable climate space

Both temperature and precipitation were important for defining the suitable climate space for bristlecone pine (Fig 2). The random forests classification tree procedure produced 2-variable

Table 2. Mean climatic conditions by lineage group. Mean annual temperature (MAT), mean temperature in the coldest (MTCM) and warmest (MTWM) months, frost free period (FFP), mean annual precipitation (MAP) and growing season precipitation (GSP) were averaged across lineages groups. Standard deviation within lineage groups are listed in parenthesis.

Lineage Group	MAT (°C)	MTCM (°C)	MTWM (°C)	FFP (days)	MAP (mm)	GSP (mm)
1	2.95 (2.5)	-8.49 (1.9)	14.55 (2.9)	56.49 (29.8)	445.08 (226.6)	261.81(86)
2	-0.25 (1.9)	-11.01 (2.1)	11.82 (1.6)	37.92 (10.6)	584.16 (124.8)	321.55 (62.8)
3	6.23 (2.9)	-4.22 (2.5)	17.40 (3.5)	91.89 (34.5)	466.43 (133.8)	314.54 (72.9)
San Francisco Peaks	5.00 (1.9)	-3.97 (1.2)	15.78 (2.2)	72.82 (14.5)	604.95 (117.5)	285.03 (42)

<https://doi.org/10.1371/journal.pone.0193481.t002>




Fig 2. Variable importance plots for the forests used to define the bristlecone pine climate space. Variables with a large (a) mean decrease in accuracy are more important for classification. (b.) The node range for variables show the range in node values that denote presence (black) and absence (grey) of bristlecone pine for each variable.

<https://doi.org/10.1371/journal.pone.0193481.g002>

algorithms that accurately predicted the presence of bristlecone pine (Accuracy = 99%; Fig 2). Most of the errors were due to erroneous predictions of presence (i.e., errors of commission = 98% of errors). Models for all lineage groups together and lineage groups 2 and 3 included the interaction between mean annual precipitation and the sum of days > 5°C (MAPDD5) and the summer precipitation balance (SMRP). For lineage group 1, the summer precipitation balance and the first freeze day (FDAY) defined the suitable climate space. The model for the San Francisco Peaks differed from the other lineage groups and did not contain a temperature parameter. The summer precipitation balance and the summer-winter temperature differential / growing season precipitation (TDGSP) defined the suitable climate space for the San Francisco Peaks. A common thread across models was a high mean decrease in accuracy for SMRP (Fig 3). Variable importance was second highest for MAPDD5, followed by FDAY, and was the lowest for TDGSP. Variable importance was low and differences in node range was small for the San Francisco Peaks model, which may reflect the low sample size available for the analysis (Fig 2).

The bristlecone pine suitable climate space was concentrated in Colorado and New Mexico (71%; Fig 3). Although the range defined by Little [31] for bristlecone pine identified general areas where the species was found, this coverage overlapped just 20% with the current climate space. Of the 399 plots in Colorado, New Mexico, and Arizona with bristlecone pine present, 98% of those plots were within the suitable climate space. Under current conditions, there was a 30% overlap in the suitable climate space of all lineage groups (Fig 4). The majority of the climate space for all lineage groups combined intersected the area dominated by lineage group 3 (90%), 70% intersected lineage group 2, and 30% intersected lineage group 1. A large portion of the current climate space for all lineage groups combined also intersected data from the San Francisco Peaks (70%). Across all lineage groups, the majority of plots that contained




Fig 3. The bristlecone pine range (Little 1971; red) and the current suitable climate space (black) overlapped 20%. Plot level data (grey) showed that areas where bristlecone pine was present (blue; $n = 399$) was in agreement (98% accuracy) with the suitable climate space.

<https://doi.org/10.1371/journal.pone.0193481.g003>

bristlecone pine were on National Forest land (84%) along with most of the current suitable climate space (74%).

Climate change projections

Changes in mean annual temperature (MAT; 4 ± 0.001 °C) and precipitation (MAP; 29 ± 0.08 mm) co-occur with shifts in maximum temperature (MMAX; 3.3 ± 0.001 °C), MAPDD5 (481 ± 0.51 mm), and the summer precipitation balance (SMRP; 8.4 ± 0.03 mm) in the 2090 climate layers. There is also a 21 day (± 0.02) delay in the Julian date of the first freezing temperature in autumn and on average, growing season precipitation increases 7.4 mm (± 0.05) in 2090 compared to current conditions.

Our models project a 74% decrease in the bristlecone pine suitable climate space in 2090 (Figs 3 and 4). Although the climate space declines substantially, overlap between the three lineage groups is maintained at 26%. Comparing the 2090 model for all lineage groups with models for individual lineage groups we estimate that, the future climate space for the San




Fig 4. The current climate space (black) versus the 2090 climate space (red) for (a.) lineage group 1, (b.) lineage group 2, (c.) lineage group 3, and (d.) the San Francisco Peaks. The current distribution of each lineage group is outlined in blue in each respective map.

<https://doi.org/10.1371/journal.pone.0193481.g004>

Francisco Peaks expands in 2090 and occupies the greatest area (99% of the 2090 climate space defined by all lineage group model), followed by lineage group 3 (35%), lineage group 2 (2%), and lineage group 1 (<1%). The 2090 climate space for lineage group 1 is compressed in the southern portion of the current distribution (Fig 4a). While we project substantial declines in the 2090 climate space for all three lineage groups in the species' core distribution, the climate space associated with the San Francisco Peaks expands (Fig 5b). The majority of the bristlecone pine 2090 climate space overlaps with the current bristlecone pine forest (90%), while just 10% occurs in new areas (Figs 4 and 5a). Small gains at high elevations (>3000 m) are projected for lineage groups 1 and 2 (Fig 6b). Between an elevation of 2000–2750 m there is expansion in the geographic distribution of the climate space for lineage group 3, and the climate space associated with the San Francisco Peaks population expands at 1000–2700 m (Fig 6b). The future projection of the San Francisco Peak source may however be an over estimate as it was not constrained by temperature; the lack of a temperature parameter in the San Francisco Peaks




Fig 5. The frequency of (a) contraction and (b) expansion in bristlecone pine climate space by lineage group and elevation in 2090.

<https://doi.org/10.1371/journal.pone.0193481.g005>

model may have been an outcome of the narrow distribution of this disjunct population. Similar to patterns observed in the current climate space, the majority of the 2090 climate space (94%) continues to occur on National Forest land.

Shifts in species and forest types

Under current conditions, the climate space for upper subalpine forests has the greatest overlap with the bristlecone pine climate space (81%), followed by the climate space for lower subalpine forests (68%), species from the montane zone (67%) and pinyon pine woodlands (6%). All groups' climate space overlapped with 45% of the current bristlecone pine climate space. The climate space for the lower elevation forest types was most frequent between 2500 and 3000 m under current conditions (Fig 6). In 2090, the climate space for all forest types declines by (13%) throughout the southern Rockies and distributions shift towards higher elevations (Fig 6). The greatest shifts in elevational distributions of forest type climate space occurs in the




Fig 6. The density of the current (black) and future (red) climate space by elevation for bristlecone pine (model for all lineage groups), upper subalpine, lower subalpine, montane zone, and pinyon pine forests in the Southern Rockies.

<https://doi.org/10.1371/journal.pone.0193481.g006>

montane zone (i.e., ponderosa pine and aspen) and in lower subalpine species (i.e., limber pine and lodgepole pine) (Fig 6). By 2090 the bristlecone pine climate space is entirely shared with the climate space of other forest types. The distribution for lower elevation species' climate space shifts up in elevation, while bristlecone pine climate space is not present and

overlap declines for all forest types climate space except for pinyon pine (+21%). The 2090 climate space for lower elevation species is more frequent between 3000m and 3500m, which coincides with declines in 2090 bristlecone pine climate space.

Discussion

This research used bristlecone pine population genetics [22], geographic distribution, and suitable climate space to evaluate the species' potential vulnerability to climate change in 2090. We hypothesized that (1) the future climate space for bristlecone pine would decline; (2) the climate space for lineage group 3 would be more extensive than the climate space for groups 1 or 2; and (3) the climate space for lower elevation species would encroach on the area that is currently occupied by bristlecone pine. Results support all three hypotheses, indicating that significant reductions in the bristlecone pine climate space and the expansion of the climate space for lower elevation species into higher elevations may reduce the future bristlecone pine range.

Abiotic versus biotic pressure

One general pattern of range limitation is the relative importance of abiotic versus biotic limiting factors in ecological gradients [55]. Currently, abiotic limitations (i.e., maximum temperatures, seasonal precipitation patterns, and disturbance regimes) constrain upward expansion of bristlecone pine forests [56], while biotic limitations (i.e., competition from lower elevation species) have a greater influence on persistence at mid-elevations and limit downward expansion. Like most other species [55], one direction is physically stressful (higher elevation) and the other is biologically stressful (lower elevation) for bristlecone pine. Although climate change is likely to reduce upward physical stress (e.g., minimum temperature), this is expected to be offset by a corresponding increase in competitive stress from lower elevation species that will represent a larger portion of the species composition in the subalpine zone (Fig 6) [57].

Bristlecone pine appears to be excluded from climate space specific to other species (Fig 5). While competitive interactions are not explicitly modeled here, overlapping climate space for co-occurring and neighboring tree species might reflect competitive pressures (Fig 6) [58,59]. Suitable 2090 climate space for bristlecone pine exhibits a bimodal distribution along the elevation gradient in the future presumably due to persistence at lower elevations in the Southern Rocky Mountains and exclusion at the mid-elevations by changes in climatic conditions that favor other lower elevation species. This pattern can be seen currently along the eastern margins of South Park and the Sawatch range in central Colorado, where bristlecone pine coexists with ponderosa pine at low elevations, is absent from portions of the upper montane and subalpine zones, then reoccurs near alpine treeline [60]. Historical patterns of species distributions along the elevational gradient further suggests the role of competition in the distribution of five-needle pines [61]. Pollen and charcoal data from the Yellowstone region suggest that five-needle pine species, now restricted to higher elevations, were abundant across broader elevational ranges during a period when Engelmann spruce and lodgepole pine were less common [62]. These shifts have strong implications for changes in fire regimes, vulnerability to pest and pathogens, and altered C dynamics across the Southern Rockies.

Climate change and disturbance regimes

Climate change is expected to increase the number of fires at high elevations in the southern Rockies [63,64], resulting in a decline in the abundance of older forests [29]. As an early seral species, bristlecone pine forest structure is currently thought to be the result of long fire return intervals, time since fire, and dispersal limitations and modified by smaller intra-stand disturbances [29,30]. Shifts in the location of suitable climate, the increase in lower elevation species

presence that have different fire regimes, and recovery patterns in bristlecone pine forest might alter the role of fire in promoting bristlecone pine forests in the future. With these shifts in forest composition, fire return intervals are likely to be greatly reduced (i.e. increasing fire frequency), which might promote novel forest development throughout the current bristlecone pine range.

An increase in fire frequency may promote regeneration of bristlecone pine, increasing the adaptive capacity of the species [65]. Diversifying age-class structure, particularly through increased abundance of younger cohorts, may serve the dual purposes of 1) facilitating more rapid selection for adaptive traits [17,66,67], and 2) ensuring the presence of many small-diameter individuals less vulnerable to drought stress, and more likely to survive mountain pine beetle (*Dendroctonus ponderosae* Hopkins) attack [60]. However, an increase in fire frequency and a smaller climate space suitable for bristlecone pine at the high elevations may reduce the likelihood of bristlecone pine trees achieving exceptional longevity (>1500 years) in the future.

Bristlecone pine is confronted by another directional selection threat—the lethal disease white pine blister rust (WPBR). The southern Rocky Mountains are at the infection front of the non-native pathogen *Cronartium ribicola* J. C. Fisch. that causes WPBR [68]. Bristlecone pine is susceptible to WPBR [69,70] and the disease is expected to continue to spread through much of the bristlecone pine distribution [71]. Currently, the disease has only been found on bristlecone pine in south-central Colorado [72] in the area that contains lineage group 3 [22]. Genetic resistance to WPBR is not distributed uniformly within the five-needle pine species, including bristlecone pine [22], and how disease resistance traits are associated with abiotic stress tolerances is unclear but may be significant [73,74]. Consequently, the climate space of five-needle pines may be shifting under directional selection for resistance to WPBR [73,74], further complicating climate change projections. Range-wide genetic collections (seed and tissues) of bristlecone pine have been made and these efforts are the first proactive coordinated range-wide genetic collection design and forest health assessment for a threatened, but not yet heavily impacted, tree species [21]. The 2090 climate space projections presented here provide further insights to guide proactive management to increase the resilience of the threatened populations [66] and consider assisted migration of WPBR resistant genotypes to areas with suitable 2090 climate space.

Recent threats from the expanding distribution of native bark beetle infestations [75] may also affect future tree species range distributions. This may be of particular concern if a greater proportion of bristlecone pines distribution is restricted to lower elevations in the future as projected by the results of this study (Fig 6). However, the other species that may be excluding bristlecone from the higher elevation climate spaces in the future also have uncertain futures in the presence of expanding and intensifying bark beetle epidemics.

Limitations to understanding climate change impacts on species range shifts

The climate spaces of specific genetic lineage groups for bristlecone pine shifted differently under a climate change scenario, with proportional representation of each climate space changing in the future. The implication of this shift on genetic diversity patterns is complex. Populations can respond to environmental change by moving to a new area corresponding to environmental conditions for which they are adapted [15], or through phenotypic plasticity and/or genetic adaptation to the new conditions, or combinations of these responses [17]. The inferences made in the current study to future species distribution and genetic structure assume only unimpeded population migration and no other evolutionary processes.

Persistence, via plasticity or adaptation, on sites with climates that are no longer suitable is not considered in the model, suggesting that the projected range contraction for the 2090 bristlecone pine climate space may be unrealistically severe. Due to the stress tolerance and long life span of bristlecone pine, individual trees may persist in suboptimal environments for centuries, yet without regeneration populations are not adaptable or sustainable into the future. Plant species distributions that are in climatic disequilibrium might lead to an increased potential for extinction within all or portions of species' ranges. This, combined with long fire return intervals, could be the cause of the lack of regeneration in some bristlecone pine forests [29]. The low genetic diversity, delayed reproductive maturity, protracted regeneration dynamic, and slow growth of bristlecone pine suggests a low adaptive potential to rapidly changing conditions [22].

Conclusions

Although the availability of bristlecone pine climate space is projected to decline substantially in 2090, availability of suitable climate space for lineage group 3 and for populations from the San Francisco Peaks suggest that these populations are likely to be the most suitable for the future climate. While these populations may be of particular management interest, current populations of lineage groups whose 2090 climate space contracts (lineage group 1 and 2) may be of particular conservation interest. Considering that the majority of the current and future suitable climate space exists on Forest Service land, the future of bristlecone pine forests is essentially within the jurisdiction of the United States Forest Service. With management intervention, assisted migration is possible and this analysis provides insights into suitable genetic lineage groups (i.e. lineage group 3 and populations from the San Francisco Peaks) that may be considered for source material. Although this approach provides an important and novel step beyond models that assume a species is genetically homogenous, further integration of known evolutionary processes are needed in climate change response modeling [14].

Supporting information

S1 Fig. (a.) Differences in elevation between true and public Forest Inventory and Analysis (FIA) plot locations. (b.) A comparison of the bristlecone pine climate space defined by true (yellow) and public FIA plots combined with supplemental plots. (TIF)

Acknowledgments

The authors would like to acknowledge the excellent support provided by Nicholas Crookston and the Rocky Mountain Research Station. We also extend thanks to Kristen Pelz for FIA support and Peter Z. Fulè for providing location information for bristlecone pine in the San Francisco Peaks ($n = 23$ plots) [34]. Any opinions, findings, conclusions, or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the authors affiliation.

Author Contributions

Conceptualization: Anna W. Schoettle.

Data curation: Anna W. Schoettle, Jonathan D. Coop.

Formal analysis: Sparkle L. Malone.

Funding acquisition: Sparkle L. Malone, Anna W. Schoettle.

Investigation: Anna W. Schoettle, Jonathan D. Coop.

Methodology: Sparkle L. Malone.

Project administration: Anna W. Schoettle.

Supervision: Anna W. Schoettle.

Validation: Sparkle L. Malone.

Writing – original draft: Sparkle L. Malone.

Writing – review & editing: Sparkle L. Malone, Anna W. Schoettle, Jonathan D. Coop.

References

1. Oldfield S, Lusty C, Nackineven A. The world list of threatened trees [Internet]. Cambridge: WCMC, IUCN; 1998. http://125.235.8.196:8080/dspace/handle/CEID_123456789/7191
2. Sturrock RN, Frankel SJ, Brown AV, Hennon PE, Kliejunas JT, Lewis KJ, et al. Climate change and forest diseases. *Plant Pathol.* 2011; 60: 133–149. <https://doi.org/10.1111/j.1365-3059.2010.02406.x>
3. Savage M, Mast JN. How resilient are southwestern ponderosa pine forests after crown fires? *Can J For Res.* 2005; 35: 967–977. <https://doi.org/10.1139/x05-028>
4. McDowell NG, Allen CD. Darcy's law predicts widespread forest mortality under climate warming. *Nat Clim Chang Nature Research.* 2015; 5: 669–672. Available from: <https://www.nature.com/articles/nclimate2641>
5. Kelly AE, Goulden ML. Rapid shifts in plant distribution with recent climate change. *Proc Natl Acad Sci U S A.* 2008; 105: 11823–11826. <https://doi.org/10.1073/pnas.0802891105> PMID: 18697941
6. Randin CF, Engler R, Normand S, Zappa M, Zimmerman NE, Pearman PB, et al. Climate change and plant distribution: local models predict high-elevation persistence. *Glob Chang Biol.* 2009; 15: 1557–1569. <https://doi.org/10.1111/j.1365-2486.2008.01766.x>
7. McDowell NG, Coops NC, Beck PSA, Chambers JQ, Gangodagamage C, Hicke JA, et al. Global satellite monitoring of climate-induced vegetation disturbances. *Trends Plant Sci.* 2015; 20: 114–123. <https://doi.org/10.1016/j.tplants.2014.10.008> PMID: 25500552
8. Woodward FI, Williams BG. Climate and plant distribution at global and local scales. In: Colin Prentice I, van der Maarel E, editors. *Theory and Models in Vegetation Science.* 1987. pp. 189–197. https://doi.org/10.1007/978-94-009-4061-1_19
9. Parmesan C, Yohe G. A globally coherent fingerprint of climate change impacts across natural systems. *Nature.* 2003; 421: 37–42. <https://doi.org/10.1038/nature01286> PMID: 12511946
10. Parmesan C. Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol Evol Syst.* 2006; 37: 637–669. Available from: <http://www.jstor.org/stable/30033846>
11. Smithers BV, North MP, Millar CI, Latimer AM. Leap-frog in slow-motion: divergent responses of tree species and life stages to climatic warming in Great Basin sub-alpine forests. *Glob Chang Biol.* 2017; <https://doi.org/10.1111/gcb.13881> PMID: 28850759
12. Breshears DD, Huxman TE, Adams HD, Zou CB, Davison JE. Vegetation synchronously leans upslope as climate warms. *Proc Natl Acad Sci U S A.* 2008; 105: 11591–11592. <https://doi.org/10.1073/pnas.0806579105> PMID: 18697950
13. Williams JW, Jackson ST, Kutzbach JE. Projected distributions of novel and disappearing climates by 2100 AD. *Proc Natl Acad Sci U S A.* 2007; 104: 5738–5742. <https://doi.org/10.1073/pnas.0606292104> PMID: 17389402
14. Alberto FJ, Aitken SN, Alía R, González-Martínez SC, Hänninen H, Kremer A, et al. Potential for evolutionary responses to climate change—evidence from tree populations. *Glob Chang Biol.* 2013; 19: 1645–1661. Available: <http://onlinelibrary.wiley.com/doi/10.1111/gcb.12181/full> PMID: 23505261
15. Kueppers LM, Conlisk E, Castanha C, Moyes AB, Germino MJ, de Valpine P, et al. Warming and provenance limit tree recruitment across and beyond the elevation range of subalpine forest. *Glob Chang Biol.* 2017; 23: 2383–2395. <https://doi.org/10.1111/gcb.13561> PMID: 27976819
16. Kremer A, Potts BM, Delzon S. Genetic divergence in forest trees: understanding the consequences of climate change. *Funct Ecol.* 2014; 28: 22–36. <https://doi.org/10.1111/1365-2435.12169>

17. Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S. Adaptation, migration or extirpation: climate change outcomes for tree populations. *Evol Appl*. 2008; 1: 95–111. <https://doi.org/10.1111/j.1752-4571.2007.00013.x> PMID: 25567494
18. Beniston M. Mountain weather and climate: A general overview and a focus on climatic change in the Alps. *Hydrobiologia*. 2006; 562: 3–16. <https://doi.org/10.1007/s10750-005-1802-0>
19. Pearson RG, Dawson TP, Berry PM, Harrison PA. SPECIES: A spatial evaluation of climate impact on the envelope of species. *Ecol Modell*. 2002; 154: 289–300. [https://doi.org/10.1016/S0304-3800\(02\)00056-X](https://doi.org/10.1016/S0304-3800(02)00056-X)
20. Brunstein FC, Yamaguchi DK. The oldest known Rocky Mountain Bristlecone pines (*Pinus aristata* Engelm.). *Arct Alp Res*. 1992; 24: 253–256. <https://doi.org/10.2307/1551666>
21. Schoettle AW, Coop JD. Range-wide conservation of *Pinus aristata*: a genetic collection with ecological context for proactive management today and resources for tomorrow. *New Forests*. 2017; 48: 181–199. <https://doi.org/10.1007/s11056-017-9570-z>
22. Schoettle AW, Goodrich AB, Hipkins V, Richards C, Kray J. Geographic patterns of genetic variation and population structure in *Pinus aristata*, Rocky Mountain bristlecone pine. *Can J For Res*. 2012; 42: 23–37. <https://doi.org/10.1139/x11-152>
23. Ackerly DD. Community assembly, niche conservatism, and adaptive evolution in changing environments. *Int J Plant Sci*. 2003; 164: S165–S184. <https://doi.org/10.1086/368401>
24. Carrara PE, McGeehin JP. Evidence of a higher late-Holocene treeline along the Continental Divide in central Colorado. *Holocene*. 2015; 25: 1829–1837. Available from: <http://journals.sagepub.com/doi/abs/10.1177/0959683615591353>
25. Danby RK, Hik DS. Variability, contingency and rapid change in recent subarctic alpine tree line dynamics. *J Ecol*. 2007; 95: 352–363. <https://doi.org/10.1111/j.1365-2745.2006.01200.x>
26. Houghton JT, Ding YDJG, Griggs DJ, Noguer M, van der Linden PJ, Dai X, et al. Climate change 2001: the scientific basis. The Press Syndicate of the University of Cambridge; 2001.
27. Critchfield WB, Little EL. Geographic Distribution of the Pines of the World [Internet]. Washington, D. C.: U.S. Department of Agriculture, Forest Service; 1966. <https://archive.org/details/geographicdistri991crit>
28. Schoettle AW. Ecological roles of five-needle pine in Colorado: potential consequences of their loss. In: Snieszko RA et al, editors. Breeding and genetic resources of five-needle pines: growth, adaptability and pest resistance. USDA Forest Service, Rocky Mountain Research Station; Fort Collins, CO: 2004. Proceedings No.: RMRS-P-32. Pp 124–135. <http://www.treesearch.fs.fed.us/pubs/52826>
29. Baker WL. Structure, Disturbance, and Change in the Bristlecone Pine Forests of Colorado, U.S.A. *Arct Alp Res*. 1992; 24: 17–26. <https://doi.org/10.2307/1551316>
30. Brown PM, Schoettle AW. Fire and stand history in two limber pine (*Pinus flexilis*) and Rocky Mountain bristlecone pine (*Pinus aristata*) stands in Colorado. *Int J Wildland Fire*. 2008; 17: 339–347. <https://doi.org/10.1071/WF06159>
31. Little EL. Atlas of United States Trees: conifers and important hardwoods [Internet]. Washington, DC: Forest Service; 1971. Report No.: Miscellaneous Publication 1146. <https://market.android.com/details?id=book-mlovAAAAYAAJ>
32. Bechtold WA, Patterson PL, Editors. The enhanced forest inventory and analysis program-national sampling design and estimation procedures [Internet]. USDA Forest Service, Southern Research Station; Asheville, NC: 2005. Report No.: SRS-GTR-80. <http://www.treesearch.fs.fed.us/pubs/download/20371.pdf>
33. O'Connell BM, LaPoint EB, Turner JA, Ridley T, Pugh SA, Wilson AM, et al. The Forest Inventory and Analysis database: Database description and user guide version 6.0 for phase 2. US Department of Agriculture Forest Service. 2014;6.0. <https://www.fia.fs.fed.us/library/database-documentation/>
34. Cocke AE, Fulé PZ, Crouse JE. Forest change on a steep mountain gradient after extended fire exclusion: San Francisco Peaks, Arizona, USA. *J Appl Ecol*. 2005; 42: 814–823. <https://doi.org/10.1111/j.1365-2664.2005.01077.x>
35. Crookston NL. Plant species and climate profile predictions [Internet]. Rocky Mountain Research Station, Moscow Forestry Sciences Laboratory; 2012. <http://charcoal.cnre.vt.edu/climate/species/>
36. Gibson J, Moisen G, Frescino T, Edwards TC. Using publicly available Forest Inventory Data in climate-based models of tree species distribution: examining effects of true versus altered location coordinates. *Ecosystems*. 2014; 17: 43–53. <https://doi.org/10.1007/s10021-013-9703-y>
37. Guisan A, Thuiller W. Predicting species distribution: offering more than simple habitat models. *Ecol Lett*. 2005; 8: 993–1009. <https://doi.org/10.1111/j.1461-0248.2005.00792.x>
38. Kearney M, Porter WP. Mapping the fundamental niche: physiology climate, and the distribution of a nocturnal lizard. *Ecology*. 2004; 85: 3119–3131. <https://doi.org/10.1890/03-0820>

39. Hutchinson GE. Concluding remarks. Cold Spring Harbor symposium on quantitative biology. 1957; 22: 415–427.
40. Bruening JM, Tran TJ, Bunn AG. Fine-scale modeling of bristlecone pine treeline position in the Great Basin, USA. *Environmental Research Letters*. 2017; 12: 014008. Available from: <http://iopscience.iop.org/article/10.1088/1748-9326/aa5432/meta>
41. Breiman L. Random Forests. *Mach Learn*. 2001; 45: 5–32. <https://doi.org/10.1023/A:1010933404324>
42. Liaw A, Wiener M. Classification and regression by randomForest. *R news*. ai2-s2-pdfs.s3.amazonaws.com; 2002;2: 18–22. <http://ai2-s2-pdfs.s3.amazonaws.com/6e63/3b41d93051375ef9135102d54fa097dc8cf8.pdf>
43. R Core Team. The R project for statistical computing [Internet]. 2014. www.r-project.org
44. Rehfeldt GE, Crookston NL, Warwell MV, Evans JS. Empirical analyses of plant-climate relationships for the western United States. *International Journal of Plant Sciences*. 2006; 167: 1123–1150.
45. Fyfe JC, Flato GM. Enhanced climate change and its detection over the Rocky Mountains. *J. Climate*. 1999; 12: 230–243. <https://doi.org/10.1175/1520-0442-12.1.230>
46. Warwell MV, Rehfeldt GE, Crookston NL. Modeling species' realized climatic niche space and predicting their response to global warming for several western forest species with small geographic distributions [Internet]. USDA Forest Service; Pacific Northwest Research Station; Portland OR: 2010. Report No.: PNW-GTR-802. Pp 171–182. <http://www.treeseearch.fs.fed.us/pubs/download/37041.pdf>
47. Crookston NL, Rehfeldt GE, Warwell MV. Using Forest Inventory and Analysis data to model plant-climate relationships. *Proceedings of the Seventh Annual Forest Inventory and Analysis Symposium*. USDA Forest Service, Northern Research Station: 2005. Report No.: WO-GTR-77, Pp 243–250. https://www.nrs.fs.fed.us/pubs/gtr/gtr_wo077/gtr_wo077_243.pdf
48. Warwell MV, Rehfeldt GE, Crookston NL. Modeling contemporary climate profiles of whitebark pine (*Pinus albicaulis*) and predicting responses to global warming [Internet]. USDA Forest Service: 2006. Report No.: R6-NR-FHP-2007-01. <https://pdfs.semanticscholar.org/e1d5/ef37b3e8f9a80683aa51b682f5f496b3b35a.pdf#page=149>
49. Rehfeldt GE, Ferguson DE, Crookston NL. Aspen, climate, and sudden decline in western USA. *For Ecol Manage*. 2009; 258: 2353–2364. <https://doi.org/10.1016/j.foreco.2009.06.005>
50. Sáenz-Romero C, Rehfeldt GE, Crookston NL, Duval P, St-Amant R, Beaulieu J, et al. Spline models of contemporary, 2030, 2060 and 2090 climates for Mexico and their use in understanding climate-change impacts on the vegetation. *Clim Change*. 2010; 102: 595–623. <https://doi.org/10.1007/s10584-009-9753-5>
51. Rehfeldt GE, Crookston NL, Sáenz-Romero C, Campbell EM. North American vegetation model for land-use planning in a changing climate: a solution to large classification problems. *Ecol Appl*. 2012; 22: 119–141. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22471079> PMID: 22471079
52. Joyce DG, Rehfeldt GE. Climatic niche, ecological genetics, and impact of climate change on eastern white pine (*Pinus strobus* L.): Guidelines for land managers. *For Ecol Manage*. 2013; 295: 173–192. <https://doi.org/10.1016/j.foreco.2012.12.024>
53. Leathwick JR, Austin MP. Competitive interactions between tree species in New Zealand's old-growth indigenous forests. *Ecology*. 2001; 82: 2560–2573. <https://doi.org/10.2307/2679936>
54. Anderson RP, Townsend Peterson A, Gomez-Laverde M. Using niche-based GIS modeling to test geographic predictions of competitive exclusion and competitive release in South American pocket mice. *Oikos*. 2002; 98: 3–16.
55. Brown JH, Stevens GC, Kaufman DM. The geographic range: size, shape, boundaries, and internal structure. *Annu Rev Ecol Syst*. 1996; 27: 597–623. Available from: <http://annualreviews.org/doi/abs/10.1146/annurev.ecolsys.27.1.597>
56. Körner C. The nutritional status of plants from high altitudes. *Oecologia*. 1989; 81: 379–391. <https://doi.org/10.1007/BF00377088> PMID: 28311193
57. Körner C. *Alpine treelines: functional ecology of the global high elevation tree limits*. Basel, Switzerland: Springer; 2012.
58. Chase JM, Leibold MA. *Ecological niche: linking classical and contemporary approaches*. Chicago: The University of Chicago Press; 2003.
59. Broennimann O, Fitzpatrick MC, Pearman PB, Petitpierre B, Pellissier L, Yoccoz NG, et al. Measuring ecological niche overlap from occurrence and spatial environmental data. *Glob Ecol Biogeogr*. 2012; 21: 481–497. <https://doi.org/10.1111/j.1466-8238.2011.00698.x>
60. Coop JD, Schoettle AW. Fire and high-elevation, five-needle pine (*Pinus aristata* & *P. flexilis*) ecosystems in the Southern Rocky Mountains: what do we know? Keane RE et al, eds. *The future of high-elevation, five-needle white pines in Western North America: Proceedings of the High Five Symposium*.

- USDA Forest Service, Rocky Mountain Research Station; Fort Collins: 2011. Report No.: RMRS-P-63. Pp 164–175. http://www.fs.fed.us/rm/pubs/rmrs_p063.html
61. Whitlock C, Colombaroli D, Conedera M, Tinner W. Land-use history as a guide for forest conservation and management. *Conserv Biol*. 2017; Available from: <http://onlinelibrary.wiley.com/doi/10.1111/cobi.12960/full>
62. Iglesias V, Krause TR, Whitlock C. Complex response of white pines to past environmental variability increases understanding of future vulnerability. *PLoS One*. 2015; 10: e0124439. <https://doi.org/10.1371/journal.pone.0124439> PMID: 25885810
63. McKenzie D, Gedalof Z, Peterson DL. Climatic change, wildfire, and conservation. *Conservation*. 2004; Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1523-1739.2004.00492.x/full>
64. Littell JS, McKenzie D, Peterson DL, Westerling AL. Climate and wildfire area burned in western U.S. ecoprovinces, 1916–2003. *Ecol Appl*. 2009; 19: 1003–1021. <https://doi.org/10.1890/07-1183.1> PMID: 19544740
65. Schoettle AW, Klutsch JG, Sniezko RA. Integrating regeneration, genetic resistance, and timing of intervention for the long-term sustainability of ecosystems challenged by non-native pests—a novel proactive approach. In: Sniezko RA, et al., editors. *Proceedings of the fourth international workshop on the genetics of host-parasite interactions in forestry: Disease and insect resistance in forest trees*. USDA Forest Service, Pacific Southwest Research Station Albany, CA: 2012. Report No.: PSW-GTR-240 pp. 112–123. <http://www.treeseearch.fs.fed.us/pubs/download/42527.pdf>
66. Schoettle AW, Sniezko RA. Proactive intervention to sustain high-elevation pine ecosystems threatened by white pine blister rust. *J Forest Res*. 2007; 12: 327–336. <https://doi.org/10.1007/s10310-007-0024-x>
67. Field SG, Schoettle AW, Klutsch JG, Tavener SJ, Antolin MF. Demographic projection of high-elevation white pines infected with white pine blister rust: a nonlinear disease model. *Ecol Appl*. 2012; 22: 166–183. Available: <https://www.ncbi.nlm.nih.gov/pubmed/22471082> PMID: 22471082
68. Burns KS, Schoettle AW, Jacobi WR, Mahalovich MF. Options for the management of white pine blister rust in the Rocky Mountain Region [Internet]. USDA Forest Service, Rocky Mountain Research Station; Fort Collins, CO: 2008. Report No.: RMRS-GTR-206. http://www.fs.fed.us/rm/pubs/rmrs_gtr206.html
69. Blodgett JT, Sullivan KF. First Report of White Pine Blister Rust on Rocky Mountain Bristlecone Pine. *Plant Dis. Scientific Societies*; 2004; 88: 311–311. <https://doi.org/10.1094/PDIS.2004.88.3.311A>
70. Schoettle AW, Sniezko RA, Kegley A, Burns KS. Preliminary overview of the first extensive rust resistance screening tests of *Pinus flexilis* and *Pinus aristata*. In: Keane RE et al., editors. *The future of high-elevation, five-needle white pines in Western North America: proceedings of the high five symposium*. USDA Forest Service, Rocky Mountain Research Station; Fort Collins, CO: 2011. Proceedings No.: RMRS-P-63. Pp 265–269. <https://www.fs.usda.gov/treeseearch/pubs/38235>
71. Howell B, Burns KS, Kearns HSJ, Witcosky JJ, Cross FJ. Biological evaluation of a model for predicting presence of white pine blister rust in Colorado based on climatic variable and susceptible white pine species distribution. USDA Forest Service, Rocky Mountain Region; Lakewood, CO; 2006. Report No.; Biological Evaluation R2-06-04.
72. Burns KS. White pine blister rust surveys in the Sangre de Cristo and Wet Mountains of Southern Colorado. USDA Forest Service, Rocky Mountain Region; Lakewood CO: 2006. Report No.; Biological Evaluation R2-06-05 https://www.researchgate.net/profile/Anna_Schoettle/publication/237418495_White_Pine_Blister_Rust_Surveys_in_the_Sangre_de_Cristo_and_Wet_Mountains_of_Southern_Colorado/links/00b7d5310212fe5031000000.pdf
73. Vogan PJ, Schoettle AW. Selection for resistance to white pine blister rust affects the abiotic stress tolerances of limber pine. *For Ecol Manage*. 2015; 344: 110–119. <https://doi.org/10.1016/j.foreco.2015.01.029>
74. Vogan PJ, Schoettle AW. Carbon costs of constitutive and expressed resistance to a non-native pathogen in limber pine. *PLoS One*. 2016; 11: e0162913. <https://doi.org/10.1371/journal.pone.0162913> PMID: 27706249
75. Bentz BJ, Régnière J, Fettig CJ, Hansen EM, Hayes JL, Hicke JA, et al. Climate change and bark beetles of the Western United States and Canada: direct and indirect effects. *Bioscience*. 2010; 60: 602–613. <https://doi.org/10.1525/bio.2010.60.8.6>