

Development of molecular diagnostic methods to distinguish Acerola species for quality assurance of food, dietary supplements and natural health products

Subramanyam Ragupathy

ragu@uoguelph.ca

University of Guelph

Arunachalam Thirugnanasambandam

University of Guelph

Varathan Vinayagam

University of Guelph

Ragupathy Sneha

American University of Antigua

Steven G. Newmaster

University of Guelph

Article

Keywords: Barbados cherry, DNA identification, NMR fingerprints, metabolite spectra, adulteration, food security

Posted Date: May 15th, 2025

DOI: <https://doi.org/10.21203/rs.3.rs-6372499/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Scientific Reports on August 4th, 2025. See the published version at <https://doi.org/10.1038/s41598-025-12408-6>.

Abstract

Acerola (Barbados cherries) has become a highly traded superfruit because it contains many phytonutrients and is a good source of vitamin C. The fruits of *Malpighia glabra*, and *M. emarginata* are utilized in food products, dietary supplements and natural health products. However, there are differences among the fruit of *Malpighia* species with respect to phytochemicals, nutrient value and clinical research. Furthermore, there is evidence of adulteration with other fruit such as cherries (*Prunus* spp.). Unfortunately, conventional morphological examination does not distinguish acerola fruit species. Furthermore, no published methods are available to distinguish the fruits of these species including chemical and DNA based techniques. This risk to quality assurance (QA) is increased when considering processed berries into juice or powdered ingredients of which are the most common source for manufactures. This lack of QA methods also increases the risk of adulteration with cheaper fruit from other species. The goal of this research is to provide orthogonal molecular methods to authenticate Acerola fruit ingredients and discuss the benefits and constraints of these two different methods. This research supports quality assurance (QA) programs with fit-for-purpose methods for verifying the authenticity of acerola species ingredients from suppliers.

Introduction

The superfruit market has had significant growth in the marketplace putting demand on suppliers. Consumer demand for superfoods such as superfruit is increasing and attracting the attention of multiple industry sectors including food products, dietary supplements and natural health products ¹. Industry has responded to this demand with marketing strategies for superfood ingredients in their products that promote considerable health benefits within the human body and play an important role in the progression of degenerative diseases ^{2,3}. The superfruit market alone was valued at 134.8 billion USD in 2024 and is expected to grow over 200 billion USD in the next decade (FMI 2025). This is partly driven by the demand from contemporary consumers that value a healthier and environmentally friendly source of nutrition of which superfoods are highly nutritional with bioavailable nutrients with documented bioactivity within the body due to the concentration of nutrients and bioactive ingredients ⁴. Few fruits meet the criteria of high nutrient value with notable nutrients such as vitamin C and can be sourced at a reasonable cost by large manufactures that focus on global markets.

Acerola berries are designated as a superfruit that can be sustainably grown by producers and sourced in large quantities in the marketplace ⁵. Acerola berries are also known as Barbados cherry and West Indian cherry and are grown commercially as small trees or shrubs. Taxonomically there are two species (*Malpighia glabra* L.; *Malpighia emarginata* DC.) from the family Malpighiaceae that contains about 45 species ^{6,7}. There is mention in the literature of another species *Malpighia puniceifolia* L. which is a synonym for *Malpighia glabra*. The fruit of these two species is undistinguishable as a red fleshy drupe that produces a juicy acidic pulp. The common names are used inconsistently for the respective species causing some confusion when sourcing acerola ingredients from the global supply chain ⁸. *Malpighia*

glabra is native to the Americas, including Mexico, Central America, South America, and the Caribbean with more recent introduction and cultivation in USA (Florida, California, Texas), Cuba, Jamaica, and Puerto Rico. *Malpighia emarginata* is originally from Mexico in the Yucatán peninsula, but is now cultivated in Central America, the Caribbean, and South America as far south as Peru^{9,10} Brazil is the world's largest producer of acerola berries with most of the production in the northeast states of Pernambuco, Paraíba, Bahia, and Ceará^{5,11}. Both species have been introduced for cultivation in Southeast Asia and southern India. In India, cultivation is reported in backyard gardens across the states of Tamil Nadu, Karnataka, Kerala and Maharashtra. More recently production has been recorded in the Andaman and Nicobar Islands¹². Production is relatively small in the southern USA. The largest markets for acerola products include United States, Germany, France, and Japan.

Acerola fruit has considerable nutritional and health benefits. There is a plethora of research on acerola to support pharmacological, medicinal and nutritional claims of which have been well documented in several scientific reviews^{5,13,14}. Acerola health benefits include high antioxidant capacity^{6,15,16}, antitumor¹⁷, antimutagenic^{18–23}, antidiabetic²⁴, hepatoprotective²⁵ and functional properties like skin anti-aging²⁶, skin whitening, and multidrug resistant reversal activity^{11,27}. This body of research is supported by many biochemistry studies that have recorded significant phytonutrients such as flavonoids, anthocyanins, carotenoids, phenolics, benzoic acid derivatives and phenylpropanoids within comprehensive scientific reviews^{28,29}. The presence of these bioactive compounds provides multiple hypothetical mechanisms to support the numerous health claims^{27,28}. This includes various biological activities using in vitro and in vivo models in which probable mechanisms have been determined¹³. One of the most studied areas of research underpins the claim that acerola is one of the richest natural sources of ascorbic acid up to 100 times more than that of oranges and lemons^{13,30–32}. It has been reported that the vitamin C of acerola is more easily absorbed than other forms including synthetic ascorbic acid^{27,33}. Acerola berries are consumed and prepared as a nutritional drink and traditional remedy in Brazil. In Europe (e.g., France, Germany, and Hungary) the fruit is utilized as juice, whilst in multiple forms within the in the United States where it is consumed as a rich source of ascorbic acid for vitamin C within supplements and natural health products²⁷.

There is a lack of published research on quality assurance tools in the acerola industry. These QA tools are needed to ensure species ingredient authenticity and prevent adulteration with other cheaper fruit species. Morphological techniques are inadequate because they cannot differentiate among the species of *Malpighia*. DNA-based markers have become a popular method for identifying and verifying botanical ingredients. This is because the genetic makeup of each species is unique, regardless of factors such as sample age, physiological condition, environmental growth conditions, harvesting methods, storage, and processing³⁴. The DNA extracted from the fruit, leaves, stems or roots of plants all carry the same genetic sequences of which can be stored for long periods of time as they are stable³⁵. Multiple DNA regions have been utilized to develop specific markers for identifying and authenticating botanical ingredients^{36,37}. Several recent reviews provide the benefits and challengers of using DNA-based

methods for quality assurance of botanical ingredients^{38–41}. The use of chemical authentication has been used extensively for quality assurance of botanical ingredients⁴². Traditional analytical chemistry methods are well established and the advantages and limitations are discussed in numerous publications^{43–45}. More advanced methods have been utilized providing rigorous approach to quality assurance of botanical ingredients that overcomes the challenges of traditional analytical methods^{46–48}. The development of orthogonal molecular methods for specific species ingredients has eliminated the challenges of DNA methods (e.g., degraded DNA in processed samples) and presented alternative metabolomic methods such as nuclear magnetic resonance (NMR)^{49–54}. The use of phytochemical analysis and metabolomics methods such as chemical profiling by NMR are the fit-for-purpose in identifying the active ingredients labelled on herbal products^{55,56}. Analytical chemistry and DNA methods are not fit-for-purpose when evaluating complex samples, such as herbal dietary supplements that can include a mixture of several plant species in one product^{36,54,57}. NMR methods overcome the admixtures barrier and have a wide range of applications in food and botanical industry, as ingredients authentication and quantification, including active ingredient quantification and differentiating the geographic origins ingredients^{58–65}. These molecular methods need to be assessed in the context of specific species ingredients to determine fit-for-purpose in the quality assessment of sourced botanical ingredients.

The objective of this study is to develop of Nuclear Magnetic Resonance Fingerprints and DNA sequence methods to distinguish Acerola species (*Malpighia glabra*, *M. emarginata*) for quality assurance of Food, dietary supplements and natural health products. More specifically we used two methods including, 1) DNA sequencing the ITS nuclear region for 16 samples and, 2) 1H-NMR metabolite fingerprinting to analyze 16 samples consisting of 10 samples of two Acerola species (*Malpighia glabra*, *M. emarginata*) and 6 samples of common adulterant cherry (*Prunus spp.*) species. The validation of genomics and metabolomics approaches in this study will ensure quality assurance for both product identity and purity.

Materials and Methods

Botanical Samples This study develop two separate orthogonal methods using 32 Samples including, 1) DNA sequencing of 16 samples representing two acerola species (*Malpighia glabra*, *M. emarginata*) of which 13 were leaf samples and 3 were berry samples (Table S1), and 2) NMR metabolite fingerprinting of 16 samples representing 10 berry samples of two acerola species (*Malpighia glabra*, *M. emarginata*) and 6 berry samples of cherry representing two species (*Prunus cerasus* L.; *Prunus avium* (L.) L.) (Table S2). Only berry samples were used for NMR metabolite fingerprinting representing the form of acerola product in the supply chain because NMR fingerprints are known to be different among leaves and fruit^{66,67}. It is known that DNA sequences for specific regions such as ITS do not change for different tissues such as leaves and fruit. Therefore, we used mostly leaves in this study for DNA sequencing. However, we did sequence five vouchers (e.g., leaves and berries) from the NMR study including samples for each *Malpighia* species to verify that the berries used for NMR metabolite fingerprinting were the correct species (Table S1). We did not attempt to sequence the cherry samples due to the known challenges of

barcoding cherry (*Prunus* spp.) in the published literature including ITS and other regions^{68–70}. The identification of the herbarium voucher samples was confirmed by qualified taxonomist on staff at the University of Guelph. All samples are housed at the Natural Health Products Research Alliance, University of Guelph.

Genomic DNA extraction Genomic DNA from the samples were extracted using the Nucleospin Plant II kit (Macherey- Nagel GmbH & Co. KG, Düren, Germany) to get high-quality DNA. DNA extractions were conducted using 100 mg of each sample, following the manufacturer's instructions. DNA quantification was carried out with the QubitTM 3.0 Fluorometer (Invitrogen, Carlsbad, CA).

DNA Amplification of ITS The ITS region was used because of its ability to discriminate con generic species^{3,71–73}. PCR was performed using species-specific ITS primer pairs as described by Fazekas et al. (2012)⁷⁴. The reaction mixtures (20 µL) contained 1 U AmpliTaq Gold Polymerase, GeneAmp buffer II (100 mM Tris-HCl pH 8.3 and 500 mM KCl), 2.5 mM MgCl₂, 0.2 mM dNTPs, 0.1 mM of each primer, and 20 ng of template DNA.

DNA sequencing of ITS region The ITS amplified products were sequenced in both directions following the protocols established at the University of Guelph Genomics Facility (www.uoguelph.ca/~genomics). The products from each specimen were purified using Sephadex columns and analyzed on a ABI 3730 sequencer (Applied Biosystems). Bidirectional sequence reads were acquired for all the PCR products. The sequences were assembled using CodonCode Aligner version 11.0.1 (CodonCode Corp) and manually aligned with BioEdit version 7.0.9. Subsequently, the aligned sequences were processed using Clustal W⁷⁵. The genetic distances were calculated using the Kimura2Parameter (K2P) model in Mega11⁷⁶. 18 DNA sequences from GenBank were used to assess the specificity of the ITS region and compare interspecific variation among the two species of *Malpighia* including, *M. glabra*, *M. emarginata*. These 18 sequences were used in a BLAST analysis within GenBank to assess the specificity of these sequences to other *Malpighia* species and that of any potential adulterants such as *Prunus* species.

NMR Sample Preparation Samples for NMR analysis were prepared by weighing 300 mg of finely ground plant tissue (Table S2). To ensure complete homogenization, the plant material was ground in liquid nitrogen using a mortar and pestle, then dissolved in a solvent mixture of 90% regular methanol and 10% deuterated methanol (CD₃OD) for its broad solubility range, which aids in NMR acquisition^{77,78}. Each sample was prepared in triplicate for consistency, sonicated in a water bath at room temperature for 10 minutes to enhance extraction, and centrifuged at 6000 rpm for 5 minutes. Finally, 650 µL of the clear supernatant was transferred into a 5 mm Wilmad® NMR tube for spectral acquisition.

NMR Spectral Acquisition The proton (¹H) NMR spectra were recorded on a Bruker Avance III 400 MHz spectrometer, which features a 5 mm Broadband Inverse (BBI) probe designed for room temperature use. We used a proton NOESY (Nuclear Overhauser Effect Spectroscopy) pulse sequence with pre-saturation to improve the results. This method included dual solvent suppression for both water and methanol, helping to reduce any interference from the solvent signals. The key acquisition parameters

included: **Spectrometer frequency:** 400.3 MHz; **Number of scans:** 64; **Acquisition time:** 2.27 seconds; **Relaxation delay:** 12.73 seconds; **Spectral width:** 8223.68 Hz. The use of dual solvent suppression was utilized to improve the spectral quality by reducing peak overlap from residual solvent signals, enhancing metabolite detection in complex biological matrices⁷⁹. Temperature stabilization was maintained at 300 K throughout data acquisition.

NMR Data Processing and Analysis In this study, NMR spectral data were processed with TopSpin 3.6.3 for consistency and accuracy, using automated phase and baseline corrections and referencing the tetramethylsilane (TMS) peak at 0.00 ppm for calibration. A rectangular binning approach with a bin width of 0.01 ppm was utilized over a chemical shift range from -1 to 12 ppm, while residual solvent peaks corresponding to water (4.75–5.06 ppm) and methanol (3.16–3.45 ppm) were excluded to minimize spectral distortion. Each spectrum was normalized via a scaling method wherein values below the mean intensity were set to zero, and intensities above the mean were scaled to fit within a range of 1 to 100. For statistical analysis, we performed multivariate analyses, including Hierarchical Clustering Analysis (HCA) and Hierarchical Clustering on Principal Components (HCP), based on an Euclidean dissimilarity matrix while employing Ward's clustering method in R⁸⁰. We converted spectral intensities into chemical fingerprints, which allowed us to use hierarchical clustering to distinguish between different botanical species. This NMR-based metabolomics approach provides reliable differentiation and ensures high reproducibility for future analyses.

Results

DNA Extraction and Sequencing. The DNA extraction protocols yielded good quality, high molecular weight genomic DNA from the dried berry and leaf samples of *M. glabra* and *M. emarginata*. The procedure yielded 40–60 ng of DNA per ng/ul of leaf tissue and for a whole dried berries and powders yielded (2 to 4.5 ng/ul). The amplification of the complete ITS region (comprising ITS1, the 5.8S rRNA gene, and ITS2) was successfully achieved using the universal primers ITS5 (forward) and ITS4 (reverse). This process generated an amplicon of approximately 700 bp for all three species investigated.

Direct sequencing of amplicon yielded a ~ 700bp ITS sequence for *M. glabra* and *M. emarginata*. GenBank accession numbers are recorded (Table S1). Searches and BLAST analysis in GenBank indicated that the sequences of *M. emarginata* matched that of those in GenBank and that those sequences for *M. glabra* were novel and added to the GenBank library. Alignment of all the sequences revealed that there was enough interspecific variation to distinguish all the samples of *M. glabra* from *M. emarginata*. Interspecific among the two species was 1% including seven species specific single nucleotide polymorphisms (SNPs) within the 700 base pair nucleotide sequence. Sequence variation indicated that the ITS1, 5.8S rRNA gene and ITS2 regions of each respective species were unique. BLAST analysis revealed that all the acerola species sequences did not match any other potential adulterant species including cherry (*Prunus* spp.).

NMR Fingerprinting Metabolite diversity among the NMR fingerprints was considerable. Hierarchical clustering analysis (HCA) of the ^1H NMR spectral data was used to assess metabolite diversity within a heatmap of metabolite diversity (Figure: 1). The resulting dendrogram clearly distinguishes the samples of *M. glabra* and *M. emarginata* highlighting their distinct metabolite profiles. However, *M. glabra* exhibits closer metabolic resemblance to *Prunus* spp. compared to *M. emarginata*, suggesting a greater risk of potential adulteration. The hierarchical clustering of cherry samples groups respective *Prunus* species from acerola species based on their metabolic composition. Clusters labeled A and D correspond to *M. glabra* and *M. emarginata*, respectively, while clusters B and C include *Prunus* spp. The distinct clustering of *M. emarginata* underscores its unique metabolite profile, which is markedly different from *M. glabra* and *Prunus* species. This separation suggests that while *M. glabra* may share some chemical characteristics with *Prunus*, *M. emarginata* remains phytochemically distinct.

Comparative ^1H NMR Spectral Analysis: *Prunus* spp. vs. *Malpighia* spp. ^1H NMR spectra of *M. emarginata*, *Prunus cerasus*, and *Prunus avium* demonstrate clear differences in metabolite distribution (Figure: S1). Although some spectral overlaps exist, key variations are observed in regions associated with carbohydrates (3.0–5.5 ppm), organic acids (2.0–3.0 ppm), and aromatic compounds (6.0–8.0 ppm). The most notable differences include:

- The prominence of organic acids such as **malic acid** (2.4–3.0 ppm) in *M. emarginata*, which is significantly reduced or absent in *Prunus* spp.
- Higher concentrations of **ascorbic acid** (4.0–4.3 ppm) in Acerola, confirming its known vitamin C content (de Souza et al., 2002).
- The presence of **ferulic acid** (6.5–7.5 ppm) in *M. emarginata*, a unique authentication marker absent in *Prunus* spp.

These spectral differences validate the use of ^1H NMR fingerprinting as a reliable method for species authentication, ensuring that *Malpighia* species can be accurately distinguished from potential adulterants such as cherry (*Prunus* spp.).

Metabolite Assignments and Authentication Markers Elucidation of specific metabolites identifies differences among the key metabolite assignments for *M. glabra* and *M. emarginata*, respectively (Figure: 2 & 3). The assigned peaks correspond to major bioactive compounds that play a role in species differentiation (Figure: 2 & 3). The presence of ferulic acid and amino acids in *M. emarginata* sets it apart from *M. glabra* and *Prunus* spp. (Figure: 2 & 3).

The metabolites identified in *M. glabra* include α -D-glucose (5.2–5.4 ppm), β -D-glucose (4.6–4.8 ppm), fructose (4.0–4.5 ppm), ascorbic acid (4.0–4.3 ppm), and malic acid (2.4–3.0 ppm). In comparison, *M. emarginata* metabolites encompass ferulic acid (6.5–7.5 ppm), notably absent in *Prunus* spp., along with α -glucose (5.2–5.4 ppm), β -glucose (4.6–4.8 ppm), fructose (4.0–4.5 ppm), and malic acid (2.4–3.0 ppm), the latter serving as a distinguishing marker. Additionally, amino acids such as valine, leucine, and isoleucine (0.8–1.5 ppm) are specifically identified in *M. emarginata*.

Discussion

One of the impediments in the acceptance of herbal formulations in the medical and pharmaceutical community is the lack of standardization and quality control. This is due to unacceptable risks to human health and unjustified health claims. It is challenging to develop robust quality control parameters using analytical chemical methods because of the complex nature and inherent variability of the chemical ingredients and molecular constituents used in supplements and natural health products, (World Health Organization 2000). These challenges including that of other conventional pharmacognosy methods such as morphology, microscopy, DNA-based and phytochemical analysis have led to the demand for more robust, fit-for-purpose methods of authentication of botanical species ingredients^{3,81–83}. This is driven by consumer demand that seek natural remedies and nutrition that support health lifestyles without the use of drugs and synthetic health supplements⁸⁴.

There are considerable challenges in authentication of acerola species ingredients from the commercial supply chain. Our experience and collaboration with industry members working directly with procuring *M. glabra* and *M. emarginata* materials indicates that these species cannot be distinguished based on their morphology or histology. Conventional analytical chemistry methods are often providing inconclusive reports. For example, high performance thin layer chromatography (HPTLC) analysis of acerola berries of *M. glabra* and *M. emarginata* often have identical or closely overlapping profiles with similar or highly variable acerolin content depending on product processing used by different suppliers (NHPRA members pers. Comm.). More recently the industry has moved to trading fruit as dried extracts of which there is a lack of standard methods for quality assurance protocols such as reference standards and product species verification. This problem is intensified when considering that acerola is considered a superfruit in high demand due to its high vitamin C content of which is highly perishable in transit and storage^{85,86}. Dehydrating acerola juice using a spray dryer has become a common solution in the supply industry to extend the product's shelf life and subsequent profit margins^{87,88}. The NHP community has repeatedly voice there is a gap in acerola quality assurance methods and need for research on the development of new methods.

This study provides two orthogonal molecular methods for the authentication of acerola species ingredients procured in the industry supply chain. The first method utilized common DNA sequencing. Several studies have used ITS sequences as genetic markers for many botanical species^{3,71–73,89}. The sequence variation of ribosomal RNA gene repeats is believed to rapidly turn over thus providing considerable interspecific variation among congeneric species⁹⁰. Universal PCR primers designed from highly conserved regions flanking the ITS region are available and are relatively small in size (600–700 bp) with high copy number provide relative ease of amplification of the ITS region. In our study we had successful sequencing and found sufficient variation to distinguish the two commercial acerola species. However, the materials we used contained considerable amounts of DNA unlike that of highly processed extracts, which have considerably less DNA presenting challenges for PCR and sequencing^{34,91}. Another consideration for extract materials is the presence of PCR amplification inhibitors such as concentrated

phenolic compounds, acidic polysaccharides and pigments, which can be overcome by choosing appropriate DNA extraction methods that can reduce or eliminate PCR inhibitors³⁵. Further research is needed to develop species specific probes based on unique SNPs for *M. glabra* and *M. emarginata* that could be used within specific QPCR methods. This DNA approach has been successfully used for species specific verification of highly processed supplements in the form of leaf or berry extracts have reported mini sequence-based authentication of several botanical species worked well in botanical extracts^{50,53}. ITS sequences were unique to commercial acerola species and not potential adulterants such as cherry species. However, a fit-for-purpose standard operating aimed at authenticating target species ingredients and preventing adulteration would require full ITS sequencing, which requires considerable resources (e.g., time, cost). If the materials are highly processed extracts, then targeted QPCR would be required including specific QPCR assays for all the potential adulterants of which resource intensive (e.g., time costs).

In this study, ¹H Nuclear Magnetic Resonance (NMR) metabolomic fingerprinting combined with chemometric clustering distinguished highly all samples of two Acerola species, *M. glabra* and *M. emarginata*, from potential adulterants belonging to the *Prunus* genus (*Prunus cerasus* and *Prunus avium*). The results provide a robust metabolic fingerprint that can serve as a scientific authentication tool based on primary and secondary metabolite composition. ¹H NMR fingerprinting, coupled with multivariate chemometric analysis, provides a powerful authentication tool for differentiating botanical species. Unlike chromatographic techniques, which can be manipulated by introducing synthetic analogs, NMR-based authentication relies on the holistic metabolite spectrum, making it more resistant to adulteration⁷⁹. The authentication of commercial acerola is particularly significant due to its high economic value and susceptibility to adulteration with cherry derivatives. The hierarchical clustering of metabolites indicates that while *M. glabra* exhibits some spectral resemblance to *Prunus* spp., *M. emarginata* remains chemically distinct. By integrating metabolite-specific peak ratios, particularly malic acid and ascorbic acid content, a more rigorous authentication framework is established. This approach provides both a species ingredient identification tools while providing verification of specific bioactive metabolites that are of value to health minded consumers. The elucidation of specific bioactive metabolites is particularly valuable for product claims and specific ingredients on product labels.

Conclusion

Two methods provide novel tools for the quality assurance of acerola species procured in the industry supply chain. DNA methods have several limitations for berry extracts and may be cost prohibitive if the goal is to detect species adulteration. NMR fingerprinting can be used for berry extracts and has the benefit of authenticating acerola species ingredients whilst detecting any adulterants in a single analysis. The resulting NMR spectra also provide quantitative estimates of bioactive phytochemicals negating the need for further analytical chemistry methods. This study highlights the application of NMR fingerprinting as a critical quality control measure in the natural health product and food industry.

Declarations

Acknowledgements

We thank Natural health product research alliance (NHPRA), University of Guelph for supporting this study. We would like to thank the Genomics facility, University of Guelph for their help in carrying out experiments in their lab and also for their timely help and assistance whenever needed. Thomas Henry is greatly acknowledged for his help in the wet lab.

Data availability statement

All data supporting the findings of this study are included in the article and supplementary material. Any additional inquiries can be directed to the corresponding author.

Funding source

Funding for this research was provided by the Natural Health Product Research Alliance, University of Guelph, grant number 053312.

Author contributions:

Sub.R. and S.G.N conceived the project and experiments; Sub.R and Sne. R. carried out the DNA molecular work and analysis; V.V. Sne. R. and A.T. carried out the NMR metabolite lab work and analysis; Sub.R. led the manuscript writing, and all other authors contributed to writing, editing, reading and accepting the final version of the manuscript.

Competing Interests: The authors declare no competing interests.

Additional Information:

The authors confirm that Dr Ragupathy is the trained taxonomist who identified all plants. All the vouchers are recorded within Table S1 and the specimens were deposited at the Natural Health Products Research Alliance, the University of Guelph. The plants used in this study are common, commercially available species that do not require any permits/permissions/licenses.

References

1. Laurindo, L. F. et al. Health benefits of acerola (*Malpighia* spp) and its by-products: A comprehensive review of nutrient-rich composition, pharmacological potential, and industrial applications. *Food Bioscience* vol. 62 Preprint at (2024). <https://doi.org/10.1016/j.fbio.2024.105422>
2. Uma Maheswari, M., Arumugam, T. & Lincy Sona, C. Fruits and vegetables as Superfoods: Scope and demand. ~ 119 ~ *Pharma Innov. J.* **10**, 119–129 (2021).

3. Li, D. Z. et al. Comparative analysis of a large dataset indicates that internal transcribed spacer (ITS) should be incorporated into the core barcode for seed plants. *Proc. Natl. Acad. Sci. U S A.* **108**, 19641–19646 (2011).
4. Fernández-Ríos, A. et al. A critical review of superfoods from a holistic nutritional and environmental approach. *Journal of Cleaner Production* vol. 379 Preprint at (2022).
<https://doi.org/10.1016/j.jclepro.2022.134491>
5. Prakash, A. & Baskaran, R. Acerola, an untapped functional superfruit: a review on latest frontiers. *Journal of Food Science and Technology* vol. 55 3373–3384 Preprint at (2018).
<https://doi.org/10.1007/s13197-018-3309-5>
6. Mezadri, T., Villaño, D., Fernández-Pachón, M. S., García-Parrilla, M. C. & Troncoso, A. M. Antioxidant compounds and antioxidant activity in acerola (*Malpighia emarginata* DC.) fruits and derivatives. *J. Food Compos. Anal.* **21**, 282–290 (2008).
7. Righetto, A. M., Netto, F. M. & Carraro, F. Chemical composition and antioxidant activity of juices from mature and immature acerola (*Malpighia emarginata* DC). *Food Sci. Technol. Int.* **11**, 315–321 (2005).
8. McGuffin, M., Tucker, A. & L., A. Y. *J. T. K. Herbs of Commerce* (American Herbal Products Association, 2000).
9. Duke, J. A. *CRC Handbook of Alternative Cash Crops* (CRC, 1993).
10. Rezende, Y. R. R. S., Nogueira, J. P. & Narain, N. Comparison and optimization of conventional and ultrasound assisted extraction for bioactive compounds and antioxidant activity from agro-industrial acerola (*Malpighia emarginata* DC) residue. *LWT* **85**, 158–169 (2017).
11. De Aparecida, S. et al. Antioxidant activity, ascorbic acid and total phenol of exotic fruits occurring in Brazil. *Int. J. Food Sci. Nutr.* **60**, 439–448 (2009).
12. Singh, D. R. R. West Indian Cherry ñ A lesser known fruit for nutritional security. in (2006).
13. Belwal, T. et al. Phytopharmacology of Acerola (*Malpighia* spp.) and its potential as functional food. *Trends in Food Science and Technology* vol. 74 99–106 Preprint at (2018).
<https://doi.org/10.1016/j.tifs.2018.01.014>
14. Laurindo, L. F. et al. Health benefits of acerola (*Malpighia* spp) and its by-products: A comprehensive review of nutrient-rich composition, pharmacological potential, and industrial applications. *Food Bioscience* vol. 62 Preprint at (2024). <https://doi.org/10.1016/j.fbio.2024.105422>
15. Paz, M. et al. Brazilian fruit pulps as functional foods and additives: Evaluation of bioactive compounds. *Food Chem.* **172**, 462–468 (2015).
16. do Rufino, M. Acerola and cashew apple as sources of antioxidants and dietary fibre. *Int. J. Food Sci. Technol.* **45**, 2227–2233 (2010).
17. Motohashi, N. et al. Biological activity of Barbados cherry (Acerola fruits, fruit of *Malpighia emarginata* DC) extracts and fractions. *Phytotherapy Research* vol. 18 212–223 Preprint at (2004).
<https://doi.org/10.1002/ptr.1426>

18. Leffa, D. D. et al. Corrective effects of acerola (*Malpighia emarginata* DC.) juice intake on biochemical and genotoxic parameters in mice fed on a high-fat diet. *Mutat. Res. - Fundamental Mol. Mech. Mutagen.* **770**, 144–152 (2014).
19. Horta, R. N. et al. Protective effects of acerola juice on genotoxicity induced by iron in vivo. *Genet. Mol. Biol.* **39**, 122–128 (2016).
20. Düsman, E., Almeida, I. V., Tonin, L. T. D. & Vicentini, V. E. P. In vivo antimutagenic effects of the barbados cherry fruit (*Malpighia glabra* linnaeus) in a chromosomal aberration assay. *Genet. Mol. Res.* **15**, (2016).
21. Da Silva Nunes, R. et al. Genotoxic and antigenotoxic activity of acerola (*malpighia glabra* L.) extract in relation to the geographic origin. *Phytother. Res.* **27**, 1495–1501 (2013).
22. Düsman, E. et al. Radioprotective Effect of the Barbados Cherry (*Malpighia Glabra* L.) against Radiopharmaceutical Iodine-131 in Wistar Rats in Vivo. (2014).
<http://www.biomedcentral.com/1472-6882/14/41>
23. da Silva Nunes, R. et al. Antigenotoxicity and Antioxidant Activity of Acerola Fruit (*Malpighia glabra* L.) at Two Stages of Ripeness. *Plant Foods Hum. Nutr.* **66**, 129–135 (2011).
24. Hanamura, T., Mayama, C., Aoki, H., Hirayama, Y. & Shimizu, M. Antihyperglycemic effect of polyphenols from acerola (*Malpighia emarginata* DC.) fruit. *Biosci. Biotechnol. Biochem.* **70**, 1813–1820 (2006).
25. El-Hawary, S. S., El-Fitany, R. A., Mousa, O. M., Salama, A. A. A. & El Gedaily, R. A. Metabolic profiling and in vivo hepatoprotective activity of *Malpighia glabra* L. leaves. *J. Food Biochem.* **45**, (2021).
26. Costa, A. et al. Clinical, biometric and ultrasound assessment of the effects of daily use of a nutraceutical composed of lycopene, acerola extract, grape seed extract and Biomarine Complex in photoaged human skin. *Bras. Dermatol.* **87**, 52–61 (2012).
27. Delva, L. & Goodrich-Schneider, R. Antioxidant activity and antimicrobial properties of phenolic extracts from acerola (*Malpighia emarginata* DC) fruit. *Int. J. Food Sci. Technol.* **48**, 1048–1056 (2013).
28. de Miskinis, A. S., Nascimento, R. & Colussi, R. L. Á. Bioactive compounds from acerola pomace: A review. *Food Chemistry* vol. 404 Preprint at (2023).
<https://doi.org/10.1016/j.foodchem.2022.134613>
29. Vilvert, J. C., de Freitas, S. T., Veloso, C. M. & Amaral, C. L. F. Genetic Diversity on Acerola Quality: A Systematic Review. *Brazilian Archives of Biology and Technology* vol. 67 Preprint at (2024).
<https://doi.org/10.1590/1678-4324-2024220490>
30. de Assis, S. A., Lima, D. C. & de Faria Oliveira, O. M. M. Activity of pectinmethylesterase, pectin content and vitamin C in acerola fruit at various stages of fruit development. *Food Chem.* **74**, 133–137 (2001).
31. dos Santos, C. P. et al. Transcriptome analysis of acerola fruit ripening: insights into ascorbate, ethylene, respiration, and softening metabolisms. *Plant. Mol. Biol.* **101**, 269–296 (2019).

32. Fernandes, F. A. N., Santos, V. O., Gomes, W. F. & Rodrigues, S. Application of High-Intensity Ultrasound on Acerola (*Malpighia emarginata*) Juice Supplemented with Fructooligosaccharides and Its Effects on Vitamins, Phenolics, Carotenoids, and Antioxidant Capacity. *Processes* 11, (2023).
33. Assis, S. A., De, Fernandes, F. P., Martins, A. B. G., Oliveira, O. M. M. D. F. & Acerola Importance, culture conditions, production and biochemical aspects. *Fruits* vol. 63 93–101 Preprint at (2008). <https://doi.org/10.1051/fruits:2007051>
34. Ragupathy, S. et al. Exploring DNA quantity and quality from raw materials to botanical extracts. *Heliyon* 5, (2019).
35. Yip, P. Y., Chau, C. F., Mak, C. Y. & Kwan, H. S. DNA methods for identification of Chinese medicinal materials. *Chinese Medicine* vol. 2 Preprint at (2007). <https://doi.org/10.1186/1749-8546-2-9>
36. Newmaster, S., Ragupathy, S. & Kress, W. J. Authentication of Medicinal Plant Components in North America's NHP Industry Using Molecular Diagnostic Tools. in *Medicinal and Aromatic Plants of North America* (ed. Máthé, Á.) 325–339 Springer International Publishing, Cham, (2020). [10.1007/978-3-030-44930-8_13](https://doi.org/10.1007/978-3-030-44930-8_13)
37. Kesanakurti, P. et al. Development of hydrolysis probe-based qpcr assays for panax ginseng and panax quinquefolius for detection of adulteration in ginseng herbal products. *Foods* 10, (2021).
38. Yu, J. et al. Progress in the use of DNA barcodes in the identification and classification of medicinal plants. *Ecotoxicol. Environ. Saf.* 208, (2021).
39. Raclariu-Manolică, A. C. & de Boer, H. J. Chapter 8 - DNA barcoding and metabarcoding for quality control of botanicals and derived herbal products. in *Evidence-Based Validation of Herbal Medicine (Second Edition)* (ed. Mukherjee, P. K.) 223–238 (Elsevier, 2022). <https://doi.org/10.1016/B978-0-323-85542-6.00004-4>
40. Travadi, T. et al. Advancing Herbal Product Authentication: A Comprehensive Review Of DNA-Based Approach For Quality Control And Safety Assurance. *Educational Adm. Theory Practices* doi:10.53555/kuey.v30i6(s).5308. (2024).
41. Lanubile, A., Stagnati, L., Marocco, A. & Busconi, M. DNA-based techniques to check quality and authenticity of food, feed and medicinal products of plant origin: A review. *Trends in Food Science and Technology* vol. 149 Preprint at (2024). <https://doi.org/10.1016/j.tifs.2024.104568>
42. Muyumba, N. W., Mutombo, S. C., Sheridan, H., Nachtergaeel, A. & Duez, P. Quality control of herbal drugs and preparations: The methods of analysis, their relevance and applications. *Talanta Open.* 4, (2021).
43. Ichim, M. C. & Booker, A. Chemical Authentication of Botanical Ingredients: A Review of Commercial Herbal Products. *Frontiers in Pharmacology* vol. 12 Preprint at (2021). <https://doi.org/10.3389/fphar.2021.666850>
44. Abraham, E. J. & Kellogg, J. J. Chemometric-Guided Approaches for Profiling and Authenticating Botanical Materials. *Frontiers in Nutrition* vol. 8 Preprint at (2021). <https://doi.org/10.3389/fnut.2021.780228>

45. Klein-Junior, L. C. et al. Quality Control of Herbal Medicines: From Traditional Techniques to State-of-the-art Approaches. *Planta Medica* vol. 87 964–988 Preprint at (2021). <https://doi.org/10.1055/a-1529-8339>
46. Salo, H. M. et al. Authentication of berries and berry-based food products. *Compr. Rev. Food Sci. Food Saf.* **20**, 5197–5225 (2021).
47. García-Pérez, P., Becchi, P. P., Zhang, L., Rocchetti, G. & Lucini, L. Metabolomics and chemometrics: The next-generation analytical toolkit for the evaluation of food quality and authenticity. *Trends in Food Science and Technology* vol. 147 Preprint at (2024). <https://doi.org/10.1016/j.tifs.2024.104481>
48. Ichim, M. C., Scotti, F. & Booker, A. Quality evaluation of commercial herbal products using chemical methods. *Critical Reviews in Food Science and Nutrition* vol. 64 4219–4239 Preprint at (2024). <https://doi.org/10.1080/10408398.2022.2140120>
49. Urumarudappa, S. K. J. et al. DNA barcoding and NMR spectroscopy-based assessment of species adulteration in the raw herbal trade of *Saraca asoca* (Roxb.) Willd, an important medicinal plant. *Int. J. Legal Med.* **130**, 1457–1470 (2016).
50. Kesanakurti, P., Thirugnanasambandam, A., Ragupathy, S. & Newmaster, S. G. Genome skimming and NMR chemical fingerprinting provide quality assurance biotechnology to validate Sarsaparilla identity and purity. *Sci. Rep.* **10**, (2020).
51. Shirahata, T. et al. Metabolic fingerprinting for discrimination of DNA-authenticated *Atractylodes* plants using ¹H NMR spectroscopy. *J. Nat. Med.* **75**, 475–488 (2021).
52. Alberts, P. S. F. & Meyer, J. J. M. Integrating chemotaxonomic-based metabolomics data with DNA barcoding for plant identification: A case study on south-east African Erythroxylaceae species. *South. Afr. J. Bot.* **146**, 174–186 (2022).
53. Ragupathy, S., Thirugnanasambandam, A., Vinayagam, V. & Newmaster, S. G. Nuclear Magnetic Resonance Fingerprints and Mini DNA Markers for the Authentication of Cinnamon Species Ingredients Used in Food and Natural Health Products. *Plants* **13**, (2024).
54. Ragupathy, S. et al. Flower Species Ingredient Verification Using Orthogonal Molecular Methods. *Foods* **13**, (2024).
55. Wang, X. et al. Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: Systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ (Online)* **349**, (2014).
56. Martinez-Farina, C. F. et al. Chemical barcoding: A nuclear-magnetic-resonance-based approach to ensure the quality and safety of natural ingredients. *J. Agric. Food Chem.* **67**, 7765–7774 (2019).
57. Grazina, L., Amaral, J. S. & Mafra, I. Botanical origin authentication of dietary supplements by DNA-based approaches. *Comprehensive Reviews in Food Science and Food Safety* vol. 19 1080–1109 Preprint at (2020). <https://doi.org/10.1111/1541-4337.12551>
58. Osman, A. et al. Quality Consistency of Herbal Products: Chemical Evaluation. in *Progress in the Chemistry of Organic Natural Products 122: Botanical Dietary Supplements and Herbal Medicines*

- (ed Kinghorn, A. D.) 163–219 (Springer Nature Switzerland, Cham, doi:10.1007/978-3-031-26768-0_2. (2023).
59. Bharti, S. K. & Roy, R. Quantitative ¹H NMR spectroscopy. *TrAC - Trends in Analytical Chemistry* vol. 35 5–26 Preprint at (2012). <https://doi.org/10.1016/j.trac.2012.02.007>
60. Sobolev, A. P., Ingallina, C., Spano, M., Di Matteo, G. & Mannina, L. NMR-Based Approaches in the Study of Foods. *Molecules* vol. 27 Preprint at (2022). <https://doi.org/10.3390/molecules27227906>
61. Esslinger, S., Riedl, J. & Fauhl-Hassek, C. Potential and limitations of non-targeted fingerprinting for authentication of food in official control. *Food Res. Int.* **60**, 189–204 (2014).
62. Lolli, V. & Caligiani, A. How nuclear magnetic resonance contributes to food authentication: current trends and perspectives. *Current Opinion in Food Science* vol. 58 Preprint at (2024). <https://doi.org/10.1016/j.cofs.2024.101200>
63. Wishart, D. S. Metabolomics: applications to food science and nutrition research. *Trends in Food Science and Technology* vol. 19 482–493 Preprint at (2008). <https://doi.org/10.1016/j.tifs.2008.03.003>
64. Spiteri, M. et al. Fast and global authenticity screening of honey using ¹H-NMR profiling. *Food Chem.* **189**, 60–66 (2015).
65. Giraudeau, P. Quantitative NMR spectroscopy of complex mixtures. *Chem. Commun.* **59**, 6627–6642 (2023).
66. Holmes, E., Tang, H., Wang, Y. & Seger, C. The Assessment of Plant Metabolite Profiles by NMR-Based Methodologies. (2006). 10.1055/s-2006-946682'Published946682'946682'Published
67. Mahmud, I., Chowdhury, K. & Boroujerdi, A. PTC&B Tissue-Specific Metabolic Profile Study of *Moringa Oleifera* L. Using Nuclear Magnetic Resonance Spectroscopy. *Plant Tissue Cult. & Biotech* vol. 24 (2014).
68. Rosario, L. H. et al. DNA barcoding of the solanaceae family in puerto rico including endangered and endemic species. *J. Am. Soc. Hortic. Sci.* **144**, 363–374 (2019).
69. Sayed, H. A., Mostafa, S., Haggag, I. M. & Hassan, N. A. DNA Barcoding of *Prunus* Species Collection Conserved in the National Gene Bank of Egypt. *Mol. Biotechnol.* **65**, 410–418 (2023).
70. Kress, W. J., Erickson, D. L. A. & Two-Locus Global DNA Barcode for Land Plants: The Coding *rbcl* Gene Complements the Non-Coding *trnH-psbA* Spacer Region. *PLoS One* **2**, (2007).
71. Baldwin, B. G. et al. The *its* Region of Nuclear Ribosomal DNA: A Valuable Source of Evidence on Angiosperm Phylogeny. *Ann. Mo. Bot. Gard.* **82**, 247 (1995).
72. Kress, W. J., Wurdack, K. J., Zimmer, E. A., Weigt, L. A. & Janzen, D. H. Use of DNA Barcodes to Identify Flowering Plants. www.pnas.org/cgi/doi/10.1073/pnas.0503123102 (2005).
73. Cheng, T. et al. Barcoding the kingdom Plantae: New PCR primers for ITS regions of plants with improved universality and specificity. *Mol. Ecol. Resour.* **16**, 138–149 (2016).
74. Fazekas, A. J., Kuzmina, M. L., Newmaster, S. G. & Hollingsworth, P. M. DNA Barcoding Methods for Land Plants. in 223–252 (2012). 10.1007/978-1-61779-591-6_11

75. Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673–4680 (1994).
76. Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol. Biol. Evol.* **38**, 3022–3027 (2021).
77. Beckonert, O. et al. Metabolic profiling, metabolomic and metabonomic procedures for NMR spectroscopy of urine, plasma, serum and tissue extracts. *Nat. Protoc.* **2**, 2692–2703 (2007).
78. Nicholson, J. K., Foxall, P. J. D., Spraul, M., Farrant, R. D. & Lindon J. C. 750 MHz ¹H and ¹H-¹³C NMR Spectroscopy of Human Blood Plasma. *Anal Chem* **67**, 793–811 (1995).
79. Emwas, A. H. M. The Strengths and Weaknesses of NMR Spectroscopy and Mass Spectrometry with Particular Focus on Metabolomics Research. in 161–193 (2015). 10.1007/978-1-4939-2377-9_13
80. Lê, S., Josse, J. & Husson, F. FactoMineR: An R Package for Multivariate Analysis. *J. Stat. Softw.* **25**, (2008).
81. Smillie, T. J. & Khan, I. A. A comprehensive approach to identifying and authenticating botanical products. *Clinical Pharmacology and Therapeutics* vol. 87 175–186 Preprint at (2010). <https://doi.org/10.1038/clpt.2009.287>
82. Mukherjee, P. K., Bahadur, S., Chaudhary, S. K., Kar, A. & Mukherjee, K. *Chapter 1 - Quality Related Safety Issue-Evidence-Based Validation of Herbal Medicine Farm to Pharmain* 1–28 (Elsevier Inc, 2015). 10.1016/B978-0-12-800874-4.00001-5
83. Govindaraghavan, S. & Sucher, N. J. Quality assessment of medicinal herbs and their extracts: Criteria and prerequisites for consistent safety and efficacy of herbal medicines. *Epilepsy Behav.* **52**, 363–371 (2015).
84. Dubey, N. K., Kumar, R. & Tripathi, P. Global promotion of herbal medicine: India's opportunity. *Curr. Sci.* **86**, (2003).
85. de Medeiros, F. G. M., Pereira, G. B. C., Pedrini, M. R. S., Hoskin, R. T. & Nunes, A. O. Evaluation of the environmental performance of the production of polyphenol-rich fruit powders: A case study on acerola. *J. Food Eng.* **372**, (2024).
86. Gomes, B. T. et al. Acerola byproducts microencapsulated by spray and freeze-drying: The effect of carrier agent and drying method on the production of bioactive powder. *Int. J. Food Eng.* **20**, 347–356 (2024).
87. Fonseca, M. T. et al. Improving the stability of spray-dried probiotic acerola juice: A study on hydrocolloids' efficacy and process variables. *Food Bioprod. Process.* **147**, 209–218 (2024).
88. Coelho, B. E. S. et al. Production and characterization of powdered acerola juice obtained by atomization. *Acta Scientiarum - Technol.* **47**, (2025).
89. Walsh, K. B. & Ragupathy, S. Mycorrhizal colonisation of three hybrid papayas (*Carica papaya*) under mulched and bare ground conditions. *Aust J. Exp. Agric.* **47**, 81 (2007).

90. Dubouzet, J. G. & Shinoda, K. Phylogenetic analysis of the internal transcribed spacer region of Japanese *Lilium* species. *Theor. Appl. Genet.* **98**, 954–960 (1999).

91. Faller, A. C. et al. DNA quality and quantity analysis of *camellia sinensis* through processing from fresh leaves to a green tea extract. *J. AOAC Int.* **102**, 1798–1807 (2019).

Figures




Figure 1

Hierarchically clustering of NMR fingerprints.

¹H NMR Spectrum of *Malpighia emarginata* - NHPRA230722-Malp_emar




Figure 2

Bioactive molecule elucidation of *M. glabra*.

¹H NMR Spectrum of *Malpighia glabra* - NHPRA221212-Malp_glab




Figure 3

Bioactive molecule elucidation of *M. emarginata*.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplement.docx](#)