

Comprehensive analysis of WUSCHEL-related Homeobox family in *Chosenia arbutifolia* (Pall.) A. Skv.: Characterization, Structure, evolutionary insights, and Expression Profiles

Pu Wang

wangpu@njfu.edu.cn

Jiangsu Academy of Forestry

Jiwei zheng

Jiangsu Academy of Forestry

Zhongyi jiao

Jiangsu Academy of Forestry

Ruifang huang

Jiangsu Academy of Forestry

Weiwei wang

Jiangsu Academy of Forestry

Hongling wang

Jiangsu Academy of Forestry

Dezong Sui

Jiangsu Academy of Forestry

Xudong He

Jiangsu Academy of Forestry

Research Article

Keywords: WOX, *Chosenia arbutifolia*, Expression Profiles, evolutionary

Posted Date: May 5th, 2026

DOI: <https://doi.org/10.21203/rs.3.rs-9273330/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Comprehensive analysis of WUSCHEL-related Homeobox family in *Chosenia arbutifolia* (Pall.) A. Skv.: Characterization, Structure, evolutionary insights, and Expression Profiles

Pu Wang^{123a}, Jiwei zheng^{123a}, Zhongyi jiao¹²³, Ruifang huang¹²³, Weiwei wang¹²³, Hongling wang¹²³, Dezong Sui¹²³, Xudong He^{123*}

¹Jiangsu Academy of Forestry, Nanjing, 211153, China

2. Key Laboratory of National Forestry and Grassland Administration on Willow, Jiangsu Academy of Forestry, Nanjing 211153, China;

3. Willow Nursery of the Jiangsu Provincial Platform for Conservation and Utilization of Agricultural Germplasm, Jiangsu Academy of Forestry, Nanjing 211153, China

^a These authors contributed equally to this work.

* Correspondence: hxd_519@163.com;

Abstract

WUSCHEL-related Homeobox (WOX) is a key transcription factor regulating plant meristems, and plays an important role in plant growth, development and stress resistance regulation. To analyze the molecular characteristics and functional potential of the *CaWOX* gene family in *Chosenia arbutifolia* (Pall.) A. Skv, genome-wide identification and bioinformatics analysis were performed in poplar in this study, and 14 *CaWOX* genes were identified. The 14 *CaWOX* genes were distributed on 11 chromosomes with 8 tandem duplications. Phylogenetic analysis divided the 14 *CaWOX* proteins into 4 conserved subgroups, and members in the same subgroup shared highly conserved motifs and gene structures. The *CaWOX* family was highly evolutionarily conserved among closely related species in Salicaceae, but showed weak collinearity with *Arabidopsis thaliana*; cis-elements related to meristem development, hormone response and abiotic stress were enriched in the promoter regions. Tissue development analysis revealed that *CaWOX3/4/5/9/11* participated in leaf development and maintenance of physiological functions in mature leaves;

CaWOX1/6/10/14 regulated root development; and *CaWOX12* mediated secondary growth of stems in *C. arbutifolia*. This study clarified the evolutionary pattern and functional differentiation of the *CaWOX* family in *C. arbutifolia*, providing a key theoretical basis and gene resources for subsequent functional verification of this gene family, as well as growth regulation, wood property improvement and stress-resistant molecular breeding of Salicaceae trees.

Keywords:

WOX, *Chosenia arbutifolia*, Expression Profiles, evolutionary

Introduction

The WOX (WUSCHEL-related Homeobox) gene family encodes plant-specific transcription factors belonging to the Homeobox (HB) superfamily[1]. As a unique class of transcription factors in plants, WOX proteins contain a conserved Homeodomain (HD) consisting of 60-66 amino acid residues[2, 3]. This domain functions in binding specific DNA sequences and folds spatially into a three-dimensional structure comprising three α -helices, known as the helix-loop-helix-turn-helix (HLHTH) motif. Beyond the HD, members of the WUS clade possess a unique WUS-box motif in the carboxy-terminal region, with the consensus sequence T-L-X-L-F-P-X-X (specifically TLELFPLH)[4]. This WUS-box is crucial for WUS function, regulating the stem cell population in the Shoot Apical Meristem (SAM) and controlling floral development. Notably, members of the WUS clade have a dipeptide motif at the start of the WUS-box, whereas WOX proteins from other clades show variations at these positions[5, 6]. Some WOX family members also contain an EAR-like (ERF-associated amphiphilic repression) domain and the repression-associated WUS-box domain (with conserved sequences LxLxLx and TLxLFP, respectively) at their C-termini. Recent research highlights that various of the *WOX* family play significant roles at different stages of plant development[7, 8].

In *Arabidopsis thaliana*, the *WOX* family comprises 15 members, while in rice, there are 12 members. Based on the conservation of protein sequences, the *WOX* family is phylogenetically divided into three clades: the Modern/WUS clade (including WUS

and *AtWOX1-7*), the Intermediate clade (including *AtWOX8*, *AtWOX9*, *AtWOX11*, and *AtWOX12*), and the Ancient clade (including *AtWOX10*, *AtWOX13*, and *AtWOX14*). For instance, *AtWUS* and *AtWOX5* are expressed in the stem and root apical meristems, respectively, maintaining meristem cell activity and exhibiting functional redundancy in root stem cell regulation[9]. *AtWOX4* and *AtWOX14* co-regulate vascular cambium differentiation, ensuring vascular bundle and stem cell maintenance, while *AtWOX6* inhibits primordium differentiation. *AtWOX11* and *AtWOX12* are induced in the root cambium and promote adventitious root formation[10].

The *WOX* gene family is pivotal throughout plant development. *WOX1* has a conserved function in promoting lateral organ growth in species like pea, *A. thaliana*, cucumber, and rice[11]. In *A. thaliana*, *WOX2* and *WOX8* exhibit functional redundancy in regulating apical embryo patterning[12]. *WOX3* is key for cell proliferation during lateral organ growth in *Medicago truncatula* and *A. thaliana* [13, 14]. Studies show that *WOX4* role in promoting vascular stem cell proliferation/maintenance is conserved across *A. thaliana*, rice, and poplar[15]. *WOX5* is expressed in the Quiescent Center and involved in maintaining the root stem cell niche[16-18]. Silencing of *WOX6* delays primordium differentiation and maturation, and regulates ovule patterning in *A. thaliana*. In rice, *WOX6* is auxin-responsive and important for linking shoot gravitropism with tiller angle control. In *A. thaliana*, *WOX7* suppresses lateral root formation in a sugar-dependent manner[19], *WOX9* is essential for meristem growth and development and positively regulates *WUS* expression, and *WOX11* plays a key role in shoot development[20-22]. Furthermore, *WOX11* and its homolog *WOX12* are involved in adventitious root formation in rice and *Liriodendron*[23-25]. Research also indicates that *A. thaliana WOX13* promotes fruit development and coordinates patterning, while *WOX14* regulates vascular tissue proliferation[26].

Given the critical roles of the *WOX* gene family in plant growth, development, and developmental plasticity in response to the environment, research on this family has been conducted in multiple species. However, it has not yet been reported in *Chosenia arbutifolia* (Pall.) A. Skv. *C. arbutifolia*, a member of the Salicaceae family, possesses pendulous male catkins, and both male and female flowers lack nectaries, resembling

features of the genus polar. This makes it scientifically valuable for studying Salicaceae phylogeny[27]. Characterized by fast growth, an elegant form, and strong ecological benefits, it is often found along riverbanks and streams, serving as an excellent species for bank protection.

Considering the impact of plant morphogenesis on productivity and the influence of *WOX* family members on embryogenesis, organ formation, and environmentally responsive plasticity, identifying key *WOX* genes affecting various meristem activities in *C. arbutifolia* could provide candidate target genes for molecular breeding aimed at trait improvement or for the molecular screening of high-yield variants. As the *WOX* gene family plays vital roles in processes like plant cell development and tissue regeneration, this study will employ bioinformatics methods to identify and systematically analyze the *WOX* gene family in *C. arbutifolia*. The aim is to provide theoretical support for subsequent genetic transformation and regeneration studies in this species.

Material and methods

Genome-wide Identification of *CaWOX* Genes

The genomic files, GFF3 annotation file, and CDS sequence file for *C. arbutifolia* were downloaded from the NCBI database[27]. Protein sequences of the *A. thaliana* *WOX* gene family were obtained from the TAIR database. A blastp search was performed against all protein sequences from the *C. arbutifolia* genome, and genes with an e-value < 0.001 were selected for preliminary identification. Candidate genes were further manually analyzed and confirmed to contain conserved domains using the CDD database. The chromosomal locations of the identified *WOX* genes were retrieved from the GFF3 file using their gene IDs. The physicochemical properties of the proteins, including relative molecular weight, theoretical isoelectric point (pI), instability index, and grand average of hydropathicity (GRAVY), were analyzed using the ExPASy online tool. Subcellular localization was predicted using Cell-PLoc 2.0.

Characterization, Chromosomal Distribution, and Phylogenetic Analysis of *CaWOX* Genes

The protein sequences encoded by *WOX* genes from *C. arbutifolia*, *Salix suchowensis*, *A. thaliana*, and *Populus trichocarpa* were downloaded from the Phytozome database. Multiple sequence alignment of these *WOX* protein sequences was performed using ClustalW [28] within the MEGA11 software[29]. A phylogenetic tree was subsequently constructed using the Neighbor-Joining (NJ) method with 1000 bootstrap replicates based on the alignment results. The tree was visualized and edited using Evolview.

Gene Structure and Conserved Motif Analysis of *CaWOX* Genes

Gene structure information (exon-intron organization) for the *CaWOX* genes was extracted from the *C. arbutifolia* genome GFF3 file based on gene IDs and visualized. Protein sequences of *CaWOX* members were retrieved using their protein IDs. Conserved motifs were analyzed using the MEME software[30] suite with parameters set to identify 10 motifs, a maximum motif width of 50, and a minimum motif width of 6. The identified motifs were annotated using meme-v4.12.0.

Analysis of Cis-acting Elements and Collinearity Relationships of *CaWOX* Genes

The promoter regions (2,000 bp upstream of the transcription start site) of *CaWOX* genes were extracted. Cis-acting regulatory elements within these promoter sequences were predicted and analyzed using the PlantCARE database and visualized with the GSDS online tool. Interspecific collinearity analysis was conducted among four species (*A. thaliana*, *P. trichocarpa*, *C. arbutifolia*, and *S. suchowensis*) using McscanX. Tandemly duplicated gene pairs within the *CaWOX* family were identified based on screening criteria. The Ka/Ks ratio for each identified tandem duplicate gene pair was calculated using the Ka/Ks Calculator function in TBtools software[31, 32]. A circos plot was generated to illustrate the tandem duplication events[33].

Plant Growth Conditions

Branches were collected from superior individuals of *Chosenia arbutifolia* (Peter Simon Pallas) A. Skvortsov growing in Manjiang Town, Fusong County, Jilin Province, China (41°47'10.55" N, 127°55'56.13" E). The specimen materials of *Chosenia arbutifolia* were stored in the Chinese virtual herbarium (<https://www.cvh.ac.cn/index.php>). The specimen number is BJFC00076636 and the identifier is Zhixiang Zhang. The branches were stored and hydroponically cultured in the laboratory of Jiangsu Academy of Forestry until leaves emerged. *C. arbutifolia* were grown in a greenhouse at the Jiangsu Academy of Forestry. The growth conditions were maintained at 23°C, 74% relative humidity, with a photoperiod of 16 hours light / 8 hours dark. After approximately one month of growth, different tissues were harvested. All collected samples were immediately frozen in liquid nitrogen and stored at -80°C until use.

Total RNA Extraction and Quantitative Real-Time PCR (qRT-PCR)

Analysis

Total RNA was extracted from samples using an RNA extraction kit (TIANGEN, Beijing). cDNA was synthesized from the RNA using the Evo M-MLV Reverse Transcription Kit (Accurate Biology, China). Specific primers for the *CaWOX* gene family were designed using Primer5, validated, and the optimal primer pairs were selected for subsequent use. qRT-PCR was performed using SYBR Green Premix Pro Taq HS (Accurate Biology) on a qPCR instrument with the following program: 95°C for 10 min, followed by 40 cycles of 95°C for 30 s, 60°C for 30 s, and 72°C for 30 s. The reference gene used was PtActin (Accession number: XM-006370951.1). Relative gene expression levels were calculated using the $2^{-\Delta\Delta CT}$ method. Three independent biological replicates were performed for each treatment, with three technical replicates for each biological sample.

Results

Identification and Analysis of the *CaWOX* Gene Family

Based on BLASTP search results and CDD domain analysis, a total of 14 *CaWOX* genes were identified in the *C. arbutifolia* genome (Table 1). Following the *WOX* gene nomenclature rules established in *Arabidopsis thaliana*, these 14 genes were designated *CaWOX1* to *CaWOX14*. The 14 *CaWOX* genes are distributed across 11 chromosomes: Chromosomes 4, 9, 10, 11, 12, 13, 14, 15, and 19 each contain 1 gene; Chromosome 2 contains 2 genes; Chromosome 5 contains the highest number with 3 genes. The CDS lengths of the 14 *CaWOX* genes range from 642 bp (*CaWOX2*) to 1221 bp (*CaWOX9*), encoding proteins with amino acid lengths between 213 and 406 aa. The relative molecular weights of the proteins range from 24,481.4 Da to 45,121.3 Da. Among the 14 *CaWOX* proteins, 8 have theoretical isoelectric points (pI) between 4.87 and 6.91, classifying them as acidic proteins; 6 have theoretical pI values between 7.66 and 9.99, classifying them as basic proteins. Subcellular localization prediction indicates that all 14 *CaWOX* proteins are localized to the Nucleus.

Table1 *CaWOX* Gene Family

GN	GID	CID	C	PL	CDS	MW	PI	NE	SL
CaWOX1	Salix001816	Salix001816.1	Chr02	215	648	24600.6	4.87	3	Nucleus
CaWOX2	Salix002763	Salix002763.1	Chr02	213	642	24481.4	9.7	3	Nucleus
CaWOX3	Salix006034	Salix006034.1	Chr04	391	1176	43212.9	7.86	3	Nucleus
CaWOX4	Salix008073	Salix008073.1	Chr05	250	753	28278.2	5.59	3	Nucleus
CaWOX5	Salix008176	Salix008176.1	Chr05	266	801	30264.7	6.29	3	Nucleus
CaWOX6	Salix009306	Salix009306.1	Chr05	217	654	24731.7	5.66	3	Nucleus
CaWOX7	Salix014938	Salix014938.1	Chr09	245	738	27878	7.66	2	Nucleus
CaWOX8	Salix017014	Salix017014.1	Chr10	309	930	34793.8	9.99	4	Nucleus
CaWOX9	Salix018617	Salix018617.1	Chr11	406	1221	45121.3	7.86	4	Nucleus
CaWOX10	Salix019842	Salix019842.1	Chr12	365	1098	41117.1	6.52	4	Nucleus
CaWOX11	Salix021133	Salix021133.1	Chr13	255	768	28051.9	5.95	2	Nucleus
CaWOX12	Salix021874	Salix021874.1	Chr14	217	654	24976.8	8.48	3	Nucleus
CaWOX13	Salix023427	Salix023427.1	Chr15	383	1152	42941.1	6.91	4	Nucleus
CaWOX14	Salix029936	Salix029936.1	Chr19	235	708	25850.5	5.74	2	Nucleus

NOTE: GN:Gene Name. GID: GeneID. CID: MrnaId. Chr.PL: length of Pro in AA. MW: Molecular Weights (MV). PI: Protein Isoelectric Points. NE: Number of Exons. SL: subcellular localization.

Phylogenetic Analysis of the *CaWOX* Gene Family

A phylogenetic tree was constructed using 15 *A. thaliana* *WOX* genes, 18 *P. trichocarpa* *WOX* genes, 16 *S. suchowensis* *WOX* genes, and the 14 *CaWOX* genes, as shown in the figure 1. According to the clustering pattern of the *A. thaliana* *WOX* gene family, the 63 *WOX* genes were divided into three major clades. Clade 1 contains 36 genes, Clade 2 contains 12 genes, and Clade 3 contains 15 genes. *WOX* genes from all four species are distributed across each clade, indicating the relatively conserved evolution of the *WOX* gene family in plants. The 14 *CaWOX* genes are clustered into these clades as follows: Clade 1 contains the largest number (7 members): *CaWOX2*, *CaWOX5*, *CaWOX7*, *CaWOX8*, *CaWOX10*, *CaWOX12*, and *CaWOX13*. Clade 2 contains 4 members: *CaWOX3*, *CaWOX9*, *CaWOX11*, and *CaWOX14*. Clade 3 contains 3 members: *CaWOX1*, *CaWOX4*, and *CaWOX6*.





Figure 1. Phylogenetic tree of the *WOX* gene family across different species. The evolutionary relationships among *WOX* genes from *A. thaliana*, *P. trichocarpa*, *C. arbutifolia*, and *S. suchowensis* are shown.

Conserved Motif Analysis and Gene Structure Analysis of the Gene Family

The results from the MEME software analysis indicate that the number and arrangement of conserved protein motifs within the 14 CaWOX protein sequences exhibit a degree of similarity and conservation (Figure 2). All 14 CaWOX protein sequences contain motif 1 and motif 2. Within Clade 3, the three CaWOX protein sequences share identical types and numbers of conserved motifs: motif 1, motif 2, motif 4, and motif 6. In Clade 1, the seven CaWOX protein sequences show some variation in their conserved motif composition. CaWOX2 and CaWOX12 share


identical motif composition and order: motif 1, motif 2, motif 5, motif 8. CaWOX5 and CaWOX7 share identical motif composition and order: motif 2, motif 3, motif 8. CaWOX10 and CaWOX13 share identical motif composition and order: motif 1, motif 2, motif 8, motif 10. In Clade 2, the conserved motifs also show specific patterns. CaWOX3 and CaWOX9 share identical motif composition and order: motif 1, motif 2, motif 3, motif 7, motif 9. CaWOX11 and CaWOX14 share identical motif composition and order: motif 1, motif 2, motif 3. Analysis of the gene structures (exon-intron organization) of the 14 *CaWOX* genes reveals a relatively high degree of similarity among genes within the same clade. In contrast, notable differences in both motif composition and gene structure are observed between different clades. This suggests that the *CaWOX* genes may have undergone functional differentiation in *C. arbutifolia*, potentially participating in distinct physiological processes within the plant.



Analysis of Cis-acting Elements in Gene Promoter Regions

Cis-acting elements located in promoter regions can regulate gene expression by interacting with trans-acting factors. Analysis of these elements can reveal a gene's potential responsiveness to various biotic and abiotic stresses. Examination of the 2,000 bp upstream sequences indicates that the *CaWOX* gene family can be clearly clustered into two distinct evolutionary clades, with significant genetic differentiation

characteristics between them (Figure 3). The first clade includes *CaWOX2*, *CaWOX12*, *CaWOX7*, *CaWOX8*, *CaWOX10*, and *CaWOX13*. High bootstrap support values, often reaching 100%, among genes within this clade suggest a reliable and close evolutionary relationship. The second clade comprises *CaWOX1*, *CaWOX4*, *CaWOX6*, *CaWOX3*, *CaWOX9*, *CaWOX11*, and *CaWOX14*, whose evolutionary relationships appear more stable and conserved. The promoter regions of the *CaWOX* genes contain a rich variety of cis-acting elements with specific distribution patterns, primarily falling into three functional categories. Abiotic Stress Response Elements: These include the low-temperature-responsive LTR, drought-responsive MBS, and dehydration/salt stress-responsive DRE elements. For instance, the *CaWOX1* promoter contains both LTR and DRE elements, while *CaWOX4* and *CaWOX5* promoters are enriched with MBS elements, suggesting these genes may be involved in adaptive responses to multiple abiotic stresses. Hormone Signaling Response Elements: These involve ABA-responsive ABRE, jasmonate-responsive CGTCA/TGACG-motif, and gibberellin-responsive P-box elements. For example, the *CaWOX2* promoter contains both ABRE and CGTCA-motif, and *CaWOX6* contains the gibberellin-responsive P-box, indicating this gene family may participate in physiological process regulation through hormone signaling pathways. Growth and Development-Related Elements: These include meristem expression-related CAT-box and endosperm development-related GCN4_motif. Specifically, the *CaWOX2* promoter contains a CAT-box, and *CaWOX13* contains a GCN4_motif, reflecting the potential roles of some genes in plant growth and differentiation. Further correlation between phylogeny and element distribution reveals that genes within the same evolutionary clade share more similar compositions of cis-acting elements, suggesting that closely related genes tend to have more consistent functional regulation patterns. In contrast, genes from different clades show significant differences in element types, corresponding to distinct functional differentiations. In summary, the evolutionary relationships within the *CaWOX* gene family are well-defined. Through diverse promoter cis-acting elements, these genes are likely broadly involved in abiotic stress responses, hormone signal transduction, and growth regulation. This analysis provides crucial clues for the subsequent targeted



257 Ka/Ks Analysis of the *CaWOX* Gene Family

258 The results of the Ka/Ks analysis for the 14 *CaWOX* genes showed that all
 259 homologous gene pairs have Ka/Ks ratios less than 1, ranging from 0.145 to 0.391
 260 (Supplementary data). The pair *CaWOX1-CaWOX6* has the smallest Ka/Ks value
 261 (0.145), while *CaWOX11-CaWOX14* has the largest (0.391). This indicates that
 262 homologous genes within the *CaWOX* family have undergone purifying selection
 263 during evolution, with varying intensities of selective pressure across different gene
 264 pairs. Regarding sequence characteristics, the effective lengths of homologous gene
 265 pairs range from 630 to 1143 bp, with *CaWOX3-CaWOX9* having the longest effective
 266 length (1143 bp). The average number of synonymous sites (AverageS-sites) ranges
 267 from 133.92 to 265.67, and the average number of non-synonymous sites (AverageN-
 268 sites) ranges from 493.42 to 877.33. For all gene pairs, the frequency of synonymous
 269 substitutions (pS) is higher than that of non-synonymous substitutions (pN), further
 270 supporting the role of purifying selection in maintaining the functional stability of
 271 *CaWOX* genes.

Collinearity Analysis

Analysis of tandem duplication among the 14 *CaWOX* genes revealed a dispersed distribution pattern (Figure 4A), with genes not clustered on a few chromosomes but rather widely distributed across multiple chromosomes including Chr02, Chr04, Chr11, Chr12, and Chr15. Chr04 appears to be an enriched region, harboring multiple family members such as *CaWOX3*, *CaWOX6*, and *CaWOX5*. A total of 8 gene pairs were identified as tandem duplications. Significant collinearity relationships were observed across several chromosomes: genes on Chr04 (*CaWOX3*, *CaWOX6*) show collinearity with genes on Chr15 (*CaWOX13*) and Chr12 (*CaWOX10*). Similarly, *CaWOX9* on Chr11 forms collinear pairs with *CaWOX* genes on chromosomes such as Chr04 and Chr12. These findings indicate that the *CaWOX* gene family in *C. arbutifolia* has experienced gene duplication events during genome evolution, which is a core mechanism driving the expansion of this gene family. The collinear regions are often located on dispersed, non-homologous chromosomes, suggesting that genes were relocated following duplication events during chromosomal evolution. This dispersed genomic distribution provides a structural basis for the functional differentiation of *CaWOX* genes. In summary, the *CaWOX* gene family in *C. arbutifolia* expanded through duplication events, and its dispersed chromosomal distribution and extensive collinearity relationships provide crucial chromosomal-level evidence for future investigations into the functional specialization and evolutionary origins of this gene family. Interspecific collinearity analysis was performed among the genomes of four species: *P. trichocarpa*, *A. thaliana*, *S. suchowensis*, and *C. arbutifolia* (Figure 4B). The collinearity results for the *CaWOX* gene family were plotted to obtain insights into its evolutionary history across these lineages. The results show a significant positive correlation between phylogenetic relatedness and the degree of collinearity conservation. The densest network of collinear connections is observed between *C. arbutifolia* and *P. trichocarpa*, with multiple chromosomes of *C. arbutifolia* forming numerous collinear pairs with most chromosomes of *P. trichocarpa*. This reflects the highest degree of evolutionary conservation within the *WOX* gene family between these

two species. The number of collinear connections between *C. arbutifolia* and *A. thaliana* is intermediate, reflecting their family-level phylogenetic relationship. In contrast, collinear connections between *C. arbutifolia* and *S. suchowensis* are relatively sparse, which may be attributed to their greater phylogenetic distance and consequently lower gene conservation. Furthermore, *CaWOX* genes can form collinear associations with homologous genes from multiple species, indicating that the core sequences and functions of this gene family have retained a degree of evolutionary conservation throughout plant evolution. In conclusion, this cross-species collinearity analysis not only confirms the evolutionary conservation of the *CaWOX* gene family but also clarifies the influence of phylogenetic relationships on gene collinearity. It provides crucial evolutionary evidence for inferring the potential biological functions of *CaWOX* genes in *C. arbutifolia* by leveraging known functions from model species.





Figure 4. Chromosomal collinearity analysis of the *WOX* gene family. (A) Intra-species collinearity plot among chromosomes of *C. arbutifolia*. (B) Inter-species collinearity plot showing chromosomes of four species: *A. thaliana*, *P. trichocarpa*, *C. arbutifolia*, and *S. suchowensis*. Red connecting lines represent collinear gene pairs between different species, illustrating conserved evolutionary relationships of *WOX* genes across species.

Expression Profiles of *CaWOX* Genes

To investigate the roles of *CaWOX* genes in the growth and development of *C. arbutifolia*, we analyzed their differential expression across various tissues. Quantitative real-time PCR (qRT-PCR) was performed to systematically analyze the

expression patterns of 10 *CaWOX* family genes across five different tissues, namely upper stem, lower stem, root, young leaf, and mature leaf (Figure 5). The results revealed that *CaWOX* family genes exhibited significant tissue-specific expression characteristics, with distinct expression profiles of different members during the development of various plant organs. In leaf tissues, *CaWOX3*, *CaWOX4*, *CaWOX5*, *CaWOX9*, and *CaWOX11* all showed dominant expression in mature leaves, with their relative expression levels in mature leaves significantly higher than those in other tissues including stems and roots ($P < 0.05$). Among them, *CaWOX3*, *CaWOX5*, and *CaWOX11* also maintained high expression levels in young leaves, indicating that these genes may be widely involved in the whole process of leaf initiation, development, and maturation in *C. arbutifolia*. In root tissues, *CaWOX1*, *CaWOX6*, *CaWOX10*, and *CaWOX14* displayed a prominent root-specific high expression pattern. Their relative expression levels in roots were the highest among all tested tissues, and were significantly higher than those in stem and leaf tissues ($P < 0.05$), suggesting that these four genes play a core regulatory role in root growth, development, and morphogenesis. In stem tissues, only *CaWOX12* exhibited stem-specific dominant expression. The expression level of this gene was the highest in the lower stem, followed by the upper stem, and was significantly downregulated in root and leaf tissues ($P < 0.05$). In contrast, the expression levels of other *CaWOX* family members were generally low in stem tissues, indicating that *CaWOX12* may be a key member of the *WOX* family that mediates stem growth and development. In summary, the expression patterns of different members of the *CaWOX* family are clearly differentiated in a tissue-specific manner, with each member showing unique tissue-specific expression characteristics during the development of roots, stems, and leaves. These findings provide a key expression-level basis for the subsequent functional characterization of *CaWOX* family genes in plant organ morphogenesis.



Discussion

The *WOX* gene family represents a class of plant-specific transcription factors characterized by a conserved homeodomain (HD), making it a focal point in plant developmental biology and molecular genetics. As a key regulator in plant growth, development, and stress responses, deciphering its evolutionary patterns, structural features, and expression profiles is fundamental to understanding its function. However, this family had not been previously identified or analyzed in the woody species *C. arbutifolia*. In this study, a BLASTP search against the *C. arbutifolia* database led to the identification of 14 *CaWOX* genes. Comparative analysis shows that while the number of *WOX* genes varies across species. 15 in *A. thaliana*, 13 in rice, 18 in flax, and 12 in grape. the overall count remains relatively stable[11]. This suggests a high degree of evolutionary conservation within the *WOX* family across the plant kingdom.

Variations in protein physicochemical properties hint at potential functional diversification[34, 35]. The *CaWOX* members exhibit significant differences in amino acid length (213–406 AA) and molecular weight (24,481.4–45,121.3 Da). For instance, *CaWOX9* (406 AA) has a substantially higher molecular weight than *CaWOX2* (213 AA), a disparity that may correspond to the gain or loss of functional domains. The range in isoelectric points (pI 5.59–9.99) could influence protein stability within the nucleus and its affinity for DNA or other interacting proteins. For example, the relatively basic *CaWOX2* (pI=9.7) and the acidic *CaWOX4* (pI=5.59) may function in different pH microenvironments or distinct regulatory pathways. Subcellular localization predictions indicate that all *CaWOX* proteins are localized to the nucleus, which is entirely consistent with their core function as transcription factors. This nuclear localization ensures their capacity to bind DNA regulatory regions, forming the structural basis for their transcriptional activity and further underscoring the evolutionary conservation of this family's function.

The core structural feature of the *WOX* family is the HD domain, composed of 60–66 amino acids responsible for binding DNA regulatory elements. Beyond this, members can be further classified based on additional domains such as the WUS-box, EAR repression motif, and VLWV motif[36–38]. Phylogenetically, the family is typically divided into three clades: the ancient clade, found in basal plants like mosses and ferns; the Intermediate clade, exclusive to angiosperms; and the WUS clade, also angiosperm-specific and containing distinctive domains like the WUS-box. This structural differentiation among clades lays the groundwork for their functional diversity. Our phylogenetic analysis, which included *WOX* genes from *P. trichocarpa* and *Arabidopsis thaliana*, investigated the origin and evolution of *CaWOX* genes. The cross-species phylogenetic tree grouped *CaWOX* genes with those from *A. thaliana*, *P. trichocarpa*, and *Eutrema salsugineum* into distinct evolutionary groups with high bootstrap support. This reflects both the evolutionary conservation of the *WOX* family in higher plants where conserved core domains like the HD maintain basic regulatory functions and the species-specific expansion of *CaWOX* in *C. arbutifolia*. Intra-species collinearity analysis in *C. arbutifolia* revealed collinear gene pairs across multiple

chromosomes, confirming that segmental or whole-genome duplication events are the primary drivers of *CaWOX* family expansion. The subsequent dispersed chromosomal distribution of these duplicated genes provides a genomic structural basis for their functional divergence.

Gene structure and conserved motif analyses further support the functional correlations implied by phylogenetic grouping[39, 40]. Genes within the same evolutionary clade share highly similar motif compositions and exon-intron structures. Motif1 corresponds to the conserved HD, the core element for DNA binding and transcriptional regulation, ensuring basic functional consistency within the clade. Unique motifs and differential exon distributions likely serve as structural markers of functional divergence following gene duplication. These unique motifs may participate in novel protein interactions or regulatory networks, enabling the *CaWOX* family to adapt to specific growth requirements of *C. arbutifolia*. Variations in exon number directly reflect structural diversification. Some members contain only 2 exons, while others like *CaWOX10* and *CaWOX9* possess 4 exons. This variation may be related to alternative splicing events or domain reorganization, where a greater number of exons could confer increased structural diversity, allowing participation in more complex regulatory networks.

Analysis of the promoter regions of the *CaWOX* gene family revealed enrichment for three core categories of cis-acting elements. Hormone-responsive elements, such as ABA-responsive ABRE, jasmonate-responsive CGTCA/TGACG-motif, and gibberellin-responsive P-box, indicating broad involvement in plant hormone signaling pathways. The presence of ABRE elements in the promoters of *CaWOX1* and *CaWOX4* suggests potential roles in responding to stresses like drought via ABA signaling. Abiotic stress-responsive elements, including low-temperature-responsive LTR, drought-responsive MBS, and dehydration/salt-responsive DRE, highlighting the potential function of *CaWOX* genes in stress adaptation in *C. arbutifolia*. The presence of MBS elements in *CaWOX5* and *CaWOX3* promoters aligns with recently reported drought-resistance functions of the *WOX* family. Growth and development-related elements, such as the meristem-specific CAT-box and the endosperm development-

related GCN4_motif, corroborating the classic growth-regulatory roles of the *WOX* family. For example, CAT-box elements in the promoters of *CaWOX2* and *CaWOX12* closely match their high expression in root meristems, suggesting a role in root meristem maintenance.

Correlating phylogeny with element distribution shows that *CaWOX* genes within the same evolutionary clade share more consistent compositions and distributions of promoter cis-elements[41, 42]. For instance, *CaWOX1*, *CaWOX4*, and *CaWOX6*, clustered in one clade, all contain both hormone-responsive and stress-responsive elements, indicating conserved regulatory pathways among evolutionarily related genes. In contrast, genes from different clades show significant divergence in element types, the *CaWOX2* clade is enriched for meristem-related CAT-box elements, while the *CaWOX3* clade contains more stress-responsive elements. This differentiation corresponds to functional specialization, an outcome of gene duplication adapting to diverse regulatory demands. The diversity of promoter cis-elements further explains the functional complexity of the *CaWOX* family. For example, the *CaWOX12* promoter contains both CAT-box and LTR elements, implying a dual role in root meristem development and cold stress response, showcasing its potential for "growth-stress response" coordination. The specific distribution of these elements within the promoter region may influence signal sensitivity and regulatory efficiency. The characteristics of *CaWOX* promoter cis-elements not only reflect the relationship between evolutionary conservation and functional divergence but also reveal the family potential as a regulatory hub connecting growth and development with stress response and integrating hormonal, stress, and growth signals to build a multi-dimensional regulatory network, providing a molecular foundation for *C. arbutifolia* to adapt to complex environments and coordinate growth.

Gene duplication events, including tandem and segmental duplications, are closely associated with gene family expansion and evolution[43-45]. The evolution of the *WOX* family follows a pattern of species differentiation and gene duplication". Lower plants possess only 1–2 *WOX* members, while higher angiosperms show significant expansion, driven primarily by whole-genome and segmental duplications. The expansion of the

Arabidopsis *WOX* family is linked to ancient hexaploidization events, and the collinear characteristics of poplar *WOX* genes also confirm the role of duplication. Concurrently, highly conserved homologous relationships exist between *WOX* families of different species, reflecting their evolutionary conservation. In this study, the 14 *CaWOX* genes are unevenly distributed across 11 chromosomes. The clustering of *CaWOX1/2* on Chr02 and *CaWOX4/5/6* on Chr05 aligns with the conclusion that "gene duplication drives family expansion": members on the same chromosome likely originated from segmental duplication events, subsequently diverging in physicochemical properties and exon structures to form specialized functions, adapting to the regulatory needs of different developmental stages or environmental conditions in *C. arbutifolia*. Regarding tandem duplication, we identified 8 pairs of duplicated genes within the *CaWOX* family in the poplar genome, indicating that gene duplication has provided impetus for *CaWOX* evolution, promoting the evolutionary process of these genes in *C. arbutifolia*. Interspecific collinearity analysis with Arabidopsis, rice, and *S. suchowensis* genomes revealed significant differences in the number of tandemly duplicated genes between the *CaWOX* family and *WOX* families of other species. Notably, the collinear connections between *C. arbutifolia* and *P. trichocarpa* were the most dense, while connections with Arabidopsis were relatively sparse. This pattern is highly consistent with phylogenetic relatedness, indicating that the conservation of *WOX* genes decreases with increasing phylogenetic distance. Based on this, known functions of *WOX* genes in closely related species like *P. trichocarpa* can be used to infer the core regulatory roles of *CaWOX* genes, providing targeted references for subsequent functional validation.

The WUSCHEL-related homeobox family is a plant-specific family of homeobox transcription factors that play critical roles in core biological processes, including meristem homeostasis maintenance, organ morphogenesis, and temporal regulation of growth and development in plants[2]. In this study, we characterized the spatial expression profiles of 10 *CaWOX* family genes across root, stem, and leaf tissues of *C. arbutifolia*, which exhibited highly tissue-specific expression patterns. *CaWOX3*, *CaWOX4*, *CaWOX5*, *CaWOX9*, and *CaWOX11* showed high expression levels in leaf

tissues, with expression patterns highly consistent with the reported functions of their homologous genes. *WOX* family members are well-recognized key regulators of plant leaf development[46]. In *Arabidopsis thaliana*, *AtWOX3* determines leaf margin morphogenesis by regulating the activity of marginal meristems, while *AtWOX9* is involved in leaf primordium initiation and the regulation of cell proliferation in young leaves[13]. In the present study, the expression levels of the above five genes in mature leaves were significantly higher than those in other tissues, and *CaWOX3*, *CaWOX5*, and *CaWOX11* also maintained high expression levels in young leaves. This result suggests that these *CaWOX* genes are not only involved in the early developmental processes of *C. arbutifolia*, including leaf primordium initiation and cell proliferation and differentiation in young leaves, but may also function in maintaining the physiological functions of mature leaves. Given the biological characteristics of *C. arbutifolia* as a deciduous tree species, mature leaves serve as the core organs for photosynthetic carbon fixation and stress response in the plant. Therefore, in addition to regulating leaf morphogenesis, these leaf-specific *CaWOX* genes may also participate in maintaining photosynthetic homeostasis and mediating abiotic stress responses in mature leaves by regulating the expression of downstream target genes. This finding provides a new direction for subsequent studies on their application value in improving photosynthetic efficiency and stress resistance breeding of forest trees. *CaWOX1*, *CaWOX6*, *CaWOX10*, and *CaWOX14* exhibited root-specific high expression, which is highly consistent with the conserved regulatory function of the *WOX* family in root development. The root system is the core organ of forest trees for water and nutrient uptake, as well as the perception of soil environmental signals, and *WOX* transcription factors are key regulators for maintaining root apical meristem homeostasis and the initiation of lateral and adventitious roots[47, 48]. Studies in model plants have confirmed that *AtWOX11* and *AtWOX12* synergistically regulate adventitious root initiation, and *AtWOX5* sustains continuous root growth by maintaining cell viability in the root apical quiescent center[49-51]. In this study, the expression levels of these four *CaWOX* genes in roots were significantly higher than those in aboveground tissues, indicating that these genes are core regulators of root

development in *C. arbutifolia* and may determine root architecture by modulating root apical meristem activity and lateral root formation. Notably, the root developmental status of *C. arbutifolia* directly determines the drought and barren tolerance of the plant. These root-specific *CaWOX* genes are not only regulators of root morphogenesis, but may also serve as key targets mediating stress adaptation in forest trees. This is consistent with our previous finding that *WOX* family genes are involved in drought response from transcriptome analysis of drought-stressed *C. arbutifolia*, and provides core candidate genes for subsequent dissection of the synergistic regulatory mechanism between root development and stress resistance in forest trees.

In stem tissues, only *CaWOX12* exhibited stem-specific dominant expression, with a significantly higher expression level in the lower stem than in the upper stem. This expression pattern is highly matched with the secondary growth characteristics of woody plants. Unlike herbaceous plants, the stems of woody plants undergo continuous secondary growth, a process in which vascular cambium activity drives xylem and phloem differentiation and ultimately wood formation[52, 53]. Furthermore, the lower stem of the plant has a more vigorous secondary growth process and a higher degree of lignification. Previous studies have shown that *WOX* family members play important regulatory roles in vascular development and secondary cell wall biosynthesis in woody plants. For example, *PtoWOX4* in *Populus* regulates secondary growth by modulating vascular cambium activity[1, 54]. In this study, *CaWOX12* was specifically highly expressed in the lower stem but had extremely low expression levels in roots and leaves, suggesting that this gene may be a key transcription factor regulating stem secondary growth, vascular tissue differentiation, and lignification in *C. arbutifolia*. This result provides an important research target for subsequent dissection of the molecular mechanism of wood formation and molecular breeding for wood property improvement in Salicaceae tree species.

The spatial expression patterns of different *CaWOX* family members in *C. arbutifolia* showed significant functional differentiation, and such tissue-specific expression characteristics are a core manifestation of subfunctionalization during the evolution of the gene family[55, 56]. During long-term evolution, different copies of

the *CaWOX* family acquired tissue-specific cis-regulatory elements, thus differentiating into specific functions that separately regulate the development of root, stem, and leaf organs, and forming a transcriptional regulatory network with clear functional division and synergistic regulation of plant growth and development[57, 58]. The tissue expression characteristics of the *CaWOX* family identified in this study lay a solid foundation for subsequent functional validation of these genes via technologies such as gene editing and genetic transformation, and also provide an important theoretical basis for the regulation of growth and development, as well as molecular breeding for stress resistance and wood property improvement of *C. arbutifolia* and even other Salicaceae trees.

Conclusion

In this study, 14 members of the *CaWOX* gene family were identified in *C. arbutifolia*, and the evolutionary characteristics and functional differentiation patterns of this family were systematically elucidated. Phylogenetic analysis classified the *CaWOX* proteins into four conserved subgroups, and members within the same subgroup exhibited highly similar gene structures and conserved motifs, confirming subfunctionalization during the family's evolution. Collinearity analysis verified that whole-genome duplication constituted the core driving force for gene expansion in this family; furthermore, the *CaWOX* family displayed extremely high conservation among closely related species in Salicaceae, but weak collinearity with *Arabidopsis thaliana*, reflecting species-specific evolution after the divergence of monocotyledons and dicotyledons. Cis-element analysis revealed that the promoter regions of these genes were rich in elements associated with hormone response, stress response, and meristem regulation, indicating their dual functions in regulating organ development and stress adaptation. Tissue expression analysis further uncovered distinct functional differentiation: *CaWOX3/4/5/9/11* were predominantly expressed in leaves and participated in leaf development and physiological homeostasis maintenance; *CaWOX1/6/10/14* exhibited root-specific high expression and regulated root growth and development; *CaWOX12* was specifically highly expressed in stems and mediated

stem secondary growth and lignification. This study lays a solid foundation for subsequent functional verification of *CaWOX* family genes in *C. arbutifolia*, and also provides vital theoretical support and high-quality gene resources for growth regulation, wood property improvement, and stress-resistant molecular breeding of Salicaceae trees.

Funding

This work was financially supported by the National Natural Science Foundation of China (Grant No. 31670662) and the Independent Scientific Research Project of Jiangsu Academy of Forestry (Grant No. ZZKY202101).

Author contributions

P.W. designed the experiments and drafted and prepared the manuscript. J.Z., R.H., W.W., and H.W. reviewed and edited the manuscript. D.S. and X.H. conceived and funded the study. All authors read and approved the final version of the manuscript.

Competing Interests

The authors declare no known competing financial interests or personal relationships that could influence the work reported in this paper.

Acknowledgements

Not applicable.

Abbreviations

Not applicable.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Data will be made available on request.

Data availability

The genomic data supporting this study are publicly available to ensure accessibility and reproducibility. Genome sequences of *C. arbutifolia* and *S. suchowensis* can be accessed in the NCBI database under accession number ASM2516995v1 and ASM2903076v1. Reference genome sequences and annotations for *A. thaliana*, *P. trichocarpa*, were obtained from Phytozome V14(<https://phytozome-next.jgi.doe.gov/info>).

References

1. Zhang M, Chen X, Lou X, Zhang Y, Han X, Yang Q, Tong Z, Zhang J: Identification of WUSCHEL-related homeobox (WOX) gene family members and determination of their expression profiles during somatic embryogenesis in *Phoebe bournei*. *For Res (Fayettev)* 2023, 3:5.
2. Jha P, Ochatt SJ, Kumar V: WUSCHEL: a master regulator in plant growth signaling. *Plant Cell Reports* 2020, 39(4):431-444.
3. Hanada K, Sawada Y, Kuromori T, Klausnitzer R, Saito K, Toyoda T, Shinozaki K, Li W-H, Hirai MY: Functional Compensation of Primary and Secondary Metabolites by Duplicate Genes in *Arabidopsis thaliana*. *Molecular biology and evolution* 2010, 28(1):377-382.
4. van der Graaff E, Laux T, Rensing SA: The WUS homeobox-containing (WOX) protein family. *Genome Biology* 2009, 10(12):248.
5. Zhao Y, Hu Y, Dai M, Huang L, Zhou D-X: The WUSCHEL-Related Homeobox Gene WOX11 Is Required to Activate Shoot-Borne Crown Root Development in Rice. *The Plant Cell* 2009, 21(3):736-748.
6. Jha P, Ochatt SJ, Kumar V: WUSCHEL: a master regulator in plant growth signaling. *Plant Cell Rep* 2020, 39(4):431-444.
7. Ikeda M, Mitsuda N, Ohme-Takagi M: *Arabidopsis* WUSCHEL Is a Bifunctional Transcription Factor That Acts as a Repressor in Stem Cell Regulation and as an Activator in Floral Patterning. *The Plant Cell* 2009, 21(11):3493-3505.
8. Alvarez JM, Bueno N, Cañas RA, Avila C, Cánovas FM, Ordás RJ: Analysis of

-
- 620 the WUSCHEL-RELATED HOMEBOX gene family in *Pinus pinaster*: New
621 insights into the gene family evolution. *Plant Physiology and Biochemistry* 2018,
622 123:304-318.
- 623 9. Eeda SK, Werr W: Transcription of the WUSCHEL-RELATED HOMEBOX 4
624 gene in *Arabidopsis thaliana*. *Gene Expr Patterns* 2020, 38:119150.
- 625 10. Liu J, Sheng L, Xu Y, Li J, Yang Z, Huang H, Xu L: WOX11 and 12 are involved
626 in the first-step cell fate transition during de novo root organogenesis in
627 *Arabidopsis*. *Plant Cell* 2014, 26(3):1081-1093.
- 628 11. Zhang X, Zong J, Liu J, Yin J, Zhang D: Genome-Wide Analysis of WOX Gene
629 Family in Rice, Sorghum, Maize, *Arabidopsis* and Poplar. *Journal of Integrative*
630 *Plant Biology* 2010, 52(11):1016-1026.
- 631 12. Lie C, Kelsom C, Wu X: WOX2 and STIMPY-LIKE/WOX8 promote cotyledon
632 boundary formation in *Arabidopsis*. *The Plant Journal* 2012, 72(4):674-682.
- 633 13. Yang Y, Ren Z, Li L, Li Y, Han Y, Liu Y, Cao H: WOX2 functions redundantly
634 with WOX1 and WOX4 to positively regulate seed germination in *Arabidopsis*.
635 *Planta* 2024, 259(4):83.
- 636 14. Gong W, Bak DT, Wendrich JR, Weijers D, Laux T: CDC48A, an interactor of
637 WOX2, is required for embryonic patterning in *Arabidopsis thaliana*. *Plant Cell*
638 *Reports* 2024, 43(7):174.
- 639 15. Hirakawa Y, Kondo Y, Fukuda H: TDIF Peptide Signaling Regulates Vascular
640 Stem Cell Proliferation via the WOX4 Homeobox Gene in *Arabidopsis* *The*
641 *Plant Cell* 2010, 22(8):2618-2629.
- 642 16. Pang A, Li Y, He C, Liu C, Jin H, Pi L, Yang Y: A WOX5/7–SCRReciprocal
643 Feedback Loop in Middle Cell Layer Drives Callus Proliferation. *Plants* 2026,
644 15(2):210.
- 645 17. Yang Y, Liu C, Yu Y, Ran G, Zhai N, Pi L: WUSCHEL RELATED HOMEBOX5
646 and 7 maintain callus development by promoting cell division in *Arabidopsis*.
647 *Plant Science* 2024, 346:112133.
- 648 18. Tian H, Wabnik K, Niu T, Li H, Yu Q, Pollmann S, Vanneste S, Govaerts W, Rolčik
649 J, Geisler M *et al*: WOX5–IAA17 Feedback Circuit-Mediated Cellular Auxin

-
- 650 Response Is Crucial for the Patterning of Root Stem Cell Niches in Arabidopsis.
651 *Molecular Plant* 2014, 7(2):277-289.
- 652 19. Kong D, Hao Y, Cui H: The WUSCHEL Related Homeobox Protein WOX7
653 Regulates the Sugar Response of Lateral Root Development in Arabidopsis
654 *thaliana*. *Molecular Plant* 2016, 9(2):261-270.
- 655 20. Baesso B, Chiatante D, Terzaghi M, Zenga D, Nieminen K, Mahonen AP, Siligato
656 R, Helariutta Y, Scippa GS, Montagnoli A: Transcription factors PRE3 and
657 WOX11 are involved in the formation of new lateral roots from secondary growth
658 taproot in *A. thaliana*. *Plant Biol (Stuttg)* 2018, 20(3):426-432.
- 659 21. Jiang W, Zhou S, Zhang Q, Song H, Zhou D-X, Zhao Y: Transcriptional regulatory
660 network of WOX11 is involved in the control of crown root development,
661 cytokinin signals, and redox in rice. *Journal of Experimental Botany* 2017,
662 68(11):2787-2798.
- 663 22. Xu M, Xie W, Huang M: Two WUSCHEL-related HOMEBOX genes,
664 PeWOX11a and PeWOX11b, are involved in adventitious root formation of poplar.
665 *Physiologia Plantarum* 2015, 155(4):446-456.
- 666 23. Ran N, Liu S, Qi H, Wang J, Shen T, Xu W, Xu M: Long Non-Coding RNA
667 lncWOX11a Suppresses Adventitious Root Formation of Poplar by Regulating the
668 Expression of PeWOX11a. *International Journal of Molecular Sciences* 2023,
669 24(6):5766.
- 670 24. Liao J, Deng B, Cai X, Yang Q, Hu B, Cong J, Zhang Y, Wang G, Xin G, Li Y *et*
671 *al*: Time-course transcriptome analysis reveals regulation of Arabidopsis seed
672 dormancy by the transcription factors WOX11/12. *J Exp Bot* 2023, 74(3):1090-
673 1106.
- 674 25. Liao J, Deng B, Cai X, Yang Q, Hu B, Cong J, Zhang Y, Wang G, Xin G, Li Y *et*
675 *al*: Time-course transcriptome analysis reveals regulation of Arabidopsis seed
676 dormancy by the transcription factors WOX11/12. *Journal of Experimental*
677 *Botany* 2022, 74(3):1090-1106.
- 678 26. Jiang D, Zhang X, Luo L, Li T, Chen H, Ma N, Fu L, Tian P, Mao F, Lü P *et al*:
679 Cytosine Methylation Changes the Preferred Cis-Regulatory Configuration of

-
- 680 Arabidopsis WUSCHEL-Related Homeobox 14. *International Journal of*
681 *Molecular Sciences* 2025, 26(2):763.
- 682 27. He X, Wang Y, Lian J, Zheng J, Zhou J, Li J, Jiao Z, Niu Y, Wang W, Zhang J *et*
683 *al*: The whole-genome assembly of an endangered Salicaceae species: *Chosenia*
684 *arbutifolia* (Pall.) A. Skv. *Gigascience* 2022, 11.
- 685 28. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam
686 H, Valentin F, Wallace IM, Wilm A, Lopez R *et al*: Clustal W and Clustal X version
687 2.0. *Bioinformatics* 2007, 23(21):2947-2948.
- 688 29. Tamura K, Stecher G, Kumar S: MEGA11: Molecular Evolutionary Genetics
689 Analysis Version 11. *Molecular biology and evolution* 2021, 38(7):3022-3027.
- 690 30. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW,
691 Noble WS: MEME SUITE: tools for motif discovery and searching. *Nucleic Acids*
692 *Res* 2009, 37(Web Server issue):W202-208.
- 693 31. Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, Xia R: TBtools: An
694 Integrative Toolkit Developed for Interactive Analyses of Big Biological Data.
695 *Mol Plant* 2020, 13(8):1194-1202.
- 696 32. Chen C, Wu Y, Li J, Wang X, Zeng Z, Xu J, Liu Y, Feng J, Chen H, He Y *et al*:
697 TBtools-II: A "One for All, All for One" Bioinformatics Platform for Biological
698 Big-data Mining. *Molecular Plant* 2023, 16.
- 699 33. Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ,
700 Marra MA: Circos: an information aesthetic for comparative genomics. *Genome*
701 *Res* 2009, 19(9):1639-1645.
- 702 34. Zhu T, Moschou PN, Alvarez JM, Sohlberg JJ, von Arnold S: WUSCHEL-
703 RELATED HOMEBOX 8/9 is important for proper embryo patterning in the
704 gymnosperm Norway spruce. *Journal of Experimental Botany* 2014, 65(22):6543-
705 6552.
- 706 35. Alvarez JM, Sohlberg J, Engström P, Zhu T, Englund M, Moschou PN, von Arnold
707 S: The WUSCHEL-RELATED HOMEBOX 3 gene PaWOX3 regulates lateral
708 organ formation in Norway spruce. *New Phytologist* 2015, 208(4):1078-1088.
- 709 36. Sheng L, Hu X, Du Y, Zhang G, Huang H, Scheres B, Xu L: Non-canonical

-
- 710 WOX11-mediated root branching contributes to plasticity in Arabidopsis root
711 system architecture. *Development* 2017, 144(17):3126-3133.
- 712 37. Wang H, Niu H, Li C, Shen G, Liu X, Weng Y, Wu T, Li Z: WUSCHEL-related
713 homeobox1 (WOX1) regulates vein patterning and leaf size in *Cucumis sativus*.
714 *Horticulture Research* 2020, 7(1):182.
- 715 38. Shafique Khan F, Zeng R-F, Gan Z-M, Zhang J-Z, Hu C-G: Genome-Wide
716 Identification and Expression Profiling of the WOX Gene Family in *Citrus*
717 *sinensis* and Functional Analysis of a CsWUS Member. *International Journal of*
718 *Molecular Sciences* 2021, 22(9):4919.
- 719 39. Rathour M, Sharma A, Kaur A, Upadhyay SK: Genome-wide characterization and
720 expression and co-expression analysis suggested diverse functions of WOX genes
721 in bread wheat. *Heliyon* 2020, 6(12):e05762.
- 722 40. Wen J, Li PF, Ran F, Guo PC, Zhu JT, Yang J, Zhang LL, Chen P, Li JN, Du H:
723 Genome-wide characterization, expression analyses, and functional prediction of
724 the NPF family in *Brassica napus*. *BMC Genomics* 2020, 21(1):871-888.
- 725 41. Riccucci E, Vanni C, Vangelisti A, Fambrini M, Giordani T, Cavallini A, Mascagni
726 F, Pugliesi C: Genome-Wide Analysis of WOX Multigene Family in Sunflower
727 (*Helianthus annuus* L.). *International Journal of Molecular Sciences* 2023,
728 24(4):3352.
- 729 42. Tang L, He Y, Liu B, Xu Y, Zhao G: Genome-Wide Identification and
730 Characterization Analysis of WUSCHEL-Related Homeobox Family in Melon
731 (*Cucumis melo* L.). *International Journal of Molecular Sciences* 2023,
732 24(15):12326.
- 733 43. Akbulut SE, Okay A, Aksoy T, Aras ES, Büyük İ: The genome-wide
734 characterization of WOX gene family in *Phaseolus vulgaris* L. during salt stress.
735 *Physiology and Molecular Biology of Plants* 2022, 28(6):1297-1309.
- 736 44. Lynch M: Gene Duplication and Evolution. *Science* 2002, 297(5583):945-947.
- 737 45. Shen JY, Su TH, Yu DQ, Tan SJ, Zhang YE: Evolution by gene duplication: in the
738 era of genomics. *Yi Chuan* 2025, 47(2):147-171.
- 739 46. Abubakar AS, Wu Y, Chen F, Zhu A, Chen P, Chen K, Qiu X, Huang X, Zhao H,

-
- 740 Chen J *et al*: Comprehensive Analysis of WUSCEL-Related Homeobox Gene
741 Family in Ramie (*Boehmeria nivea*) Indicates Its Potential Role in Adventitious
742 Root Development. *Biology* 2023, 12(12):1475.
- 743 47. Willig J-J, Guarneri N, van Loon T, Wahyuni S, Astudillo-Estévez IE, Xu L,
744 Willemsen V, Goverse A, Sterken MG, Lozano-Torres JL *et al*: Transcription
745 factor WOX11 modulates tolerance to cyst nematodes via adventitious lateral root
746 formation. *Plant Physiology* 2024, 195(1):799-811.
- 747 48. Deng Y, Zhu Y, Su W, Zhang M, Liao W: Transcription factor WUSCHEL-related
748 homeobox (WOX) underground revelations: Insights into plant root development.
749 *Plant Physiology and Biochemistry* 2025, 224:109928.
- 750 49. Zhou S, Jiang W, Long F, Cheng S, Yang W, Zhao Y, Zhou D-X: Rice
751 Homeodomain Protein WOX11 Recruits a Histone Acetyltransferase Complex to
752 Establish Programs of Cell Proliferation of Crown Root Meristem. *The Plant Cell*
753 2017, 29(5):1088-1104.
- 754 50. Liu J, Hu X, Qin P, Prasad K, Hu Y, Xu L: The WOX11-LBD16 Pathway Promotes
755 Pluripotency Acquisition in Callus Cells During De Novo Shoot Regeneration in
756 Tissue Culture. *Plant Cell Physiol* 2018, 59(4):734-743.
- 757 51. Liu J, Sheng L, Xu Y, Li J, Yang Z, Huang H, Xu L: WOX11 and 12 Are Involved
758 in the First-Step Cell Fate Transition during de Novo Root Organogenesis in
759 Arabidopsis *The Plant Cell* 2014, 26(3):1081-1093.
- 760 52. Tanaka H, Hashimoto N, Kawai S, Yumoto E, Shibata K, Tameshige T, Yamamoto
761 Y, Sugimoto K, Asahina M, Ikeuchi M: Auxin-Induced WUSCHEL-RELATED
762 HOMEODOMAIN13 Mediates Asymmetric Activity of Callus Formation upon
763 Cutting. *Plant and Cell Physiology* 2022, 64(3):305-316.
- 764 53. Liu R, Wen S-S, Sun T-T, Wang R, Zuo W-T, Yang T, Wang C, Hu J-J, Lu M-Z,
765 Wang L-Q: PagWOX11/12a positively regulates the PagSAUR36 gene that
766 enhances adventitious root development in poplar. *Journal of Experimental*
767 *Botany* 2022, 73(22):7298-7311.
- 768 54. Sun R, Zhang X, Ma D, Liu C: Identification and Evolutionary Analysis of Cotton
769 (*Gossypium hirsutum*) WOX Family Genes and Their Potential Function in

-
- 770 Somatic Embryogenesis. *International Journal of Molecular Sciences* 2023,
771 24(13):11077.
- 772 55. Wang D, Qiu Z, Xu T, Yao S, Zhang M, Cheng X, Zhao Y, Ji K: Identification and
773 Expression Patterns of WOX Transcription Factors under Abiotic Stresses in *Pinus*
774 *massoniana*. *International Journal of Molecular Sciences* 2024, 25(3):1627.
- 775 56. Khan FS, Goher F, Hu CG, Zhang JZ: WUSCHEL-related homeobox (WOX)
776 transcription factors: key regulators in combating abiotic stresses in plants.
777 *Horticulture Advances* 2024, 2(1):2.
- 778 57. Leonardo B, Emanuela T, Letizia MM, Antonella M, Marco M, Fabrizio A,
779 Beatrice BM, Adriana C: Cadmium affects cell niches maintenance in *Arabidopsis*
780 *thaliana* post-embryonic shoot and root apical meristem by altering the expression
781 of WUS/WOX homolog genes and cytokinin accumulation. *Plant Physiology and*
782 *Biochemistry* 2021, 167:785-794.
- 783 58. Jiang Y, Liang J, Wang C, Tan L, Kawano Y, Nagawa S: Stem Cell Factors BAM1
784 and WOX1 Suppressing Longitudinal Cell Division of Margin Cells Evoked by
785 Low-Concentration Auxin in Young Cotyledon of *Arabidopsis*. *International*
786 *Journal of Molecular Sciences* 2025, 26(10):4724.
787

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarydata.docx](#)