MycoKeys 121: 1–20 (2025)
DOI: 10.3897/mycokeys.121.154055

Research Article

The polyphasic approach reveals two new species and two new records of *Nigrospora* (Apiosporaceae, Amphisphaeriales) associated with *Aquilaria sinensis* from China

Shiyu Zhang[1,2,3,4], Junfu Li[2,3], Hongbo Jiang[2,3], Shuang Ye[5], Ausana Mapook[4], Jianchu Xu[2,3], Kevin D. Hyde[4], Prapassorn Damrongkool Eungwanichayapant[1]

1 *School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand*
2 *Center for Mountain Futures, Kunming Institute of Botany, Kunming 650201, China*
3 *CIFOR-ICRAF China Program, World Agroforestry (ICRAF), Kunming 650201, China*
4 *Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand*
5 *Tea Research Institute, Yunnan Academy of Agricultural Sciences, Kunming 650201, China*

Corresponding authors: Junfu Li (lijunfu@mail.kib.ac.cn); Prapassorn Damrongkool Eungwanichayapant (prapassorn@mfu.ac.th)

Abstract

This study isolated endophytic fungi of the genus *Nigrospora* from *Aquilaria sinensis* (Chinese agarwood) in Guangxi, China. Through rigorous morphological comparisons and molecular phylogenetic analyses, we have identified two new species (*N. guangxiensis* and *N. pubeiensis*) and two new host-recorded species (*N. oryzae* and *N. camelliae-sinensis*). The morphological analysis revealed that the new species align with the genus-level definition of *Nigrospora* in colony morphology, conidiophores, and conidial characteristics. However, they were distinguished from known species by finer features such as conidial dimensions and conidiogenous structures. For instance, conidiophores in *Nigrospora* are typically differentiated from hyphae, characterized by dark pigmentation, which is consistent with broader taxonomic descriptions. Molecular phylogeny, based on concatenated datasets of the ITS, *TEF1-α*, and *TUB2* loci, showed that *N. pubeiensis* is most closely related to *N. chinensis*, while *N. guangxiensis* forms a distinct basal clade relative to these two groups. Additionally, *N. oryzae* and *N. camelliae-sinensis* were first recorded on *A. sinensis*, expanding their known host ranges. The study enriches the taxonomic framework of *Nigrospora* by providing novel data from China, emphasizing the importance of integrating morphological and molecular approaches in fungal systematics. It also underscores the ecological diversity of fungal-host interactions in agarwood-producing trees, a topic explored in related research on endophytic fungi. These findings have significant implications for our understanding of fungal diversity and their ecological roles in agarwood-producing trees, and they pave the way for further research in this area.

Key words: 2 new species, asexual fungi, endophyte, Hyphomycetes, multigene phylogeny, Sordariomycetes

OPEN ACCESS

This article is part of:
Exploring the Hidden Fungal Diversity: Biodiversity, Taxonomy, and Phylogeny of Saprobic Fungi
Edited by Samantha C. Karunarathna, Danushka Sandaruwan Tennakoon, Ajay Kumar Gautam

Academic editor:
Samantha C. Karunarathna
Received: 27 March 2025
Accepted: 10 June 2025
Published: 21 August 2025

Citation: Zhang S, Li J, Jiang H, Ye S, Mapook A, Xu J, Hyde KD, Eungwanichayapant PD (2025) The polyphasic approach reveals two new species and two new records of *Nigrospora* (Apiosporaceae, Amphisphaeriales) associated with *Aquilaria sinensis* from China. MycoKeys 121: 1–20. https://doi.org/10.3897/mycokeys.121.154055

Copyright: © Shiyu Zhang et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Species of *Nigrospora* Zimm. are pathogens, endophytes, or saprobes on various plant hosts in different habitats (Wang et al. 2017). This genus belongs to Apiosporaceae K.D. Hyde, J. Fröhl., Joanne E. Taylor & M.E. Barr (Hyde et al. 2024). *Nigrospora* was introduced by Zimmerman (1902) for *N. panici* Zimm., isolated from the dead leaves of *Panicum amphibium* Steud, in Indonesia. It was known from the asexual characters because of its black opaque conidial spores borne from grey conidiophores. Mason (1927) classified four species occurring on monocotyledons, *viz.*, *N. oryzae* (Berk. & Broome) Petch, *N. arundinacea* (Cooke & Massee) Potl., *N. sacchari* (Speg.) E.W. Mason, and *N. sphaerica* (Sacc.) E.W. Mason, by their black-spored hyphomycetous characters and further indicated *N. javanica* Palm as a synonym of *N. panici*. Species of *Nigrospora* were delimited based on morphological characteristics until it was found that some key morphological characteristics overlapped between phylogenetically distinct species, such as conidial dimensions (Kirk et al. 2008). To address this issue, Wang et al. (2017) analyzed recent species in the genus *Nigrospora* and introduced twelve novel species based on morphological characters, host associations, and a multigene combined phylogeny of ITS, *TEF1-α*, and *TUB2*. They further placed this genus within the Apiosporaceae (Xylariales). Moreover, Hao et al. (2020) reported *N. gorlenkoana* Novobr., *N. oryzae*, *N. osmanthi* Mei Wang & L. Cai, *N. rubi* Mei Wang & L. Cai, and *N. sphaerica* with 13 novel host associations, including economically important crops from China. They confirmed the genus in the family Apiosporaceae based on multi-gene phylogenetic analyses. Liu et al. (2024) isolated *N. anhuiensis* Y. Liu & F. He as a pathogen from rice in Anhui Province, China. Zou et al. (2024) established a novel species, *N. yunnanensis* M.T. Zou & Yong Wang, from *Juglans regia* L. in Yunnan Province, China.

The secondary metabolites of *Nigrospora* species typically exhibit good biological activities and are thus regarded as an interesting source of natural products due to their potential application value. For example, *Nigrospora sphaerica* produces hydroanthraquinones, which exhibit anti-inflammatory activity (Kuang et al. 2022). The metabolites of this species have also shown prominent herbicidal activity in the treatment of intact, greenhouse-grown plants (Fukushima et al. 1998). *Nigrospora sphaerica* and *Nigrospora camelliae-sinensis* Mei Wang & L. Cai have also been proven to be a rich source of other secondary metabolites, such as hydroanthraquinone diketopiperazines, lactones, isochromene, and cyclopentanone derivatives (Metwaly et al. 2014; Zhang et al. 2015; Wu et al. 2018; Zhu et al. 2018; Huang et al. 2021; Kuang et al. 2022). Some have been used in the production of green tea, textile dyes, and food preservatives (Xu et al. 2022).

Aquilaria sinensis (Lour.) Spreng. (Thymelaeaceae) is an important economic and medicinal material in China. The high-value, fragrant, and dark resinous heartwood of *A. sinensis* is known by its popular name "agarwood" (Liu et al. 2017; Wang et al. 2018; Tan et al. 2019; Wang et al. 2019). The damage and infection of the white wood of *A. sinensis* stimulate the plant's defense response, and subsequently, dark resin is produced to suppress further infection, resulting in the formation of agarwood (Chen et al. 2017; Chhipa et al. 2017; Faizal et al. 2020). Fungal inoculation is considered to be effective in the agarwood-producing

industry, and most of the fungi used for inoculation are endophytes from agarwood-producing trees, including *A. sinensis* (Han et al. 2014; Azren et al. 2018; Subasinghe et al.2018; Du et al. 2024). Du et al. (2022b) reviewed the biological activity and agarwood induction potential of endophytic fungi; they indicated that *Nigrospora oryzae* isolated from *A. sinensis* produced biological activity metabolites such as *β*-phenylethyl alcohol, 11-hydroxycapitulatin B, and Capitulatin B (Li et al. 2014), showing that endophytic *Nigrospora* species from *Aquilaria* play a potential application value in the agarwood-producing industry.

During the survey of endophytic fungi, five strains of *Nigrospora* were isolated and examined from *Aquilaria sinensis* in Guangxi Zhuang Autonomous Region, China, for their morphological characteristics. Phylogenetic relationships based on DNA sequence analyses from nucleotides were investigated to evaluate the taxonomic and phylogenetic status of our strains and other *Nigrospora* species using combined ITS, *TEF1-α*, and *TUB2* sequence data. Full descriptions, illustrations, and phylogenetic analysis results for the two new species and two new host records of *Nigrospora* are provided.

Materials and methods

Sampling and isolation of endophytic *Nigrospora* species

Healthy leaf samples of *Aquilaria sinensis* were collected in Qinzhou, Guangxi Zhuang Autonomous Region, China. The samples were transported to the laboratory in self-sealing plastic bags stored in insulated ice boxes(Rathnayaka et al. 2025). Leaf samples were rinsed with tap water to remove soil and debris, then aseptically cut into 0.5 cm × 0.5 cm pieces. Surface sterilization was carried out using 75% ethanol for 30 s, followed by 3% H_2O_2 for 3 min, and another 30 s treatment with 75% ethanol. The samples were then rinsed three times with sterile distilled water and dried on sterile tissue paper (Du et al. 2022a). After drying, four pieces of leaf tissue were placed on potato dextrose agar (PDA) plates and incubated at room temperature (25 °C). Fungal culture characteristics on PDA were observed after one month. Living cultures were deposited in the Kunming Institute of Botany Culture Collection, Kunming, China (KUNCC).

MycoBank numbers and Facesoffungi numbers were obtained as per the instructions in (https://www.mycobank.org/) and Jayasiri et al. (2015), respectively.

DNA extraction, PCR amplification, and sequencing

Fungal mycelia from 30-day-old PDA cultures were scraped off using a sterilized scalpel and transferred into 1.5 mL centrifuge tubes. Genomic DNA was extracted using the Biospin Fungus Genomic DNA Extraction Kit-BSC14S1 (BioFlux®, P.R. China), following the manufacturer's instructions.

DNA amplification was performed using polymerase chain reaction (PCR) targeting three loci: ITS, *TEF1-α*, and *TUB2*.

The internal transcribed spacer (ITS) region was amplified using the primer pair ITS1 and ITS4 (White et al. 1990). The partial translation elongation factor 1-alpha (*TEF1-α*) region was amplified using EF1-728F and EF2 primers (Carbone and Kohn 1999; O'Donnell et al. 1998). The beta-tubulin (*TUB2*) fragment was amplified using Bt-2a and Bt-2b (Glass and Donaldson 1995).

The PCR reaction volume was 25 μL, containing 1 μL forward primer, 1 μL reverse primer, 22 μL JPmix (Tsingke Biotech Co., Ltd.), and 1 μL of template DNA (48 ng/μL). PCR amplification conditions were as follows: initial denaturation at 98 °C for 2 min, followed by 32 cycles of denaturation at 98 °C for 15 s, annealing at 53 °C for 15 s, extension at 72 °C for 45 s, and a final extension at 72 °C for 5 min. Purification and sequencing of PCR products were carried out by Tsingke Biotech Co., Ltd. (Kunming, P.R. China). The DNA product was stored at -20 °C for long-term preservation.

Phylogenetic analyses

Phylogenetic analyses were performed following the procedures described in Dissanayake et al. (2020). Sequence quality was assessed using SnapGene v6.1. A BLAST search based on ITS, *TEF1-α*, and *TUB2* sequences was conducted to identify closely related taxa in GenBank (https://www.ncbi.nlm.nih.gov/genbank/), and corresponding sequences were downloaded. *Apiospora obovata* (LC4940) and *Ap. malaysiana* (CBS 102053) were selected as outgroup taxa (Table 1). Sequences were aligned in MEGA v7.0 using MUSCLE, with manual adjustments where necessary (Kumar et al. 2016). Phylogenetic trees were generated using both maximum likelihood (ML) and Bayesian inference (BI) methods via the CIPRES Science Gateway (Miller et al. 2010). The ML analysis was conducted using the GTRGAMMA model with 1000 bootstrap replicates. The BI analysis employed the Invgamma+5 model with 100,000 generations. Resulting trees were visualized in FigTree v1.4.2 (http://tree.bio.ed.ac.uk/software/figtree/), with bootstrap support values indicated at the nodes. The phylogram was rendered in TreeView (Page 1996), illustrated in Microsoft PowerPoint, and exported as a JPEG image using Adobe Photoshop CS5 (Adobe Systems Inc., United States). Newly generated sequences were deposited in GenBank (Table 1). Sequence alignments were submitted to TreeBASE under accession number S31751 (https://treebase.org/).

Results

Phylogenetic analyses

The combined ITS, *TEF1-α*, and *TUB2* sequence dataset comprised 77 taxa, with *Apiospora obovata* (LC4940) and *Ap. malaysiana* (CBS 102053) selected as outgroup taxa.

Bayesian inference (BI) and maximum likelihood (ML) analyses of the combined dataset were conducted to determine the placement of our new taxa, infer relationships at the intrageneric level, and resolve phylogenetic relationships within Apiosporaceae. The phylogenetic trees derived from BI and ML analyses exhibited largely similar topologies and were consistent with previous studies based on ML analysis (Li et al. 2017; Su et al. 2018). The best-scoring RAxML tree is shown in Fig. 1, with a final ML optimization likelihood value of −10322.409322 (ln). The dataset consisted of 1,595 total characters, including gaps (ITS: 1–591 bp; *TEF1-α*: 592–1170 bp; *TUB2*: 1171–1595 bp). RAxML analysis yielded 1,595 distinct alignment patterns, with 33.97% of characters undetermined or containing gaps.

Table 1. Names, culture collection numbers, hosts, and corresponding GenBank accession numbers of the taxa used in the phylogenetic analyses.

Taxa names	Culture collection number	Host	ITS	*TUB2*	*TEF1-α*
Apiospora obovata	LC4940	–	KY494696	KY705166	KY705095
Ap. malaysiana	CBS 102053	–	NR120273	KF144988	KF145030
Nigrospora aurantiaca(T)	CGMCC3.18130 = LC7302	*Nelumbo* sp. (leaf)	KX986064	KY019465	KY019295
N. aurantiaca	LC7034	*Musa paradisiaca*	KX986093	KY019598	KY019394
N. bambusae(T)	CGMCC3.18327 = LC7114	Bamboo (leaf)	KY385307	KY385319	KY385313
N. bambusae	LC7244	Bamboo (leaf)	KY385306	KY385320	KY385314
N. bambusae	LC7245	Bamboo (leaf)	KY385305	KY385321	KY385315
N. camelliae-sinensis	LC2710	*Castanopsis* sp.	KX985957	KY019484	KY019310
N. camelliae-sinensis	LC3287	*Camellia sinensis*	KX985975	KY019502	KY019323
N. camelliae-sinensis	LC3496	*Camellia sinensis*	KX985985	KY019510	KY019327
N. camelliae-sinensis(T)	CGMCC3.18125 = LC3500	*Camellia sinensis*	KX985986	KY019460	KY019293
N. camelliae-sinensis	LC6684	*Camellia sinensis*	KX986046	KY019570	KY019449
N. camelliae-sinensis	**KUNCC23-16748**	***Aquilaria sinensis***	**PQ553689**	**PQ613611**	**PQ613606**
N. chinensis	LC2696	*Lindera aggregata*	KX985947	KY019474	KY019424
N. chinensis	LC3493	*Camellia sinensis*	KX985984	KY019509	KY019434
N. chinensis	LC4433	*Castanopsis* sp.	KX986013	KY019536	KY019436
N. chinensis	LC4558	Unknown host plant	KX986020	KY019543	KY019441
N. chinensis(T)	CGMCC3.18127 = LC4575	*Machilus breviflora*	KX986023	KY019462	KY019422
N. chinensis	LC4660	–	–	KY019548	KY019445
N. chinensis	LC6631	*Camellia sinensis*	KX986043	KY019569	KY019448
N. chinensis	LC6851	Unknown host plant	KX986049	KY019579	KY019450
N. gorlenkoana(T)	CBS 480.73	*Vitis vinifera*	KX986048	KY019456	KY019420
N. gorlenkoana	JZB3230001	*Cirsium setosum*	MN495939	MN549381	MN54464
N. guangxiensis	**KUNCC23-16747**	***Aquilaria sinensis***	**PQ553688**	**PQ613610**	**PQ613605**
N. guilinensis	LC7301	*Vitis vinifera*	KX986063	KY019608	KY019404
N. guilinensis(T)	CGMCC3.18124 = LC3481	*Nelumbo* sp. (stem)	KX985983	KY019459	KY019292
N. hainanensis(T)	CGMCC3.18129 = LC7030	*Musa paradisiaca* (leaf)	KX986091	KY019464	KY019415
N. hainanensis	LC6979	*Musa paradisiaca* (leaf)	KX986079	KY019586	KY019416
N. hainanensis	LC7031	*Musa paradisiaca*	KX986092	KY019597	KY019417
N. hainanensis	LC7042	*Musa paradisiaca* (leaf)	KX986094	KY019599	KY019418
N. lacticolonia(T)	CGMCC3.18123 = LC3324	*Camellia sinensis*	KX985978	KY019458	KY019291
N. lacticolonia	LC7009	*Musa paradisiaca* (leaf)	KX986087	KY019594	KY019454
N. musae(T)	CBS 319.34	*Musa paradisiaca* (fruit)	KX986076	KY019455	KY019419
N. musae	LC6385	*Camellia sinensis*	KX986042	KY019567	KY019371
N. oryzae	LC6761	*Oryza sativa*	KX986056	KY019574	KY019376
N. oryzae	LC7297	*Nelumbo* sp. (leaf)	KX985936	KY019605	KY019400
N. oryzae	LC2693	*Neolitsea* sp.	KX985944	KY019471	KY019299
N. oryzae	LC2707	*Rhododendron simiarum*	KX985954	KY019481	KY019307
N. oryzae	LC4338	*Camellia* sp.	KX986008	KY019532	KY019349
N. oryzae	LC4961	*Pittosporum illicioides*	KX986031	KY019553	KY019358
N. oryzae	LC5243	Submerged wood	KX986033	KY019555	KY019360
N. oryzae	LC6923	*Oryza sativa* L.	KX986051	KY019581	KY019383
N. oryzae	JZB3230002	*Phyllostachys nigra*	MN495940	–	MN544639
N. oryzae	JZB3230003	*Rudbeckia hirta*	MN495941	–	MN544640
N. oryzae	JZB3230004	*Scirpus* sp.	MN495942	MN549382	MN544641
N. oryzae	**KUNCC23-16746**	***Aquilaria sinensis***	**PQ553687**	**PQ613609**	**PQ613604**
N. osmanthi(T)	CGMCC3.18126 = LC4350	*Osmanthus* sp.	KX986010	KY019461	KY019421
N. osmanthi	LC4487	*Hedera nepalensis*	KX986017	KY019540	KY019438
N. osmanthi	JZB3230005	*Rosa chinensis*	MN495943	MN549383	MN508179
N. osmanthi	JZB3230006	*Rosa chinensis*	MN495944	MN549384	MN508180

Taxa names	Culture collection number	Host	ITS	*TUB2*	*TEF1-α*
N. osmanthi	JZB3230007	*Phragmites australis*	MN495945	MN549385	MN508181
N. osmanthi	JZB3230008	*Cirsium setosum*	MN495946	MN549386	MN508182
N. osmanthi	JZB3230009	*Phyllostachys nigra*	MN495947	MN549387	MN508183
N. osmanthi	JZB3230010	*Phyllostachys nigra*	MN495948	MN549388	MN508184
N. osmanthi	JZB3230011	*Rudbeckia hirta*	MN495949	MN549389	MN508185
N. pubeiensis	**KUNCC23-16745**	***Aquilaria sinensis***	**PQ553686**	**PQ613608**	**PQ613603**
N. pyriformis(T)	CGMCC3.18122 = LC2045	*Citrus sinensis*	KX985940	KY019457	KY019290
N. pyriformis	LC2688	*Lindera aggregata*	KX985941	KY019468	KY019297
N. pyriformis	LC2694	*Rubus reflexus*	KX985945	KY019472	KY019300
N. pyriformis	LC3099	*Camellia sinensis*	KX985971	KY019498	KY019322
N. pyriformis	LC3292	*Camellia sinensis*	KX985976	KY019503	KY019324
N. rubi(T)	CGMCC3.18326 = LC2698	*Rubus* sp.	KX985948	KY019475	KY019302
N. rubi	JZB3230012	*Fraxinus* sp.	MN495950	–	MN54464
N. sphaerica	LC7312	*Nelumbo* sp. (leaf)	KX985935	KY019618	KY019414
N. sphaerica	LC7298	*Nelumbo* sp. (leaf)	KX985937	KY019606	KY019401
N. sphaerica	LC2840	*Harpullia longipetala*	KX985965	KY019492	KY019318
N. sphaerica	LC3477	*Camellia sinensis*	KX985982	KY019508	KY019326
N. sphaerica	LC4264	*Rhododendron arboretum*	KX985993	KY019517	KY019334
N. sphaerica	LC4307	*Rhododendron arboretum*	KX986005	KY019529	KY019346
N. sphaerica	LC5901	Submerged wood	KX986034	KY019556	KY019361
N. sphaerica	LC6294	*Camellia sinensis*	KX986044	KY019565	KY019369
N. sphaerica	LC6996	*Musa paradisiaca* (leaf)	KX986085	KY019592	KY019390
N. sphaerica	JZB3230013	*Cirsium setosum*	MN495951	MN549390	MN544642
N. sphaerica	JZB3230014	*Phragmites australis*	MN495952	MN549391	MN544643
N. sphaerica	JZB3230015	*Fraxinus* sp.	MN495953	MN549392	MN544644
Nigrospora sp. 1	LC2725	*Symplocos zizyphoides*	KX985960	KY019487	KY019313
Nigrospora sp. 1	LC4566	*Lithocarpus* sp.	KX986022	KY019545	KY019354
Nigrospora sp. 2	LC6704	*Camellia sinensis*	KX986047	KY019571	KY019373
N. vesicularis	LC0322	Unknown	KX985939	KY019467	KY019296
N. vesicularis(T)	CGMCC3.18128 = LC7010	*Musa paradisiaca* (leaf)	KX986088	KY019463	KY019294
N. zimmermanii	CBS 167.26	Unknown	KY385308	KY385318	KY385312
N. zimmermanii(T)	CBS 290.62	*Saccharum officinarum* (leaf)	KY385309	KY385317	KY385311
N. zimmermanii	CBS 984.69	*Saccharum officinarum* (leaf)	KY385310	KY385322	KY385316
Nigrospora sp.	YL-2024a	–	OP677969	PP103614	PP103590
N. shadeganensis	SCUA-Saf-N27	–	PP256499	PP263821	PP263812
N. coryli	W18	*Corylus heterophylla*	PP218065	PP218065	PP461302
N. humicola	CFCC 56884	Soil	ON555686	ON557392	ON557394
N. yunnanensis	GUCC24-0008	–	PP915796	PP947937	PP947933

– Refers to the data unavailability, (T)refers to the type specimens, and the taxa produced in this study are in bold.

Nigrospora camelliae-sinensis **Mei Wang & L. Cai, in Wang, Liu, Crous & Cai, Persoonia 39: 127 (2017)**

MycoBank No: 820731

Fig. 2

Description. ***Endophytic*** from healthy leaves of *Aquilaria sinensis*. Sexual morph: Undetermined. Asexual morph: ***Hyphae*** branched, septate, hyaline, or light grey, 2–4 µm diam. ***Conidiophores*** 3–6 µm diam. micronematous, mononematous, solitary, smooth, branched or not, 0–1 septate, hyaline to subhyaline, consisting of 1–2 cells or usually reduced to conidiogenous cells. ***Conidiogenous cells*** 5.5–7.2 × 6.3–9.4 µm diam. $\bar{x}$ = 5.8 × 12.1 µm, n = 30), monoblastic, discrete, determinate, smooth, subglobose to ampulliform, subhyaline to light brown.




Figure 1. Tree generated by RAxML analysis showing the placement of *Nigrospora* species based on combined ITS, *TEF1-α*, and *TUB2* sequence data. Bootstrap support values for ML (left) ≥ 75% and Bayesian posterior probabilities (PP, right) ≥ 0.9 are indicated at the nodes. The tree is rooted with *Ap. obovata* (LC4940) and *Ap. malaysiana* (CBS 102053). Strains generated in this study are shown in red and blue bold; red indicates new species, and blue indicates new host records.

Conidia 8–13 µm diam. $\bar{x}$ = 11 µm, n = 30) solitary, smooth-walled, aseptate, spherical, light grey to black, globose or subglobose.

Cultural characteristics. Mycelium effusing on PDA within 12 hours from the edges of the surface sterilized leaf tissue piece. Colonies growing on PDA, hairy, black, reaching 9 cm in 7 days at 30 °C; mycelium partly superficial, partly immersed, slightly effuse, radially striate, with an irregular edge, black. Asexual conidia spores were formed after 25 days on PDA. Sexual spores were not formed within 60 days.

Material examined. China • Guangxi Zhuang Autonomous Region, Pubei City, in healthy living leaves of *Aquilaria sinensis*, September 25, 2020, Shiyu Zhang, GX9-3 (HKAS 134953, new host record**)**. Living culture KUNCC23-16748.

Known distributions (based on molecular data): China (Wang et al. 2017, this study).

Known hosts (based on molecular data): *Aquilaria sinensis* (this study), *Camellia sinensis* (Wang et al. 2017), *Castanopsis* sp. (Wang et al. 2017), *Musa paradisiaca* (Wang et al. 2017).




Figure 2. *Nigrospora camelliae-sinensis* (HKAS–134953, new host record). **a.** Culture on PDA for 30 days; **b–d.** Mass of conidia; **e–j.** Conidiogenous cells attached conidia; **k.** Conidiogenous cell arising from hyphae; **l–o.** Conidiophores bearing conidia; **p–t.** Conidia. Scale bars: 10 μm (**b–t**).

Notes. *Nigrospora camelliae-sinensis* has been reported to have a cosmopolitan distribution and a broad host range (Wang et al. 2017). In this study, our isolation (KUNCC23-16748), collected from *Aquilaria sinensis* in Guangxi Zhuang Autonomous Region, China, clustered in one single clade with *N. camelliae-sinensis* (94% ML, 1PP). Its morphological characteristics were in good agreement with those of *N. camelliae-sinensis*. Therefore, we regard this isolation (KUNCC23-16748)

as *N. camelliae-sinensis* collected from *Aquilaria sinensis* for the first time. The study showed that *N. camelliae-sinensis* is also characterized by having determinate and micro- and mononematous, subcylindrical conidiophores.

Nigrospora guangxiensis **S.Y. Zhang, J.F. Li & K.D. Hyde, sp. nov.**
MycoBank No: 859462
Facesoffungi Number: FoF17621
Fig. 3

Etymology. Named after the type location, Guangxi Zhuang Autonomous Region, China.

Holotype. HKAS 134952.




Figure 3. *Nigrospora guangxiensis* (HKAS–134952, holotype). **a.** Culture on PDA for 30 days; **b, c.** Mass of conidia; **d–g.** Conidiogenous cells attached conidia; **h, i.** Conidiophores; **j–s.** Conidia. Scale bars: 10 μm (**b–s**).

Table 2. Base pair comparison in ITS, *TEF1-α*, and *TUB2* sequences of *N. chinensis*, *N. pubeiensis*, and *N. guangxiensis*.

Species	Base pair comparison in ITS, *TUB2*, and *TEF1-α*		
	ITS	*TUB2*	*TEF1-α*
N. chinensis (CGMCC-38127) *vs. N. pubeiensis* (KUNCC23-16745)	3/492(0.6%)	26/365(7.1%)	0
N. chinensis (CGMCC-38127) *vs. N. guangxiensis* (KUNCC23-16747)	10/492(2.0%)	26/373(7.0%)	62/425(14.5%)
N. guangxiensis (KUNCC23-16747) *vs. N. pubeiensis* (KUNCC23-16745)	9/511(1.8%)	0	62/425(14.5%)

Description. ***Endophytic*** from healthy leaves of *Aquilaria sinensis*. Sexual morph: Undetermined. Asexual morph: ***Hyphae*** branched, septate, hyaline to pale brown, 1–3 µm diam. ***Conidiophores*** usually reduced to conidiogenous cells, which are dispersed on hyphae. ***Conidiogenous cells*** 4.5–6 × 5.3–7.3 µm diam. $\bar{x}$ = 5 × 6.8 µm, n = 30), discrete, solitary, monoblastic, determinate, subglobose, pale brown to brown colored. *Conidia* 8–14 µm diam. $\bar{x}$ = 10.5 µm, n = 30), solitary, discrete, dark brown to black, globose or subglobose.

Cultural characteristics. Mycelium effusing on PDA within 12 hours from the edges of the surface sterilized leaf tissue piece. Colonies growing on PDA, hairy, reaching 9 cm in 7 days at 30 °C; mycelium partly superficial, partly immersed, slightly effuse, radially striate, with irregular edge, initially greyish-brown, becoming black colored with age; asexual spores were formed after 25 days on PDA, and sexual spores not formed within 60 days on PDA.

Known distribution (based on molecular data): China (this study).

Known hosts. *Aquilaria sinensis* (this study).

Material examined. **China** • Guangxi Zhuang Autonomous Region, Pubei city, on healthy living leaves of *Aquilaria sinensis*, September 25, 2020, Shiyu Zhang, GX4-1 (HKAS 134952, holotype**)**; extype living culture, KUNCC23-16747.

Notes. In the multi-locus phylogenetic analyses, our strain *Nigrospora guangxiensis* (KUNCC23-16747) formed a separate branch basal (97% ML, 1PP) to *N. pubeiensis* (KUNCC23-16745) and *N. chinensis* (eight strains). *Nigrospora guangxiensis* differs from *N. pubeiensis* in having smaller conidiogenous cells (5 × 6.8 µm *vs.* 5.8 × 9.4 µm) and shorter conidia (10.5 µm vs. 11.3 µm). Additionally, its initial culture on PDA medium is greyish-brown. *Nigrospora guangxiensis* is distinct from *N. chinensis* in having smaller conidia (10.5 µm vs. 12.97 ± 1.07 µm), pale brown to brown-colored conidiogenous cells, and sterile cells, which are absent in *N. guangxiensis* (Wang et al. 2017). A nucleotide base comparison of these species is shown in the notes of *N. pubeiensis* and Table 2.

Nigrospora oryzae **(Berk. & Broome) Petch, J. Indian bot. Soc. 4: 24 (1924)**

MycoBank No: 253729

Facesoffungi Number: FoF06596

Fig. 4

Monotospora oryzae Berk. & Broome, J. Linn. Soc., Bot. 14: 99. 1873. Basionym.
Khuskia oryzae H.J. Huds., Trans. Brit. Mycol. Soc. 46: 358. 1963. Synonym.

Description. ***Endophytic*** from healthy leaves of *Aquilaria sinensis*. Sexual state: Undetermined. Asexual state: Hyphomycetes. ***Hyphae*** 1–3.5 µm diam. branched, smooth, septate, hyaline, or subhyaline. ***Conidiophores*** 2–5 µm diam. micronematous, solitary in sporodochia, subcylindrical, pale grey, smooth,




Figure 4. *Nigrospora oryzae* (HKAS—134951), new host and geographical record. **a.** Culture on PDA for 30 days; **b, c.** Mass of conidia; **d, e.** Conidiogenous cells attached conidia; **f–h.** Conidiophores bearing conidia; **i–r.** Conidia. Scale bars: 10 μm (**a–r**).

0–2-septate, branched or not, usually reduced to conidiogenous cells. ***Conidiogenous cells*** 4.7–5.2 × 6.3–7.4 μm diam. $\bar{x}$ = 5.0 × 6.9 μm, n = 50), monoblastic, solitary, determined, hyaline to subhyaline, ampliform to subspherical, discrete, smooth-walled. ***Conidia*** 8–13.5 μm diam $\bar{x}$ = 11.3 μm, n = 50) solitary, globose or subglobose, black, shiny, smooth, aseptate.

Cultural characteristics. Mycelium effusing on PDA within 12 hours from the edges of the surface-sterilized leaf tissue piece. Colonies growing on PDA, hairy, black, reaching 5 cm in 7 days at 30 °C; mycelium partly superficial, partly

immersed, slightly effuse, radially striate, with an irregular edge, dark brown to black colored; asexual spores were formed after 25 days on PDA, and sexual spores were not formed within 60 days on PDA.

Material examined. China • Guangxi Zhuang Autonomous Region, Pubei city, on healthy living leaves of *Aquilaria sinensis*, September 25, 2020, Shiyu Zhang, GX2-3 (HKAS 134951, new host record**)**. Living culture KUNCC23-16746.

Known distribution (based on molecular data): Australia (Barkat et al. 2016), Bangladesh (Begum et al. 2018), China (Sun et al. 2011; Chen and Kirschner2018; He et al. 2019; Zhang et al. 2019, this study), Iran (Hashemian et al. 2014), Italy (Lorenzini et al. 2016), Kazakhstan (Eken et al. 2016), Pakistan (Alam et al. 2017), Sri Lanka (Wang et al. 2017), and Thailand (Hyde et al. 2020).

Known hosts (based on molecular data): *Acer truncatum* (Sun et al. 2011), *Actinidia* sp. (Li et al. 2018), *Aloe vera* (Alam et al. 2017), *Aquilaria sinensis* (This study), *Castanopsis* sp. (Wang et al. 2017), *Cephalotaxus sinensis* (Wang et al. 2017), *Citrullus lanatus* (Chen and Kirschner 2018), *Citrus reticulata* (Wang et al. 2017), *Cleyera japonica* (Wang et al. 2017), *Daphniphyllum macropodum* (Wang et al. 2017), *Daphniphyllum oldhamii* (Wang et al. 2017), *Gossypium hirsutum* (Zhang et al. 2019), *Hamamelis mollis* (Wang et al. 2017), *Nelumbo nucifera* (Chen and Kirschner 2018), *Nelumbo* sp. (Wang et al. 2017), *Neolitsea* sp. (Wang et al. 2017), *Oryza sativa* (Wang et al. 2017), *Osmanthus fragrans* (Wang et al. 2017), *Osmanthus* sp. (Wang et al. 2017), *Pennisetum americanum* (Hashemian et al. 2014), *Pentactina rupicola* (Wang et al. 2017), *Photinia serrulata* (He et al. 2019), *Phyllostachys heterocycla* (Hyde et al. 2020), *Rhododendron simiarum* (Wang et al. 2017), *Rhododendron* sp. (Wang et al. 2017), *Rubus reflexus* (Wang et al. 2017), *Rubus* sp. (Wang et al. 2017), *Symplocos zizyphoides* (Wang et al. 2017), *Ternstroemia* sp. (Wang et al. 2017), *Triticum aestivum* (Eken et al. 2016; Barkat et al. 2016), *Tutcheria microcarpa* (Wang et al. 2017), *Vaccinium corymbosum* (Zhang et al. 2019), *Vitis vinifera* (Lorenzini et al. 2016).

Notes. *Nigrospora oryzae* has been reported to have a cosmopolitan distribution and a broad host range (Wang et al. 2017). In this study, our isolate (KUNCC23-16746) clustered in the same clade (100% ML, 1 PP) as *N. oryzae*, and its morphological characteristics were consistent with those of *N. oryzae* (Wang et al. 2017). Therefore, we regard this isolate (KUNCC23-16746) as *N. oryzae*, which was collected from *Aquilaria sinensis* in the Guangxi Zhuang Autonomous Region, China, for the first time.

Nigrospora pubeiensis **S.Y. Zhang, J.F. Li & K.D. Hyde, sp. nov.**

MycoBank No: 859460

Facesoffungi Number: FoF17622

Fig. 5

Etymology. Named after the location from which it was collected, Pubei, Guangxi Zhuang Autonomous Region, China.

Holotype. HKAS 134950.

Description. ***Endophytic*** from healthy leaves of *Aquilaria sinensis*. Sexual morph: Undetermined. ***Asexual morph*: *Hyphae*** 3–6 µm diam. branched, smooth, septate, hyaline, or subhyaline. ***Conidiophores*** 1–4 µm diam, micronematous, solitary,




Figure 5. *Nigrospora pubeiensis* (HKAS–134950, holotype). **a.** Culture on PDA for 30 days; **b.** Mass of conidia; **c, d.** Conidiogenous cells attached conidia; **e–i.** Conidiophores; **j.** Conidiophores bearing conidia; **k–t.** Conidia. Scale bars: 10 μm (**b–t**).

subcylindrical, hyaline to pale grey, smooth, 0–2-septate, branched or not, usually reduced to conidiogenous cells. ***Conidiogenous cells*** 5–6.3 × 6.3–11.3 μm diam. ($\bar{x}$ = 5.8 × 9.4 μm, n = 20), discrete, solitary, monoblastic, determinate, subglobose, straight, smooth, hyaline or dark brown. ***Conidia*** 6–14 μm diam ($\bar{x}$ = 11.3 μm, n = 20), solitary, granular, dark brown to black, globose, or subglobose.

Cultural characteristics. Mycelium effusing on PDA within 12 hours from the edges of the surface-sterilized leaf tissue piece. Colonies growing on PDA, hairy, black, reaching 9 cm in 7 days at 30 °C; mycelium partly superficial, partly immersed, slightly effuse, radially striate, with irregular edge, initially grayish white, becoming black colored with age; asexual spores were formed after 25 days on PDA, and sexual spores not formed within 60 days on PDA.

Known distribution (based on molecular data): China (this study).

Known hosts. *Aquilaria sinensis* (this study).

Material examined. China • Guangxi Zhuang Autonomous Region, Pubei City, in healthy living leaves of *Aquilaria sinensis*, September 25, 2020, Shiyu Zhang, GX2-2 (HKAS 134950, ***holotype***). Ex-type living culture at KUNCC23-16745.

Notes. *Nigrospora pubeiensis* is described herein as a new species, which resembles other species in the genus in having granular, dark brown to black, globose, or subglobose conidia, but differs in its short conidiophores, sometimes reduced to conidiogenous cells. The strain representing *N. pubeiensis* (KUNCC23-16745) clustered in a supported clade (100% ML, 1 PP) and is closely related to *N. guangxiensis* (KUNCC23-16747) and *N. chinensis* (CGMCC-38127). However, *N. pubeiensis* is distinct in nucleotide base pair comparison with *N. guangxiensis* (KUNCC23-16747) across the ITS gene region (9/511 bp, 1.8% difference, no gap) and the *TEF1-α* gene region (62/425 bp, 14.5% difference, no gap), *TUB2* (0); it differs with *N. chinensis* (CGMCC-38127) across the ITS gene region (3/492 bp, 0.6% difference, no gap) and the *TUB2* gene region (26/365 bp, 7.1% difference, no gap), *TEF1-α* (0). They also have different conidiophore structures. Compared with the photoplate from Wang et al. (2017), the conidiophore of *N. chinensis* has an ampulliform structure, but *N. pubeiensis* does not. Based on distinct morphological characteristics and phylogenetic support, we regard this isolation (KUNCC23-16745) as a new species, *N. pubeiensis*, which was collected from *Aquilaria sinensis* in Guangxi Province, China.

Discussion

The discovery of two novel *Nigrospora* species (*N. pubeiensis* and *N. guangxiensis*), along with new host records for *N. oryzae* and *N. camelliae-sinensis*, highlights the adaptability of *Nigrospora* to diverse ecological niches. The broad host range of *Nigrospora* on *Aquilaria* spp. may be linked to its ability to exploit host-specific chemical environments. For instance, endophytic fungi in *A. sinensis* often interact with host-derived secondary metabolites, such as sesquiterpenes and chromones, which are abundant in agarwood resin (Li et al. 2025). These compounds may create selective pressures that favor fungal lineages capable of tolerating or metabolizing them (Li et al. 2025). Evidence from other studies shows that *Nigrospora* species produce bioactive metabolites (e.g., antimicrobial compounds), which could confer competitive advantages in colonizing new hosts or resisting host defenses (Metwaly et al. 2014; Kuang et al. 2022).

Ecologically, the occurrence of *Nigrospora* on *A. sinensis* aligns with observations that endophytic fungal communities shift dynamically during agarwood formation. For example, fungal diversity decreases as resinous compounds accumulate, suggesting a trade-off between host defense and fungal survival (Li et al. 2025). Similar patterns have been observed in *Fusarium*-induced agarwood, where dominant fungi such as *Colletotrichum* thrive in resin-rich tissues (Fang et al. 2024). This implies that *Nigrospora* may occupy niches where host metabolites are less inhibitory or where their secondary metabolites counteract host defenses.

Evolutionarily, the divergence of *Nigrospora* lineages may reflect co-adaptation with *Aquilaria* hosts. Multi-gene phylogenies (e.g., ITS, *TEF1*, and *TUB2*) reveal that fungal speciation often correlates with host specificity (Tibpromma et al. 2021). For instance, *Nemania* species isolated from *A. sinensis* produce sesquiterpenoids identical to those found in agarwood, suggesting a symbiotic relationship in which the fungi contribute to resin biosynthesis (Tibpromma et al. 2021). Analogously, *Nigrospora* may have evolved metabolic pathways that enable coexistence with *Aquilaria*'s chemical defenses, thus facilitating colonization across diverse hosts.

Acknowledgments

Li Junfu thanks the Yunnan Provincial Department of Human Resources and Social Security and the Yunnan Provincial Postdoctoral Orienteering Training and Funding Program (E33O386261), entitled "Investigation of Fungal Diversity and Fungal Resources in the Dry Valley Area of Honghe, Yunnan Province." We thank two anonymous reviewers and Prof. Saowaluck Tibpromma for their valuable suggestions to improve the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

No use of AI was reported.

Funding

The research is supported by the Yunnan Department of Science and Technology of China (Grant Nos: 202302AE090023 and 202303AP140001).

Author contributions

Shiyu Zhang, Prapassorn Damrongkool Eungwanichayapant and Junfu Li conceived the study. Shiyu Zhang, Junfu Li and Shuang Ye performed the experiments. Shiyu Zhang, Hongbo Jiang and Junfu Li analyzed the data. Shiyu Zhang and Junfu Li drafted the manuscript. Ausana Mapook, Jianchu Xu, Kevin D. Hyde and Prapassorn Damrongkool Eungwanichayapant reviewed and edited the manuscript. All authors contributed to the article and approved the submitted version.

Author ORCIDs

Shiyu Zhang https://orcid.org/0009-0001-4174-6103
Hongbo Jiang https://orcid.org/0000-0001-9655-4662
Kevin D. Hyde https://orcid.org/0000-0002-2191-0762
Prapassorn Damrongkool Eungwanichayapant https://orcid.org/0000-0001-8005-4137

Data availability

All of the data that support the findings of this study are available in the main text.

References

Alam MW, Rehman A, Saira M, Kan NA, Aslam S, Fiaz M, Muhammad S (2017) First report of leaf spots in *Aloe vera* caused by *Nigrospora oryzae* in Pakistan. Plant Disease 101(5): 841. https://doi.org/10.1094/PDIS-11-16-1684-PDN

Azren PD, Lee SY, Emang D, Mohamed R (2018) History and perspectives of induction technology for agarwood production from cultivated *Aquilaria* in Asia: A review. Journal of Forestry Research 30(1): 1–11. https://doi.org/10.1007/s11676-018-0627-4

Barkat EH, St J, Hardy GE, Ren Y, Calver M, Bayliss KL (2016) Fungal contaminants of stored wheat vary between Australian states. Australas. Australasian Plant Pathology 45(6): 621–628. https://doi.org/10.1007/s13313-016-0449-9

Begum M, Hamza A, Tanny T, Das KC, Mahmud MT, Salimullah M, Alam I (2018) First report of leaf spot disease in Aloe vera caused by *Nigrospora oryzae* in Bangladesh. Plant disease 102(7): 1461. https://doi.org/10.1094/PDIS-11-17-1857-PDN

Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. Mycologia 3: 553–556. https://doi.org/10.1080/00275514.1999.12061051

Chen KL, Kirschner R (2018) Fungi from leaves of lotus (*Nelumbo nucifera*). Mycological Progress 17(1–2): 275–293. https://doi.org/10.1007/s11557-017-1324-y

Chen XY, Sui C, Liu YY, Yang Y, Liu PW, Zhang Z, Wei JH (2017) Agarwood formation induced by fermentation liquid of *Lasiodiplodia theobromae*, the dominating fungus in wounded wood of *Aquilaria sinensis*. Current Microbiology 74(4): 460–468. https://doi.org/10.1007/s00284-016-1193-7

Chhipa H, Chowdhary K, Kaushik N (2017) Artificial production of agarwood oil in *Aquilaria* sp. by fungi: A review. Phytochemistry Reviews 16(5): 835–860. https://doi.org/10.1007/s11101-017-9492-6

Dissanayake AJ, Bhunjun CS, Maharachchikumbura SSN, Liu JK (2020) Applied aspects of methods to infer phylogenetic relationships amongst fungi. Mycosphere 11(1): 2652–2676. https://doi.org/10.5943/mycosphere/11/1/18

Du TY, Karunarathna SC, Hyde KD, Mapook A, Wariss HM, Aluthwattha ST, Wang YH, Mortimer PE, Xu JC, Tibpromma S (2022a) The endophytic fungi of *Aquilaria sinensis* from southern China. Fungal Biotec 2: 1–15.

Du TY, Dao CJ, Mapook A, Stephenson SL, Elgorban AM, Al-Rejaie S, Suwannarach N, Karunarathna SC, Tibpromma S (2022b) Diversity and biosynthetic activities of agarwood associated fungi. Diversity 14(3): 211. https://doi.org/10.3390/d14030211

Du TY, Tibpromma S, Hyde KD, Wang YH, Chomnunti P, Chukeatirote E, Lu WH, Mapook A, Dai DQ, Wijesundara DSA, Elgorban AM, Suwannarach N, Kumla J, Moussa IM, Wang HH, Karunarathna SC (2024) The fungal strain promotes rapid agarwood resin production with medicinally accepted agarotetrol level. Food Bioscience 62: 105535. https://doi.org/10.1016/j.fbio.2024.105535

Eken C, Spanbayev A, Tulegenova Z, Yechshzhanov T (2016) First report of *Nigrospora oryzae* on wheat in Kazakhstan. Plant Disease 100(4): 861. https://doi.org/10.1094/PDIS-08-15-0915-PDN

Faizal A, Azar AWP, Turjaman M, Esyanti RR (2020) *Fusarium solani* induces the formation of agarwood in *Gyrinops versteegii* (Gilg.) Domke branches. Symbiosis 81(1): 15–23. https://doi.org/10.1007/s13199-020-00677-w

Fang X, Li X, Zhang Q, Hu H, Hong Z, Liu X, Cui Z, Xu D (2024) Physiological and endophytic fungi changes in grafting seedlings of Qi-Nan clones (*Aquilaria sinensis*). Forests 15: 106. https://doi.org/10.3390/f15010106

Fukushima T, Tanaka M, Gohbara M, Fujimori T (1998) Phytotoxicity of three lactones sacchari from *Nigrospora sacchari*. Phytochemistry 48(4): 625–630. https://doi.org/10.1016/S0031-9422(97)01023-6

Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. Applied and Environmental Microbiology 61: 1323–1330. https://doi.org/10.1128/aem.61.4.1323-1330.1995

Han XM, Liang L, Zhang Z, Li XJ, Yang Y, Meng H, Gao ZH, Xu YH (2014) Study of production of sesquiterpenes of *Aquilaria sinensis* stimulated by *Lasiodiplodia theobromae*. Zhongguo Zhongyao Zazhi 39: 192–196. [In Chinese]

Hao YY, Aluthmuhandiram Janith VS, Chethana KWT, Manawasinghe IS, Li XH, Liu M, Hyde KD, Phillips AJL, Zhang W (2020) *Nigrospora* species associated with various hosts from Shandong Peninsula, China. Mycobiology 48(3): 169–183. https://doi.org/10.1080/12298093.2020.1761747

Hashemian KT, Jahani M, Zare R, Mirzaee MR (2014) First report of *Nigrospora* leaf spot on *Pennisetum americanum*. Iranian Journal of Plant Pathology 96(3): 606.

He BY, Cernava T, He HD, Li HX, Chen XY, Yang H (2019) First report of leaf spots on *Photinia serrulata* caused by *Nigrospora oryzae* in China. Plant Disease 103(9): 2480. https://doi.org/10.1094/PDIS-03-19-0431-PDN

Huang DY, Nong XH, Zhang YQ, Xu W, Sun LY, Zhang T, Chen GY, Han CR (2021) Two new 2,5-diketopiperazine derivatives from mangrove-derived endophytic fungus *Nigrospora camelliae-sinensis* S30. Natural Product Research 36(14): 3651–3656. https://doi.org/10.1080/14786419.2021.1878168

Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo A, Chethana KWT, Clericuzio M, et al. (2017) Fungal diversity notes 603–708: taxonomic and phylogenetic notes on genera and species. Fungal Divers 87: 1–235.

Hyde KD, de Sliva NI, Jeewon R, Bhat DJ, Phookamsak R, Doilom M, Boonmee S, Jayawardena RS, Maharachchikumbura SSN, Senanayake IC, Manawasinghe IS, Liu NG, Abeywickrama PD, Chaiwan N, Karunarathna A, Pem D, Lin CG, Sysouphanthong P, Luo ZL, Wei DP, Wanasinghe DN, Norphanphoun C, Tennakoon DS, Samarakoon MC, Jayasiri SC, Jiang HB, Zeng XY, Li JF, Wijesinghe SN, Devadatha B, Goonasekara ID, Brahmanage RS, Yang EF, Aluthmuhandiram JVS, Dayarathne MC, Marasinghe DS, Li WJ, Dissanayake LS, Dong W, Huanraluek N, Lumyong S, Liu JK, Karunarathna SC, Jones EBG, Al-Sadi AM, Xu JC, Harishchandra D, Sarma VV, Bulgakov TS (2020) AJOM new records and collections of fungi: 1–100. Asian Journal of Mycology 3(1): 22–294.

Hyde KD, Noorabadi MT, Thiyagaraja V, He MQ, Johnston PR, Wijesinghe SN, Armand A, Biketova AY, Chethana KWT, Erdoğdu M, Ge ZW, Groenewald JZ, Hongsanan S, Kušan I, Leontyev DV, Li DW, Lin CG, Liu NG, Maharachchikumbura SSN, Matočec N, May TW, McKenzie EHC, Mešić A, Perera RH, Phukhamsakda C, Piątek M, Samarakoon MC, Selcuk F, Senanayake IC, Tanney JB, Tian Q, Vizzini A, Wanasinghe DN, Wannasawang N, Wijayawardene NN, Zhao RL, Abdel-Wahab MA, Abdollahzadeh J, Abeywickrama PD, Abhinav, Absalan S, Acharya K, Afshari N, Afshan NS, Afzalinia S, Ahmadpour SA, Akulov O, Alizadeh A, Alizadeh M, Al-Sadi AM, Alves A, Alves VCS, Alves-Silva G, Antonín V, Aouali S, Aptroot A, Apurillo CCS, Arias RM, Asgari, B, Asghari R, Assis DMA, Assyov B, Atienza V, Aumentado HDR, Avasthi S, Azevedo E, Bakhshi M, Bao DF, Baral HO, Barata M, Barbosa KD, Barbosa RN, Barbosa FR, Baroncelli R, Barreto GG, Baschien C, Bennett RM, Bera I, Bezerra JDP, Bhunjun CS, Bianchinotti MV, Błaszkowski J, Boekhout T, Bonito GM, Boonmee S, Boonyuen N, Bortnikov FM, Bregant C, Bundhun D, Burgaud G, Buyck B, Caeiro MF, Cabarroi-Hernández M, Cai M Feng, Cai L, Calabon MS, Calaça FJS, Callalli M, Câmara MPS, Cano-Lira J, Cao B, 5162 Carlavilla JR, Carvalho A, Carvalho TG, Castañeda-Ruiz RF, Catania MDV, Cazabonne J, Cedeño-Sanchez M, Chaharmiri-Dokhaharani S, Chaiwan N, Chakraborty N, Cheewankoon R, Chen C, Chen J, Chen Q, Chen YP, Chinaglia S, Coelho-Nascimento CC, Coleine C, CostaRezende DH, Cortés-Pérez A, Crouch, JA, Crous PW, Cruz RHSF, Czachura P, Damm U, Darmostuk V, Daroodi Z, Das K, Das K, Davoodian N, Davydov EA, et al. (2024) The 2024 Outline of Fungi and fungus-like taxa. Mycosphere 15(1): 5146–6239. https://doi.org/10.5943/mycosphere/15/1/25

Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang KA, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsanan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The Faces of Fungi database: Fungal names linked with morphology, phylogeny and human impacts. Fungal Diversity 74(1): 3–18. https://doi.org/10.1007/s13225-015-0351-8

Kirk PM, Cannon PF, Minter DW, et al. (2008) Dictionary of the Fungi 10th edn. CABI Bioscience, UK.

Kuang QX, Luo Y, Lei LR, Guo WX, Li XA, Wang YM, Huo XY, Liu MD, Zhang Q, Feng D, Huang LJ, Wang D, Gu YC, Deng Y, Guo DL (2022) Hydroanthraquinones from *Nigrospora sphaerica* and their anti-inflammatory activity uncovered by transcriptome analysis. Journal of Natural Products 85(6): 1474–1485. https://doi.org/10.1021/acs.jnatprod.1c01141

Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. Molecular biology and evolution 33(7): 1870–1874. https://doi.org/10.1093/molbev/msw054

Li DL, Chen YC, Pan QL, Tao MH, Zhang WM (2014) A new eudesmane sesquiterpene from *Nigrospora oryzae*, an endophytic fungus of *Aquilaria sinensis*. Records of Natural Products 8: 330–333.

Li JF, Phookamsak R, Jeewon R, Bhat DJ, Mapook A, Camporesi E, Shang Q-J, Chukeatirote E, Bahkali AH, Hyd KD (2017) Molecular taxonomy and morphological characterization reveal new species and new host records of *Torula species* (Torulaceae, Pleosporales). Mycological Progress 16: 447–461. https://doi.org/10.1007/s11557-017-1292-2

Li L, Pan H, Chen MY, Zhang SJ, Zhong CH (2018) First report of *Nigrospora oryzae* causing brown/black spot disease of kiwifruit in China. Plant Disease 102(1): 243. https://doi.org/10.1094/PDIS-06-17-0833-PDN

Li M, Hong Z, Wang S, Xu D, Yang Z, Li Z, Hu H, Li S (2025) Characterization and Analysis of 2-(2-Phenylethyl)chromone Derivatives and Sesquiterpenoids from Agarwood of Four "Qi-Nan" Clones (*Aquilaria sinensis*) with Different Induction Times. Molecules 30(2): 352. https://doi.org/10.3390/molecules30020352

Liu YY, Wei JH, Gao ZH, Zhang Z, Lyu JC (2017) A review of quality assessment and grading for agarwood. Chinese Herbal Medicines 9(1): 22–30. https://doi.org/10.1016/S1674-6384(17)60072-8

Liu Y, An J, Safdar A, Shen Y, Sun Y, Shu W, Tan X, Zhu B, Xiao J, Schirawski J, He F, Zhou G (2024) Identification and characterization of *Nigrospora* species and a novel species, *Nigrospora anhuiensis*, causing black leaf spot on rice and wild rice in the Anhui Province of China. Journal of Fungi 10: 156. https://doi.org/10.3390/jof10020156

Lorenzini M, Cappello MS, Logrieco A, Zapparoli G (2016) Polymorphism and phylogenetic species delimitation in filamentous fungi from predominant mycobiota in withered grapes. International Journal of Food Microbiology 238: 56–62. https://doi.org/10.1016/j.ijfoodmicro.2016.08.039

Mason EW (1927) On species of the genus *Nigrospora Zimmermann* recorded on monocotyledons. Transactions of the British Mycological Society 12: 152–165. https://doi.org/10.1016/S0007-1536(27)80011-2

Metwaly AM, Kadry HA, El-Hela AA, Mohammad AI, Ma G, Cutler SJ, Ross SA (2014) Nigrosphaerin A a new isochromene derivative from the endophytic fungus *Nigrospora sphaerica*. Phytochemistry Letters 7: 1–5. https://doi.org/10.1016/j.phytol.2013.09.001

Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Gateway Computing Environments Workshop (GCE), New Orleans, LA, USA, 2010, 1–8. https://doi.org/10.1109/GCE.2010.5676129

O'Donnell K, Kistler HC, Cigelnik E, Ploetz RC (1998) Multiple evolutionary origins of the fungus causing Panama disease of banana: Concordant evidence from nuclear and mitochondrial gene genealogies. Proceedings of the National Academy of Sciences of the United States of America 95(5): 2044–2049. https://doi.org/10.1073/pnas.95.5.2044

Page RDM (1996) TreeView: An application to display phylogenetic trees on personal computers. CABIOS 12: 357–358. https://doi.org/10.1093/bioinformatics/12.4.357

Rathnayaka AR, Tennakoon DS, Jones GE, Wanasinghe DN, Bhat DJ, Priyashantha AH, Stephenson SL, Tibpromma S, Karunarathna SC (2025) Significance of precise documentation of hosts and geospatial data of fungal collections, with an emphasis on plant-associated fungi. New Zealand Journal of Botany 63(2–3): 462–489. https://doi.org/10.1080/0028825X.2024.2381734

Su XJ, Luo ZL, Jeewon R, Bhat DJ, Bao DF, Li WL, et al. (2018) Morphology and multigene phylogeny reveal new genus and species of Torulaceae from freshwater habitats in northwestern Yunnan, China. Mycological Progress 17(5): 531–545. https://doi.org/10.1007/s11557-018-1388-3

Subasinghe SMCUP, Hitihamu HID, Fernando KMEP (2018) Use of two fungal species to induce agarwood resin formation in *Gyrinops walla*. Journal of Forestry Research 30(2): 721–726. https://doi.org/10.1007/s11676-018-0654-1

Sun X, Guo LD, Hyde KD (2011) Community composition of endophytic fungi in *Acer truncatum* and their role in decomposition. Fungal Diversity 47: 85–95. https://doi.org/10.1007/s13225-010-0086-5

Tan CS, Isa NM, Ismail I, Zainal Z (2019) Agarwood induction: Current developments and future perspectives. Frontiers in Plant Science 10: 122. https://doi.org/10.3389/fpls.2019.00122

Tibpromma S, Karunarathna SC, Du T Y, Xu J, Mortimer PE, Zhang L, Wang Y H, Phukhamsakda C, Suwannarach N, Rachakunta M (2021) Volatile Constituents of Endophytic Fungi Isolated from *Aquilaria sinensis* with descriptions of two new species of *Nemania*. Life 11: 363. https://doi.org/10.3390/life

Wang M, Liu F, Crous PW, Cai L (2017) Phylogenetic reassessment of *Nigrospora*: Ubiquitous endophytes, plant and human pathogens. Persoonia 39: 118–142. https://doi.org/10.3767/persoonia.2017.39.06

Wang S, Yu ZX, Wang CH, Wu CM, Guo P, Wei JH (2018) Chemical constituents and pharmacological activity of agarwood and *Aquilaria* plants. Molecules 23(2): 342. https://doi.org/10.3390/molecules23020342

Wang Y, Lin S, Zhao L, Sun X, He W, Zhang Y, Dai YC (2019) *Lasiodiplodia* spp. associated with *Aquilaria crassna* in Laos. Mycological Progress 18(5): 683–701. https://doi.org/10.1007/s11557-019-01481-7

White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications 18: 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Wu Z, Xie Z, Wu M, Li X, Li W, Ding W, She Z, Li C (2018) New antimicrobial Cyclopentenones from *Nigrospora sphaerica* ZMT05, a fungus derived from *Oxya chinensis*

thunber. Journal of Agricultural and Food Chemistry 66(21): 5368–5372. https://doi.org/10.1021/acs.jafc.8b01376

Xu T, Song Z, Hou Y, Liu S, Li X, Yang Q, Wu S (2022) Secondary metabolites of the genus *Nigrospora* from terrestrial and marine habitats: Chemical diversity and biological activity. Fitoterapia Sep; 161: 105254. https://doi.org/10.1016/j.fitote.2022.105254

Zhang QH, Tian L, Sun ZL, Fang S, Cai GL, Wang YJ, Pei YH (2015) Two new secondary metabolites from the marine-derived fungus *Nigrospora sphaerica.* Journal of Asian Natural Products Research 17(5): 497–503. https://doi.org/10.1080/10286020.2015.1009899

Zhang LQ, Jiang S, Meng JJ, An HS, Zhang XY (2019) First report of leaf spot caused by *Nigrospora oryzae* on blueberry in Shanghai, China. Plant Disease 103(9): 2473–2474. https://doi.org/10.1094/PDIS-02-19-0242-PDN

Zhu XW, Chen JQ, Zhu SR, He YY, Ding WJ, Li CY (2018) Two new compounds from *Nigrospora sphaerica* ZMT05, a fungus derivated from *Oxya chinensis* thunber. Natural Product Research 32(20): 2375–2381. https://doi.org/10.1080/14786419.2017.1413566

Zimmerman A (1902) Ueber einige an tropischen Kulturpflanzen beobachtete Pilze III. Zentralblatt für Bakteriologie. Parasitenkunde 8: 216–221.

Zou M, Al-Otibi F, Hyde KD, Wang Y, Pan XJ (2024) New *Helminthosporium* (Massarinaceae, Dothideomycetes) and *Nigrospora* (Incertae sedis, Sordariomycetes) species associated with walnut (*Juglans regia* L.) in China. MycoKeys 109: 265–284. https://doi.org/10.3897/mycokeys.109.133431