

Factors influencing pollinator abundance in Indigenous coffee farms of the Nilgiris, Western Ghats, India

Manju Vasudevan Sharma (✉ mann.vasu@gmail.com)

Keystone Foundation

Poomima Viswanathan

Keystone Foundation

Robert Leo

Keystone Foundation

Barbara Gemmill-Herren

Prescott College

Christos Mammides

Frederick University

Hien T. Ngo

Food and Agriculture Organization of the United Nations

Research Article

Keywords: Bees, ecosystem services, tree cover, forest edge, coffee management, biodiversity conservation

Posted Date: January 19th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2485017/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Wild pollinators are shown to be declining in many parts of the world where data and evidence exist; trends could be similar in other regions, but data and evidence are lacking. Land-use change is recognized as the top driver of biodiversity loss, including pollinator loss. In this study, we focused on coffee plantations in Indigenous land holdings in the Nilgiri Biosphere Reserve in the Western Ghats of India, where changing agricultural practices and reducing tree shade diversity and/or changing tree cover type may threaten pollinator communities. We assessed pollinator abundance, through scan sampling of flowers, in ten coffee farms – five of which had (*Grevillea robusta*) silver oak as shade trees and five of which had native tree species. We then evaluated the combined effect of (a) tree cover type, (b) distance from the forest edge, and (c) area under coffee cultivation on the abundance of four dominant coffee pollinators (*Apis dorsata*, *A. cerana*, *A. florea*, and *Tetragonula iridipennis*) and the abundance of *Xylocopa* sp., which is also known as a coffee pollinator. We found that the abundances of all five species were associated positively with the area under coffee cultivation. The abundance of *A. cerana* and *T. iridipennis* were also associated with the distance from the forest edge; the closer a farm to the forest, the more individuals of *A. cerana* and *T. iridipennis* were found visiting coffee flowers on the farm. Lastly, we found no statistically significant relationship between the abundances of the five species examined and tree cover type (either mixed native forest trees or silver oak (*G. robusta*)). The absence of a pattern may have been driven partly by our relatively small sample size since the abundances of *A. cerana*, *A. florea*, and *T. iridipennis* were on average higher in farms with native tree species. Our results suggest that maintaining forests near coffee systems increases insect pollinator abundance (i.e., delivery of pollination services) in the case of *A. cerana* and *T. iridipennis* and maintaining forest cover with native tree species composition plays a role in supporting pollinator habitats as well as providing foraging resources.

Implications for insect conservation

In working with Indigenous land holdings in India, our results show that forests and land use (area under cultivation) play a key role in maintaining bee pollinators in coffee agroecosystems and further investigation is needed to clarify the role of tree cover type and insect populations.

Introduction

It is widely accepted that pollination services, provided by a diversity of pollinators, are important for global food security, ecosystem integrity, nutrition and health security, and human well-being (Chaplin-Kramer et al., 2014; Huttenhower et al., 2012; IPBES, 2016; Klein et al., 2018; Ollerton, 2021; Potts et al., 2016). Pollination is an indisputably critical ecosystem service, with approximately 90% of our wild-growing plant communities (Ollerton et al., 2011) and 75% of our leading global crop types dependent on pollinators, to varying degrees (IPBES, 2016; Klein et al., 2007). This is no exception for India, which, together with China and the USA, accounts for approximately 30% of the total global cropland (FAO, 2020); important crops for India include mangoes, fresh vegetables, tomatoes, soybeans, eggplant,

cardamom, coffee, and others — which are all pollinator-dependent (Abraham et al., 2013; Agarwal et al., 2013; Basu et al., 2011; Bhattacharya & Basu, 2018; Cooley & Vallejo-Marín, 2021; Garibaldi et al., 2021; Huda et al., 2015; Krishnan et al., 2012).

Studies from various parts of the world have suggested that bees are on the decline (Biesmeijer et al., 2006; Cameron & Sadd, 2020; Gallai et al., 2009; IPBES, 2016; Koh et al., 2016; Orr et al., 2021; Potts et al., 2016). Yet, data and evidence are sparse from some regions. The largest drivers of biodiversity loss are land- and sea-use change, direct exploitation of nature, climate change, invasive alien species and pollution (IPBES, 2016), which unsurprisingly, match the drivers of pollinator decline (Dicks et al., 2021; Goulson et al., 2015). Dicks et al. (2021) found changes in land cover and configuration to be important drivers of pollinator loss in most regions worldwide; for the Asia-Pacific region, the study found pesticides to be among the top three drivers of pollinator loss, which differs from the global ranking of drivers for biodiversity loss overall (i.e., not the same relative rank of importance). In terms of land configuration, native forests are known to play a beneficial role in wild bee populations (Aizen & Feinsinger, 2003; Ghazoul et al., 1998; Schrader et al., 2018), largely because native trees and older trees provide suitable nesting substrates for bees such as *Apis dorsata* (Thomas et al., 2009) and *Tetragonula* species (Bhatta et al., 2019). Proximity to forests or semi-natural sites increases pollinator species richness and abundance within agricultural systems (Blanche et al., 2006; Blanche & Cunningham, 2005; Chacoff & Aizen, 2006; Free, 1993; Klein et al., 2003; Kremen et al., 2002, 2004; Morandin & Winston, 2006; Ricketts, 2004; Viswanathan et al., 2020, 2020; Widhiono & Sudiana, 2016). Natural forests and semi-natural plots play an important role for pollinators and their delivery of pollination services in several important cash crops, including coffee and spices.

One of the most important global agricultural commodities is coffee (Khoury et al., 2014), worth approximately \$40 billion/yr in production value alone and significantly more when considering retail value (Rhiney et al., 2021); coffee prices reached an average of \$190.82 US cents/lb (July 2022). The coffee industry involves upwards of twenty-five million coffee farmers and producers (Donald, 2004; Jha et al., 2011; Pham et al., 2019). Coffee production occupies over 10 million hectares of land (FAO, 2021). India is the seventh top coffee producer in the world (Sänger, 2018), making coffee an important cash crop for India (Bhattacharya, 2014), being the seventh top coffee grower in the world, India produces approximately 3% of the total global coffee export, equivalent to approximately 334,000 metric tons (estimated for 2020–2021, Coffee Board of India), which mainly comes from smallholder farmers (Chengappa et al., 2017; DaMatta et al., 2007; Ranjan Jena & Grote, 2017), some of whom are Indigenous forest-dwelling tribes. In the Western Ghats of India, coffee is traditionally grown under an agroforestry landscape with a diversity of native shade trees or intercropped with pepper, but in recent years, some coffee farms have shifted to a more intensified cultivation system (Chengappa et al., 2017). Specifically, in the coffee-growing slopes of Nilgiri Biosphere Reserve, which spans across the States of Tamil Nadu, Karnataka, and Kerala, the characteristic coffee agro-forest is densely planted with coffee bushes under an evergreen forest patch close to human habitation. However, some coffee growers have shifted to using an exotic fast-growing monocrop of silver oak (*Grevillea robusta*) trees (Bali et al., 2007) to shade their coffee crops. The impact of this shift on pollinator communities is unclear.

For coffee, fruit set is known to increase with cross-pollination by bees – as shown in the Indonesian coffee agroecosystems (Klein et al., 2003); fruit set increased by approximately 12.3% in *C. arabica* and 16% in *C. robusta* (Klein et al., 2003) with “open pollination” treatment where all insects could access coffee flowers compared to “control” where insect pollinators were excluded from the treatment. Pollination is important not only for the quantity of coffee production but also the quality; sufficient coffee pollination results in double-seeded full berries (fruit), while a pollination deficit leads to peaberries (i.e., the development of one ovule into a single-seeded fruit) (Classen et al., 2014; Ghazoul & Krishnan, 2012; Krishnan et al., 2012). Previous studies on pollinators and coffee production in other regions have shown that coffee fruit set increases when coffee is grown near rainforests, even fragments (De Marco & Coelho, 2004; Klein et al., 2003; Ricketts, 2004; Ricketts et al., 2004), since many wild bees nest in natural habitats – large tree branches and cavities, rocks, etc. (Aizen & Feinsinger, 2003; Cunningham, 2000; Ghazoul et al., 1998). In a recent global meta-analysis of *Coffea arabica* systems, Moreaux et al. (2022) found animal pollination to be important for increasing fruit set by approximately 18%, yet when looking at habitat features such as forest cover, distance to nearby forest and forest canopy density, the study found that forest cover and distance to open forest were not to be correlated to bee species richness and fruit set – which does not support previous research (e.g., Klein, 2009; Ricketts, 2004) highlighting the need for further research on this topic.

The Western Ghats of India is a biodiversity hotspot (Myers et al., 2000) and a UNESCO World Heritage Site with “high geological, cultural and aesthetic values” but has undergone extensive loss in forest cover and biodiversity owing to land-use change (Ambinakudige & Choi, 2009; Daniels et al., 1995; Gupta et al., 2014; Kale et al., 2016). Data has been documented the last 60 years for honey bee stocks (i.e., hives) for India and the trend has been increasing (approx. 5 million hives in 1961 to almost 13 million hives in 2021) at the same time, data are lacking on the status and trends of wild bees, which are important for coffee production (Pannure, 2016; Veddeler et al., 2008). Previous studies have cited *Apis dorsata*, the giant Asian honey bee, as the main pollinator of coffee in the Kodagu district (Karnataka); Krishnan et al. (2012) found *A. dorsata* individuals comprising almost 60% of floral visits recorded on coffee (*C. canephora*), which was associated with increased coffee fruit production by 50% (Krishnan et al., 2012).

There is a knowledge gap regarding the impacts of reducing shade tree diversity or changing tree cover type on bee pollinators and their pollination services to coffee crops in the Western Ghats. In a study in Kodagu, “a coffee-growing region where a high percentage of native tree cover is still intact to provide shade for coffee plants” (Bhagwat et al., 2005) shade tree diversity has been shown to enhance coffee production and quality of *C. canephora* while reducing incidences of the coffee berry borer *Hypothenemus hampei* (Nesper et al., 2017).

We assessed bee abundance in small holdings of *Coffea arabica* grown under two types of tree cover – native forest trees and silver oak (*G. robusta*). Based on previous work (Barrios et al., 2018; Nesper et al., 2017; Prado et al., 2018), we predicted that coffee farms with native tree cover would offer better delivery of pollination services compared to those with only silver oak (*G. robusta*); bee abundance was used as proxy indicator of pollination services. In order to determine bee abundance, only insect visitors observed

foraging on pollen or nectar were considered potential pollinators. We quantified bee abundance as a function of (a) tree cover type, (b) distance from the nearest forest edge, and (c) area under coffee cultivation. We examined the influence of these three factors on abundance of *Apis dorsata*, *A. cerana*, *A. florea*, *Tetragonula iridipennis* and *Xylocopa* sp., the dominant pollinators of coffee. Although *Xylocopa* has been reported as a minor pollinator in several studies (Krishnan et al., 2012; Prado et al., 2019; Samnegård et al., 2016), their efficiency in pollination is believed to be high in comparison to social bees (Klein et al., 2003).

Methods

Study sites

The Nilgiri Biosphere Reserve (NBR) is one of the top twenty-five global hotspots of biodiversity in the world based on the level of endemic fauna and flora (Baskaran et al., 2012) and within India is the "largest protected ecologically sensitive areas" (Srinivasan et al., 2021) the NBR is composed of protected forests and human-dominated landscapes, including that of tea, coffee and eucalyptus plantations, orchards and vegetable farms. We chose ten coffee-growing farms (*Coffea arabica*) during the coffee flowering season of February to May (2014), which were part of the agricultural land belonging to the Irula and Kurumba Indigenous communities of the NBR (Fig. 1). These lands are located within evergreen and semi-evergreen forests composed of native tree species such as *Syzygium cuminii*, *Bombax* sp., *Dalbergia* sp., *Toona ciliata*, *Canarium strictum*, *Albizia lebbek*, *Lannea coromandelica*, *Sapindus emarginatus*, and *Anogeissus latifolia*. *Bombax* sp. and *Artocarpus* sp. (LR, pers. obs.) were more common closer to human settlements, having spread from planted trees. We based our study design on the protocol developed by FAO and Vaissière et al., (2011) for testing pollination deficits in crops. Tree cover type (mixed vs. silver oak) was the treatment tested. Half of the farms (n = 5) had coffee with only the silver oak (*G. robusta*) species for shade, and the other half of the farms (n = 5) had coffee with a composition of > 5 native tree species for shade. All of the latter (mixed) were similar to each other in terms of tree species composition. The area under coffee production ranged from 0.5 ha to 1 ha, and the sites were at a minimum distance of 210 m from each other and a maximum of 1097 m (average = 564 m; standard deviation = 312 m) to eliminate issues that could arise due to spatial autocorrelation. All sites were selected based on similar topography, soil, slope, and exposure; in addition, all sites were managed in a similar way, with the exception of the independent variable tested (tree cover type) to minimize the effects of confounding factors. There were no managed bee colonies or bee boxes in any of the study sites. Distances from the nearest forest edge and tree cover type were recorded at each crop site.

Floral density and bee activity

Four trained samplers collected data from the ten sites in rotation. Floral density was measured and standardized in four random plots at each site during the coffee bloom. Floral density has been an important predictor of bee abundance (Hamblin et al., 2018; Mahon & Hodge, 2022) although other studies have not shown an effect of floral density on pollinator visits (Elliott & Irwin, 2009; Essenberg,

2012). In order avoid any effect of floral density on bees within plots, we estimated number of flowers per transect and made sure that each transect had approximately 250 flowers. Flowers were sampled on two random branches of two adjacent coffee bushes in a given plot. No plot was sampled twice. At each site, six transects were walked four times a day (0800h, 1000h, 1300h, 1500h), and scan sampling for bees visiting coffee flowers, or floral units (approximately $n = 250$ flowers per transect, using a counter) was carried out to assess pollinator density/abundance (Vaissière et al., 2011). To measure pollinator diversity and abundance, we sampled insects visiting flowers in 5-minute sampling slots and walked in transects that were roughly 20–25 m long. Sweep netting was also done to assess the diversity and abundance of insects. The diversity counts were done after density counts. Only visitors foraging on pollen or nectar were considered potential pollinators.

Data Analyses

We used the abundance of each of the five pollinator species (*A. dorsata*, *A. cerana*, *A. florea*, *Tetragonula iridipennis*, and *Xylocopa* sp.) as the response variable and (a) tree cover type (native trees vs. silver oak), (b) distance from the forest edge, and (c) area under coffee cultivation, as the explanatory variables. To assess these variables' potential influence on each pollinator species, we ran a generalized linear mixed regression model with a Poisson error distribution. We used the identity of the sites as a random effect to account for the fact that each site was surveyed multiple times ($n = 24$) during the flowering season. We also used the date at which each survey was conducted as a random effect to account for any temporal similarities, e.g., due to weather conditions. To estimate the amount of variance explained by each model, we used the “r.squaredGLMM” function in the “MuMin” package in R; for each model, we calculated the corresponding marginal and conditional R^2 values. The marginal R^2 value represents the amount of variance explained by the explanatory variables (i.e., the fixed effects), while the conditional R^2 value corresponds to the amount of variance explained by the whole model, i.e., including the random effects. To confirm there was no collinearity, we used the “vif” function in the “car” package (Fox & Weisberg, 2019) to measure the variance inflation factor (VIF) of each of the three variables in our models. In all cases, VIF was < 2 ; therefore, we retained all variables in the analyses. To confirm that our data were not spatially autocorrelated, we used the “ape” package (Paradis & Schliep, 2019) to calculate the Moran's I value of each model's residuals. In all five models, the p-value was > 0.05 , suggesting no spatial autocorrelation. All analyses were conducted using the R statistical software (R Core Team, 2022).

Results

A total of 24 species of insects were documented visiting the coffee flowers in the NBR during scan sampling. This included those within the orders Hymenoptera (10 spp. of bees and one unidentified wasp), Diptera (5 spp. of Syrphid flies, one sp. in Bombyliidae, one sp. in Ulidiidae and one sp. unidentified), Coleoptera (1 sp. each in Cantharidae, Elateridae and Curculionidae) and Hemiptera (1 sp. of Pyrrhocoridae).

The abundances of *A. dorsata*, *A. florea*, and *Xylocopa* sp. were mostly associated with the size of the coffee farm, meaning that the larger the size, the more individuals were found visiting the coffee flowers (Tables 1 and 2). The abundances of *A. cerana* and *T. iridipennis* were associated with the size of the farm and the distance from the nearest forest (Fig. 3; Tables 1 and 2). The larger the farm and the closer it was to the forest, the more individuals of *A. cerana* and *T. iridipennis* were found visiting coffee flowers. The relationship between tree cover type and flower visitation was not statistically significant for any of the five species examined (Tables 1 and 2), although the abundances, from scan sampling of *A. cerana*, *A. florea*, and *T. iridipennis* were on average higher in farms with native tree species.

Table 1

Results of the generalized linear mixed models showing the relationship between the three variables examined and the abundances of *A. dorsata*, *A. cerana*, and *A. florea*. Statistically significant effects in bold.

	<i>Apis dorsata</i>			<i>Apis cerana</i>			<i>Apis florea</i>		
<i>Variables</i>	β	<i>std. Error</i>	<i>p</i>	β	<i>std. Error</i>	<i>p</i>	β	<i>std. Error</i>	<i>p</i>
(Intercept)	-5.42	2.04	0.008	-1.34	0.72	0.065	-6.49	1.77	< 0.001
treecover [Silveroak]	1.06	0.79	0.184	-0.19	0.22	0.388	-0.49	0.70	0.487
Area	4.99	2.53	0.049	3.91	0.72	< 0.001	9.37	2.26	< 0.001
Dist	4.56	2.59	0.079	-3.79	0.84	< 0.001	1.10	2.31	0.633
R ² Marginal / R ² Conditional	0.141 / 0.997			0.236 / 0.983			0.319 / 0.995		

Table 2

Results of the generalized linear mixed models showing the relationship between the three variables examined and the abundances of *Tetragonula iridipennis* and *Xylocopa* sp. Statistically significant effects in bold.

	<i>Tetragonula iridipennis</i>			<i>Xylocopa</i> sp.		
<i>Variables</i>	β	<i>std. Error</i>	<i>p</i>	β	<i>std. Error</i>	<i>p</i>
(Intercept)	-6.01	0.97	<0.001	-13.36	4.22	0.002
treecover [Silveroak]	-0.19	0.33	0.558	0.30	1.60	0.854
Area	9.84	1.21	<0.001	12.04	5.20	0.020
Dist	-2.83	1.32	0.033	1.91	5.60	0.733
R ² Marginal / R ² Conditional	0.578 / 0.994			0.306 / 0.949		

Discussion

Changes in land cover and configuration, land management, and pesticides have been identified as the top three drivers threatening pollinator communities across all regions of the world (Dicks et al., 2021). Surrounding natural or semi-natural landscapes are important for the preservation of nature and related ecosystem services (also referred to as nature's contributions to people (Díaz et al., 2018) such as pollination (Julier & Roulston, 2009; Millennium Ecosystem Assessment, 2005; Ricketts, 2004). This is supported by most studies examining the impacts of habitat fragmentation and isolation on pollinator richness and abundance (e.g., (Aizen & Feinsinger, 1994a, 1994b; Blanche et al., 2006; Carvalheiro et al., 2010; Klein et al., 2003; Klein et al., 2003; Perfecto & Vandermeer, 2002; Ricketts, 2004; Steffan-Dewenter & Tscharntke, 1999) but not all (Moreaux et al., 2022).

Our study contributes some empirical evidence that forest fragments are important for pollinators including bees. Forest areas and riparian strips (both small and large) provide potential nesting sites for some bee species. Krishnan et al., (2012) while studying coffee farms (*Coffea canephora*) in Kodagu, South India, demonstrated adjacent forest fragments (0.3–20 ha) positively impacted the number of insect pollinators, dominated by *Apis dorsata*. They found in their study that the giant honey bee *Apis dorsata* – known to nest in forested areas, was the main pollinator of coffee (58% of the total coffee flower visitors). In our study, there was a weaker link between nearby or adjacent forests and *A. dorsata*, although proximity from the forest was correlated to *A. cerana* and *T. iridipennis* bee species abundances. This may indicate that our study required another study period (i.e., more than one sampling season) or more replicates of study sites. Future research may wish to collect insect specimens to be able to identify them to species level, beyond morphospecies identifications, in order to determine diversity of insect visitors to each farm. In addition, future research to monitor insects could consider standardized or multiple methods in order to determine pollinator communities more accurately. For example, our study used pollination scan sampling to determine pollinator density, meaning “an insect will be recorded or not

depending on whether it is present at the time a given flower is first seen” (FAO, 2011) and we complemented this with sweep net sampling, however the sweep net sampling results were not used in our analysis. Other studies use slightly different methods to monitor pollinators on coffee such as flower visits consisting of bees observed to both carry pollen and contact the coffee flower stigma (Boreux et al., 2013) or pollinator observations on selected branches/flowers (Krishnan et al., 2012) – this difference in insect pollinator monitoring precludes a direct comparison of pollinator communities in coffee systems. Our study did support, in part, the findings of the meta-analysis done by Moreaux et al. (2022), where forest proximity was not a strong factor in determining bee richness or in our case, *A. dorsata*, *A. florea*, and *Xylocopa* sp bee abundance. Further research is needed to study the importance of distance from forest fragments, forest fragment size, and forest fragment/habitat quality in influencing nearby pollinator communities for coffee agroecosystems. These factors are hard to isolate (Boreux et al., 2013).

Lastly, the importance of natural forest fragments to coffee agroecosystems and coffee pollinators may be diminished if the shade tree diversity within coffee farms is sufficient in providing nesting sites to pollinators – this relationship has not been sufficiently studied within Indian coffee systems. Shade coffee systems grown under traditional or rustic management have shown potential to have the capacity to house equivalent levels of biodiversity compared to those of adjacent forests (e.g., arthropods) in other regions such as Latin America (Perfecto et al., 1996; Philpott et al., 2008 etc.). Not only can diversity (heterogeneity) of shade trees be of importance but the type (species) of shade tree can also be important for maintaining insect or other faunal diversity in those systems.

There is evidence that a diversity of native tree cover for shade helps maintain floral and faunal diversity (Badrinarayanan et al., 2001; Bhagwat et al., 2005; Dolia et al., 2008; Kapoor, 2008) and within coffee agroecosystems, this diversity supports pollinator communities (Belavadi, 2020). In a recent study, 107 species of shade trees were found in coffee systems in the Kodagu district (Belavadi, 2020). Trees such as *Schefflera venulose* and *S. wallachiana* have their flowering season following coffee; this sequential blooming supports pollinators with a stable source of nectar and pollen (i.e., provisioning of foraging sources during non-flowering season) with added benefits of providing microclimates for better fruit maturation (Krishnan et al., 2017). Planting indigenous tree species such as *Dalbergia latifolia*, *Ficus racemosa*, and *A. fraxinifolius* could have multiple benefits (Maheswarappa et al., 2022; Nath et al., 2016) including those for soil (e.g., improved soil quality through litter fall or reduced soil erosion (Dhanya et al., 2014), timber, firewood, and birds and pollinators. In the coffee-growing areas of Kodagu, Karnataka, there are 280 tree species which could be used as native shade tree species (Ambinakudige & Sathish, 2009; Maheswarappa et al., 2022), yet larger-scale coffee farmers tend to prefer the exotic silver oak species (*G. robusta*) within their coffee farms (Maheswarappa et al., 2022). There are incentives to use *G. robusta* (Nath et al., 2011), as it is fast-growing and holds the lure of timber market. There have also been cited negative aspects to having *G. robusta* as shade trees such as their impact on berry dry weight and coffee quality (Boreux et al., 2018). In our study, having only silver oak (*G. robusta*) was correlated with decreased abundance for three of the five pollinator species (Tables 1 and 2), albeit not statistically significant, suggesting that their abundance was, on average lower in silver oak-covered coffee crops compared to those sites that had native shade trees. As noted above, a larger sample size with more

replicates may have shown a stronger significant relationship between pollinators and tree cover type. We recommend that future studies investigate this relationship further. As landscape elements (size of farm and proximity to forest) were shown in this study to have impacts on key pollinator abundances, these factors should be of importance to farming communities. Our study encourages coffee growers in the region to consider diverse native shade trees for tree cover rather than the non-native silver oak (*G. robusta*) for coffee systems, and to support forest conservation within the Reserve.

In general, more research, funding, and attention are needed in areas where pollinators and pollination are understudied, such as Asia. Coffee production (quality and quantity) is important for farmers within the Western Ghats and across India. Improving and stabilizing coffee production will in turn improve the livelihoods and well-being of farmers and their families and communities. While it is recognized that countries within Asia lack pollination studies compared to other regions such as North America and Europe (Archer et al., 2014; Klein et al., 2018; Orr et al., 2021; Teichroew et al., 2017) there are some initiatives starting that could help promote and support more pollinator and pollination-related research in Asia, including India. The Indian Pollinator Initiative, established in 2021, is run by academics in the field of pollination ecology. The Indian Pollinator Initiative has three main objectives: 1) to advance and promote studies on pollinators and pollination, 2) to provide a platform to exchange findings, ideas, methods and concepts to enhance collaboration, and 3) to encourage and promote education related to pollinators or pollination (more information can be found on <https://inpollin.com/>). There have been several attempts to start an Asian Pollinator Initiative – which is not yet formally established due to the COVID-19 pandemic and the disruption in international travel. A continuation of these efforts is envisioned, and a second meeting is planned to take place in early 2023. This endeavor aimed to have pollinator ecologists from the region meet to discuss how the Asian network could be established, connected, and further developed. The Indigenous Pollinator Network (IPN), supported by the Indigenous Partnership for Agrobiodiversity and Food Sovereignty has various organizations such as the Keystone Foundation (India), Slowfood International (Italy), Kivulini Trust (Kenya), and the Ogiek Peoples Development Project (Kenya) as part of it; this initiative hopes to expand its operations in the coming years (more information can be found at <https://www.theindigenouspartnership.org/pollinators-network>). Future research and support for such initiatives could increase the understanding of pollinators' role in important productive systems and, ultimately, their conservation.

Declarations

Acknowledgements

We are grateful to the Food and Agricultural Organization of the United Nations (UN FAO) for a grant under the GEF/UNEP/FAO Global Pollination Project (GEF Project ID: 2123), to support this study. We thank the Keystone Foundation for their collaboration and partnership in this project and the broader International Pollinator Initiative. We would like to acknowledge Aidin Niamir for early comments on support and advice on the analysis of data. We are grateful to Febina, Kuttan, M Chandran (Late), Sumin George, Sivraj, Palani, Justin, Mahadesh for assistance during data collection. Our biggest thanks and

sincere gratitude to the cooperation and co-production of knowledge with the farmers and villagers of our field sites.

Funding

The project was supported by Food and Agricultural Organization of the United Nations (UN FAO) for a grant under the GEF/UNEP/FAO Global Pollination Project (GEF Project ID: 2123).

Author information

Authors and Affiliations

- Keystone Foundation, PB 35, Groves Hill Road Kotagiri 643 217 Nilgiris District, Tamil Nadu India
- Prescott College, 220 Grove Avenue Prescott, AZ, USA
- Nature Conservation Unit, Frederick University, 7 Yianni Frederickou Str., Nicosia, 1036, Cyprus
- Food and Agriculture Organization of the United Nations, Viale delle Terme di Caracalla, 00153 Roma RM, Italy

Contributions

M. V. Sharma, B. Gemmill-Herren, R. Leo and H. T. Ngo designed the project and field sampling. P. Viswanathan and R. Leo implemented the field sampling and monitoring. M. V. Sharma and H. T. Ngo wrote the main manuscript text. C. Mammides carried out the majority of statistical analyses and reviewed the entire manuscript and added invaluable discussion and guidance to the manuscript.

Corresponding author

Correspondence to Manju Vasudevan Sharma.

Ethics declarations

Conflict of interest

The authors have no conflicts of interest.

References

1. Abraham R, Purushothaman S, Devy S (2013) Conservation and Coffee Production: Creating Synergies in Kodagu, Karnataka. In S. Purushothaman & R. Abraham (Eds.), *Livelihood Strategies in Southern India: Conservation and Poverty Reduction in Forest Fringes* (pp. 89–107). Springer India. https://doi.org/10.1007/978-81-322-1626-1_6
2. Agarwal DK, Billore SD, Sharma AN, Dupare BU, Srivastava SK (2013) Soybean: Introduction, Improvement, and Utilization in India—Problems and Prospects. *Agricultural Res* 2(4):293–300.

<https://doi.org/10.1007/s40003-013-0088-0>

3. Aizen MA, Feinsinger P (1994a) Forest Fragmentation, Pollination, and Plant Reproduction in a Chaco Dry Forest, Argentina. *Ecology* 75(2):330–351. <https://doi.org/10.2307/1939538>
4. Aizen MA, Feinsinger P (1994b) Habitat Fragmentation, Native Insect Pollinators, and Feral Honey Bees in Argentine 'Chaco Serrano'. *Ecol Appl* 4(2):378–392. <https://doi.org/10.2307/1941941>
5. Aizen MA, Feinsinger P (2003) Bees Not to Be? Responses of Insect Pollinator Faunas and Flower Pollination to Habitat Fragmentation. In: Bradshaw GA, Marquet PA (eds) *How Landscapes Change: Human Disturbance and Ecosystem Fragmentation in the Americas*. Springer, Berlin Heidelberg, pp 111–129. https://doi.org/10.1007/978-3-662-05238-9_7
6. Ambinakudige S, Choi J (2009) Global coffee market influence on land-use and land-cover change in the Western Ghats Of India. *Land Degrad Dev* 20(3):327–335. <https://doi.org/10.1002/ldr.921>
7. Ambinakudige S, Sathish BN (2009) Comparing tree diversity and composition in coffee farms and sacred forests in the Western Ghats of India. *Biodivers Conserv* 18(4):987–1000. <https://doi.org/10.1007/s10531-008-9502-5>
8. Archer CR, Pirk CWW, Carneiro LG, Nicolson SW (2014) Economic and ecological implications of geographic bias in pollinator ecology in the light of pollinator declines. *Oikos* 123(4):401–407. <https://doi.org/10.1111/j.1600-0706.2013.00949.x>
9. Badrinarayanan S, Krishnaswamy J, Lele S, Chandrashekara K (2001) Consequences of forest conversion to coffee plantation on litter beetle and ant communities. In *K.N. Ganeshiah, R.U. Shaanker, and K.S. Bawa, (Eds.), Proceedings of International Conference on Tropical Ecosystems: Structure, Diversity and Human Welfare* (pp. 162–166). Oxford-IBH Publishing Co. http://atree.org/sites/default/files/pubs/slele/journal-publications/slele_5_forest-coffee_and_litter_insects-summary.pdf
10. Bali A, Kumar A, Krishnaswamy J (2007) The mammalian communities in coffee plantations around a protected area in the Western Ghats, India. *Biol Conserv* 139(1):93–102. <https://doi.org/10.1016/j.biocon.2007.06.017>
11. Barrios E, Valencia V, Jonsson M, Brauman A, Hairiah K, Mortimer PE, Okubo S (2018) Contribution of trees to the conservation of biodiversity and ecosystem services in agricultural landscapes. *Int J Biodivers Sci Ecosyst Serv Manage* 14(1):1–16. <https://doi.org/10.1080/21513732.2017.1399167>
12. Baskaran N, Anbarasan U, Agoramoorthy G (2012) India's biodiversity hotspot under anthropogenic pressure: A case study of Nilgiri Biosphere Reserve. *J Nat Conserv* 20(1):56–61. <https://doi.org/10.1016/j.jnc.2011.08.004>
13. Basu P, Bhattacharya R, Ianetta P (2011) A decline in pollinator dependent vegetable crop productivity in India indicates pollination limitation and consequent agro-economic crises. *Nat Precedings*. <https://doi.org/10.1038/npre.2011.6044.1>
14. Belavadi VV (2020) Chapter 9: Benefits of planning shade-tree cultivation to favour pollinators. In *Towards sustainable crop pollination services: Measures at field, farm and landscape scales* (pp. 93–104). FAO. <https://doi.org/10.4060/ca8965en>

15. Bhagwat SA, Kushalappa CD, Williams PH, Brown ND (2005) The Role of Informal Protected Areas in Maintaining Biodiversity in the Western Ghats of India. *Ecol Soc* 10(1):8
16. Bhatta CP, Gonzalez VH, Mayes D, Simões M, Smith DR (2019) Nesting biology and niche modelling of *Tetragonula iridipennis* (Smith) (Hymenoptera: Apidae, Meliponini) in Nepal. *J Apic Res* 58(4):501–511. <https://doi.org/10.1080/00218839.2019.1614729>
17. Bhattacharya B (2014) Local History of a Global Commodity: Production of Coffee in Mysore and Coorg in the Nineteenth Century. *Indian Hist Rev* 41(1):67–86. <https://doi.org/10.1177/0376983614521734>
18. Bhattacharya R, Basu P (2018) Pollinator Limitation and Crop Production: Experimental Observations on Few Economically Important Vegetable Crops in West Bengal, India. *Proceedings of the Zoological Society*, 71(1), 88–91. <https://doi.org/10.1007/s12595-016-0189-4>
19. Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP, Potts SG, Kleukers R, Thomas CD, Settele J, Kunin WE (2006) Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313(5785):351–354. <https://doi.org/10.1126/science.1127863>
20. Blanche KR, Ludwig JA, Cunningham SA (2006) Proximity to rainforest enhances pollination and fruit set in orchards. *J Appl Ecol* 43(6):1182–1187. <https://doi.org/10.1111/j.1365-2664.2006.01230.x>
21. Blanche R, Cunningham SA (2005) Rain Forest Provides Pollinating Beetles for Atemoya Crops. *J Econ Entomol* 98(4):1193–1201. <https://doi.org/10.1603/0022-0493-98.4.1193>
22. Boreux V, Krishnan S, Cheppudira KG, Ghazoul J (2013) Impact of forest fragments on bee visits and fruit set in rain-fed and irrigated coffee agro-forests. *Agric Ecosyst Environ* 172:42–48. <https://doi.org/10.1016/j.agee.2012.05.003>
23. Cameron SA, Sadd BM (2020) Global Trends in Bumble Bee Health. *Ann Rev Entomol* 65(1):209–232. <https://doi.org/10.1146/annurev-ento-011118-111847>
24. Carvalheiro LG, Seymour CL, Veldtman R, Nicolson SW (2010) Pollination services decline with distance from natural habitat even in biodiversity-rich areas. *J Appl Ecol* 47(4):810–820. <https://doi.org/10.1111/j.1365-2664.2010.01829.x>
25. Chacoff NP, Aizen MA (2006) Edge effects on flower-visiting insects in grapefruit plantations bordering premontane subtropical forest. *J Appl Ecol* 43(1):18–27. <https://doi.org/10.1111/j.1365-2664.2005.01116.x>
26. Chaplin-Kramer R, Dombeck E, Gerber J, Knuth KA, Mueller ND, Mueller M, Ziv G, Klein A-M (2014) Global malnutrition overlaps with pollinator-dependent micronutrient production. *Proceedings of the Royal Society B: Biological Sciences*, 281(1794), 20141799. <https://doi.org/10.1098/rspb.2014.1799>
27. Chengappa PG, Devika CM, Rudragouda CS (2017) Climate variability and mitigation: Perceptions and strategies adopted by traditional coffee growers in India. *Climate Dev* 9(7):593–604. <https://doi.org/10.1080/17565529.2017.1318740>

28. Classen A, Peters MK, Ferger SW, Helbig-Bonitz M, Schmack JM, Maassen G, Schleuning M, Kalko EKV, Böhning-Gaese K, Steffan-Dewenter I (2014) Complementary ecosystem services provided by pest predators and pollinators increase quantity and quality of coffee yields. *Proceedings of the Royal Society B: Biological Sciences*, 281(1779), 20133148. <https://doi.org/10.1098/rspb.2013.3148>
29. Cooley H, Vallejo-Marín M (2021) Buzz-Pollinated Crops: A Global Review and Meta-analysis of the Effects of Supplemental Bee Pollination in Tomato. *J Econ Entomol* 114(2):505–519. <https://doi.org/10.1093/jee/toab009>
30. Cunningham SA (2000) Depressed pollination in habitat fragments causes low fruit set. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 267(1448), 1149–1152. <https://doi.org/10.1098/rspb.2000.1121>
31. DaMatta FM, Ronchi CP, Maestri M, Barros RS (2007) Ecophysiology of coffee growth and production. *Braz J Plant Physiol* 19:485–510. <https://doi.org/10.1590/S1677-04202007000400014>
32. Daniels RJR, Gadgil M, Joshi NV (1995) Impact of Human Extraction on Tropical Humid Forests in the Western Ghats Uttara Kannada, South India. *J Appl Ecol* 32(4):866–874. <https://doi.org/10.2307/2404826>
33. De Marco P, Coelho FM (2004) Services performed by the ecosystem: Forest remnants influence agricultural cultures' pollination and production. *Biodivers Conserv* 13(7):1245–1255. <https://doi.org/10.1023/B:BIOC.0000019402.51193.e8>
34. Dhanya B, Sathish BN, Viswanath S, Purushothaman S (2014) Ecosystem services of native trees: Experiences from two traditional agroforestry systems in Karnataka, Southern India. *Int J Biodivers Sci Ecosyst Serv Manage* 10(2):101–111. <https://doi.org/10.1080/21513732.2014.918057>
35. Díaz S, Pascual U, Stenseke M, Martín-López B, Watson RT, Molnár Z, Hill R, Chan KMA, Baste IA, Brauman KA, Polasky S, Church A, Lonsdale M, Larigauderie A, Leadley PW, van Oudenhoven APE, van der Plaats F, Schröter M, Lavorel S, ..., Shirayama Y (2018) Assessing nature's contributions to people. *Science* 359(6373):270–272. <https://doi.org/10.1126/science.aap8826>
36. Dicks LV, Breeze TD, Ngo HT, Senapathi D, An J, Aizen MA, Basu P, Buchori D, Galetto L, Garibaldi LA, Gemmill-Herren B, Howlett BG, Imperatriz-Fonseca VL, Johnson SD, Kovács-Hostyánszki A, Kwon YJ, Lattorff HMG, Lungharwo T, Seymour CL, ..., Potts SG (2021) A global-scale expert assessment of drivers and risks associated with pollinator decline. *Nat Ecol Evol* 5(10) Article 10. <https://doi.org/10.1038/s41559-021-01534-9>
37. Dolia J, Devy MS, Aravind NA, Kumar A (2008) Adult butterfly communities in coffee plantations around a protected area in the Western Ghats, India. *Anim Conserv* 11(1):26–34. <https://doi.org/10.1111/j.1469-1795.2007.00143.x>
38. Donald PF (2004) Biodiversity Impacts of Some Agricultural Commodity Production Systems. *Conserv Biol* 18(1):17–37
39. Elliott SE, Irwin RE (2009) Effects of flowering plant density on pollinator visitation, pollen receipt, and seed production in *Delphinium barbeyi* (Ranunculaceae). *Am J Bot* 96(5):912–919. <https://doi.org/10.3732/ajb.0800260>

40. Essenberg CJ (2012) Explaining Variation in the Effect of Floral Density on Pollinator Visitation. *Am Nat* 180(2):153–166. <https://doi.org/10.1086/666610>
41. Esther Julier H, Roulston TH (2009) Wild Bee Abundance and Pollination Service in Cultivated Pumpkins: Farm Management, Nesting Behavior and Landscape Effects. *J Econ Entomol* 102(2):563–573. <https://doi.org/10.1603/029.102.0214>
42. FAO (2020) *World Food and Agriculture—Statistical Yearbook 2020*. <https://doi.org/10.4060/cb1329en>
43. FAO (2021) *Food and agriculture Organization of the United Nations. FAOSTAT (database)*.
44. Fox J, Weisberg S (2019) *An {R} Companion to Applied Regression* (3rd ed.). Sage. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
45. Free JB (1993) *Insect Pollination of Crops*. Academic Press
46. Gallai N, Salles J-M, Settele J, Vaissière BE (2009) Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecol Econ* 68(3):810–821. <https://doi.org/10.1016/j.ecolecon.2008.06.014>
47. Garibaldi LA, Schulte LA, Nabaes Jodar DN, Carella G, D. S., Kremen C (2021) Time to Integrate Pollinator Science into Soybean Production. *Trends Ecol Evol* 36(7):573–575. <https://doi.org/10.1016/j.tree.2021.03.013>
48. Ghazoul J, Krishnan S (2012) Importance of bees, shade trees and forests in coffee production in Kodagu. *Indian Coffee* 76(1):10–12
49. Ghazoul J, Liston KA, Boyle TJB (1998) Disturbance-induced density-dependent seed set in *Shorea siamensis* (Dipterocarpaceae), a tropical forest tree. *J Ecol* 86(3):462–473. <https://doi.org/10.1046/j.1365-2745.1998.00270.x>
50. Goulson D, Nicholls E, Botías C, Rotheray EL (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* 347(6229). <https://doi.org/10.1126/science.1255957>
51. Gupta S, Gupta S, Aithal BH, Ramchandra TV (2014) Land use Dynamics in central and southern Western Ghats. *LAKE 2014: Conference on Conservation and Sustainable Management of Wetland Ecosystems in Western Ghats. Sahyadri Conservation Series 47, In, ETR 87*, 199–205. http://wgbis.ces.iisc.ernet.in/energy/lake2014/proceedings/23_Shweta_LAKE2014_R.pdf
52. Hamblin AL, Youngsteadt E, Frank SD (2018) Wild bee abundance declines with urban warming, regardless of floral density. *Urban Ecosyst* 21(3):419–428. <https://doi.org/10.1007/s11252-018-0731-4>
53. Huda AN, Salmah MRC, Hassan AA, Hamdan A, Razak MNA (2015) Pollination Services of Mango Flower Pollinators. *J Insect Sci* 15(1):113. <https://doi.org/10.1093/jisesa/iev090>
54. Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, Chinwalla AT, Creasy HH, Earl AM, FitzGerald MG, Fulton RS, Giglio MG, Hallsworth-Pepin K, Lobos EA, Madupu R, Magrini V, Martin JC, Mitreva M, Muzny DM, Sodergren EJ, The Human Microbiome Project Consortium (2012) Structure,

- function and diversity of the healthy human microbiome. *Nature* 486(7402) Article 7402.
<https://doi.org/10.1038/nature11234>
55. IPBES (2016) In: Potts SG, Imperatriz-Fonseca VL, Ngo HT (eds) The assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES)
 56. Jha S, Bacon CM, Philpott SM, Rice RA, Méndez VE, Läderach P (2011) A Review of Ecosystem Services, Farmer Livelihoods, and Value Chains in Shade Coffee Agroecosystems. In: Campbell WB, Lopez Ortiz S (eds) Integrating Agriculture, Conservation and Ecotourism: Examples from the Field. Springer Netherlands, pp 141–208. https://doi.org/10.1007/978-94-007-1309-3_4
 57. Kale MP, Chavan M, Pardeshi S, Joshi C, Verma PA, Roy PS, Srivastav SK, Srivastava VK, Jha AK, Chaudhari S, Giri Y (2016) Land-use and land-cover change in Western Ghats of India. *Environ Monit Assess* 188(7):387 & Krishna Murthy, Y. V. N. <https://doi.org/10.1007/s10661-016-5369-1>
 58. Kapoor V (2008) Effects of rainforest fragmentation and shade-coffee plantations on spider communities in the Western Ghats, India. *J Insect Conserv* 12(1):53–68.
<https://doi.org/10.1007/s10841-006-9062-5>
 59. Khoury CK, Bjorkman AD, Dempewolf H, Ramirez-Villegas J, Guarino L, Jarvis A, Rieseberg LH, Struik PC (2014) Increasing homogeneity in global food supplies and the implications for food security. *Proceedings of the National Academy of Sciences*, 111(11), 4001–4006.
<https://doi.org/10.1073/pnas.1313490111>
 60. Klein AM, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B-Biological Sciences*, 274(1608), 303–313. <https://doi.org/10.1098/rspb.2006.3721>
 61. Klein A, Steffan-Dewenter I, Tscharntke T (2003) Fruit set of highland coffee increases with the diversity of pollinating bees. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1518), 955–961. <https://doi.org/10.1098/rspb.2002.2306>
 62. Klein A-M (2009) Nearby rainforest promotes coffee pollination by increasing spatio-temporal stability in bee species richness. *For Ecol Manag* 258(9):1838–1845.
<https://doi.org/10.1016/j.foreco.2009.05.005>
 63. Klein A-M, Boreux V, Fornoff F, Mupepele A-C, Pufal G (2018) Relevance of wild and managed bees for human well-being. *Curr Opin Insect Sci* 26:82–88. <https://doi.org/10.1016/j.cois.2018.02.011>
 64. Klein A-M, Steffan-Dewenter I, Tscharntke T (2003) Pollination of *Coffea canephora* in relation to local and regional agroforestry management. *J Appl Ecol* 40(5):837–845.
<https://doi.org/10.1046/j.1365-2664.2003.00847.x>
 65. Koh I, Lonsdorf EV, Williams NM, Brittain C, Isaacs R, Gibbs J, Ricketts TH (2016) Modeling the status, trends, and impacts of wild bee abundance in the United States. *Proceedings of the National Academy of Sciences*, 113(1), 140–145. <https://doi.org/10.1073/pnas.1517685113>

66. Kremen C, Williams NM, Bugg RL, Fay JP, Thorp RW (2004) The area requirements of an ecosystem service: Crop pollination by native bee communities in California: Area requirements for pollination services to crops. *Ecol Lett* 7(11):1109–1119. <https://doi.org/10.1111/j.1461-0248.2004.00662.x>
67. Kremen C, Williams NM, Thorp RW (2002) Crop pollination from native bees at risk from agricultural intensification. *Proceedings of the National Academy of Sciences*, 99(26), 16812–16816. <https://doi.org/10.1073/pnas.262413599>
68. Krishnan S, Kushalappa CG, Shaanker RU, Ghazoul J (2012) Status of pollinators and their efficiency in coffee fruit set in a fragmented landscape mosaic in South India. *Basic Appl Ecol* 13(3):277–285. <https://doi.org/10.1016/j.baae.2012.03.007>
69. Maheswarappa V, Vasudeva R, Hegde R (2022) Circa situm conservation of coffee agroforests: Farmer's perception in Kodagu landscape of Karnataka, India. *Trop Ecol* 63(1):104–112. <https://doi.org/10.1007/s42965-021-00141-w>
70. Mahon N, Hodge S (2022) Evaluating the effects of observation period, floral density, and weather conditions on the consistency and accuracy of timed pollinator counts. *J Pollinat Ecol* 31:124–138. [https://doi.org/10.26786/1920-7603\(2022\)699](https://doi.org/10.26786/1920-7603(2022)699)
71. Millennium Ecosystem Assessment (2005) *Ecosystems and Human Well-Being: Synthesis*. Island Press
72. Morandin LA, Winston ML (2006) Pollinators provide economic incentive to preserve natural land in agroecosystems. *Agric Ecosyst Environ* 116(3–4):289–292. <https://doi.org/10.1016/j.agee.2006.02.012>
73. Moreaux C, Meireles DAL, Sonne J, Badano EI, Classen A, González-Chaves A, Hipólito J, Klein A-M, Maruyama PK, Metzger JP, Philpott SM, Rahbek C, Saturni FT, Sritongchuay T, Tscharntke T, Uno S, Vergara CH, Viana BF, Strange N, Dalsgaard B (2022) The value of biotic pollination and dense forest for fruit set of Arabica coffee: A global assessment. *Agric Ecosyst Environ* 323:107680. <https://doi.org/10.1016/j.agee.2021.107680>
74. Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403(6772) Article 6772. <https://doi.org/10.1038/35002501>
75. Nath CD, Schroth G, Burslem DFRP (2016) Why do farmers plant more exotic than native trees? A case study from the Western Ghats, India. *Agric Ecosyst Environ* 230:315–328. <https://doi.org/10.1016/j.agee.2016.05.013>
76. Nesper M, Kueffer C, Krishnan S, Kushalappa CG, Ghazoul J (2017) Shade tree diversity enhances coffee production and quality in agroforestry systems in the Western Ghats. *Agric Ecosyst Environ* 247:172–181. <https://doi.org/10.1016/j.agee.2017.06.024>
77. Ollerton J (2021) *Pollinators and Pollination: Nature and Society*. Pelagic Publishing Limited
78. Ollerton J, Winfree R, Tarrant S (2011) How many flowering plants are pollinated by animals? *Oikos* 120(3):321–326. <https://doi.org/10.1111/j.1600-0706.2010.18644.x>
79. Orr MC, Hughes AC, Chesters D, Pickering J, Zhu C-D, Ascher JS (2021) Global Patterns and Drivers of Bee Distribution. *Curr Biol* 31(3):451–458e4. <https://doi.org/10.1016/j.cub.2020.10.053>

80. Pannure A (2016) Bee pollinators decline: Perspectives from India. *Int Res J Nat Appl Sci* 3(5):1–10
81. Paradis E, Schliep K (2019) ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35(3):526–528
82. Perfecto I, Rice RA, Greenberg R, Van der Voort ME (1996) Shade Coffee: A Disappearing Refuge for Biodiversity: Shade coffee plantations can contain as much biodiversity as forest habitats. *Bioscience* 46(8):598–608. <https://doi.org/10.2307/1312989>
83. Perfecto I, Vandermeer J (2002) Quality of Agroecological Matrix in a Tropical Montane Landscape: Ants in Coffee Plantations in Southern Mexico. *Conserv Biol* 16(1):174–182. <https://doi.org/10.1046/j.1523-1739.2002.99536.x>
84. Pham Y, Reardon-Smith K, Mushtaq S, Cockfield G (2019) The impact of climate change and variability on coffee production: A systematic review. *Clim Change* 156(4):609–630. <https://doi.org/10.1007/s10584-019-02538-y>
85. Philpott SM, Arendt WJ, Armbrrecht I, Bichier P, Diestch TV, Gordon C, Greenberg R, Perfecto I, Reynoso-Santos R, Soto-Pinto L, Tejeda-Cruz C, Williams-Linera G, Valenzuela J, Zolotoff JM (2008) Biodiversity Loss in Latin American Coffee Landscapes: Review of the Evidence on Ants, Birds, and Trees. *Conserv Biol* 22(5):1093–1105. <https://doi.org/10.1111/j.1523-1739.2008.01029.x>
86. Potts SG, Imperatriz-Fonseca V, Ngo HT, Aizen MA, Biesmeijer JC, Breeze TD, Dicks LV, Garibaldi LA, Hill R, Settele J, Vanbergen AJ (2016) Safeguarding pollinators and their values to human well-being. *Nature* 540(7632):220–229. <https://doi.org/10.1038/nature20588>
87. Prado SG, Collazo JA, Irwin RE (2018) Resurgence of specialized shade coffee cultivation: Effects on pollination services and quality of coffee production. *Agric Ecosyst Environ* 265:567–575. <https://doi.org/10.1016/j.agee.2018.07.002>
88. Prado SG, Collazo JA, Stevenson PC, Irwin RE (2019) A comparison of coffee floral traits under two different agricultural practices. *Sci Rep* 9(1). Article 1. <https://doi.org/10.1038/s41598-019-43753-y>
89. R Core Team (2022) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL [i>https://www.R-project.org/](https://www.R-project.org/).
90. Ranjan Jena P, Grote U (2017) Fairtrade Certification and Livelihood Impacts on Small-scale Coffee Producers in a Tribal Community of India. *Appl Economic Perspect Policy* 39(1):87–110. <https://doi.org/10.1093/aepp/ppw006>
91. Rhiney K, Guido Z, Knudson C, Avelino J, Bacon CM, Leclerc G, Aime MC, Bebbler DP (2021) Epidemics and the future of coffee production. *Proceedings of the National Academy of Sciences*, 118(27), e2023212118. <https://doi.org/10.1073/pnas.2023212118>
92. Ricketts TH (2004) Tropical Forest Fragments Enhance Pollinator Activity in Nearby Coffee Crops. *Conserv Biol* 18(5):1262–1271. <https://doi.org/10.1111/j.1523-1739.2004.00227.x>
93. Ricketts TH, Daily GC, Ehrlich PR, Michener CD (2004) Economic value of tropical forest to coffee production. *Proceedings of the National Academy of Sciences*, 101(34), 12579–12582. <https://doi.org/10.1073/pnas.0405147101>

94. Samnegård U, Hambäck PA, Lemessa D, Nemomissa S, Hylander K (2016) A heterogeneous landscape does not guarantee high crop pollination. *Proceedings. Biological Sciences*, 283(1838), 20161472. PubMed. <https://doi.org/10.1098/rspb.2016.1472>
95. Sängner C (n.d.). *State of the global coffee market*. United Nations Conference on Trade and Development, 10th Multi-Year Expert Meetings on Commodities and Development, Geneva, Switzerland. https://unctad.org/system/files/non-official-document/MYEM2018_Christoph%20Saenger_25042018.pdf
96. Schrader J, Franzén M, Sattler C, Ferderer P, Westphal C (2018) Woody habitats promote pollinators and complexity of plant–pollinator interactions in homegardens located in rice terraces of the Philippine Cordilleras. *Paddy Water Environ* 16(2):253–263. <https://doi.org/10.1007/s10333-017-0612-0>
97. Srinivasan K, Anand S, Bilyaminu H, Haritha (2021) Assessment of Dynamic Land System in Nilgiri Biosphere Reserve Using MODIS Derived Temporal Data Sets during 2001 to 2018. *Int J Environ Clim Change* 11(6):132–149. <https://doi.org/10.9734/IJECC/2021/v11i630430>
98. Steffan-Dewenter I, Tscharntke T (1999) Effects of habitat isolation on pollinator communities and seed set. *Oecologia* 121(3):432–440. <https://doi.org/10.1007/s004420050949>
99. Teichroew JL, Xu J, Ahrends A, Huang ZY, Tan K, Xie Z (2017) Is China's unparalleled and understudied bee diversity at risk? *Biol Conserv* 210:19–28. <https://doi.org/10.1016/j.biocon.2016.05.023>
100. Thomas SG, Varghese A, Roy P, Bradbear N, Potts SG, Davidar P (2009) Characteristics of trees used as nest sites by *Apis dorsata* (Hymenoptera, Apidae) in the Nilgiri Biosphere Reserve, India. *J Trop Ecol* 25(5):559–562. <https://doi.org/10.1017/S026646740900621X>
101. Vaissière B, Freitas B, Gemmill-Herren B (2011) *Protocol to Detect and Assess Pollination Deficits in Crops: A Handbook for its Use*
102. Veddeler D, Olschewski R, Tscharntke T, Klein A-M (2008) The contribution of non-managed social bees to coffee production: New economic insights based on farm-scale yield data. *Agroforest Syst* 73(2):109–114. <https://doi.org/10.1007/s10457-008-9120-y>
103. Viswanathan P, Mammides C, Roy P, Sharma MV (2020) Flower visitors in agricultural farms of Nilgiri Biosphere Reserve: Do forests act as pollinator reservoirs? *J Apic Res* 59(5):978–987. <https://doi.org/10.1080/00218839.2020.1762994>
104. Widhiono I, Sudiana E (2016) Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet. *Biosaintifika: J Biology Biology Educ* 8(2):148–154

Figures



Figure 1

Map of study area and farm sites. The Nilgiri Biosphere Reserve (NBR) and the ten coffee-growing farms (*Coffea arabica*) belonging to the Irula and Kurumba Indigenous communities of the NBR.




Figure 2

The abundances of *A. dorsata* and *A. florea* were associated with the size of the farm, while the abundances of *A. cerana* were associated with the size of the farm and the distance from the nearest forest edge.




Figure 3

The abundances of *Tetragonula iridipennis* were associated with the size of the farm and the distance from the nearest forest edge, while the abundances of *Xylocopa* sp. were only associated with the size of the farm.