

1 **Running head:** Plant evolution in thermal soils

2 **The impact of microclimate and soil on the ecology and evolution of an arctic plant**

3

4 Niklas J. Wickander¹, Pil U. Rasmussen², Bryndís Marteinsdóttir^{3,4}, Johan Ehrlén¹ and Ayco
5 J. M. Tack¹

6

7 ¹Department of Ecology, Environment and Plant Sciences, Stockholm University, Sweden

8 ²The National Research Centre for the Working Environment, 105 Lersø Parkallé, DK-2100
9 Copenhagen, Denmark

10 ³The Soil conservation Service of Iceland, Gunnarsholt, Hella, Iceland (Present address)

11 ⁴Institute of Life and Environmental Sciences, University of Iceland, Reykjavík, Iceland

12

13 **ORCID IDs**

14 Niklas: <https://orcid.org/0000-0003-0892-5801>

15 Pil: <https://orcid.org/0000-0003-0607-4230>

16 Ayco: <https://orcid.org/0000-0002-3550-1070>

17 Bryndís: <https://orcid.org/0000-0003-3779-7327>

18

19 **Keywords:** *Bistorta vivipara*, geothermal soil, local adaptation, performance, phenotypic
20 plasticity, soil temperature

21

22 **Abstract**

23 The arctic and alpine regions are predicted to experience one of the highest rates of climate
 24 change, and the arctic vegetation is expected to be especially sensitive to such changes.
 25 Understanding the ecological and evolutionary responses of arctic plant species to changes in
 26 climate is therefore a key objective. Geothermal areas, where temperature gradients naturally
 27 occur over small spatial scales, and without many of the confounding environmental factors
 28 present in latitudinal and other gradient studies, provide a natural experimental setting to
 29 examine the impact of temperature on the response of arctic-alpine plants to increasing
 30 temperatures. To test the ecological and evolutionary response of the circumpolar alpine
 31 bistort (*Bistorta vivipara*) to temperature, we collected plant material and soil from areas with
 32 low, intermediate, and high soil temperatures and grew them in all combinations at three
 33 different temperatures. At higher experimental soil temperatures, sprouting was earlier, and
 34 plants had more leaves. Sprouting was earlier in soil originating from intermediate
 35 temperature and plants had more leaves when grown in soil originating from low
 36 temperatures. We did not find evidence of local adaptation or genetic variation in reaction
 37 norms among plants originating from areas with low, intermediate, and high soil temperature.
 38 Our findings suggest that the alpine bistort has a strong plastic response to warming, but that
 39 differences in soil temperature have not resulted in genetic differentiation. The lack of an
 40 observed evolutionary response may, for example, be due to the absence of temperature-
 41 mediated selection on *B. vivipara*, or high levels of gene flow balancing differences in
 42 selection. When placed within the context of other studies, we conclude that arctic-alpine
 43 plant species often show strong plastic responses to spring warming, while evidence of
 44 evolutionary responses varies among species.

45

46 **Introduction**

47 Warming in the Arctic is predicted to be particularly strong (Pachauri et al. 2015). This
 48 increase in temperature may have a major impact on the phenology, growth and reproduction
 49 of arctic plants, whose life cycles are strongly influenced by seasonal shifts in temperature
 50 (Henry and Molau 1997, Callaghan et al. 2004, Gornish and Prather 2014, Semenchuk et al.
 51 2015). To predict the long-term effects of increasing temperatures on the arctic flora, it is
 52 important to understand both immediate and evolutionary responses. For example, plant
 53 individuals may respond plastically and advance development in response to increasing
 54 spring temperatures (Anderson et al. 2012). At the same time, overall increases in
 55 temperatures might lead to changes in how strong the optimal response of development to
 56 temperature should be, and to evolutionary responses in reaction norms (Valdés et al. 2018).
 57 To study the response of arctic plants to increasing temperature, studies commonly use
 58 experiments with open-top climate chambers (Arft et al. 1999, Wolkovich et al. 2012), or
 59 sample plants growing along altitudinal (Frei et al. 2014) and latitudinal gradients
 60 (Toftegaard et al. 2016). Such methods have limitations, with the time-scale of experimental
 61 work often too short to observe evolutionary responses, and changes in temperature along
 62 altitudinal and latitudinal gradients often occurring alongside changes in other environmental
 63 factors. Given these limitations, geothermal systems, where radiation heating causes small
 64 scale variation in temperature but not in other environmental variables (O’Gorman et al.
 65 2014, Sigurdsson et al. 2016), offer an alternative setting where plastic and evolutionary
 66 responses to temperature can be explored (Gudmundsdottir et al., 2011; O’Gorman et al.,
 67 2014; Sigurdsson et al., 2016; Valdés et al., 2018). For example, Iceland has many naturally
 68 occurring geothermal areas, where volcanic activity close to the surface warms water and soil
 69 through radiation heating (Gudmundsdottir et al. 2011, Zakharova and Spichak 2012). In

70 such geothermal areas, plants of the same species frequently grow side-by-side in heated and
71 non-heated soil.

72 Plasticity allows plants to grow under variable conditions and to respond to changes
73 in the environment (Agrawal 2001). For example, plants often germinate earlier and grow
74 larger at higher temperatures (Anderson et al. 2012). The optimal plastic response of plants is
75 likely to differ depending on their local environmental conditions (Grether 2005). If there is
76 genetic variation in plasticity, we would thus expect selection to result in reaction norms that
77 are locally adapted (Valdés et al. 2018). For example, plants that have evolved in a low-
78 temperature environment may rapidly germinate, flush leaves or flower in response to
79 warming in spring, whereas plants from a high-temperature environment may respond more
80 slowly to spring warming. Indeed, previous studies have found that plants from geothermally
81 heated areas have an earlier spring phenology than plants from non-heated areas in the field,
82 while the pattern is reversed when grown in a common environment, a phenomenon referred
83 to as counter-gradient variation (Conover and Schultz 1995, Valdés et al. 2018).

84 The plastic responses and local adaptation of plants in geothermal landscapes may not
85 only be in response to the direct effect of warming, but also due to indirect effects of
86 warming on the local environment. One such potentially important indirect effect is the soil
87 environment (Wittenmayer and Merbach 2005, Bardgett and van der Putten 2014). There is
88 evidence for plant local adaptation to both abiotic (Bradshaw 1952, Joshi et al. 2001, Brady et
89 al. 2005) and biotic soil characteristics (Mursinoff and Tack 2017). Warmer soils often have
90 higher nutrient levels, which are frequently limited in the arctic environment (Sigurdsson et
91 al. 2016). Moreover, both beneficial and antagonistic soil biota, like fungi, bacteria and
92 archaea, might be more diverse in warmer soils (Tedersoo et al. 2014, Rasmussen et al. 2018,
93 2020).

94 The aim of this study was to investigate the effects of soil heating on plastic and
 95 evolutionary responses of the plant *Bistorta vivipara*. For this, we carried out a three-way
 96 multifactorial greenhouse experiment where we grew bulbils (i.e. asexually produced axillary
 97 buds), collected from locations with low, intermediate and high temperatures, in soils from
 98 the same three types of locations, under low, intermediate and high soil temperatures in a
 99 climate chamber. We tested the following specific hypotheses:

- 100 1) Experimental heating of soil results in a higher probability of sprouting, earlier
 101 sprouting, and increased size
- 102 2) The bulbils planted in soils originating from heated areas have a higher probability of
 103 sprouting, sprout earlier, and grow larger due to higher availability of mineralized
 104 nutrients
- 105 3) Selection has resulted in a counter-gradient pattern where bulbils originating from
 106 heated sites show a weaker growth response to temperature than bulbils originating
 107 from non-heated sites
- 108 4) Genetic differences among plants originating from different soil temperatures
 109 correspond to local adaptation, i.e. plants perform better under environmental
 110 conditions that correspond to conditions at their site of origin
- 111 5) Both the abiotic and biotic soil environment influence the ecological and evolutionary
 112 response of plants to heating

113

114

Material and methods

Study system

Geothermal landscapes, where magma gets close to the surface and heats soils and ground water, can act as hotspots of local adaptation (O’Gorman et al. 2014). Notably, within these landscapes the temperature can vary strongly at small spatial scales, with temperatures differing by several or tens of degrees Celsius between locations separated by only a few meters (Gudmundsdottir et al. 2011). The plant and soil microbial communities in these locations are frequently strongly affected by spatial variation in soil temperature (Friberg et al. 2009, Zakharova and Spichak 2012). At the same time, variation in soil nutrient concentrations across the temperature gradient is relatively limited, yet availability increases with rising temperatures in arctic ecosystems through mineralization of locked nutrients (Weintraub and Schimel 2003, O’Gorman et al. 2014, Semenchuk et al. 2015, Sigurdsson et al. 2016).

The alpine bistort (*Bistorta vivipara*) is a perennial herb that occurs in many parts of the northern hemisphere (Jonsell and Karlsson 2000). In Iceland, *B. vivipara* can be found growing naturally in a wide range of temperature conditions (Kristinsson 2001), including geothermal areas. The herb primarily spreads clonally by bulbils, and only rarely produces fertile seeds (Jonsell and Karlsson 2000). Bulbils are asexually produced bulb-like structures that replace the lower inflorescence of *B. vivipara*. Bulbils have no specialized dispersal structures, and often start growing within tens of centimeters from the mother plant (Bills et al. 2015).

Origin of plants and soil

We collected plants and soil from Hengladalir, a geothermally active valley located a few tens of kilometers east of Reykjavik, Iceland. Within this area, we identified three locations

with no heating (soil temperatures c. 13°C), three locations with slight heating (14-17°C) and three locations with strong heating (25-30°C). From each location, we collected soil up to a depth of c. 10 cm, which matches the average root depth of *B. vivipara*. From the soil, we removed the litter layer and larger rocks and roots. From each location, we further collected bulbils from one to three *B. vivipara* plants (henceforth referred to as ‘mother plants’), resulting in seven mother plants from each temperature.

Experimental design

To investigate the effect of temperature and soil on plant performance, we designed a multifactorial experiment where bulbils collected at different temperatures (3 levels) were grown in soils collected at different temperatures (3 levels), and the soils were then experimentally heated to low, intermediate and high temperatures, to simulate the natural variation in soil temperature at the geothermal area. This resulted in 27 different treatment combinations. For each treatment combination, we had seven 82 mm diameter pots (330 ml) planted with two bulbils. We distributed bulbils from each mother plant equally across treatment combinations. As maternal investment in bulbils may affect growth, we recorded bulbil weight prior to planting.

We experimentally heated the plant pots in water baths. The water baths were uniformly heated using a combination of aquarium heaters (Easyheater 75W, AQUAEL, Poland) and water circulation pumps (Compact 300, EHEIM, Germany). In total there were nine water baths, three at each temperature level: i) low temperature (13°C), ii) intermediate (18°C) and iii) high (30°C). In each water bath, we placed 21 plant pots. Water baths were placed in a climate chamber set to 13°C and a day length of 16 h.

To avoid microbial contamination, we sealed the plant pots, but filled the lower 2.5 cm with gravel and water-absorbing cloth to allow for drainage. To allow water to flow

below the plant pot, we glued three metal washers under each plant pot. To avoid positional effects, the location of the plant pots was regularly randomized within the three water baths belonging to the same temperature treatment.

To disentangle the effect of soil biota from that of the abiotic soil environment (e.g. physical or chemical structure) on the probability and timing of bulbil sprouting, we set up a complementary experiment using sterile soil. For this, we applied only the low and high temperatures, resulting in eight treatment combinations and 56 plant pots. For sterilization, soil was autoclaved for 1 h at 121°C.

Plant performance measures

Plants were monitored every other day for sprouting. The day of sprouting was determined as the number of days from planting until the first sign of sprouting was observed. Plant height and number of leaves were recorded on days 84 and 126, and the length and width of the longest leaf were measured on days 97 and 159. Given the elliptical shape of the leaf, we calculated leaf size using the formula for the area of an ellipse:

$$Leaf\ size = \left(\frac{leafwidth}{2} * \frac{leaflength}{2} \right) * \pi$$

Analysis

Data were analyzed using the generalized linear mixed model framework as implemented in R v. 3.6.1 (R Core Team 2019). We tested for significance using the *Anova* function in the *car* package (Fox and Weisberg 2019).

To test for the impact of soil temperature and origin of plants and soil on bulbil sprouting and plant growth, we modelled sprouting (0/1), day of sprouting, plant height, number of leaves and leaf size as a function of original soil temperature, original bulbil temperature, experimental soil temperature and their two- and three-way interactions. Bulbil

weight was included as a covariate. To account for variation among mother plants, we included the random factor ‘*Mother plant*’. As we planted two bulbils in each plant pot, we further included the random factor ‘*Plant pot*’ which was nested under ‘*Mother plant*’. For probability of sprouting, we used a binomial distribution with a logit link; for day of sprouting, plant height and leaf size, we used a normal distribution with an identity link; and for the number of leaves, we used a Poisson distribution with a log link.

We *a priori* planned to test for local adaptation in those cases where we found an interaction between original bulbil temperature and original soil temperature, or between original bulbil temperature and experimental soil temperature (Kawecki and Ebert 2004, Blanquart et al. 2013), using a contrast statement as implemented with the function *emmeans* in the R-package *emmeans* (Lenth 2019).

An effect of temperature of the soil at the collection site on the probability of sprouting and the date of sprouting could be due to either the abiotic or biotic soil characteristics. To disentangle the role of the soil abiotic and biotic environment, we first created a data set with data from both field and sterile soil experiment, where we only retained those treatment combinations present in both experiments (i.e. only ‘low’ and ‘high’ temperatures). We then modelled the affected plant traits as a function of original soil temperature, original bulbil temperature, experimental soil temperature and soil sterilization, including all two- and three-way interactions. As above, we included the random factors ‘*Mother plant*’ and ‘*Plant pot*’ (nested under ‘*Mother plant*’). Given the many terms in the full model, we then identified the minimum adequate model using backward selection, using a p-value of 0.05 as a threshold (Crawley 2013).

Results

Bulbils were more likely to sprout and sprouted earlier when grown at higher soil temperatures (Fig 1AB, Table 1). Plants had the smallest leaves when grown at intermediate soil temperature when measured 97 days after planting, whereas plants grown at higher soil temperatures had more leaves when plants were measured 126 days after planting (Fig. 1CD, Table 1). Bulbils sprouted earlier in soil originating from areas with intermediate temperature than in soil originating from low or high temperatures (Table 1, Fig. 2A). Plants had a higher number of leaves at 126 days after planting when grown in soil from a low temperature origin (Fig. 2B, Table 1). At day 159, the end of the experiment, leaf size was interactively affected by experimental soil temperature and original soil temperature: plants had larger leaves when grown in soil originating from intermediate temperature, but only when the experimental soil temperature was either low or high (Fig. S1). Plant height increased with bulbil weight, but was not affected by any of the experimental factors (Table 1).

Plants originating from areas with low, intermediate, and high soil temperature did not differ in probability of sprouting, day of sprouting or growth traits (Table 1). We also did not detect any evidence of local adaptation in terms of an effect of the interaction between original bulbil temperature and original or experimental soil temperature on sprouting probability, day of sprouting or plant growth (Table 1).

Bulbil sprouting was later in the sterile soil than in the field soil, and sprouting was earlier with experimental soil heating (Fig. S2, Table S1). We further detected a weak interactive effect between original bulbil temperature, experimental soil temperature and soil sterilization, suggesting that bulbils of a high temperature origin sprouted later than expected when growing in soil originating from an area with low temperature, but only when the soil was sterilized.

Discussion

We used a multi-factorial experiment to investigate the impact of microclimate and soil on the immediate and evolutionary response of the alpine bistort. As expected, higher experimental soil temperature advanced bulbil sprouting and led to plants with more leaves. In contrast to our prediction, sprouting was earliest in soil originating from locations with intermediate temperature, and the number of leaves was highest when plants were grown in soil from a low temperature origin. We found no evidence for either genetic variation in phenotypic plasticity or local adaptation. The microbial community in the soil advanced bulbil sprouting, with some weak evidence for delayed sprouting of plants originating from high temperature soils when they were grown at low soil temperature in the absence of soil biota.

Plants sprouted earlier and had more leaves with higher experimental soil temperature. This is consistent with previous studies that have shown an advance of spring phenology and flowering, as well as increased plant growth, with increasing temperature (Arft et al. 1999, Valdés et al. 2018). For example, flowering phenology has been shown to advance with heating in geothermal areas (Anderson et al. 2012, Valdés et al. 2018), and plant biomass is higher in *Arctagrostis latifolia* and *Carex bigelowii* when grown at higher soil temperature (Marchand et al. 2005). Taken together, these findings suggest that the circumpolar *B. vivipara* will start growing earlier, and grow larger, with climate change.

Surprisingly, the bulbils sprouted earliest in soil from the intermediate temperature origins, and plants had the highest number of leaves when grown in soil from low temperature origins. This did not match our expectation that the higher mineralization rates, and associated higher availability of nutrients, in soils from high temperature would result in earlier sprouting and increased performance. Previous studies have shown that nutrients are frequently a limiting factor in arctic ecosystems and increased temperatures enhance nutrient

availability for plants and increase growth (Henry and Molau 1997, Weintraub and Schimel 2003, Semenchuk et al. 2015). Moreover, warmer soil is often associated with a more diverse soil community and higher soil biotic activity (Friberg et al. 2009, Zakharova and Spichak 2012). One explanation for the lack of the expected pattern is that nutrient variation across temperature gradients is limited in geothermal areas (O’Gorman et al. 2014, Sigurdsson et al. 2016), and even if mineralization is higher in soils in heated areas, this may not result in higher availability of nutrients when the soil is exposed to lower temperatures. Importantly, our findings illustrate that warming-mediated changes in the soil environment can have idiosyncratic effects on plant phenology and plant traits.

We found no evidence for genetic variation in phenotypic plasticity or local adaptation of plants to spatial variation in soil temperature. This contrasts to previous studies, which found adaptive patterns of phenotypic plasticity by comparing plants grown in the field and common garden (Valdés et al. 2018), and local adaptation using reciprocal transplant experiments (Conover and Schultz 1995, Joshi et al. 2001, Anderson et al. 2012). For example, Valdés et al. (2018) found that the arctic plant *Cerastium fontanum* flowered earlier when growing in heated areas than in non-heated areas in the field, but that individuals from heated areas flowered later than individuals from non-heated areas when growing in a common garden. The lack of counter-gradient variation and local adaptation in *B. vivipara* might have three non-mutually exclusive explanations. First, selection on the investigated traits may be weak or absent. Second, *B. vivipara* mostly reproduces clonally, and the lack of recombination may slow down the rate of local adaptation compared to sexually reproducing plant species. Indeed, sexually reproducing species used in other studies showed adaptive patterns of counter-gradient variation (Anderson et al. 2012, Valdés et al. 2018). Finally, gene flow may swamp the effect of natural selection, but as dispersal distances are low in *B. vivipara*, this appears less likely to explain the absence of an evolutionary response to

temperature in this species. To tease apart these explanations, future studies could use quantitative genetic approaches to assess additive genetic variation for the investigated traits, as combined with a genetic or experimental assessment of gene flow.

Interestingly, bulbil sprouting was slower in the sterile soil than in the field soil, indicating that the soil microbial community stimulates bulbil sprouting. The effect of soil biota matches previous studies, which found that soil organisms can advance phenology and influence plant growth (Wagner et al. 2014, Rasmussen et al. 2017, 2020). The delayed sprouting of bulbils originating from high temperature when grown at low soil temperature, but only when the soil was sterilized, may indicate that soil biota plays a role in the evolutionary response of plants to soil heating. However, we have no insights into the putative adaptive nature of this pattern.

Conclusions

Our findings highlight that the circumpolar alpine bistort has a strong plastic response to temperature. We did not find an evolutionary response in the form of counter-gradient variation or local adaptation. This contrasts with previous reports from other plant species in geothermal areas, and along latitudinal and altitudinal gradients, and may imply that the relative importance of plastic vs genetic responses is highly variable among plant species. Such variable responses, in combination with underlying mechanisms, have profound implications for our understanding of how the arctic vegetation will respond to a changing climate. For example, some plant species may lack genetic variation for selection to act on, and thereby show maladaptive plastic responses to changes in temperature. Other plant species may have additive genetic variation, but not show an evolutionary response in the current context due to high rates of gene flow. Still other plant species may have a relatively weak plastic response but show local adaptation across small spatial scales. Overall, we hope

that future studies will test for the ecological and evolutionary response of plants to changes in temperature across a broad range of plant species and environmental clines, and unravel the underlying mechanisms, which will allow for the development of eco-evolutionary models to predict the response of the arctic vegetation to changes in climate. This will give insights in the threats posed by climate change to the arctic vegetation and help inform about potential mitigation actions.

Declarations

We thank Jordbruksverket for the import permit (6.5.18-5502/16) for the soil and plant material used in this study. We thank Nia Sigrún Perron for help in the field. This research was supported by a grant from the Swedish Research Council (2015-03993 to AJMT) and funding from the University of Iceland postdoctoral grant to B.M. The authors declare no conflict of interest. NJW, PUR, JE and AJMT designed the experiment. BM collected field materials, and NJW carried out the experiment and analyzed the data. NJW wrote the manuscript, with contributions from all authors.

References

- Agrawal, A. A. 2001. Phenotypic plasticity in the interactions and evolution of species. - science 294: 321–326.
- Anderson, J. T. et al. 2012. Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in response to climate change. - Proc. R. Soc. B Biol. Sci. 279: 3843–3852.
- Arft, A. M. et al. 1999. Responses of tundra plants to experimental warming: meta-analysis of the international tundra experiment. - Ecol. Monogr. 69: 491–511.
- Bardgett, R. D. and van der Putten, W. H. 2014. Belowground biodiversity and ecosystem functioning. - Nature 515: 505–511.
- Bills, J. W. et al. 2015. Environmental and genetic correlates of allocation to sexual reproduction in the circumpolar plant *Bistorta vivipara*. - Am. J. Bot. 102: 1174–1186.
- Blanquart, F. et al. 2013. A practical guide to measuring local adaptation (D Ebert, Ed.). - Ecol. Lett. 16: 1195–1205.
- Bradshaw, A. D. 1952. Populations of *Agrostis tenuis* resistant to lead and zinc poisoning. - Nature 169: 1098–1098.
- Brady, K. U. et al. 2005. Evolutionary ecology of plant adaptation to serpentine soils. - Annu. Rev. Ecol. Evol. Syst. 36: 243–266.
- Callaghan, T. V. et al. 2004. Responses to projected changes in climate and uv-b at the species level. - AMBIO J. Hum. Environ. 33: 418–435.
- Conover, D. O. and Schultz, E. T. 1995. Phenotypic similarity and the evolutionary significance of countergradient variation. - Trends Ecol. Evol. 10: 248–252.
- Crawley, M. J. 2013. The R book. - Wiley.
- Fox, J. and Weisberg, S. 2019. An R compaion to applied regression. - Sage.

352 Frei, E. R. et al. 2014. Plant population differentiation and climate change: responses of
353 grassland species along an elevational gradient. - *Glob. Change Biol.* 20: 441–455.

354 Friberg, N. et al. 2009. Relationships between structure and function in streams contrasting in
355 temperature. - *Freshw. Biol.* 54: 2051–2068.

356 Gornish, E. S. and Prather, C. M. 2014. Foliar functional traits that predict plant biomass
357 response to warming (F de Bello, Ed.). - *J. Veg. Sci.* 25: 919–927.

358 Grether, G. F. 2005. Environmental change, phenotypic plasticity, and genetic compensation.
359 - *Am. Nat.* 166: E115–E123.

360 Gudmundsdottir, R. et al. 2011. Effects of temperature regime on primary producers in
361 Icelandic geothermal streams. - *Aquat. Bot.* 95: 278–286.

362 Henry, G. H. R. and Molau, U. 1997. Tundra plants and climate change: the international
363 tundra experiment (ITEX). - *Glob. Change Biol.* 3: 1–9.

364 2000. Lycopodiaceae to Polygonaceae (B Jonsell and T Karlsson, Eds.). - The Bergius
365 Foundation [u.a.].

366 Joshi, J. et al. 2001. Local adaptation enhances performance of common plant species. - *Ecol.*
367 *Lett.* 4: 536–544.

368 Kawecki, T. J. and Ebert, D. 2004. Conceptual issues in local adaptation. - *Ecol. Lett.* 7:
369 1225–1241.

370 Kristinsson, H. 2001. A guide to the flowering plants and ferns of Iceland. - *Mál og menning.*

371 Lenth, R. 2019. emmeans: Estimated Marginal Means, a Least-Square Means.

372 Marchand, F. L. et al. 2005. Performance of high arctic tundra plants improved during but
373 deteriorated after exposure to a simulated extreme temperature event. - *Glob. Change*
374 *Biol.* 11: 2078–2089.

375 Mursinoff, S. and Tack, A. J. M. 2017. Spatial variation in soil biota mediates plant
376 adaptation to a foliar pathogen. - *New Phytol.* 214: 644–654.

377 O’Gorman, E. J. et al. 2014. Climate change and geothermal ecosystems: natural laboratories,
378 sentinel systems, and future refugia. - *Glob. Change Biol.* 20: 3291–3299.

379 2015. Climate change 2014: synthesis report (RK Pachauri, L Mayer, and Intergovernmental
380 Panel on Climate Change, Eds.). - Intergovernmental Panel on Climate Change.

381 R Core Team 2019. R: A Language and Environment for Statistical Computing. - R
382 Foundation for Statistical Computing.

383 Rasmussen, P. U. et al. 2017. Plant and insect genetic variation mediate the impact of
384 arbuscular mycorrhizal fungi on a natural plant-herbivore interaction: AMF and
385 genetic variation. - *Ecol. Entomol.* 42: 793–802.

386 Rasmussen, P. U. et al. 2018. Multiscale patterns and drivers of arbuscular mycorrhizal
387 fungal communities in the roots and root-associated soil of a wild perennial herb. -
388 *New Phytol.* 220: 1248–1261.

389 Rasmussen, P. U. et al. 2020. The impact of elevated temperature and drought on the ecology
390 and evolution of plant–soil microbe interactions (B Singh, Ed.). - *J. Ecol.* 108: 337–
391 352.

392 Semenchuk, P. R. et al. 2015. Deeper snow alters soil nutrient availability and leaf nutrient
393 status in high Arctic tundra. - *Biogeochemistry* 124: 81–94.

394 Sigurdsson, B. D. et al. 2016. Geothermal ecosystems as natural climate change experiments:
395 The ForHot research site in Iceland as a case study. - *Icel. Agric. Sci.* 29: 53–71.

396 Tedersoo, L. et al. 2014. Global diversity and geography of soil fungi. - *Science* 346:
397 1256688.

398 Toftegaard, T. et al. 2016. Variation in plant thermal reaction norms along a latitudinal
399 gradient - more than adaptation to season length. - *Oikos* 125: 622–628.

400 Valdés, A. et al. 2018. A natural heating experiment: phenotypic and genotypic responses of
401 plant phenology to geothermal soil warming. - *Glob. Change Biol.*: gcb.14525.

402 Wagner, M. R. et al. 2014. Natural soil microbes alter flowering phenology and the intensity
403 of selection on flowering time in a wild *Arabidopsis* relative (J Klironomos, Ed.). -
404 Ecol. Lett. 17: 717–726.

405 Weintraub, M. N. and Schimel, J. P. 2003. Interactions between carbon and nitrogen
406 mineralization and soil organic matter chemistry in arctic tundra soils. - Ecosystems
407 6: 129–143.

408 Wittenmayer, L. and Merbach, W. 2005. Plant responses to drought and phosphorus
409 deficiency: contribution of phytohormones in root-related processes. - J. Plant Nutr.
410 Soil Sci. 168: 531–540.

411 Wolkovich, E. M. et al. 2012. Warming experiments underpredict plant phenological
412 responses to climate change. - Nature 485: 494–497.

413 Zakharova, O. K. and Spichak, V. V. 2012. Geothermal fields of Hengill volcano, Iceland. -
414 J. Volcanol. Seismol. 6: 1–14.

415

Table 1 The impact of original soil temperature (OST), original bulbil temperature (OBT) and experimental soil temperature (EST), as well as bulbil weight, on the performance of *Bistorta vivipara* in field soil. Shown are P-values, with significant values in bold. The number of replicates for each response variable is given within parentheses.

	Original soil temperature (OST)	Original bulbil temperature (OBT)	Experimental soil temperature (EST)	Bulbil weight	OST × OBT	OST × EST	OBT × EST	OST × OBT × EST
Sprouting success (n = 378)	0.856	0.849	<0.001	0.164	0.121	0.212	0.276	0.313
Day of sprouting (n = 240)	<0.001	0.094	<0.001	0.072	0.935	0.080	0.456	0.445
Plant height on day 84 (n = 217)	0.792	0.088	0.338	<0.001	0.790	0.106	0.753	0.468
Plant height on day 126 (n = 77)	0.401	0.648	0.529	0.030	0.561	0.337	0.623	0.102
Leaf number on day 84 (n = 217)	0.291	0.514	0.177	0.561	0.874	0.785	0.816	0.992
Leaf number on day 126 (n = 77)	0.012	0.118	0.004	0.984	0.939	0.772	0.559	0.999
Leaf size on day 97 (n = 162)	0.066	0.296	0.043	0.067	0.538	0.652	0.050	0.571
Leaf size on day 159 (n = 48)	0.162	0.924	0.055	0.521	0.296	0.025	0.074	-

419

Figure legends

Figure 1. The impact of experimental soil temperature on bulbil sprouting and performance of *Bistorta vivipara*. Shown are the impact of experimental soil temperature on A) probability of sprouting, B) day of sprouting, C) leaf size at 97 days after planting (DAP) and D) number of leaves at 126 days after planting. Shown are means $\pm$ standard errors.

Figure 2. The impact of original soil temperature on bulbil sprouting and performance of *Bistorta vivipara*. Shown are the impact of experimental soil temperature on A) day of sprouting and B) the number of leaves at 126 days after planting (DAP). Shown are means $\pm$ standard errors.

Figure 1.




Figure 2.

