



Genome-Wide Identification and Expression Analysis of Terpene Synthase Genes in *Cymbidium faberi*

Qian-Qian Wang¹, Meng-Jia Zhu², Xia Yu¹, Yuan-Yang Bi¹, Zhuang Zhou^{1,3}, Ming-Kun Chen¹, Jiating Chen¹, Diyang Zhang¹, Ye Ai¹, Zhong-Jian Liu^{1,3,4*} and Siren Lan^{12*}

¹Key Laboratory of National Forestry and Grassland Administration for Orchid Conservation and Utilization at College of Landscape Architecture, Fujian Agriculture and Forestry University, Fuzhou, China, ²College of Forestry, Fujian Agriculture and Forestry University, Fuzhou, China, ³Zhejiang Institute of Subtropical Crops, Zhejiang Academy of Agricultural Sciences, Wenzhou, China, ⁴Institute of Vegetable and Flowers, Shandong Academy of Agricultural Sciences, Jinan, China

OPEN ACCESS

Edited by:

Xinchao Wang,
Tea Research Institute,
Chinese Academy of Agricultural
Sciences (CAAS), China

Reviewed by:

Xinlu Chen,
The University of Tennessee,
Knoxville, United States
Mingle Wang,
Huazhong Agricultural University,
China

*Correspondence:

Zhong-Jian Liu
zjliu@fafu.edu.cn
Siren Lan
lkzx@fafu.edu.cn

Specialty section:

This article was submitted to
Crop and Product Physiology,
a section of the journal
Frontiers in Plant Science

Received: 02 August 2021

Accepted: 01 November 2021

Published: 25 November 2021

Citation:

Wang Q-Q, Zhu M-J, Yu X, Bi Y-Y,
Zhou Z, Chen M-K, Chen J, Zhang D,
Ai Y, Liu Z-J and Lan S (2021)
Genome-Wide Identification and
Expression Analysis of Terpene
Synthase Genes in *Cymbidium faberi*.
Front. Plant Sci. 12:751853.
doi: 10.3389/fpls.2021.751853

Terpene synthases (TPSs) are essential for forming terpenes, which play numerous functional roles in attracting pollinators, defending plants, and moderating the interaction between plants. TPSs have been reported in some orchids, but genome-wide identification of terpenes in *Cymbidium faberi* is still lacking. In this study, 32 putative TPS genes were classified in *C. faberi* and divided into three subfamilies (TPS-a, TPS-b, and TPS-e/f). Motif and gene structure analysis revealed that most *CfTPS* genes had the conserved aspartate-rich DDxxD motif. TPS genes in the TPS-a and TPS-b subfamilies had variations in the RRX₈W motif. Most *cis*-elements of *CfTPS* genes were found in the phytohormone responsiveness category, and MYC contained most of the numbers associated with MeJA responsiveness. The *Ka/Ks* ratios of 12/13 *CfTPS* gene pairs were less than one, indicated that most *CfTPS* genes have undergone negative selection. The tissue-specific expression patterns showed that 28 genes were expressed in at least one tissue in *C. faberi*, and TPS genes were most highly expressed in flowers, followed by leaves and pseudobulbs. In addition, four *CfTPS* genes were selected for the real-time reverse transcription quantitative PCR (RT-qPCR) experiment. The results revealed that *CfTPS12*, *CfTPS18*, *CfTPS23*, and *CfTPS28* were mainly expressed in the full flowering stage. *CfTPS18* could convert GPP to β -myrcene, geraniol, and α -pinene *in vitro*. These findings of *CfTPS* genes of *C. faberi* may provide valuable information for further studies on TPSs in orchids.

Keywords: terpenes, terpene synthase, orchids, floral scent, expression analysis, *Cymbidium*

INTRODUCTION

Terpenes, which contain isoprene (C₅), monoterpenes (C₁₀), sesquiterpenes (C₁₅), and diterpenes (C₂₀), constitute the most prominent family of volatile components in plants (Nagegowda, 2010; Tholl, 2015). They play significant roles in attracting pollinators, defending plants against herbivores and pathogens, and moderating the interaction between plants (Wagner and Elmadfa, 2003; Dudareva et al., 2006). Terpenoids are dominant compounds in orchid floral scents (Ramya et al., 2018). For example, sesquiterpenes are the major floral scent compounds in *Cymbidium goeringii* (Ramya et al., 2019). The monoterpenes linalool, ocimene, and linalool

oxide are the major scent components of *Vanda* Mini Palmer flowers (Mohd-Hairul et al., 2010). The floral scent in orchids is closely related to flower development stages and is vital for pollination ecology (Ramya et al., 2018).

Isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) are precursors of terpene formation, and they are generated by cytosolic mevalonate acid (MVA) and the plastid methylerythritol phosphate (MEP) pathways (Dudareva et al., 2006). A plastid prenyltransferase synthesizes geranyl diphosphate (GPP), and the second type of plastid prenyltransferase produces geranylgeranyl diphosphate (GGPP) from the condensation of IPP and DMAPP. The condensation of DMAPP and IPP forms farnesyl diphosphate (FPP) in the cytosol (Figure 1). Plants have terpene synthases (TPSs) that catalyze monoterpene formation, sesquiterpene formation, and diterpene formation from GPP, FPP, and GGPP, respectively (Pichersky et al., 2006; Vranová et al., 2013).

Most full-length TPSs contain two conserved domains defined in PFAM: PF01397 (N-terminal) containing a conserved RRX_8W motif and PF03936 (C-terminal) containing a $DDxxD$ motif and NSE/DTE motif (El-Gebali et al., 2019; Jiang et al., 2019). In addition, TPSs can be divided into seven major categories: TPS-a, TPS-b, TPS-c, TPS-d, TPS-e/f, TPS-g, and TPS-h (Chen et al., 2011). TPS-a, TPS-b, and TPS-g categories are angiosperm-specific. The TPS-a category can encode sesquiterpene synthases of these three categories and can be further classified into monocot-specific TPS-a-1 and dicot-specific TPS-a-2 groups. In recent reports, all characterized TPSs in TPS-b with RRX_8W motifs are monoterpene synthases. The TPS-g category can encode monoterpene synthases without the RRX_8W motif. In addition, TPS-d is a gymnosperm-specific category that encodes monoterpenes, sesquiterpenes, and diterpenes. TPS-e/f can encode kaurene or copalyl diphosphate synthases that function in gibberellic acid synthesis in vascular plants. TPS-c may represent the ancestral category, and TPS-h only appears in *Selaginella moellendorffii* in recent reports (Chen et al., 2011; Li et al., 2012).

Genome-wide TPS families have been identified in *Arabidopsis thaliana*, *Vitis vinifera*, *Malus domestica*, *Camellia sinensis*, *Glycine max*, *Daucus carota*, and *Dendrobium officinale* (Aubourg et al., 2002; Martin et al., 2010; Nieuwenhuizen et al., 2013; Liu et al., 2014; Keilwagen et al., 2017; Yu et al., 2020; Zhou et al., 2020). Terpenoids are predominant components in orchid flower volatiles, but genome-wide TPS identification is limited in orchids (Ramya et al., 2018; Yu et al., 2020). Orchidaceae comprises five subfamilies, and in recent reports, a few TPS genes have been identified in *Apostasia shenzhenica* in Apostasioideae subfamily; *Vanilla planifolia* in Vanilloideae subfamily; and *Phalaenopsis equestris* and *Dendrobium catenatum* in Epidendroideae subfamily

(Huang et al., 2021). However, genome-wide TPS genes identification in Orchidoideae was limited.

Cymbidium faberi is a plant of Orchidaceae with a long history of cultivation due to its characteristic flower fragrance and beautiful flower shape (Ramya et al., 2018). There are over 100 compounds in the *C. faberi* floral scent, and some terpenes have been identified in the blooming *C. faberi* flowers by headspace vapors (Omata et al., 1990). This study first analyzed the classification, phylogenetics, and expression patterns of TPS genes in *C. faberi*. Our results will provide valuable information for further studies on *C. faberi* and other orchids.

MATERIALS AND METHODS

Plant Materials

The wild *C. faberi* collected in this study were cultivated in the greenhouse at Forest Orchid Garden in Fujian Agriculture and Forestry University (Fuzhou, Fujian Province, China) under natural light and temperatures. The temperature was about 20–25°C. Flowers, leaves, and pseudobulbs of *C. faberi* were sampled at the flowering stage. The buds about 1 cm before anthesis, semi-open flowers about 3 cm, and fully open flowers were also used in this study. All samples of *C. faberi* were frozen in liquid nitrogen for storage at –80°C until use.

Identification of *Cf*TPS Genes in the *C. faberi* Protein Database

Two domains – PF01397 representing the TPS N-terminal domain and PF03936 representing the TPS C-terminal domain from PFAM¹ – were used as queries to search the *C. faberi* protein database (El-Gebali et al., 2019). The *C. faberi* genome data will be published separately. An HMM search (built-in Tbttools) was used in this study with an e-value cut at 10^{-3} . To avoid missing potential TPS genes, TPS sequences from *A. thaliana* in the TAIR database² were also used to screen the *C. faberi* protein database using BLASTP (built-in Tbttools; Chen et al., 2018). The candidate TPS genes were checked manually by Pfam to verify putative full-length TPS genes, and TPS genes lacking either PF03936 or PF01397 were excluded. The grand average of hydrophobicity (GRAVY), molecular weight (Mw), isoelectric points (pI), aliphatic index (AI), and instability index (II) of the TPS proteins were predicted by the ExPASy database (Artimo et al., 2012).³ Subcellular localization was predicted by Plant-mPloc (Chou and Shen, 2010),⁴ AtSubP (Kaundal et al., 2010),⁵ and Ploc-mPlant (Cheng et al., 2017).⁶ Terzyme⁷ and BLATP⁸ were used to predict gene function (Priya et al., 2018).

Abbreviations: AI, Aliphatic index; bp, Base pair; DMAPP, Dimethylallyl diphosphate; FPP, Farnesyl diphosphate; FPKM, Fragments per kilobase of transcript per million fragments mapped; GGPP, Geranylgeranyl diphosphate; GPP, Geranyl diphosphate; GRAVY, Grand average of hydrophobicity; II, Instability index; IPP, Isopentenyl diphosphate; MeJA, Methyl jasmonate; MEP, Methylerythritol phosphate; Mw, Molecular weight; MVA, Mevalonate acid; NCBI, National Center for Biotechnology Information; NJ, Neighbor-joining; pI, Isoelectric point; RT-qPCR, Real-time reverse transcription quantitative PCR; TPS, Terpene synthase.

¹<http://pfam.xfam.org/>

²<https://www.arabidopsis.org/>

³<https://www.expasy.org/>

⁴<http://www.csbio.sjtu.edu.cn/bioinf/plant-multi/>

⁵<http://bioinfo3.noble.org/AtSubP/>

⁶<https://www.swmath.org/software/25191>

⁷<http://www.nipgr.ac.in/terzyme.html>

⁸<http://blast.ncbi.nlm.nih.gov>




FIGURE 1 | The pathway of plant terpenes biosynthesis. Terpenes are synthesized from two precursors generated by the cytosolic mevalonate acid (MVA) and plastid methylerythritol phosphate (MEP) pathways. G3P, glyceraldehyde 3-phosphate; IPP, isopentenyl diphosphate; DMAPP, dimethylallyl diphosphate; GPP, geranyl diphosphate; GPPS, GPP synthase; FPP, farnesyl diphosphate; FPPS, FPP synthase; GGPP, geranylgeranyl diphosphate; GGPPS, GGPP synthase; and TPS, terpene synthase.

Motifs and Gene Structure Analysis

Conserved motifs in the *C. faberi* TPS sequences were employed and analyzed by MEME software⁹ with default parameters (Bailey et al., 2009). We identified 20 motifs in this study. The exon-intron structure of the sequences was determined using GSDS software (Hu et al., 2015).¹⁰

Phylogenetic Analysis of TPS Genes

The transcriptomes of TPS sequences from *P. equestris* and *A. shenzhenica* were downloaded from their genome databases. TPS sequences from *S. moellendorffii* were downloaded from NCBI.¹¹ TPS sequences from *Picea abies* were downloaded from UniProt,¹² and TPS protein sequences from *A. thaliana* and *Oryza sativa* were downloaded from Phytozome.¹³ All these sequences were aligned with MAFFT (Rozewicki et al., 2019). The maximum likelihood (ML) method was used for the phylogenetic tree, which was constructed with RAxML on the CIPRES Science Gateway web server (RAxML-HPC2 on XSEDE; Miller et al., 2011). Bootstrap values were 1,000 replicates with the JTT model. The most appropriate protein evolution model for the alignment was predicted by ProTest (Darrriba et al., 2011). The generated tree was redrawn and annotated by EVOLVIEW (He et al., 2016).¹⁴ The sequences of the *CfTPS* proteins used in this study are listed in **Supplementary Table S7**.

Promoter Element Analysis of TPS Genes

Tbtools software extracted the promoter sequences and 2,000 bp regions upstream of 32 *CfTPS* genes (Chen et al., 2018). Afterward, the online software PlantCARE¹⁵ was used to identify

the *cis*-active regulatory elements in the promoter regions of the *CfTPS* genes (Lescot et al., 2002).

Calculation of *Ka* and *Ks* Ratios

Gene pairs with similar genetic relationships were selected based on the phylogenetic tree. DNAMAN software was used to select the gene pairs with a consistency greater than 60%. Tbtools software was then used to calculate *Ka* (non-synonymous rate), *Ks* (synonymous substitution), and *Ka/Ks* (evolutionary constraint) values. Divergence time (*T*) was calculated by using the formula $T = Ks / (2 \times 9.1 \times 10^{-9}) \times 10^{-6}$ million years ago (Mya; Zhang et al., 2018). In general, *Ka/Ks* < 1.0 represents purifying or negative selection, *Ka/Ks* = 1.0 represents neutral selection, and *Ka/Ks* > 1.0 represents positive selection (Zhang et al., 2006).

Transcriptome Data and GO Classification Analysis

An RNA sequencing transcriptome database of leaves, pseudobulbs, petals, sepals, labellums, and gynostemium was established to study the expression patterns of *CfTPS* genes. Fragments per kilobase of transcript per million mapped reads (FPKM) values of *CfTPS* genes were used to evaluate translation abundance. DESeq was used to conduct gene differential expression analysis, and gene ontology (GO) classification analysis was performed based on the differentially expressed gene analysis. The heatmaps of *CfTPS* expression patterns were drawn by Tbtools software, and the color in the heatmap was expressed as the log₂-transformed expression levels of each *CfTPS* gene (Chen et al., 2018).

Extraction of RNA and RT-qPCR Analysis

RNA was isolated from flowers of *C. faberi* at the flowering stage using the Biospin Plant Total RNA Extraction Kit (Bioer Technology, Hangzhou, China). First-strand DNA was synthesized with TransScript® All-in-One First-Strand cDNA Synthesis SuperMix for quantitative PCR (qPCR; TransGen Biotech, Beijing, China). TransScript® All-in-One First-Strand cDNA

⁹<http://meme-suite.org/>

¹⁰<http://gsds.gao-lab.org/>

¹¹<https://www.ncbi.nlm.nih.gov/genbank/>

¹²<https://www.UniProt.org/>

¹³<https://phytozome.jgi.doe.gov/>

¹⁴<https://evolgenius.info/>

¹⁵<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>

Synthesis SuperMix for qPCR was also used to remove genomic DNA. The real-time reverse transcription quantitative PCR (RT-qPCR) primers of *CfTPS* were designed by Primer Premier 5 software and can be found in **Supplementary Table S12**. Primer specificity was confirmed using Primer-BLAST on the NCBI website.¹⁶ PerfectStart™ Green qPCR SuperMix (TransGen Biotech, Beijing, China) was used for RT-qPCR analysis. The *C. faberi* reference gene GAPDH (GenBank Accession: JX560732; Tian et al., 2020) was used as the internal control and quantified by the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). There were three biological replicates in the RT-qPCR analysis.

***CfTPS18* Enzyme Assays**

The ORF of *CfTPS18* was synthesized and ligated into pET28a vector. Then, the recombinant plasmid was transformed into *Escherichia coli* BL21 (DE3) pLysS cells (Transgen, China). Primers are given in **Supplementary Table S12**. The positive clones were incubated with shaking at 200 rpm, 37°C until $OD_{600}=0.6$, and then at 37°C for 3 h with shaking and 0.1 mM IPTG. Recombinant *CfTPS18* enzyme was induced at 16°C for 16 h with 0.1 mM IPTG. The precipitate was resuspended in extraction buffer [50 mM NaH_2PO_4 , 500 mM NaCl, 10% (v/v) glycerol, and pH 7.0] and disrupted with a sonicator at 200 W for 60 s. The protein was purified with Ni-NTA agarose (Clontech). Purified *CfTPS18* protein was examined by SDS-PAGE using Tris-HCl buffer (pH 7.5). The recombinant protein was performed in assay buffer [25 mM HEPES, 10 mM $MgCl_2$, 100 mM KCl, 5 mM DTT, 10% (v/v) glycerol, and 30 μ M GPP], and 30 μ g of *CfTPS18* protein at pH 7.2 and 30°C for 1 h²⁴.

GC-MS Analysis

The volatiles were exposed to SPME fiber with 50/30 μ m DVB/CAR/PDMS (divinylbenzene/carboxen/polydimethylsiloxane; Supelco Co., Bellefonte, PA, United States). The extract was analyzed using a gas chromatograph (Agilent 6890N) and mass spectrometer (Agilent 5975B, Santa Clara, CA, United States) outfitted with a silica capillary column (DB-5MS; 60 m $\times$ 0.25 mm $\times$ 0.25 μ m). The temperature program was as follows: 55°C for 2 min, 3°C min⁻¹ up to 80°C, 5°C min⁻¹ up to 180°C, 10°C min⁻¹ up to 230°C, and finally, 20°C min⁻¹ to 250°C. The ion source temperature was 230°C, and the electron energy was 70 eV. The GC-MS interface zone temperature was 250°C, and the scan range was 50–500 m/z. Reactions only added with GPP were used as the blank control. There were three biological replicates for the experiment. The retention time was compared with the NIST Mass Spectral Library.

RESULTS

TPS Gene Identification and Protein Features in *C. faberi*

To retrieve the TPS genes in *C. faberi*, two domains, PF01397 and PF03936 in the PFAM, were used to search the *C. faberi*

protein database (El-Gebali et al., 2019). BLASTP (built-in Tltools) was also used to screen the *C. faberi* protein database (Chen et al., 2018). After removing artefacts, 32 TPS genes were obtained. The deduced proteins of these genes were in the range of 115 for *CfTPS1* and *CfTPS2* to 902 amino acids for *CfTPS12* and had predicted molecular weights (Mw) in the range of 10.24 for *CfTPS12* to 103.89 KDa for *CfTPS31*. The theoretical isoelectric point (pI) values were in the range of 4.86 for *CfTPS1* and *CfTPS2* to 9.28 for *CfTPS26*, and the deduced grand average of hydrophilic (GRAVY) values were in the range of −0.358 for *CfTPS26* to 0.073 for *CfTPS6*, suggesting that most *CfTPS* proteins were hydrophilic except for *CfTPS1*, *CfTPS2*, and *CfTPS6*. Additionally, the aliphatic index (AI) of *CfTPS*-deduced proteins was in the range of 87.09 for *CfTPS21* to 107.22 for *CfTPS6*, and the instability index (II) was in the range of 28.62 for *CfTPS1* and *CfTPS2* to 52.36 for *CfTPS5*. To retrieve information on the subcellular localization of *CfTPS* proteins, three predictors were used in this study: AtSubp, Plant-mPLoc, and pLoc-mPLant (Chou and Shen, 2010; Kaundal et al., 2010; Cheng et al., 2017). The results showed that 16 *CfTPS* proteins were marked in the chloroplast, and 16 *CfTPS* proteins were marked in chloroplast or cytoplasm, indicating that these three predictors produce different results and need to be further analyzed. We also annotated 32 *CfTPS* genes using BLASTP¹⁷ and Terzyme software (**Supplementary Table S4**; Priya et al., 2018). The results showed that 11 *CfTPS*s were annotated sesquiterpene synthases, 15 *CfTPS*s were annotated monoterpene synthases, and six *CfTPS*s were annotated diterpene synthases. The secondary structure predicted by the SOPMA program revealed that the average of α -helices, random coils, extended strands, and β -turns comprised 70.86, 21.37, 4.35, and 3.63% of the structure, respectively (Geourjon and Deléage, 1995). The results showed that α -helices were predominant in all *CfTPS* proteins (**Table 1**).

Motif and Gene Structure Analysis

To understand the intron-exon structure of *CfTPS* genes, we analyzed TPS gene structure with GSDS software (Hu et al., 2015). The exons in *CfTPS* genes ranged in numbers from 2 to 15, and the results showed that most of the genes in the same category contained a similar intron-exon structure.

To further analyze the motifs of the *CfTPS* genes, we identified 20 motifs using MEME software (Bailey et al., 2009). The numbers of *C. faberi* TPS motifs ranged in length from 4 to 16. *CfTPS3* and *CfTPS23* contained the most motifs, with 16, while *CfTPS1*, *CfTPS2* had only four motifs. Motif 3 can be found in all *CfTPS* proteins except *CfTPS9* and *CfTPS16*. Motif 4 was also the most common *CfTPS* protein (28/32). Twenty-five *CfTPS* genes contained the RRX₈W motif (motif 1), and 30 *CfTPS* genes contained the DDxxD motif (motif 2; **Figure 2C**). Accordingly, different clusters have different forms of motifs. The same cluster's *CfTPS* proteins generally contained similar motifs. These results of intron-exon structure


¹⁶<https://blast.ncbi.nlm.nih.gov/Blast.cgi>

¹⁷<http://blast.ncbi.nlm.nih.gov>

TABLE 1 | A list of TPS genes in *C. faberi*, their characteristics, and functional annotation.

Gene ID ¹	Name	AA ² (aa)	pI ³	Mw ⁴ (kDa)	AI ⁵	II ⁶	GRAVY ⁷	Localization ⁸	Function ⁹
HL002300	<i>CfTPS1</i>	115	4.86	13.43	100.18	28.62	0.004	Chloroplast ^{a,b} /Cytoplasm ^c	C15
HL002301	<i>CfTPS2</i>	115	4.86	13.43	100.18	28.62	0.004	Chloroplast ^{a,b} /Cytoplasm ^c	C15
HL003318	<i>CfTPS3</i>	609	6.00	70.70	97.68	47.96	-0.274	Chloroplast ^{a,b,c}	C10
HL003423	<i>CfTPS4</i>	806	6.03	90.68	93.49	44.18	-0.114	Chloroplast ^{a,b,c}	C20
HL008810	<i>CfTPS5</i>	374	6.02	43.53	96.27	52.36	-0.294	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL012959	<i>CfTPS6</i>	321	5.45	37.26	107.22	40.93	0.073	Chloroplast ^{a,b,c}	C20
HL015149	<i>CfTPS7</i>	435	5.30	49.81	98.66	48.37	-0.109	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL015150	<i>CfTPS8</i>	448	5.43	51.45	101.25	45.47	-0.112	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL017747	<i>CfTPS9</i>	429	6.08	48.50	98.13	46.48	-0.074	Chloroplast ^{a,b,c}	C20
HL018937	<i>CfTPS10</i>	250	5.00	28.44	91.77	46.95	-0.163	Chloroplast ^{a,b,c}	C10
HL020199	<i>CfTPS11</i>	405	5.17	46.73	92.00	42.44	-0.291	Chloroplast ^{a,b,c}	C20
HL021067	<i>CfTPS12</i>	902	6.18	10.24	95.35	38.05	-0.160	Chloroplast ^{a,b} /Nucleus ^a /Cytoplasm ^c	C10
HL023892	<i>CfTPS13</i>	455	5.18	52.81	101.21	42.22	-0.072	Chloroplast ^{a,b} /Cytoplasm ^c	C20
HL024326	<i>CfTPS14</i>	501	5.91	58.85	101.00	48.05	-0.139	Chloroplast ^{a,b} /Cytoplasm ^c	C20
HL024478	<i>CfTPS15</i>	528	6.35	62.29	98.60	35.68	-0.230	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL025052	<i>CfTPS16</i>	421	5.07	49.19	89.67	45.71	-0.270	Chloroplast ^{a,b,c}	C20
HL025282	<i>CfTPS17</i>	617	6.46	69.77	93.26	52.75	-0.124	Chloroplast ^a /Mitochondrion ^b /Cytoplasm ^c	C10
HL025366	<i>CfTPS18</i>	463	5.57	54.22	106.80	49.56	-0.174	Chloroplast ^{a,b,c}	C10
HL025643	<i>CfTPS19</i>	490	5.30	57.56	98.18	39.76	-0.095	Chloroplast ^{a,b,c}	C20
HL025987	<i>CfTPS20</i>	445	5.27	51.22	97.09	50.28	-0.055	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL026777	<i>CfTPS21</i>	653	8.57	75.5	87.09	42.52	-0.257	Chloroplast ^{a,b,c}	C20
HL026987	<i>CfTPS22</i>	178	5.09	21.16	104.18	37.38	-0.237	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL027027	<i>CfTPS23</i>	569	5.30	66.31	95.63	37.83	-0.277	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL027466	<i>CfTPS24</i>	508	5.17	59.33	98.70	41.50	-0.090	Chloroplast ^{a,b} /Cytoplasm ^c	C20
HL027610	<i>CfTPS25</i>	398	6.12	46.21	99.97	52.70	-0.221	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL027633	<i>CfTPS26</i>	562	9.28	65.60	87.47	42.35	-0.358	Chloroplast ^{a,b,c}	C20
HL028595	<i>CfTPS27</i>	281	5.83	33.37	99.21	47.13	-0.312	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL029155	<i>CfTPS28</i>	311	5.53	36.76	105.65	42.67	-0.126	Chloroplast ^{a,b} /Cytoplasm ^c	C10
HL029581	<i>CfTPS29</i>	550	5.38	64.34	101.58	50.85	-0.189	Chloroplast ^{a,b,c}	C20
HL029624	<i>CfTPS30</i>	454	5.60	53.09	100.68	49.99	-0.174	Chloroplast ^{a,b} /Cytoplasm ^c	C20
HL029782	<i>CfTPS31</i>	895	6.68	103.89	88.87	40.62	-0.249	Chloroplast ^{a,c} /Unknown ^b	C20
HL030142	<i>CfTPS32</i>	321	5.38	37.77	97.84	38.32	-0.085	Chloroplast ^a /Unknown ^b /Cytoplasm ^c	C20

¹Gene ID is annotated in *C. faberi* genome.²AA, exhibits amino acid.³pI, exhibits theoretical isoelectric point.⁴Mw, exhibits molecular weight.⁵AI, exhibits aliphatic index.⁶II, exhibits instability index.⁷GRAVY, exhibits the grand average of hydrophobicity.⁸Subcellular localization depended on Plant-mPloc, AtSubP, and Ploc-mPlant, respectively (Chou and Shen, 2010; Kaundal et al., 2010; Cheng et al., 2017).⁹Gene function predicted by Terzyme website.C10, C15, and C20 represent monoterpene, sesquiterpene, and diterpene, respectively. Row data is listed in **Supplementary Tables S1–S5**.^{a, b, c}exhibit Plant-mPloc, AtSubP, and Ploc-mPlant website results, respectively.



and motif analysis verified the closeness of the phylogenetic tree in *C. faberi* (Figure 2).

Phylogenetic Analysis of *CfTPS* Genes

A phylogenetic tree was constructed to analyze the evolutionary patterns of the *CfTPS* genes. Thirty-two *CfTPS* genes were used, and TPS protein sequences from six species (*A. shenzhenica*, *P. equetris*, *O. sativa*, *A. thaliana*, *P. abies*, and *S. moellendorffii*) were used. The maximum likelihood (ML) method was used for the phylogenetic tree, which was constructed with RAxML on the CIPRES Science Gateway web server (RAxML-HPC2 on XSEDE; Miller et al., 2011). Bootstrap values were 1,000 replicates with the JTT model. The phylogenetic tree indicated that TPS proteins belonged to seven categories. This classification result was consistent with a recent study: TPS-a, TPS-b, TPS-c, TPS-d, TPS-e/f, TPS-g, and TPS-h (Chen et al., 2011). Thirty-two *CfTPS* proteins belonged to three categories according to the phylogenetic tree (Figure 3): TPS-a, TPS-b, and TPS-e/f. Of these three categories, the TPS-a and TPS-b clades contained the most members and were the most expanded categories, with 13 and 15 genes, respectively, and were consistent with other plant species, such as *A. thaliana*, *C. sinensis*, *V. planifolia*, *D. catenatum*, *P. equestris*, and *D. officinale* (Aubourg et al., 2002; Yu et al., 2020; Zhou et al., 2020; Huang et al., 2021). The remaining four TPS genes belonged to the TPS-e/f subfamily.

We aligned the multiple sequences using MAFFT to analyze the TPS RRX₈W motif in the N-terminus, DDxxD, and NSE/DTE motifs in the C-terminus (Rozewicki et al., 2019). The alignment showed that almost all the *CfTPS* proteins in the TPS-a and TPS-b subfamilies had the highly conserved aspartate-rich motif DDxxD, except *CfTPS9* and *CfTPS16* in the TPS-a

subfamily and *CfTPS17*, *CfTPS27*, and *CfTPS28* in the TPS-b subfamily (Jiang et al., 2019). TPS genes in TPS-a and TPS-b had variations in the RRX₈W and RxR motif. However, in the TPS-a and TPS-b categories, the RRX₈W motif was not found in nine *CfTPS* genes. The RRX₈W motif was absent in the TPS-e/f subfamily, and the NSE/DTE motif was found in 16 *CfTPS* genes (Figure 4). Among them, DDxxD and NSE/DTE are important in the metal-dependent ionization of the prenyl diphosphate substrate, and the RRX₈W motif is essential in the cyclization of monoterpene synthase (Bohlmann et al., 1998; Chen et al., 2011; Jiang et al., 2019). In addition, the members in the TPS-a subfamily detected in both dicot and monocot plants encode only sesquiterpenes (Jiang et al., 2019). The secondary “R” in this family is not conserved (Martin et al., 2010). TPS-b encodes monoterpenes containing the conserved RRX₈W motif (Chen et al., 2011; Jiang et al., 2019).

Promoter Analysis of *CfTPS* Genes

To retrieve the potential function of *CfTPS* genes, we obtained a 2,000-bp region upstream of the 32 *CfTPS* genes and analyzed them using the online software Plantcare (Lescot et al., 2002). In total, we found 784 *cis*-acting regulatory elements in the promoter areas of *CfTPS* genes. We classified them into three categories according to the function of these elements: plant growth and development, stress responsiveness, and phytohormone responsiveness. The plant growth and dependent category (166/784) contained nine *cis*-acting regulatory elements and consists of AAGAA motifs, As-1 elements, A1-rich elements, etc. Most of them had As-1 elements (66/166), which are related to shoot expression. The stress responsiveness category (216/874) contained ARE, DRE, LTR, ST-RE, ABRE, etc.




FIGURE 3 | Phylogenetic tree of *CFTPS* genes based on the TPS protein sequences of seven plant species. The ML method was used for the phylogenetic tree, which was constructed with RAxML on the CIPRES Science Gateway web server (RAxML-HPC2 on XSEDE; Miller et al., 2011). The bootstrap values were 1,000 replicates with the JTT model. The generated tree was redrawn and annotated by the EVOLVIEW website (He et al., 2016). The TPS family was classified into seven categories: TPS-a, TPS-b, TPS-c, TPS-d, TPS-e/f, TPS-g, and TPS-h (Chen et al., 2011). TPS protein sequences in *Cymbidium faberi* are available in **Supplementary Table S7**.

STRE (48/216) is an essential element in the promoter that is related to stress. Interestingly, most of the *cis*-elements (401/784) were in the phytohormone responsiveness category, which contained CGTCA motifs, EREs, MYC motifs, TGAGG motifs, etc. Among them, most *cis*-elements were MYC (137/401), which is associated with MeJA responsiveness (Dombrecht et al., 2007). The results indicated that the *CfTPS* gene expression patterns might be regulated by MeJA treatment (Figure 5).

Ka/Ks Analysis in *C. faberi*

The Ka/Ks ratio can show positive selection ($Ka/Ks > 1$), negative or purifying selection ($Ka/Ks < 1$), and neutral selection ($Ka/Ks = 1$) during the evolution (Zhang et al., 2006). In this study, 13 gene pairs with similar genetic relationships were selected for Ka/Ks calculation. The results showed that the

Ka/Ks ratios of 12/13 *CfTPS* genes were less than one, indicated that most *CfTPS* genes underwent negative selection (Table 2). The divergence time of 13 *CfTPS* gene pairs was in the range of 0.75 for *CfTPS1* and *CfTPS2* to 60.46 for *CfTPS3* and *CfTPS10*.

Expression Patterns in Different *C. faberi* Organs and GO Classification of TPS Genes

An RNA sequencing transcriptome database of leaves, pseudobulbs, petals, sepals, labellums, and gynostemium was established to study the expression patterns of *CfTPS* genes. Eighteen genes were expressed in both leaves and flowers, and 20 genes were expressed in the labellums. Twenty-five genes were found in sepals and 24 in gynostemium. Four genes were not found to be expressed in any of the tissues.




FIGURE 4 | RRX&W, DDxxD, and NSE/DTE motifs in the *CfTPS* protein amino acid sequences. Multiple sequence alignments were constructed by MAFFT, and Jalview software was used to visualize the sequences (Troshin et al., 2011; Rozewicki et al., 2019).

CfTPS12, *CfTPS15*, and *CfTPS23* exhibited a high level of expression in leaves and pseudobulbs. Notably, *CfTPS3*, *CfTPS12*, *CfTPS15*, *CfTPS17*, *CfTPS18*, *CfTPS28*, and *CfTPS31* displayed high transcript abundance in floral organs, suggesting that these TPS genes might be related to flower scent in *C. faberi* (Figure 6).

Gene ontology classification analysis was performed based on the differentially expressed gene analysis. According to the classification results, molecular function contained most genes, and genes were mostly enriched in lyase activity, magnesium binding, and terpene synthase activity (Figure 7).


Expression Patterns of *CfTPS* in Flowers at Three Floral Developmental Stages

Real-time reverse transcription quantitative PCR was performed to investigate the expression patterns of *CfTPS* genes in flowers at three floral developmental stages. Four putative functional genes, *CfTPS12*, *CfTPS18*, *CfTPS23*, and *CfTPS28*, which are highly expressed in floral organs, were used. They were expressed differently among the three floral stages. Interestingly, these genes were mainly expressed in the full flowering stage (Figure 8).

Notably, *CfTPS18* had the highest transcript levels during the floral developmental stages, consistent with the expression patterns displayed in Figure 5. Gene annotation indicated that *CfTPS12*, *CfTPS18*, *CfTPS23*, and *CfTPS28* likely encoded monoterpene synthases and clustered in TPS-b subfamily (Table 1).

Functional Characterization of *CfTPS18* in *E. coli*

To investigate the enzyme activity, the full-length ORF sequence of *CfTPS18* was cloned to vector pET28a and ectopically expressed in *E. coli*. Recombinant *CfTPS18* enzyme was induced with 0.1 mM isopropyl-β-D-galactopyranoside and purified with Ni-NTA agarose (Clontech). SDS-PAGE analysis of *CfTPS18* recombinant protein can be found in Supplementary Figure S1. After using GPP as substrate in the reactions, the products were analyzed by GC-MS. The result indicated that *CfTPS18* could convert GPP into β-myrcene, geraniol, and α-pinene which were validated by the NIST Mass Spectral Library (Figure 9). The blank control with only GPP added to the reactions could not produce



monoterpenes. Thus, *CfTPS18* was considered a monoterpene synthase.

DISCUSSION

Orchids are among the largest angiosperm families in angiosperms and demonstrate a diversity of epiphytic and terrestrial growth forms (Hsiao et al., 2011). *Cymbidium faberi* is one of the longest cultivated orchids planted in China and has high ornamental value due to its characteristic flower scent and beautiful flower shape (Omata et al., 1990; Hsiao et al., 2011; Sun et al., 2016). Terpenoids play an essential role in floral scent and can attract pollinating insects and defend against environmental stresses (Wagner and Elmadfa, 2003; Dudareva et al., 2006; Sun et al., 2016). Plants have enzymes called TPSs, which encode the synthesis of monoterpene (C10), sesquiterpene (C15), or diterpenes (C20) from DMAPP, GPP, and GGPP, respectively (Pichersky et al., 2006). Terpenoids are dominant in the flower scent of orchids.

In this study, we systematically retrieved and classified TPS genes in *C. faberi*.

We identified 32 *CfTPS* genes in the *C. faberi* genome according to the TPS N-terminal and C-terminal domains (Table 1). We classified them into three categories: TPS-a, TPS-b, and TPS-e/f. TPS-b was the most expanded category, which was consistent with patterns in *D. officinale*, *V. planifolia*, and *D. catenatum*, and *C. faberi* have more genes in TPS-b than *P. equestris* (Yu et al., 2020; Huang et al., 2021). However, it was not consistent with *A. thaliana*, *O. sativa*, and *S. bicolor*, which have a dominant subfamily TPS-a (Aubourg et al., 2002; Chen et al., 2011). These results suggest that orchids have more TPS genes in TPS-b than other angiosperm dicot species and are related to emit floral scent to attract pollinators (Huang et al., 2021). According to the phylogenetic tree, most of the TPS-b genes are present in dicots. TPS-a genes can be further split into monocot-specific TPS-a-1 and dicot-specific TPS-a-2 clades. The TPS gene family is a medium-sized family, and the numbers of TPS genes range from approximately 20–100 (Chen et al., 2011). For instance,

TABLE 2 | Ka/Ks analysis of TPS genes in *C. faberi*.


Gene pairs	Ka ¹	Ks ²	Ka/Ks ³ Ratio	Date (Mya)
CfTPS5- CfTPS12	0.036349	0.060903	0.596841353	3.346319222
CfTPS18- CfTPS22	0.183699	0.58801	0.31240786	32.30822426
CfTPS27- CfTPS28	0.061668	0.086265	0.71486064	4.739845716
CfTPS15- CfTPS23	0.040242	0.038395	1.048087536	2.109640277
CfTPS3- CfTPS10	0.329855	1.100284	0.299790513	60.45518349
CfTPS20- CfTPS7	0.035647	0.054369	0.655636934	2.987330286
CfTPS25- CfTPS8	0.052051	0.057777	0.900881064	3.174578759
CfTPS1- CfTPS2	0	0.01373	0	0.754415417
CfTPS13- CfTPS16	0.207258	0.585853	0.353772093	32.1897194
CfTPS11- CfTPS29	0.108769	0.203168	0.535365599	11.16305388
CfTPS14- CfTPS30	0.035698	0.0446	0.800403031	2.450539375
CfTPS24- CfTPS32	0.028484	0.045777	0.622244514	2.515215271
CfTPS21- CfTPS26	0.04343	0.044138	0.983976748	2.425138016

¹Ka, non-synonymous rate.²Ks, synonymous substitution.³Ka/Ks, evolutionary constraint.


Divergence time (T) was calculated by using the formula $T = Ks / (2 \times 9.1 \times 10^{-9}) \times 10^{-6}$ million years ago (Mya, Zhang et al., 2018). The details of Ka/Ks calculation are listed in **Supplementary Table S8**.

14 *SmTPS*, 23 *GmTPS*, 32 *AtTPS*, 34 *DoTPS*, 23 *CsTPS*, and 69 *VvTPS* were found in *S. moellendorffii*, *G. max*, *A. thaliana*, *D. officinale*, *C. sinensis*, and *V. vinifera*, respectively (Li et al., 2012; Yu et al., 2020; Zhou et al., 2020). In addition, TPS occupied 0.26 genes/M in *A. thaliana* (125 M), 0.14 genes/M in *V. vinifera* (487 M), 0.13 genes/M in *S. moellendorffii* (106 M), 0.02 genes/M in *D. officinale* (1.35 G), 0.02 genes/M in *G. max* (1.011 G), and 0.01 genes/M in *C. faberi* (3.77 G; Jaillon et al., 2007; Schmutz et al., 2010; Poczar et al., 2014; Zhang et al., 2016). These results indicate that the TPS family has undergone expansion throughout the evolutionary history of land plants, and different species may show a difference in the expansion mechanism (Chen et al., 2011; Jiang et al., 2019).

We also annotated 32 *CfTPS* genes, and the results showed that all TPSs in the TPS-a clade encode sesquiterpenes, and all TPSs in the TPS-b clade encode monoterpenes (Table 1). All the *CfTPS* in the TPS-e/f clade were annotated as diterpene synthases. This is consistent with a recent study in which most TPS genes in the TPS-a subfamily were determined to be sesquiterpene synthases. TPS-e/f encoded monoterpene, sesquiterpene, and diterpene in a recent study (Chen et al., 2011). Sesquiterpenes, diterpenes, and monoterpenes are important to emit floral scents to attract pollinators and defend against environmental stress (Huang et al., 2021).

**FIGURE 6** | Expression patterns of *CfTPS* genes in different organs. The heatmap was produced in Tbttools (Chen et al., 2018). The tissues were leaves (L), pseudobulbs (Ps), petals (Pe), sepals (Se), labellum (La), and gynostemium (Gs) in wild *C. faberi*. The fragments per kilobase of transcript per million fragments (FPKM) values are listed in **Supplementary Table S9**.

Each full-length TPS had two conserved domains, including the N-terminal domain containing the RRX_8W motif and the C-terminal domain containing two highly conserved aspartate-rich motifs: DDxxD and NSE/DTE (Starks et al., 1997; Jiang et al., 2019). DDxxD and NSE/DTE are significant in the metal-dependent ionization of the prenyl diphosphate substrate, and the RRX_8W motif is essential in the cyclization of monoterpene synthase (Bohlmann et al., 1998; Chen et al., 2011; Jiang et al., 2019). The secondary "R" in the TPS-a subfamily is not conserved (Martin et al., 2010). TPS-b contains the conserved RRX_8W motif, which is related to monoterpene formation. TPS-c does not include the DDxxD motif (Chen et al., 2011; Jiang et al., 2019). Motif RxR was also conserved in TPS-a and TPS-b subfamilies. In this study, multiple sequence alignments showed that 28/32 *CfTPS* had highly conserved DDxxD motifs, 16/32 *CfTPS* had RRX_8W motifs, 16/32 *CfTPS* had NSE/DTE motifs, and 9/32 *CfTPS* had RxR motifs. The RRX_8W motif was not found in 11 *CfTPS*s in the TPS-a and TPS-b clades, and it was absent in the TPS-e/f subfamily. The results indicated that motif loss might have appeared during family evolution in *C. faberi* and that different subfamilies have different motif features.



Different elements were observed in promoter areas of *CfTPS* genes. Most of the *cis*-elements were in the phytohormone responsiveness group, and the number of MYCs associated with MeJA responsiveness contained most of this group. The results indicated that the expression patterns of *CfTPS* may be regulated by MeJA treatment and may respond to multiple environmental stresses (Chaiprasongsuk et al., 2018). In recent studies, MeJA was shown to regulate TPS gene expression in *D. officinale* and

C. sinensis (Yu et al., 2020; Zhao et al., 2020; Zhou et al., 2020). However, this needs to be further investigated in *C. faberi*. The *Ka/Ks* ratio analysis indicated that the TPS gene family in *C. faberi* mainly underwent negative selection, making it more stable during the evolution. GO annotation analysis of *CfTPS* genes indicated that molecular function contained most genes, and genes were mostly enriched in lyase activity, magnesium binding, and terpene synthase activity. Expression pattern analysis indicated that *CfTPS*




FIGURE 8 | Real-time reverse transcription quantitative PCR (RT-qPCR) validation of transcriptomic data of the *CfTPS* genes at three flowering stages. B, budding flowers; S, semi-open flowers; and F, fully open flowers. The error bars indicate three RT-qPCR biological replicates. The values were standardized by the *C. faberi* reference gene GAPDH (GenBank Accession: JX560732; Tian et al., 2020). The expression values of *CfTPS* genes are listed in **Supplementary Table S11**, and the RT-qPCR primers of *CfTPS* are listed in **Supplementary Table S12**.




FIGURE 9 | Enzymatic assays after incubating recombinant *CfTPS18*. Reactions only performed with GPP were used as the blank control. The retention time was compared with the NIST Mass Spectral Library.

genes were mainly expressed in the floral organs of *C. faberi*, indicating that they were related to floral scent. We selected four *CfTPS* genes with the highest transcript levels in floral organs for qRT-PCR analysis at three flowering stages. The results showed that they all belonged to the

TPS-b clade and were mainly expressed in the full flowering stage. According to our annotation, four genes were all annotated as monoterpene synthases, which may play essential roles in floral scent and attracting pollinators in *C. faberi*. Enzymatic assays suggested that *CfTPS18* could convert GPP to β -myrcene, geraniol, and α -pinene. In recent reports, *DoTPS10* in *D. officinale* can convert GPP to linalool *in vitro* (Yu et al., 2020). *DoGES1* in *D. officinale* can catalyze geraniol *in vitro* and *in vivo* (Zhao et al., 2020). Linalool and geraniol belong to monoterpenes which play essential roles in floral scents. TPS that catalyze terpenes can also be found in *V. vinifera*, *M. domestica*, and *Litsea cubeba* (Martin et al., 2010; Nieuwenhuizen et al., 2013; Chen et al., 2020). To better understand terpenes production and function in *C. faberi*, more studies of expression profiles should be developed.

CONCLUSION

In this study, 32 *CfTPS* genes were identified from the genomes of *C. faberi*. We analyzed their conserved motifs, exon-intron structure, phylogeny, *Ka/Ks* ratios, and *cis*-acting regulatory elements. We also analyzed the expression patterns of *CfTPS* genes in leaves, pseudobulbs, petals, sepals, labellums, and gynostemium in wild *C. faberi*. Four putatively functional genes highly expressed in floral organs were used to analyze the expression patterns at three flowering stages. Enzymatic assays

indicated that *CfTPS18* could convert GPP to β -myrcene, geraniol, and α -pinene. The results will provide valuable information for further studies on floral scents in *C. faberi* and other orchids.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding authors.

AUTHOR CONTRIBUTIONS

SL, Z-JL, and DZ contributed to conceptualization and validation. Q-QW, M-JZ, and ZZ prepared the original draft. Q-QW, Y-YB, and XY analyzed the data. M-KC, YA, and JC contributed to the visualization. All authors contributed to the article and approved the submitted version.

REFERENCES

- Artimo, P., Jonnalagedda, M., Arnold, K., Baratin, D., Csardi, G., De Castro, E., et al. (2012). ExPASy: SIB bioinformatics resource portal. *Nucleic Acids Res.* 40, 597–603. doi: 10.1093/nar/gks400
- Aubourg, S., Lecharny, A., and Bohlmann, J. (2002). Genomic analysis of the terpenoid synthase (*AtTPS*) gene family of *Arabidopsis thaliana*. *Mol. Gen. Genomics* 267, 730–745. doi: 10.1007/s00438-002-0709-y
- Bailey, T. L., Boden, M., Buske, F. A., Frith, M., Grant, C. E., Clementi, L., et al. (2009). MEME suite: tools for motif discovery and searching. *Nucleic Acids Res.* 37, 1–7. doi: 10.1093/nar/gkp335
- Bohlmann, J., Meyer-Gauen, G., and Croteau, R. (1998). Plant terpenoid synthases: molecular biology and phylogenetic analysis. *Proc. Natl. Acad. Sci. U. S. A.* 95, 4126–4133. doi: 10.1073/pnas.95.8.4126
- Chaiprasongsuk, M., Zhang, C., Qian, P., Chen, X., Li, G., Trigiano, R. N., et al. (2018). Biochemical characterization in Norway spruce (*Picea abies*) of SABATH methyltransferases that methylate phytohormones. *Phytochemistry* 149, 146–154. doi: 10.1016/j.phytochem.2018.02.010
- Chen, C., Chen, H., He, Y., and Xia, R. (2018). TBtools, a toolkit for biologists integrating various biological data handling tools with a user-friendly interface. *bioRxiv* [Preprint]. doi: 10.1101/289660
- Chen, Y. C., Li, Z., Zhao, Y. X., Gao, M., Wang, J. Y., Liu, K. W., et al. (2020). The *Litsea* genome and the evolution of the laurel family. *Nat. Commun.* 11, 1–14. doi: 10.1038/s41467-020-15493-5
- Chen, F., Tholl, D., Bohlmann, J., and Pichersky, E. (2011). The family of terpene synthases in plants: a mid-size family of genes for specialized metabolism that is highly diversified throughout the kingdom. *Plant J.* 66, 212–229. doi: 10.1111/j.1365-3113.2011.04520.x
- Cheng, X., Xiao, X., and Chou, K. C. (2017). PLoc-mPlant: predict subcellular localization of multi-location plant proteins by incorporating the optimal GO information into general PseAAC. *Mol. BioSyst.* 13, 1722–1727. doi: 10.1039/C7MB00267J
- Chou, K. C., and Shen, H. B. (2010). Plant-mPLoc: a top-down strategy to augment the power for predicting plant protein subcellular localization. *PLoS One* 5:e11335. doi: 10.1371/journal.pone.0011335
- Darriba, D., Taboada, G. L., Doallo, R., and Posada, D. (2011). ProtTest 3: fast selection of best-fit models of protein evolution. *Bioinformatics* 27, 1164–1165. doi: 10.1093/bioinformatics/btr088
- Dombrecht, B., Gang, P. X., Sprague, S. J., Kirkegaard, J. A., Ross, J. J., Reid, J. B., et al. (2007). MYC2 differentially modulates diverse jasmonate-dependent functions in *Arabidopsis*. *Plant Cell* 19, 2225–2245. doi: 10.1105/tpc.106.048017

FUNDING

This research was funded by The National Key Research and Development Program of China (2019YFD1001000), The National Natural Science Foundation of China (no. 31870199), The Key Laboratory of National Forestry and Grassland Administration for Orchid Conservation and Utilization Construction Funds (Grant 115/118990050, 115/KJG18016A), Natural Science Foundation of Zhejiang Province (Grant nos. LY20C160005, LY19C150003), Key Research and Development Program of Zhejiang Province (Grant no. 2021C02043), and Wenzhou Agricultural New Variety Breeding Cooperative Group Project (Grant no. 2019ZX004-3).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2021.751853/full#supplementary-material>

- Dudareva, N., Negre, F., Nagegowda, D. A., and Orlova, I. (2006). Plant volatiles: recent advances and future perspectives. *Crit. Rev. Plant Sci.* 25, 417–440. doi: 10.1080/07352680600899973
- El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C., et al. (2019). The Pfam protein families database in 2019. *Nucleic Acids Res.* 47, D427–D432. doi: 10.1093/nar/gky995
- Geourjon, C., and Deléage, G. (1995). Sopma: significant improvements in protein secondary structure prediction by consensus prediction from multiple alignments. *Comput. Appl. Biosci.* 11, 681–684. doi: 10.1093/bioinformatics/11.6.681
- He, Z., Zhang, H., Gao, S., Lercher, M. J., Chen, W. H., and Hu, S. (2016). Evolvview v2: an online visualization and management tool for customized and annotated phylogenetic trees. *Nucleic Acids Res.* 44, W236–W241. doi: 10.1093/nar/gkw370
- Hsiao, Y. Y., Pan, Z. J., Hsu, C. C., Yang, Y. P., Hsu, Y. C., Chuang, Y. C., et al. (2011). Research on orchid biology and biotechnology. *Plant Cell Physiol.* 52, 1467–1486. doi: 10.1093/pcp/pcr100
- Hu, B., Jin, J., Guo, A. Y., Zhang, H., Luo, J., and Gao, G. (2015). GSDS 2.0: an upgraded gene feature visualization server. *Bioinformatics* 31, 1296–1297. doi: 10.1093/bioinformatics/btu817
- Huang, L. M., Huang, H., Chuang, Y. C., Chen, W. H., Wang, C. N., and Chen, H. H. (2021). Evolution of terpene synthases in Orchidaceae. *Int. J. Mol. Sci.* 22:6947. doi: 10.3390/ijms22136947
- Jaillon, O., Aury, J. M., Noel, B., Policriti, A., Clepet, C., Casagrande, A., et al. (2007). The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* 449, 463–467. doi: 10.1038/nature06148
- Jiang, S. Y., Jin, J., Sarojam, R., and Ramachandran, S. (2019). A comprehensive survey on the terpene synthase gene family provides new insight into its evolutionary patterns. *Genome Biol. Evol.* 11, 2078–2098. doi: 10.1093/gbe/evz142
- Kaundal, R., Saini, R., and Zhao, P. X. (2010). Combining machine learning and homology-based approaches to accurately predict subcellular localization in *Arabidopsis*. *Plant Physiol.* 154, 36–54. doi: 10.1104/pp.110.156851
- Keilwagen, J., Lehnert, H., Berner, T., Budahn, H., Nothnagel, T., Ulrich, D., et al. (2017). The terpene synthase gene family of carrot (*Daucus carota* L.): identification of QTLs and candidate genes associated with terpenoid volatile compounds. *Front. Plant Sci.* 8:1930. doi: 10.3389/fpls.2017.01930
- Lescot, M., Déhais, P., Thijs, G., Marchal, K., Moreau, Y., Van De Peer, Y., et al. (2002). PlantCARE, a database of plant *cis*-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res.* 30, 325–327. doi: 10.1093/nar/30.1.325
- Li, G., Köllner, T. G., Yin, Y., Jiang, Y., Xu, Y., Gershenzon, J., et al. (2012). Correction for Li et al., nonseed plant *Selaginella moellendorffii* has both

- seed plant and microbial types of terpene synthases. *Proc. Natl. Acad. Sci. U. S. A.* 109, 20774–20774. doi: 10.1073/pnas.1204300109
- Liu, J., Huang, F., Wang, X., Zhang, M., Zheng, R., Wang, J., et al. (2014). Genome-wide analysis of terpene synthases in soybean: functional characterization of GmTPS3. *Gene* 544, 83–92. doi: 10.1016/j.gene.2014.04.046
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25, 402–408. doi: 10.1006/meth.2001.1262
- Martin, D. M., Aubourg, S., Schouwey, M. B., Daviet, L., Schalk, M., Toub, O., et al. (2010). Functional annotation, genome organization and phylogeny of the grapevine (*Vitis vinifera*) terpene synthase gene family based on genome assembly, FLcDNA cloning, and enzyme assays. *BMC Plant Biol.* 10:226. doi: 10.1186/1471-2229-10-226
- Miller, M. A., Pfeiffer, W., and Schwartz, T. (2011). “The CIPRES science gateway: A community resource for phylogenetic analyses” in *Proceedings of the TeraGrid 2011 Conference: Extreme Digital Discovery, TG’11*.
- Mohd-Hairul, A. R., Namasivayam, P., Lian, G. E. C., and Abdullah, J. O. (2010). Terpenoid, benzenoid, and phenylpropanoid compounds in the floral scent of *Vanda Mimi palmeri*. *J. Plant Biol.* 53, 358–366. doi: 10.1007/s12374-010-9123-x
- Nagegowda, D. A. (2010). Plant volatile terpenoid metabolism: biosynthetic genes, transcriptional regulation and subcellular compartmentation. *FEBS Lett.* 584, 2965–2973. doi: 10.1016/j.febslet.2010.05.045
- Nieuwenhuizen, N. J., Green, S. A., Chen, X., Bailleul, E. J. D., Matich, A. J., Wang, M. Y., et al. (2013). Functional genomics reveals that a compact terpene synthase gene family can account for terpene volatile production in apple. *Plant Physiol.* 161, 787–804. doi: 10.1104/pp.112.208249
- Omata, A., Nakamura, S., Yomogida, K., Moriai, K., Ichikawa, Y., and Watanabe, I. (1990). Volatile components of to-Yo-ran flowers (*Cymbidium faberi* and *Cymbidium virens*). *Agric. Biol. Chem.* 54, 1029–1033. doi: 10.1080/00021369.1990.10870040
- Pichersky, E., Noel, J. P., and Dudareva, N. (2006). Biosynthesis of plant volatiles: nature’s diversity and ingenuity. *Science* 311, 808–811. doi: 10.1126/science.1118510
- Pocza, P., Cernák, I., Varga, I., and Hyvönen, J. (2014). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Genet. Resour. Crop. Evol.* 61, 796–815. doi: 10.1134/S1022795411020074
- Priya, P., Yadav, A., Chand, J., and Yadav, G. (2018). Terzyme: a tool for identification and analysis of the plant terpenome. *Plant Methods* 14, 1–18. doi: 10.1186/s13007-017-0269-0
- Ramya, M., An, H. R., Baek, Y. S., Reddy, K. E., and Park, P. H. (2018). Orchid floral volatiles: biosynthesis genes and transcriptional regulations. *Sci. Hortic.* 235, 62–69. doi: 10.1016/j.scienta.2017.12.049
- Ramya, M., Park, P. H., Chuang, Y. C., Kwon, O. K., An, H. R., Park, P. M., et al. (2019). RNA sequencing analysis of *cymbidium goeringii* identifies floral scent biosynthesis related genes. *BMC Plant Biol.* 19:337. doi: 10.1186/s12870-019-1940-6
- Rozewicki, J., Li, S., Amada, K. M., Standley, D. M., and Katoh, K. (2019). MAFFT-DASH: integrated protein sequence and structural alignment. *Nucleic Acids Res.* 47, W5–W10. doi: 10.1093/nar/gkz342
- Schmutz, J., Cannon, S. B., Schlueter, J., Ma, J., Mitros, T., Nelson, W., et al. (2010). Genome sequence of the palaeopolyploid soybean. *Nature* 463, 178–183. doi: 10.1038/nature08670
- Starks, C. M., Back, K., Chappell, J., and Noel, J. P. (1997). Structural basis for cyclic terpene biosynthesis by tobacco 5-epi-aristolochene synthase. *Science* 277, 1815–1820. doi: 10.1126/science.277.5333.1815
- Sun, Y., Wang, G., Li, Y., Jiang, L., Yang, Y., and Guan, S. (2016). De novo transcriptome sequencing and comparative analysis to discover genes related to floral development in *Cymbidium faberi* Rolfe. *Springerplus* 5:1458. doi: 10.1186/s40064-016-3089-1
- Tholl, D. (2015). Biosynthesis and biological functions of terpenoids in plants. *Adv. Biochem. Eng. Biotechnol.* 148, 63–106. doi: 10.1007/10_2014_295
- Tian, C., Liu, S., Jiang, L., Tian, S., and Wang, G. (2020). The expression characteristics of methyl jasmonate biosynthesis-related genes in *Cymbidium faberi* and influence of heterologous expression of *CjJMT* in *Petunia hybrida*. *Plant Physiol. Biochem.* 151, 400–410. doi: 10.1016/j.plaphy.2020.03.051
- Troshin, P. V., Procter, J. B., and Barton, G. J. (2011). Java bioinformatics analysis web services for multiple sequence alignment-JABAWS:MSA. *Bioinformatics* 27, 2001–2002. doi: 10.1093/bioinformatics/btr304
- Vranová, E., Coman, D., and Grissem, W. (2013). Network analysis of the MVA and MEP pathways for isoprenoid synthesis. *Annu. Rev. Plant Biol.* 64, 665–700. doi: 10.1146/annurev-arplant-050312-120116
- Wagner, K. H., and Elmadfa, I. (2003). Biological relevance of terpenoids: overview focusing on mono-, di- and tetraterpenes. *Ann. Nutr. Metab.* 47, 95–106. doi: 10.1159/000070030
- Yu, Z., Zhao, C., Zhang, G., Teixeira da Silva, J. A., and Duan, J. (2020). Genome-wide identification and expression profile of TPS gene family in *Dendrobium officinale* and the role of *DoTPS10* in linalool biosynthesis. *Int. J. Mol. Sci.* 21, 1–22. doi: 10.3390/ijms21155419
- Zhang, J., Li, Y., Liu, B., Wang, L., Zhang, L., Hu, J., et al. (2018). Characterization of the Populus Rab family genes and the function of PtRabE1b in salt tolerance. *BMC Plant Biol.* 18:124. doi: 10.1186/s12870-018-1342-1
- Zhang, Z., Li, J., Zhao, X. Q., Wang, J., Wong, G. K. S., and Yu, J. (2006). KaKs_calculator: calculating Ka and Ks through model selection and model averaging. *Genomics Proteomics Bioinformatics* 4, 259–263. doi: 10.1016/S1672-0229(07)60007-2
- Zhang, G. Q., Xu, Q., Bian, C., Tsai, W. C., Yeh, C. M., Liu, K. W., et al. (2016). The *Dendrobium catenatum* Lindl. genome sequence provides insights into polysaccharide synthase, floral development and adaptive evolution. *Sci. Rep.* 6, 1–10. doi: 10.1038/srep19029
- Zhao, C., Yu, Z., Teixeira da Silva, J. A., He, C., Wang, H., Si, C., et al. (2020). Functional characterization of a *Dendrobium officinale* geraniol synthase *DoGES1* involved in floral scent formation. *Int. J. Mol. Sci.* 21:7005. doi: 10.3390/ijms21197005
- Zhou, H. C., Shamala, L. F., Yi, X. K., Yan, Z., and Wei, S. (2020). Analysis of terpene synthase family genes in *Camellia sinensis* with an emphasis on abiotic stress conditions. *Sci. Rep.* 10:933. doi: 10.1038/s41598-020-57805-1

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Wang, Zhu, Yu, Bi, Zhou, Chen, Chen, Zhang, Ai, Liu and Lan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.