

Using Cold As An Ally: Balsam Fir (*Abies Balsamea* L., Mill) Needles And Essential Oil Kill Overwintering Ticks (*Ixodes Scapularis*)

Shelley A. Adamo (✉ sadamo@dal.ca)

Dalhousie University

Amal El Nabbout

Dalhousie University

Laura Ferguson

Dalhousie University

Jeffrey Zbarsky

Dalhousie University

Nicoletta Faraone

Acadia University

Research Article

Keywords: bacterium, synthetic, toxicity, acaricidal

Posted Date: December 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1096371/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The blacklegged tick, *Ixodes scapularis* Say, vectors *Borrelia burgdorferi*, a bacterium that causes Lyme Disease. Although synthetic pesticides can reduce tick numbers, less toxic acaricides would reduce impacts on other animals, such as pollinating insects. Natural-based pest control agents could provide an alternative because they have low environmental persistency; however, natural products struggle to provide effective control. We found a new natural acaricide, balsam fir (*Abies balsamea*) needles, that kill overwintering *I. scapularis* ticks. We extracted the essential oils from the needles, analyzed their chemical composition, and tested them for acaricidal activity. Both the oil and one of its main components, β -pinene, kill overwintering ticks. Whole balsam fir needles require several weeks to kill overwintering ticks, while the essential oil is lethal within days at low temperatures ($\leq 4^{\circ}\text{C}$). Further, low temperatures increase the efficacy of this low toxicity compound. Higher temperatures (i.e. 20°C) reduce the acaricidal effectiveness of the essential oil by 50% at 0.1% v/v. Low temperatures may promote the effectiveness of other low toxicity, natural control products. Winter is an overlooked season for tick control and should be explored as a possible time for the application of low toxicity products for successful tick management.

Introduction

The tick *Ixodes scapularis* Say is a vector of *Borrelia burgdorferi*, a bacterium that causes Lyme Disease¹. Synthetic pesticides are currently used to kill ticks and reduce the transmission of Lyme Disease². However, concerns over the deleterious effects of pesticides on the environment have spurred the development of natural, less toxic, acaricidal compounds²⁻⁴. Although natural products can kill *I. scapularis*^{3,4}, they have had limited success in the field^{2,5-7}. Some of the very attributes that reduce their environmental impact, such as low persistence, also reduce their effectiveness^{5,8}. In this paper, we demonstrate that it may be possible to boost the effectiveness of natural products by applying them during the winter season when temperatures are low. Natural products such as essential oils have greater persistence at lower temperatures⁹.

Synthetic acaricides (i.e. tick-killing compounds) are applied during the ticks' active season – i.e. spring, summer and fall⁶. However, all three stages of *I. scapularis* ticks (e.g., larvae, nymph and adult) can be found during the winter¹⁰. Low winter temperatures are stressful for terrestrial arthropods¹¹ and *I. scapularis* is susceptible to stressors such as chill injury, desiccation, and inoculative freezing; thus cold and dry winters can be lethal^{12,13}. To survive these stresses, *I. scapularis* ticks must use resources to increase the production of protective molecules, such as antifreeze proteins¹⁴. During this stressful time, ticks may be more susceptible to a second stressor (e.g. a toxic compound). Animals often respond less effectively when facing two stressors simultaneously^{15,16}. Therefore, it is plausible that compounds with minimal toxicity to ticks at higher temperatures may become more lethal in the cold.

One important behavioural strategy *I. scapularis* ticks use to survive through the winter is burrowing into leaf litter¹⁷. Leaf litter provides an insulated and more humid environment than that offered by an

exposed site¹³. Indeed, ticks overwintering under leaf litter can withstand temperatures well below 0°C^{17,18} with rates of mortality similar to those in summer conditions¹². For example, leaf litter allows ticks (nymphs¹⁰, larvae/nymphs¹⁸) to have greater than 80% survival over the winter months, although overwintering success can vary widely from year to year^{19,20}. Given that winter conditions typically persist for four to five months in Canada and parts of the American Northeast, ticks in these areas spend considerable time in contact with plant material in the leaf litter, and thus the type of leaf litter surrounding ticks could impact their winter mortality²¹. Some have suggested that coniferous forests are likely to provide less insulating leaf litter, leading to reduced overwintering survival¹³. However, the survival of engorged larval *I. scapularis* ticks was the same whether they overwintered in deciduous or coniferous wood stands²¹. Similarly there was no difference in the overwintering survival of unfed adult female ticks in a maple forest, white pine stand, or oak savannah environment, although in these studies the bottom substrate of the microcosm was Kentucky bluegrass sod (*Poa pratensis* L., Poaceae)^{19,20}. Therefore, although leaf litter is important for tick overwintering survival¹⁷, there is no direct evidence that the type of leaf litter is important.

Because of the need for inexpensive, environmentally-friendly, methods of tick control, we decided to revisit the question of whether the type of leaf litter can reduce overwintering survival in *I. scapularis*. Anecdotal observations by one of us (SAA) suggested that ticks did not successfully overwinter under balsam fir (*Abies balsamea* L. Mill, Pinaceae) needles. We also chose this particular conifer because it is not a common tree in the historical home range of *I. scapularis*. *I. scapularis* ticks may be more likely to be sensitive to compounds with which they have not co-evolved. Balsam fir is common in Eastern Canada and in the extreme northern part of the eastern half of the US, but it is rare further south (<https://tidcf.nrcan.gc.ca/en/trees/factsheet/80>), in the region thought to be the ancestral range of *I. scapularis*²². We tested the acaricidal activity of balsam fir needles, and the essential oils extracted from those needles, against overwintering *I. scapularis* ticks using both lab and field studies. We determined the essential oil composition of our local balsam fir needles, and tested whether the main essential oil plays a role in killing ticks. We also tested whether the acaricidal effect was temperature-dependent.

Results

Balsam fir needles kill overwintering ticks under both field and lab conditions

Locally collected adult ticks were placed in individual tubes with vermiculite and sand, topped with 1 cm of balsam fir needles or 1 cm of maple/oak leaf litter (approximately 2 g of leaf litter). More than 98.5% of ticks overwintering in the field in balsam fir died (69/70), but only 57.1% (40/70) died in maple/oak leaf litter (G-test=38.2, $p < 0.0001$). Balsam fir killed both male (N=34/34 dead) and female (N=35/36 dead) ticks. Ticks died in balsam fir leaf litter whether the tubes were placed in a forest environment (N=31/32 dead), or in dune grass (N=38/38 dead). Adult female ticks also died in balsam fir leaf litter when held in incubators simulating winter conditions for 12 weeks. During the simulated winter (Fig. S1), 100% of ticks

in balsam fir died (48/48), while only 20.8% (10/48) of ticks in maple/oak died (G-test=72.9, $p<0.0001$). Balsam fir needles killed ticks infected with *B. burgdorferi* as well as non-infected ticks (Field – 100% infected died (8/8) and 98.4% uninfected died (61/62); Incubator – 100% of infected (33/33) and 100% non-infected (15/15) died). The nymph stage was also killed by balsam fir needles. Lab-reared *I. scapularis* adult females and nymphs were obtained from the National Tick Rearing Facility in Oklahoma, USA and were placed in individual tubes. The incubator was set at $0^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 10 weeks. As expected, all adult females in balsam fir (5/5) died during the simulated winter, while 60% survived in maple/oak (3/5), demonstrating that both field-collected and lab-reared ticks are susceptible to balsam fir needles. Overwintering nymphs died in balsam fir needles (0/26 survived in balsam fir, 22/24 survived in maple/oak, Fisher Exact Test, $p<0.0001$).

These results could be due to differences in the insulating properties of the two leaf litters. Maple/oak leaf litter did have better insulating properties compared with balsam fir, but the differences were small. The average temperature inside both leaf litters varied between each other by only $0.5^{\circ} \pm 0.4^{\circ}\text{C}$, averaged across 3 incubators with different winter temperature regimes (see Supplementary Information). Differences in relative humidity between the two leaf litters were larger, with maple/oak having higher relative humidity on average ($5.5\% \text{ RH} \pm 0.4$ higher) than balsam fir leaf litter.

To avoid the possibility that the effect we observed was due to the small differences in leaf litter insulating properties, we used a vermiculite/sand mixture or cotton ball as the substrate at the bottom of the tubes in subsequent experiments. Grinding balsam fir needles increased their acaricidal effect in incubators simulating winter conditions (Fig. 1; Chi-square (2)=149, $p<0.0001$, difference across groups. Linear trend with time for Ground needles, Chi-square (1)= (3.26), $p=0.07$). The acaricidal effect of ground needles was not immediate, but required about 3 weeks to kill 75% of the ticks (Fig. 1). Ticks housed with intact needles were still alive 3 weeks later, suggesting that balsam fir needles require several weeks to kill ticks. However, although ticks were not killed outright by balsam fir needles, after a 2 week exposure, ticks suffered from slower righting reflexes than controls in all 3 winter incubation temperatures (Fig. S2, 1-way ANOVA, $F(5, 127)=44.35$, $p<0.0001$, post hoc comparisons using a Tukey test, 0°C , $p<0.0001$; 4°C , $p<0.0001$, Fluctuating -2° to $+2^{\circ}\text{C}$, $p<0.0001$, no differences across the 3 controls, $F(3, 58)=0.80$, $p=0.45$).

Acaricidal compounds in balsam fir needles

To determine whether balsam fir needles contained compounds that kill ticks, we fractionated ground balsam fir needles (1 g balsam fir needles ground with 10 mL double distilled water). The slurry was filtered using Whatman #4 filters. The green filtrate smelt strongly of balsam fir. This material was added to 10K MWCO centrifuge filters. The clear aqueous fraction at the bottom of the centrifuge tube had no smell, while the material that did not pass through the filter was a green colour and retained the smell of balsam fir. Three-hundred (300) μL of either the clear aqueous fraction or 300 μL of the green residue fraction was added to a sand/vermiculite mix at the bottom of a tube. The tubes, containing a single lab-reared adult female, were placed in an incubator set at 0°C for 4 weeks; 80.9% of ticks (N=21) died when exposed to the green filtrate, while fewer than 15% of ticks died when exposed to the clear filtrate (14.3%,

N=21) or water (300 μ L of water added); 11.1%, N=18; G-test(2)=26.6, $p<0.0001$). Based on these results, we explored the possibility that an essential oil may be involved in balsam fir's acaricidal effect.

First, we determined the essential oil composition of our locally collected balsam fir using gas chromatography mass spectrometry after extracting the oil by hydro-distillation. The essential oil composition of fresh balsam fir needles showed an abundance of β -pinene (Table 1; Fig. S3), as was found by others^{23,24}. To mimic the amount of balsam fir or β -pinene essential oil that would be found in 2 g of balsam fir needles (based on data from Poaty²⁴), we added 2mL of 1.75% (v/v) of balsam fir essential oil or 0.65% (v/v) of β -pinene in 1% Tween 80 (v/v) in double distilled water to a cotton ball at the bottom of each tube. One lab-reared adult female tick was placed in each tube, and the tubes were placed in incubators to simulate overwintering conditions for 3 weeks. Ticks exposed to ground balsam fir needles (positive control), β -pinene and balsam fir essential oil died, while ticks in control tubes (1% v/v Tween 80) survived (Fig. 2, Planned comparisons for contingency tables²⁵; Control significantly different than other groups, 0°C: Z=8.1; $p<0.0001$; Overall Chi-square(3)=66.3, $p<0.0001$. 12°C: Control significantly different than other groups (Z= 8.5; $p<0.0001$). Overall Chi-square (3)=73.8, $p<0.0001$).

Essential oils are known for their volatility (e.g. balsam fir²⁶). In the incubator trials described above, the tubes were sealed with parafilm to maintain high humidity. However, this method also concentrates released volatiles, increasing the concentration in the head-space. We thus tested acaricidal effect of the essential oils if the tubes were left open (i.e. entrance covered only by a fine mesh), mimicking an outdoor environment in which volatiles could dissipate. High humidity was maintained in the tubes, and the cotton remained damp throughout the 4-week trial. Mortality was greater under closed tube conditions and depended on essential oil concentration (Fig. S4; Chi-square test. Significant difference between closed and open tubes, Chi-square (1)=25.7, $p<0.00001$, Significant difference across concentrations, Chi-square (2)=35.1, $p<0.0001$). We also found that balsam fir essential oil was more toxic to ticks than β -pinene (Chi-square test. Significant difference between balsam fir essential oil (N=112) and β -pinene (N=48), Chi-square (1)=15.0, $p<0.0001$, significant difference across concentrations, Chi-square (2)=35.2, $p<0.0001$).

Given the acaricidal effects of balsam fir, we also tested whether balsam fir was a repellent using horizontal bioassays according to Faraone et al.²⁷. Balsam fir essential oil was repellent for up to 10 min at room temperature, and the effect was concentration dependent (Fig. 3, $F_{4,436}=34.65$, $p<0.001$, N=30/group). The lowest concentration that exerted a significant repellent effect for up to 10 minutes was 4% v/v (z=3.087, $p=0.002$); 86.7% of the tested ticks were repelled when exposed to 8% v/v balsam fir essential oil after 3 min of exposure (Table S1). The repellent effect decreased over time (Fig. 3).

Table 1
Compounds identified in Balsam fir needles by GS-MS.

	Compound	ID	RT (min)	% Analyte in Total Oil $\pm$ SE
1	Santen	NIST	7.131	1.50 $\pm$ 0.11
2	Tricyclene	NIST	8.368	1.11 $\pm$ 0.04
3	3-Methyl-4-heptanol	NIST	8.495	2.69 $\pm$ 0.02
4	α -pinene	S	8.726	7.09 $\pm$ 0.18
5	Camphene	S	9.205	7.98 $\pm$ 0.15
6	4-hepten-2-one	NIST	9.399	1.53 $\pm$ 0.12
7	8-hydroxylinalool	NIST	9.732	9.34 $\pm$ 0.11
8	β -pinene	S	10.017	22.5 $\pm$ 0.44
9	β -myrcene	S	10.377	0.59 $\pm$ 0.03
10	3-carene	S	10.897	10.3 $\pm$ 0.15
11	Limonene	S	11.455	3.44 $\pm$ 0.10
12	β -phellandrene	NIST	11.482	5.38 $\pm$ 0.06
13	Isoborneol	NIST	14.946	0.67 $\pm$ 0.06
14	Borneol acetate	S	17.342	18.0 $\pm$ 0.23
15	Caryophyllene	S	20.044	2.18 $\pm$ 0.16
16	Humulene	S	20.690	1.12 $\pm$ 0.15
17	β -bisabolene	NIST	21.560	1.22 $\pm$ 0.12
	Unknowns	-	-	3.6 $\pm$ 0.2

RT= Retention Time

ID= Identification Reference Source

NIST=National Institute of Standard Technology

S=Chemical standard

Effect of temperature on the acaricidal activity of balsam fir essential oil

To test whether low temperature enhanced the acaricidal effect of balsam fir essential oil, we placed adult female ticks in individual tubes in one of two incubators (2°C and 20°C). At 20°C, a higher concentration

of balsam fir essential oil was required to kill ticks compared to 2°C (Fig. 4; Significant temperature effect, Pearson Chi-square test (1)=19.64, $p<0.0001$; Significant effect of essential oil concentration, Pearson Chi-square (3)=128.7, $p<0.0001$).

Discussion

Balsam fir needles killed overwintering ticks. The essential oils extracted from these needles were also acaricidal. The main essential oil within balsam fir, β -pinene, killed ticks too, although less effectively than the whole balsam fir essential oil. Balsam fir needles required several weeks to kill ticks, while the essential oil killed ticks in days at cold temperatures. Room temperature (i.e. 20°C) reduced the tick-killing effectiveness of the essential oil (Fig. 4). Lower temperatures may have decreased the loss of the volatile compounds responsible for the tick-killing. By decreasing the loss of these compounds, the duration of the tick's exposure to the essential oils could have been increased.

Low temperatures may aid in killing ticks for reasons other than reducing the loss of volatile substances. Low temperatures slow enzyme activity¹¹ and could reduce the rate of detoxification, allowing toxic compounds to accumulate. In addition, detoxification pathways may be compromised by a shift in resources towards overwintering survival. In some insects, detoxification is impaired during the activation of other physiological systems (e.g. immune responses) due to competition for limited resources²⁸. Exposure to toxic compounds may force an over-investment in detoxification; such an over-investment may impair overwintering survival²⁹. Some insects experience oxidative stress during exposure to low temperatures³⁰. If ticks do the same, then low temperatures could compound the toxic effects of balsam fir. Additionally, the decline in mobility at low temperatures^{31,32} may limit the ability of *Ixodes* ticks to move away from toxic leaf litter. Although this was not a serious issue in this study, as ticks were confined to tubes, it would reduce the ability of ticks to exit toxic leaf litter in the field. Even though ticks are capable of some movement in the winter³², forcing ticks to expend energy in locomotion during the winter to avoid toxic leaf litter could reduce overwintering survival during this non-feeding state. Not only will ticks be forced to consume scarce reserves, they will be more vulnerable to predation, freezing and desiccation while searching for a new patch of leaf litter. In addition, ticks exposed to balsam fir needles showed evidence of toxicity even after a short exposure (i.e. a slower righting reflex). These sublethal effects could produce a reduction in motor abilities, leaving ticks unable to find hosts. Thus, even if ticks are able to leave balsam fir leaf litter, shorter exposures may nonetheless impact tick populations.

In addition to β -pinene, other components are likely to play a role in the acaricidal effect of balsam fir essential oil. Combining essential oils can enhance their efficacy³³, and balsam fir contains a number of essential oils (Table 1). Two of the main essential oils found in our locally collected balsam fir needles, α -pinene and β -pinene (Table 1), are known to be lethal to some insects³⁴. β -pinene has also been reported to repel *Amblyomma americanum* and *Rhipicephalus appendiculatus* ticks, although it has not previously been identified as an acaricide³³. In fact, α - and β -pinene were not effective at killing *I. scapularis* nymphs at any concentration in a previous study³⁵ that examined their acaricidal effect at 21°C and for 24 h. α -

pinene and β -pinene may be lethal only at colder temperatures. The standard method for testing whether plant compounds kill ticks is to incubate them for 15 days at 27°C³⁶, which could explain why some low-toxicity compounds may be overlooked.

Balsam fir essential oil showed a significant repellent effect at room temperature, but it required a relatively high concentration (8% v/v) in the horizontal assay test. The repellency effect is significantly reduced after only 10 minutes, indicating the high volatility of the essential oil (Figure 3). Other plant essential oils can repel ticks (*I. ricinus*) at lower concentrations (e.g. lavender oil, < 1% w/w³⁷). These results demonstrate the low persistence of balsam fir essential oil at room temperature.

These results suggest that balsam fir needles, and their essential oil, have development potential as an acaricide. Balsam fir essential oil has minimal toxicity to terrestrial vertebrates at the concentrations used in this study.

(https://www.praannaturals.com/downloads/msds/SDS_Organic_Essential_Oil_Fir_Balsam_Canada.pdf). A balsam fir-derived product could provide environmentally-friendly control, especially when used during the winter when ticks are most sensitive to it. Moreover, winter application of an acaricide could offer additional ecological advantages. By late fall fewer non-target organisms are likely to be exposed³⁸. Therefore, later applications should reduce negative impacts on pollinators³⁹. Late application would also reduce the phytotoxicity that can occur with some essential oils³⁹. However, the compounds in plants that are deadly to ticks probably evolved to defend plants against insect herbivores, not blood-sucking arthropods³. They have effects on ticks because of similarities in the physiology of insects and ticks³. Therefore, any development of balsam fir, or other natural product targeting the over-wintering period, should examine its effects on arthropods that over-winter in the ground, especially on soil organisms that help control ticks⁴⁰.

Whether other plants can decrease tick overwintering survival should also be explored. In previous studies that showed high overwintering success in *I. scapularis*, the leaf litter was from deciduous trees^{10,18,41}. Ticks appear to have better survival and are more abundant in deciduous forests during their active period (i.e., spring, summer and fall) than in coniferous forests^{42–45}, although not all studies find these trends^{13,46}. Nevertheless, our results suggest that some species of conifers may be a potential source of anti-tick compounds. We hypothesize it will be plants like balsam fir, with which *I. scapularis* ticks historically have had little experience. For example, few *I. scapularis* ticks are found under eastern hemlock (*Tsuga canadensis* L., Carrière)⁴⁴, suggesting another possible northern conifer with acaricidal properties.

Although low humidity can be lethal to *I. scapularis*^{47,48}, it is unlikely that the decline in tick survival in balsam fir leaf litter was because it was a poor insulator. Lindsay et al.⁴⁹ found little difference in temperature or vapour pressure deficit between white pine and maple forest leaf litter over the winter. Brunner¹⁰ found no differences in the overwintering survival of *I. scapularis* nymphs between two sites that differed in relative humidity by about 20% (one site had 40% RH, the other had 20% RH). This result

suggests that the small differences we observed in humidity between the two leaf litters in this study is unlikely to explain the very large difference in mortality.

Many people become infected with Lyme disease in their own yard; a safe, efficient acaricide could play an important role in reducing the exposure to tick bites and to the spread of these infections⁴. Even if a vaccine for Lyme disease becomes available, *I. scapularis* can vector other diseases⁵⁰, and tick control will be necessary to prevent them. Our results show that a single application of balsam fir needles, or a balsam fir-based product, could be sufficient to kill overwintering ticks. Despite winter rain and periodic snow melts, a single patch of balsam fir needles was able to kill ticks in our open tube field tests. Other studies have noted that applications of acaricides prior to re-activation of ticks in the spring can significantly reduce the numbers of early season ticks⁷, suggesting that reducing overwintering survival will also reduce numbers in the spring⁵¹. For example, Rand et al.⁵² found that a low toxicity acaricide (IC2™, made with rosemary essential oil) sprayed in October substantially reduced the number of *I. scapularis* ticks in the following spring. However, a different study applied the same low toxicity control agent (IC2™) in June and found that it had little lasting effect⁶. Possibly the application of IC2™ under cold weather conditions (i.e. late October in Maine, US) enhanced its efficacy⁵². Cold temperatures may be an important ally of low toxicity, natural control products.

Methods

Ticks

Studies used unfed nymph and adult *I. scapularis* ticks. Ticks were either locally collected in Nova Scotia (i.e. Halifax Regional Municipality, Lunenburg County or Port Joli, Nova Scotia, Canada), or were purchased from the National Tick Research and Education Resource, housed at Oklahoma State University (Stillwater OK, USA). Studies were approved by the University Committee on Laboratory Animals at Dalhousie University (protocol number: I18-06). Research on plants in this study complies with the Convention on Biological Diversity and the Convention on the Trade in Endangered Species of Wild Fauna and Flora.

Effects of balsam fir needles on overwintering *I. scapularis* in the field

Field work was conducted at the Harrison Lewis Coastal Discovery Centre (HLC), Port L'Hebert, NS (HLC43°49'04.2" N 64°53'40.4" W). *Ixodes scapularis* were collected by dragging⁵⁴ from the HLC and Thomas Raddall Provincial Park, located adjacent to the HLC. Following collection, 70 ticks were placed in individual microcosms using a design modified from Burtis⁵³. The microcosm was an open plastic tube (15.24 cm in length x 1.27 cm in width); with mesh glued to each end, secured by a plastic cap. The plastic caps had a central hole that allowed rain and snow to enter in the top and connected the soil with the open bottom of the tube. The tubes contained sand and vermiculite (75% w/w sand and 25% w/w vermiculite) to a depth of 5 cm in the tube. The tubes were placed 5 cm into the ground. On top of the

sand/vermiculite was 1 cm of either a) cut oak (Red oak (*Quercus rubra* L.) and maple (Red maple (*Acer rubrum* L.) leaves or b) 1 cm of cut balsam fir (*Abies balsamea* L., Mill.) needles. A single tick was placed into each microcosm and then the microcosm was placed under a coarse netting (NuVue® (78.7cm L x 78.7 cm W x 83.8 cm H). A data logger (HOBO MX2301A) was placed in each enclosure with a probe 5 cm underground between the microcosms; the logger was covered with a solar radiation shield (Onset RS3-B) and the logger automatically recorded temperature and relative humidity every hour. Tick vials and data loggers were placed on October 25, 2018 and retrieved on February 8, 2019. Ticks were placed in either dune grass (N=38) or in a mixed maple/spruce stand of trees (N=32) behind the dune grass.

Ticks were recovered by carefully sieving the tubes. For all the experiments described in this paper, mortality was determined by allowing ticks to return to room temperature (approximately 22°C) for 24 h. Ticks that were immobile after both a gentle exhalation by the experimenter (AEN) and probing with a paint brush, were declared dead. Both dead and living ticks were tested for the presence of *B. burgdorferi* DNA using qPCR (see Supplementary Information).

Effects of balsam fir needles on *I. scapularis* overwintering in incubators

Ticks were collected in Burnside (Nova Scotia, Canada) and maintained at 4° C for 1 to 6 weeks prior to the study. Ticks were placed in glass tubes of the same size (15.24 cm in length x 1.27 cm in width) as was used in the field study, and had the same contents, but these were sealed at the bottom and capped at the top with parafilm. They were placed in one of 3 incubators simulating winter conditions (see Supplementary Information for details, Fig. S1). Ticks were placed in incubators in late fall and removed 12 weeks later. Tick mortality was recorded, and then ticks were tested for *B. burgdorferi* (Supplementary Information).

Essential oil extraction from balsam fir needles

Fresh balsam fir stems and leaves were collected in Port Joli and Lunenburg County (Nova Scotia, Canada). Needles were manually removed from the branches and ground using an electric coffee grinder (Black+Decker). Plant material was stored at -20°C until required. Essential oil extraction through hydrodistillation was performed using a Clevenger-type apparatus as previously described (Wang et al., in review). Hexane and deionized water were used as the solvent for the balsam fir extractions. Anhydrous sodium sulfate were used for drying balsam fir oils after extraction. Essential oil was stored at -20°C.

GC-MS analysis and characterization of balsam fir essential oil

Essential oil samples and chemical standards were analyzed via gas chromatography-mass spectrometry (GC-MS) to determine composition and main constituents. Essential oils were diluted in hexane at a concentration of 50 ng/µl. Samples were analyzed using a Scion 456 Gas Chromatograph-Single Quad (GC-MS; SCION Instruments UK Ltd., Livingston, UK). A non-polar capillary column Rxi®-5silms (30 m x 0.25 mm, film thickness 0.25 mm; Restek Corporation, State College, PA, USA) linked to a Bruker mass

spectrometer (Bruker Daltonics Ltd., Coventry, UK) was used for analysis. One μL was injected at 250°C by using a Gerstel Maestro Autosampler (Gerstel, Mülheim an der Ruhr, Germany). The split ratio for chemical standards was set up as 1:100. The oven temperature was set up at 50°C for 5 minutes, then increased to 220°C at 7°C/minute, and then increased up to 280°C at 30°C/minute. Both the injector and transfer line were maintained at 250°C. Helium was used as a carrier gas at 1 mL/min rate flow. Compounds were identified using analytical standards, and NIST Mass Spectral Search Program for the NIST/EPA/NIH Mass Spectral Library Version 2.0g 2011 (Scion Instruments UK Ltd., Livingston, West Lothian, UK).

Test of the acaricidal activity of balsam fir essential oil

Lab-reared ticks were placed singly in tubes as described for the incubator study. We added 2 mL of 1.75% (v/v) of balsam fir essential oil in 1% (v/v) Tween 80 (Sigma-Aldrich, Saint Louis, MO, US) to each tube to mimic the amount of balsam fir essential oil found in 2 g of balsam fir needles²⁴. We also added 2 mL of 0.65% (v/v) β -pinene in 1% (v/v) Tween 80 to mimic the amount of β -pinene found in 2 g of balsam fir needles²⁴. Control tubes received 2 mL of 1% (v/v) Tween 80. Tubes were placed in incubators set at 0°C and 12°C for 4 weeks.

Balsam fir oils horizontal repellency bioassay using nymphs

Nymphs were stored in sealed plastic containers lined with moist Kimwipe® in the fridge at 4 °C in dark conditions prior to trials. Balsam fir essential oil was diluted to 4% (v/v) or 8% (v/v) in hexane, with hexane acting as the control. Each concentration was tested against five nymphs for a total of 6 replications (N=30). The horizontal repellency bioassay was performed as described in Faraone²⁷ with some modifications. Briefly, the experimental arena was composed of three circles: the dropping zone (i.e. inner circle), treated circle, and outer circle. 320 μL of either the 4%, 8%, or control solution was added to the treated circle. Five active nymphs were transferred to the dropping zone by using a fine paintbrush. The location of each tick was recorded at 3, 5, and 10 minutes. Immobile ticks were recorded as “non-responders”. Active ticks, which exited the dropping zone, but did not cross the treated circle, were recorded as repelled. Active ticks that crossed into the treated circle or into the outer untreated circle were recorded as non-repelled.

Effect of temperature on the effect of balsam fir essential oil

Adult lab-reared ticks were placed singly in tubes as previously described above. Based on the results of the previous experiments, we used the following balsam fir essential oil concentrations: 0.1%, 0.5% and 1% (v/v) (N=32; 16 males and 16 females/concentration) in 1% Tween 80 for each temperature. We added 2 mL of each concentration to tubes with cotton as described above. Control tubes received 2 mL of 1% Tween 80 N=(36/temperature; 18 males and 18 females). Tubes were placed in incubators set at 2°C or 20°C for 2 weeks. Mortality was assessed. The results for both sexes were pooled because they did not differ significantly (Pearson Chi-Square (1)=2.184, p=0.139).

Statistical analysis

All data were tested for the assumptions of the statistical test used. Statistical analyses for contingency tests were performed on SPSS (ver. 26). Other tests were performed using Prism (GraphPad, Version 9.20) or RStudio Version 1.1.453 (RStudio Team 2009-2018). When multiple statistical tests were performed on the same data set, the alpha criterion was adjusted using a Bonferroni correction. Most of the data were categorical and were analyzed using Pearson Chi-square with Bonferroni correction. For the repellency tests, efficacy was analyzed with generalized linear mixed effect regression (glmer) with a binomial link to model the logit of tick repelled, followed by simultaneous tests for general linear hypotheses and multiple comparisons of means (Tukey's test), using the emmeans package.

Declarations

Data Availability

Data are available on Mendeley (<https://data.mendeley.com/datasets/gmegg47rcb5/1>).

Acknowledgements

We thank the Harrison Lewis Discovery Centre (Port Joli, Nova Scotia, Canada) for hosting our field study and Thomas Raddall Provincial Park (Port Joli, Nova Scotia, Canada) for permission to drag for ticks. We thank the Public Health Agency of Canada (PHAC) for the gift of the OspA for our positive control and for sharing their standard operating procedure for the molecular identification of *B. burgdorferi*. We thank Haozhe Vincent Wang for performing repellency assays. This research was supported by a grant from the Canadian Lyme Foundation to SAA, and a MITACS grant to SAA and AEN.

Author contributions. SAA conceived and developed the hypothesis, assisted with experimental design, data analysis, running the trials and led the writing of the paper. AEN assisted with experimental design, data analysis and running trials, LF helped with formulating the hypothesis, experimental design and running trials. JF helped with running trials and experimental design. NF assisted with formulating the hypothesis, experimental design, running trials and performed the chemical study on the balsam fir. All authors contributed to the manuscript.

Competing interest statement. The authors declare no competing interests.

References

1. Kilpatrick, A. M. *et al.* Lyme disease ecology in a changing world: consensus, uncertainty and critical gaps for improving control. *Philos. Trans. R. Soc. B-Biol. Sci.* **372**, 15, doi:10.1098/rstb.2016.0117 (2017).
2. Isman, M. B. Botanical insecticides in the twenty-first century - Fulfilling their promise? *Annual Review of Entomology* **65**, 233–249 (2020).

3. Adenubi, O. T. *et al.* Pesticidal plants as a possible alternative to synthetic acaricides in tick control: A systematic review and meta-analysis. *Industrial Crops and Products* **123**, 779–806, doi:10.1016/j.indcrop.2018.06.075 (2018).
4. Eisen, L. Control of ixodid ticks and prevention of tick-borne diseases in the United States: The prospect of a new Lyme disease vaccine and the continuing problem with tick exposure on residential properties. *Ticks Tick-Borne Dis.* **12**, doi:10.1016/j.ttbdis.2021.101649 (2021).
5. Eisen, L. & Dolan, M. C. Evidence for Personal Protective Measures to Reduce Human Contact With Blacklegged Ticks and for Environmentally Based Control Methods to Suppress Host-Seeking Blacklegged Ticks and Reduce Infection with Lyme Disease Spirochetes in Tick Vectors and Rodent Reservoirs. *J. Med. Entomol.* **53**, 1063–1092, doi:10.1093/jme/tjw103 (2016).
6. Dyer, M. C., Requintina, M. D., Berger, K. A., Puggioni, G. & Mather, T. N. Evaluating the Effects of Minimal Risk Natural Products for Control of the Tick, *Ixodes scapularis* (Acari: Ixodidae). *J. Med. Entomol.* **58**, 390–397, doi:10.1093/jme/tjaa188 (2021).
7. Schulze, T. L. & Jordan, R. A. Synthetic Pyrethroid, Natural Product, and Entomopathogenic Fungal Acaricide Product Formulations for Sustained Early Season Suppression of Host-Seeking *Ixodes scapularis* (Acari: Ixodidae) and *Amblyomma americanum* Nymphs. *J. Med. Entomol.* **58**, 814–820, doi:10.1093/jme/tjaa248 (2021).
8. Bharadwaj, A., Stafford, K. C. & Behle, R. W. Efficacy and Environmental Persistence of Nootkatone for the Control of the Blacklegged Tick (Acari: Ixodidae) in Residential Landscapes. *J. Med. Entomol.* **49**, 1035–1044, doi:10.1603/me11251 (2012).
9. Pavela, R. & Sedláč, P. Post-application temperature as a factor influencing the insecticidal activity of the essential oil from *Thymus vulgaris*. *Industrial crops and products* **113**, 46–49 2018.
10. Brunner, J. L., Killilea, M. & Ostfeld, R. S. Overwintering Survival of Nymphal *Ixodes scapularis* (Acari: Ixodidae) Under Natural Conditions. *J. Med. Entomol.* **49**, 981-987, doi:10.1603/me12060 (2012).
11. Chown, S. L. & Nicolson, S. W. *Insect Physiological Ecology*. (Oxford University Press, 2004).
12. Ogden, N. H., Beard, C. B., Ginsberg, H. S. & Tsao, J. I. Possible effects of climate change on Ixodid ticks and the pathogens they transmit: Predictions and observations. *J. Med. Entomol.* **58**, 1536-1545 (2021).
13. Ballard, K. & Bone, C. Exploring spatially varying relationships between Lyme disease and land cover with geographically weighted regression. *Applied Geography* **127**, 102383 (2021).
14. Neelakanta, G., Sultana, H., Fish, D., Anderson, J. F. & Fikrig, E. *Anaplasma phagocytophilum* induces *Ixodes scapularis* ticks to express an antifreeze glycoprotein gene that enhances their survival in the cold. *Journal of Clinical Investigation* **120**, 3179-3190, doi:10.1172/jci42868 (2010).
15. Adamo, S. A. Animals have a Plan B: how insects deal with the dual challenge of predators and pathogens. *J. Comp. Physiol. B-Biochem. Syst. Environ. Physiol.* **190**, 381-390, doi:10.1007/s00360-020-01282-5 (2020).
16. Adamo, S. A. How insects protect themselves against combined starvation and pathogen challenges, and the implications for reductionism. *Comparative Biochemistry and Physiology B-Biochemistry &*

17. Linske, M. A.*et al.* Impacts of Deciduous Leaf Litter and Snow Presence on Nymphal *Ixodes scapularis* (Acari: Ixodidae) Overwintering Survival in Coastal New England, USA. *Insects* **10**, doi:10.3390/insects10080227 (2019).
18. Burtis, J. C., Fahey, T. J. & Yavitt, J. B. Survival and energy use of *Ixodes scapularis* nymphs throughout their overwintering period. *Parasitology* **146**, 781-790, doi:10.1017/s0031182018002147 (2019).
19. Lindsay, L. R.*et al.* Survival and development of *Ixodes scapularis* (acari, ixodidae) under various climatic conditions in Ontario, Canada. *J. Med. Entomol.* **32**, 143-152, doi:10.1093/jmedent/32.2.143 (1995).
20. Lindsay, L. R.*et al.* Survival and development of the different life stages of *Ixodes scapularis* (Acari: Ixodidae) held within four habitats on Long Point, Ontario, Canada. *J. Med. Entomol.* **35**, 189-199, doi:10.1093/jmedent/35.3.189 (1998).
21. Ginsberg, H. S.*et al.* Woodland type and spatial distribution of nymphal *Ixodes scapularis* (Acari: Ixodidae). *Environmental Entomology* **33**, 1266-1273, doi:10.1603/0046-225x-33.5.1266 (2004).
22. Khatchikian, C. E.*et al.* Recent and rapid population growth and range expansion of the Lyme disease tick vector, *Ixodes scapularis*, in North America. *Evolution* **69**, 1678-1689, doi:10.1111/evo.12690 (2015).
23. Pichette, A., Larouche, P. L., Lebrun, M. & Legault, J. Composition and antibacterial activity of *Abies balsamea* essential oil. *Phytotherapy Research* **20**, 371-373, doi:10.1002/ptr.1863 (2006).
24. Poaty, B., Lahlah, J., Porqueres, F. & Bouafif, H. Composition, antimicrobial and antioxidant activities of seven essential oils from the North American boreal forest. *World Journal of Microbiology & Biotechnology* **31**, 907-919, doi:10.1007/s11274-015-1845-y (2015).
25. Beasley, T. M. & Schumacker, R. E. Multiple regression approach to analyzing contingency tables: Post hoc and planned comparison procedures. *Journal of Experimental Education* **64**, 79-93, doi:10.1080/00220973.1995.9943797 (1995).
26. Canon, L., Deslauriers, A., Mshvildadze, V. & Pichette, A. Volatile compounds in the foliage of balsam fir analyzed by static headspace gas chromatography (HS-GS): an example of the spruce budworm defoliation effect in the boreal forest of Quebec, Canada. *Microchemical Journal* **110**, 587-590 (2013).
27. Faraone, N., MacPherson, S. & Hillier, N. K. Behavioral responses of *Ixodes scapularis* tick to natural products: development of novel repellents. *Experimental and Applied Acarology* **79**, 195-207, doi:10.1007/s10493-019-00421-0 (2019).
28. McMillan, L. E., Miller, D. W. & Adamo, S. A. Eating when ill is risky: immune defense impairs food detoxification in the caterpillar *Manduca sexta*. *J. Exp. Biol.* **221**, jeb173336, doi:10.1242/jeb.173336 (2018).
29. Gazave, E., Chevillon, C., Lenormand, T., Marquine, M. & Raymond, M. Dissecting the cost of insecticide resistance genes during the overwintering period of the mosquito *Culex pipiens*. *Heredity* **87**, 441-448 (2001).

30. Lalouette, L., Williams, C. M., Hervant, F., Sinclair, B. J. & Renault, D. Metabolic rate and oxidative stress in insects exposed to low temperature thermal fluctuations. *Comparative Biochemistry and Physiology A* **158**, 229-234 (2011).
31. Clark, D. D. Lower temperature limits for activity of several Ixodid ticks: Effects of body size and rate of temperature change. *J. Med. Entomol.* **32**, 449-452 (1995).
32. Carroll, J. F. & Kramer, M. Winter activity of *Ixodes scapularis* (Acari: Ixodidae) and the operation of deer-targeted tick control devices in Maryland. *J. Med. Entomol.* **40**, 238-244, doi:10.1603/0022-2585-40.2.238 (2003).
33. Quadros, D. G., Johnson, T. L., Whitney, T. R., Oliver, J. D. & Chavez, A. S.O. Plant-Derived Natural Compounds for Tick Pest Control in Livestock and Wildlife: Pragmatism or Utopia? *Insects* **11**, doi:10.3390/insects11080490 (2020).
34. Ogendo, J. *et al.* Biocontrol potential of selected plant essential oil constituents as fumigants of insect pests attacking stored food commodities. *Health* **10**, 287-318 (2011).
35. Panella, N. A., Karchesy, J., Maupin, G. O., Malan, J. C. & Piesman, J. Susceptibility of immature *Ixodes scapularis* (Acari: Ixodidae) to plant-derived acaricides. *J. Med. Entomol.* **34**, 340-345 (1997).
36. Rosado-Aguilar, J. A. *et al.* Plant products and secondary metabolites with acaricide activity against ticks. *Veterinary Parasitology* **238**, 66-76, doi:10.1016/j.vetpar.2017.03.023 (2017).
37. Jaenson, T. G. T., Carboui, S. & Palsson, K. Repellency of oils of lemon eucalyptus, geranium, and lavender and the mosquito repellent MyggA natural to *Ixodes ricinus* (Acari: Ixodidae) in the laboratory and field. *J. Med. Entomol.* **43**, 731-736, doi:10.1603/0022-2585(2006)43[731:Roole]2.0.Co;2 (2006).
38. Schulze, T. L. *et al.* Efficacy of granular deltamethrin against *Ixodes scapularis* and *Amblyomma americanum* (Acari: Ixodidae) nymphs. *J. Med. Entomol.* **38**, 344-346, doi:10.1603/0022-2585-38.2.344 (2001).
39. Elias, S. P. *et al.* Effect of a Botanical Acaricide on *Ixodes scapularis* (Acari: Ixodidae) and Nontarget Arthropods. *J. Med. Entomol.* **50**, 126-136, doi:10.1603/me12124 (2013).
40. Burtis, J. C., Yavitt, J. B., Fahey, T. J. & Ostfeld, R. S. Ticks as Soil-Dwelling Arthropods: An Intersection Between Disease and Soil Ecology. *J. Med. Entomol.* **56**, 1555-1564, doi:10.1093/jme/tjz116 (2019).
41. Burtis, J. C., Ostfeld, R. S., Yavitt, J. B. & Fahey, T. J. The Relationship Between Soil Arthropods and the Overwinter Survival of *Ixodes scapularis* (Acari: Ixodidae) Under Manipulated Snow Cover. *J. Med. Entomol.* **53**, 225-229, doi:10.1093/jme/tjv151 (2016).
42. Guerra, M. *et al.* Predicting the risk of Lyme disease: Habitat suitability for *Ixodes scapularis* in the north central United States. *Emerging Infectious Diseases* **8**, 289-297, doi:10.3201/eid0803.010166 (2002).
43. Bunnell, J. E., Price, S. D., Das, A., Shields, T. M. & Glass, G. E. Geographic Information Systems and spatial analysis of adult *Ixodes scapularis* (Acari: Ixodidae) in the Middle Atlantic region of the USA. *J. Med. Entomol.* **40**, 570-576, doi:10.1603/0022-2585-40.4.570 (2003).

44. Lubelczyk, C. B.*et al.* Habitat associations of *Ixodes scapularis* (Acari: Ixodidae) in Maine. *Environmental Entomology* **33**, 900-906, doi:10.1603/0046-225x-33.4.900 (2004).
45. Killilea, M. E., Swei, A., Lane, R. S., Briggs, C. J. & Ostfeld, R. S. Spatial dynamics of Lyme disease: A review. *Ecohealth* **5**, 167-195, doi:10.1007/s10393-008-0171-3 (2008).
46. Clow, K. M.*et al.* The influence of abiotic and biotic factors on the invasion of *Ixodes scapularis* in Ontario, Canada. *Ticks Tick-Borne Dis.* **8**, 554-563, doi:10.1016/j.ttbdis.2017.03.003 (2017).
47. Stafford, K. C. Survival of immature *Ixodes scapularis* (Acari: Ixodidae) at different relative humidities. *J. Med. Entomol.* **31**, 310-314 (1994).
48. Bertrand, M. R. & Wilson, M. L. Microclimate-dependent survival of unfed adult *Ixodes scapularis* (Acari: Ixodidae) in Nature: Life cycle and study design implications. *J. Med. Entomol.* **33**, 619-627 (1996).
49. Lindsay, L. R.*et al.* Microclimate and habitat in relation to *Ixodes scapularis* (Acari: Ixodidae) populations on Long Point, Ontario, Canada. *J. Med. Entomol.* **36**, 255-262, doi:10.1093/jmedent/36.3.255 (1999).
50. Thompson, C., Spielman, A. & Krause, P. J. Coinfecting deer-associated zoonoses: Lyme disease, babesiosis, and ehrlichiosis. *Clinical Infectious Diseases* **33**, 676-685 (2001).
51. Hayes, L. E., Scott, J. A. & Stafford, K. C. Influences of weather on *Ixodes scapularis* nymphal densities at long-term study sites in Connecticut. *Ticks Tick-Borne Dis.* **6**, 258-266, doi:10.1016/j.ttbdis.2015.01.006 (2015).
52. Rand, P. W.*et al.* Trial of a minimal-risk botanical compound to control the vector tick of Lyme disease. *J. Med. Entomol.* **47**, 695-698 (2010).
53. Burtis, J. C. Method for the Efficient Deployment and Recovery of *Ixodes scapularis* (Acari: Ixodidae) Nymphs and Engorged Larvae from Field Microcosms. *J. Med. Entomol.* **54**, 1778-1782, doi:10.1093/jme/tjx157 (2017).
54. Schulze, T. L., Jordan, R. A. & Hung, R. W. Biases associated with several sampling methods used to estimate abundance of *Ixodes scapularis* and *Amblyomma americanum* (Acari: Ixodidae). *J. Med. Entomol.* **34**, 615-623 (1997).

Figures




Figure 1

Effect of intact and ground balsam fir needles on the survival of adult female *Ixodes scapularis* exposed to 0°C - 4°C. N=27 ticks/group for Week 1 and Week 3. For Week 2, N=36 for controls and N=45 for intact or ground balsam fir needles (BF Needles – intact balsam fir needles, Ground BF – ground balsam fir needles).




Figure 2

Ground balsam fir needles, b-pinene and balsam fir essential oil kill ticks at two different temperatures (0°C and 12°C). Bars represent the percentage of dead ticks after 4 weeks. BF=Balsam Fir, BF EO=Balsam Fir Essential Oils. Control N=24/incubator, Ground BF N=16/incubator, b-pinene N=24/incubator, BF EO N=32/incubator.




Figure 3

Mean repellency (%) over time of *Ixodes scapularis* nymphs exposed to balsam fir essential oil (EO) at different concentrations (1-8% v/v). N= 30 per treatment.




Figure 4

Balsam fir essential oil killed ticks at a lower concentration at 2°C than at 20°C. The black bars represent the mortality at 2°C while the grey bars represent mortality at 20°C. N=32 for each bar except N=36 at 0%.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInfodec14.docx](#)