

Two novel species of sooty mould fungi (Capnodiales, Dothideomycetes) from China

Jipeng Sun^{1,2,3}, Entaj Tarafder⁴, Tao Chen⁵, Sinang Hongsanan⁶, Shengxi Chen¹, Chada Norphanphoun¹, Samantha C. Karunarathna⁴, Xiangyu Zeng^{1,2,3}

¹ Department of Plant Pathology, Agricultural College, Guizhou University, Guiyang 550025, China

² Guizhou Key Laboratory of Agricultural Microbiology, Guiyang 550025, China

³ Institute of Edible Mushrooms, Guizhou University, Guiyang 550025, China

⁴ Center for Yunnan Plateau Biological Resources Protection and Utilization & Yunnan International Joint Laboratory of Fungal Sustainable Utilization in South and Southeast Asia, College of Biology and Food Engineering, Qujing Normal University, Qujing 655099, China

⁵ Public Health and Health Sciences College, Guizhou Medical University, Guiyang 551113, China

⁶ Shenzhen Key Laboratory of Microbial Genetic Engineering, College of Life Science and Oceanography, Shenzhen University, Shenzhen 518060, China

Corresponding author: Xiangyu Zeng (xyzeng3@gzu.edu.cn)

Abstract

Conidiocarpus and *Polychaeton* are genera of sooty mould fungi (Capnodiales, Dothideomycetes) that grow epiphytically on the leaf surfaces of living plants and typically depend on honeydew excreted by sap-sucking insects as their primary nutrient source. Heavy colonization by these fungi may impair host photosynthesis, reduce plant vigor, and decrease the aesthetic or economic value of affected plants. In this study, two novel species, *Conidiocarpus nanshanense* and *Polychaeton cengongense*, were collected from the leaves of *Schefflera macrostachya* (large-spiked schefflera) and *Dalbergia assamica* (South China rosewood) in Guizhou and Guangdong Provinces, respectively. Species delimitation was based on detailed morphological characteristics and phylogenetic analyses of multiple loci, including ITS, LSU, *tef1-α*, and *rpb2*. Morphological descriptions and illustrations are provided, together with phylogenetic analyses inferred using maximum likelihood (ML) and Bayesian inference (BI). The results expand the known taxonomic diversity of sooty mould fungi in China and provide important morphological and molecular data that support species delimitation and classification within Capnodiales.

Key words: Capnodiaceae, fungal diversity, new species, phylogenetic analysis, taxonomy



Academic editor:

Danushka Sandaruwan Tennakoon

Received: 14 December 2025

Accepted: 3 February 2026

Published: 19 February 2026

Citation: Sun J, Tarafder E, Chen T, Hongsanan S, Chen S, Norphanphoun C, Karunarathna SC, Zeng X (2026) Two novel species of sooty mould fungi (Capnodiales, Dothideomycetes) from China. MycoKeys 129: 23–41. <https://doi.org/10.3897/mycokeys.129.182290>

Copyright: © Jipeng Sun et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Capnodiaceae (Capnodiales, Dothideomycetes), a family of epiphytic sooty mould fungi, was first established by Höhnelt (1910), with *Capnodium* designated as the type genus (Kirk et al. 2008; Lumbsch and Huhndorf 2011). The taxonomic status of the family was formally validated by Theissen (1915). Currently, this family comprises 11 genera and approximately 150 species (Lu et al. 2022; Wijayawardene et al. 2022; Yang et al. 2025). Among these genera, *Scoriadopsis*, *Limaciniaseta*, *Kosmimatomyces*, and *Hyphocapnodia* are monotypic and lack molecular data (Lu et al. 2022; Wijayawardene et

al. 2022), with only morphological descriptions available. Hughes (1976) published the first monograph on sooty mould fungi, systematically summarizing the entire group. Subsequent researchers have further refined the morphological descriptions and supplemented molecular data for species in this family based on this work (Crous et al. 2009; Chomnunti et al. 2011; Abdollahzadeh et al. 2020).

Capnodiaceae is the representative family of sooty mould fungi (Crous et al. 2007; Abdollahzadeh et al. 2020). Its key characteristics include septate, dark-brown hyphae that form a thin hyphal network on the host surface; bitunicate asci; and asexual morphs producing slender conidiomata with short or long, narrow necks (Chomnunti et al. 2011). Distinct elliptical swellings are present near the base, middle, or apex of the conidiomata, and transparent conidia are produced inside these swollen regions (Chomnunti et al. 2011; Abdollahzadeh et al. 2020). Sooty mould refers to a group of epiphytic fungi that obtain nutrients from honeydew secreted by pests such as aphids, whiteflies, and mealybugs, or from exudates produced by the leaves of specific plants. These fungi are characterized by black, patchy colonies on plant surfaces (Jouraeva et al. 2006; Hongsanan et al. 2015). Species of Capnodiaceae dominate sooty mould communities, as their hyphal layers cover the plant surface, blocking sunlight and thereby interfering with leaf photosynthesis. In addition, the hyphal layers trap heat on the leaf surface. Overall, they not only reduce photosynthetic efficiency but also increase the plant's own energy consumption, ultimately affecting plant growth and crop yields (Chomnunti et al. 2011; Abdollahzadeh et al. 2020).

Capnodiaceae fungi are widely distributed in tropical and subtropical regions (Chomnunti et al. 2011; Hongsanan et al. 2020). Nevertheless, the currently documented species diversity remains insufficient to support in-depth investigations of these fungi, particularly with respect to their ecological roles and interactions among fungi, insects, and host plants. China, despite its immense biodiversity, has so far reported only four species of Capnodiaceae, including three newly described species (*Conidiocarpus fici-septicae*, *Hyphocapnodia sichuanensis*, and *C. cinnamomeus*) and one newly recorded species (*C. caucasicus*) (Lu et al. 2022; Tennakoon et al. 2022; Thungdee et al. 2023; Tian et al. 2024; Yang et al. 2025). These taxa were described from different regions and host plants, including *C. fici-septicae* from fallen leaves of *Ficus septica* (Moraceae) (Tennakoon et al. 2022), *H. sichuanensis*, the sole species of the monotypic genus *Hyphocapnodia* from Sichuan Province (Lu et al. 2022), *C. cinnamomeus* from leaves of *Cinnamomum japonicum* (Yang et al. 2025), and *C. caucasicus* from withered leaves of *Cryptomeria japonica* in southwestern China (Tian et al. 2024). Collectively, these limited records indicate that Capnodiaceae remain markedly under-represented in China and suggest that the diversity of sooty mould fungi in this region is still poorly explored.

Consequently, further systematic studies are required to reveal additional species diversity and to clarify unresolved taxonomic issues within Capnodiaceae. In this study, we describe novel species of Capnodiaceae associated with *Schefflera macrostachya* and *Dalbergia assamica* from China. These findings expand the known geographic and host ranges of the family and provide a foundation for future studies on the diversity and ecology of sooty mould fungi.

Materials and methods

Sample collection and morphological observations

Sooty mould-infected leaves were collected from *Schefflera macrostachya* in Guangdong Province and from *Dalbergia assamica* in Guizhou Province, China, in early spring (24 March 2025) and mid-summer (14 July 2025), respectively. The samples were placed in paper envelopes, transported to the laboratory, and stored at 4 °C. Macromorphological characteristics, including colony characteristics (size, shape, elevation, margin, texture, and color) and the distribution of conidiomata, were observed and documented using a stereomicroscope (Keyence VHX-7000 Digital Microscope, Japan). For microscopic analysis, samples were prepared on slides with sterile water and examined under a compound optical microscope (Zeiss Axioscope 5, Germany) equipped with an AxioCam 208 color camera and a phase-contrast optical system. The shape, pigmentation, and dimensions of hyphae, conidiomata, ostioles, the position of the swollen region of the conidiomata, and conidia were photographed and recorded. All images for illustration and dimensional measurements were processed using Adobe Photoshop (2023, 24.0.0.59). The holotype was deposited at the Herbarium of IFRD (International Fungal Research & Development Centre; Institute of Highland Forest Science, Chinese Academy of Forestry, Kunming, China). The ex-type living culture was deposited at the Culture Collection of the Herbarium of IFRD (IFRDCC). The newly described species were registered in Index Fungorum, and the corresponding accession numbers were obtained (Index Fungorum 2026).

DNA extraction, PCR amplifications, and sequencing

DNA extraction was performed with fresh fungal mycelia cultured on potato dextrose agar (PDA) at 26 °C in darkness for 2 weeks using the BEIWO Fungal DNA Extraction Kit (Hangzhou Beiwo Medical Technology Co., Ltd., catalog No. BW-GD2416), following the manufacturer's protocol. Genomic DNA of the host plant was extracted using the Plant Genomic DNA Extraction Kit from Beijing Solarbio Science & Technology Co., Ltd. For fungal gene amplification, four molecular markers were targeted: the internal transcribed spacer (ITS) region, including 5.8S rDNA, using primers ITS1/ITS4 (White et al. 1990); the large subunit ribosomal RNA gene (LSU), using primers LR0R/LR5 (Vilgalys and Hester 1990); the translation elongation factor 1- α gene (*tef1*- α), using primers EF1-983F/EF1-2218R (Reynolds and Gilbert 2005); and the RNA polymerase II second largest subunit gene (*rpb2*), using primers RPB2-5F/RPB2-7cR (Liu et al. 1999). For the host plant, the partial *rbcL* gene was amplified using the primer pair SI_Forward and SI_Reverse, which were developed by Kress et al. (2009). The PCR reaction mixture had a total volume of 20 μ l, consisting of 17.5 μ l GoldenStar T6 Super PCR Mix Ver. 2 (1.1 $\times$), 1 μ l DNA template, 1 μ l forward primer, and 1 μ l reverse primer. PCR cycling parameters were optimized for each molecular marker, with the key annealing temperatures specified as follows: ITS and LSU were amplified with an annealing temperature of 52 °C (35 cycles); *tef1*- α at 55 °C (35 cycles); and *rpb2* with a two-step annealing protocol (60 °C for 5 cycles, followed by 54 °C for 35 cycles). Other standard PCR conditions included initial denaturation, denaturation, extension, and final hold steps, as described in the original protocols (Vilgalys and Hester 1990; White et

al. 1990; Reynolds and Gilbert 2005; Kress et al. 2009). Detailed procedures for the PCR assays used in the present study are listed in Table 1. PCR products were analyzed using 1% agarose gel, which was prepared and, after complete solidification, placed in an electrophoresis tank containing buffer. DNA samples were loaded into the wells, and electrophoresis was performed at 100 V for 30 min. The gel was visualized using a gel imaging system. PCR products were considered qualified if clear, bright bands were observed at the expected positions corresponding to the target gene fragments. Qualified PCR products were sent to Tsingke Biotechnology Co., Ltd. (Beijing) for sequencing. The resulting valid sequences were deposited in GenBank (Table 2).

Phylogenetic analyses

Gene sequencing results obtained from the biotechnology company were analyzed by examining DNA sequence peaks using BioEdit version 7.2.5 (Hall 1999). Phylogenetic analyses were performed using a one-stop software for fungal phylogeny (OFPT) (Zeng et al. 2023), which first requires preparation of reference sequence data from the NCBI database (<https://www.ncbi.nlm.nih.gov/>) and newly collected sequences. The specific workflow was as follows: gene sequences were downloaded from NCBI. Afterwards, datasets for each gene region were aligned independently using MAFFT (<https://mafft.cbrc.jp/alignment/server/>) (Kato and Standley 2013). These alignments were then trimmed using TrimAl with the “gappyout” method (Capella-Gutiérrez et al. 2009). Phylogenetic trees were constructed using the Bayesian inference (BI) method (Ronquist et al. 2012), and bootstrap (BS) values were obtained. Aligned files (NEX format) were run in MrBayes v. 3.2.6, and the selection of the optimal nucleotide substitution model was performed using MrModeltest v. 2.3. Bayesian inference was conducted using a Markov chain Monte Carlo (MCMC) approach with two independent runs of four Markov chains each. The analysis was run for 50 million generations, sampling every 100 generations, and was terminated when the average standard deviation of split frequencies between runs dropped below 0.01. In total, 1,000,000 trees were generated, of which the first 25% were discarded as burn-in prior to constructing the majority-rule consensus tree. Posterior probabilities (PP) were calculated for each clade. The resulting tree files were visualized using FigTree v. 1.4.3, and the phylogenetic trees were further edited and formatted in Adobe Illustrator CC 2024 (AI) before being saved as PDF files.

Table 1. The PCR conditions and the primers used in this study.

Gene	Primers	Sequence (5'–3')	PCR cycles	References
ITS	ITS1	TCCGTAGGTGAACCTGCGG	(95 °C: 30 s, 52 °C: 45 s, 72 °C: 60 s) × 35 cycles	White et al. (1990)
	ITS4	TCCTCCGCTTATTGATATGC		
LSU	LR0R	ACCCGCTGAACCTTAAGC	(95 °C: 30 s, 52 °C: 45 s, 72 °C: 60 s) × 35 cycles	Vilgalys and Hester (1990)
	LR5	ATCCTGAGGGAAACTTC		
<i>tef1-a</i>	EF1-983F	GCCYCCYGGHCAYCGTGAYTTYAT	(95 °C: 30 s, 55 °C: 45 s, 72 °C: 60 s) × 35 cycles	Reynolds and Gilbert (2005)
	EF1-2218R	ATGACACCRACRGCRACRGTYTG		
<i>rpb2</i>	RPB2-5F	GAYGAYMGWGATCAYTTYGG	(94 °C: 45 s, 60 °C: 45 s, 72 °C: 2 min) × 5 cycles; (94 °C: 45 s, 54 °C: 45 s, 72 °C: 2 min) × 5 cycles; (94 °C: 45 s, 54 °C: 45 s, 72 °C: 2 min) × 30 cycles	Liu et al. (1999)
	RPB2-7cR	CCCATRGCTTGYYTTRCCCAT		
<i>rbcL</i>	SL_Forward	ATGTCACCACAAACAGAGACTAAAGC	(94 °C: 30 s, 55 °C: 30 s, 72 °C: 60 s) × 35 cycles	Kress et al. (2009)
	SL_Reverse	GTAAATCAAGTCCACCRCG		

Table 2. Taxa used in the phylogenetic analysis of Capnodiaceae and their corresponding GenBank accession numbers.

Species	Strain	GenBank Accession Number				Reference
		ITS	LSU	<i>tef1-α</i>	<i>rpb2</i>	
<i>Capnodium aciculiforme</i>	CBS 892.73	-	-	GU349045	-	Schoch et al. (2009)
<i>Capnodium alfenasii</i> ^T	CBS 146151	MN749233	MN749165	MN829346	MN829260	Abdollahzadeh et al. (2020)
<i>Capnodium blackwelliae</i> ^T	CBS 133588	MN749235	-	-	-	Abdollahzadeh et al. (2020)
<i>Capnodium coartatum</i>	CPC 17779	MN749236	MN749167	MN829348	MN829262	Abdollahzadeh et al. (2020)
<i>Capnodium coartatum</i>	SDBR-CMU477	OR458575	OR458609	OR523660	-	Thungdee et al. (2023)
<i>Capnodium coffeae</i> ^T	CBS 147.52	DQ491515	GU214400	DQ471089	KT216519	Spatafora et al. (2006)
<i>Capnodium coffeicola</i> ^{ET}	MFLUCC 15-0206	KU358921	KU358920	-	-	Ariyawansa et al. (2015)
<i>Capnodium gamsii</i>	CBS 146153	MN749238	MN749168	MN829349	MN829263	Abdollahzadeh et al. (2020)
<i>Capnodium gardeniarum</i>	CPC 14327	-	GU301807	GU349054	GU371743	Schoch et al. (2009); Li et al. (2020)
<i>Capnodium neocoffeicola</i> ^T	CBS 139614	MN749242	MN749172	MN829353	MN829267	Abdollahzadeh et al. (2020)
<i>Capnodium paracoartatum</i>	MFLUCC 14-0282	NR_169716	NG_073832	-	-	Li et al. (2020)
<i>Capnodium paracoffeicola</i> ^T	CBS 139616	MN749244	MN749174	MN829355	MN829269	Abdollahzadeh et al. (2020)
<i>Chaetocapnodium insulare</i> ^T	CBS 146159	MN749248	NG_068681	-	-	Abdollahzadeh et al. (2020)
<i>Chaetocapnodium magnum</i> ^T	CBS 153154	PV583761	PV583758	PV591868	PV591871	Czachura et al. (2025)
<i>Chaetocapnodium polonicum</i> ^T	CBS 153156	PV583763	PV583760	PV591870	PV591873	Czachura et al. (2025)
<i>Chaetocapnodium siamensis</i> ^T	MFLUCC 13-0778	-	KP744479	-	-	Hongsanan et al. (2015)
<i>Chaetocapnodium tanzanicum</i> ^T	CBS 145.79	MN749253	MN749182	MN829365	MN829280	Abdollahzadeh et al. (2020)
<i>Chaetocapnodium thailandense</i> ^T	CBS 139619	MN749254	MN749183	MN829366	MN829281	Abdollahzadeh et al. (2020)
<i>Conidiocarpus caucasicus</i> ^T	GUMH 937	-	KC833050	-	-	Bose et al. (2014)
<i>Conidiocarpus cinnamomi</i> ^T	SICAU 23-0100	PP736390	PP732745	PP779591	PP782090	Yang et al. (2025)
<i>Conidiocarpus fici-septicae</i> ^T	MFLUCC 19-0072	MW063143	MW063206	-	-	Tennakoon et al. (2021)
<i>Conidiocarpus guilanensis</i> ^T	IRAN 2474C	MG906804	-	-	-	Khodaparast et al. (2020)
<i>Conidiocarpus nanshanense</i> ^T	IFRDCC 25-0001	PX668540	PX668542	PX682093	PX687868	This study
<i>Conidiocarpus siamensis</i> ^T	MFLUCC 10-0061	KU358923	JN832607	-	-	Hongsanan et al. (2015)
<i>Heteroconium citharexyl</i> ^T	CPC 13957	HM628776	HM628775	-	-	Cheewangkoon et al. (2012)
<i>Hyphocapnodia sichuanensis</i> ^T	CGMCC 3.23573	ON603981	ON603986	ON646697	ON646698	Lu et al. (2022)
<i>Leptoxylum cacuminum</i> ^T	MFLUCC 10-0059	-	JN832603	-	-	Chomnunti et al. (2011)
<i>Leptoxylum citri</i>	CBS 146162	MN749267	MN749195	MN829378	-	Abdollahzadeh et al. (2020)
<i>Leptoxylum citri</i> ^T	CBS 451.66	MN749266	KF902094	GU349039	GU371727	Abdollahzadeh et al. (2020)
<i>Leptoxylum fumago</i>	CBS 123.26	MH854862	GU214430	GU349051	-	Schoch et al. (2009)
<i>Leptoxylum glochidion</i> ^T	IFRDCC 2651	KF982307	KF982308	-	-	Yang et al. (2014)
<i>Leptoxylum kurandae</i>	CBS 129530	JF951150	JF951170	MN829379	MN829295	Crous et al. (2011)
<i>Leptoxylum madagascariense</i> ^T	CBS 124766	MH863407	MH874923	MN829380	MN829296	Abdollahzadeh et al. (2020)
<i>Phaeoxyphiella australiana</i> ^T	CBS 146169	MN749292	MN749220	MN829406	MN829322	Abdollahzadeh et al. (2020)
<i>Phaeoxyphiella phylicae</i> ^T	CBS 146170	MN749291	MN749219	MN829405	MN829321	Abdollahzadeh et al. (2020)
<i>Phragmocapnias betle</i>	CPC 17762	MN749293	MN749221	MN829407	MN829323	Abdollahzadeh et al. (2020)
<i>Phragmocapnias betle</i>	MFLUCC 10-0050	-	JN832605	-	-	Bose et al. (2014)
<i>Phragmocapnias betle</i> ^{ET}	MFLUCC 10-0053	KU358922	JN832606	-	-	Bose et al. (2014)
<i>Phragmocapnias plumeriae</i> ^T	MFLUCC 15-0205	KU358919	KU358918	-	-	Ariyawansa et al. (2015)
<i>Polychaeton cengongense</i> ^T	IFRDCC 25-0002	PX668539	PX668541	PX682092	PX682094	This study
<i>Polychaeton citri</i>	CBS 116435	GU214649	GU214469	MN829394	MN829310	Crous et al. (2009)
<i>Scolecoxyphium blechni</i> ^T	CBS 146174	MN749296	MN749224	MN829412	MN829328	Abdollahzadeh et al. (2020)
<i>Scolecoxyphium blechnicola</i> ^T	CBS 146175	MN749297	MN749225	MN829413	MN829329	Abdollahzadeh et al. (2020)
<i>Scolecoxyphium leucadendri</i> ^T	CBS 146176	MN749298	MN749226	MN829414	MN829330	Abdollahzadeh et al. (2020)
<i>Scolecoxyphium phylicae</i> ^T	CBS 146177	MN749299	MN749227	MN829415	MN829331	Abdollahzadeh et al. (2020)

Sequences and details of strains obtained in this study are shown in bold. Abbreviations: T and ET indicate type species. - indicates unavailable data or sequence.

Results

Phylogenetic analyses

The alignment included 45 strains and contained a total of 3223 characters, including gaps, with 1–445, 446–1,280, 1,281–2,332, and 2,333–3,223 corresponding to ITS, LSU, *rpb2*, and *tef1-α*, respectively. The optimal nucleotide substitution models were TIM2e+G4, TNe+R2, TIM2e+I+G4, and TIM3+F+I+G4, respectively.

The phylogenetic analysis of the concatenated dataset produced a best-scoring tree with a final maximum likelihood optimization value of -16613.553 .

The tree topology from BI was similar to that from ML. Thus, only the ML tree is shown. Capnodiaceae comprises eight genera: *Conidiocarpus*, *Phragmocapnias*, *Capnodium*, *Hyphocapnodia*, *Polychaeton*, *Chaetocapnodium*, *Heteroconium*, and *Leptoxyphium*. The genera *Conidiocarpus* (6 representative sequences), *Phragmocapnias* (4), *Capnodium* (12), *Hyphocapnodia* (1), *Polychaeton* (2), *Chaetocapnodium* (6), *Heteroconium* (1), and *Leptoxyphium* (8) form a monophyletic group. In this study, the two new species belong to different genera, with *Conidiocarpus* sister to *Phragmocapnias* and *Polychaeton* sister to *Capnodium* (Fig. 1).




Figure 1. Phylogenetic tree generated from ML analysis based on the concatenated datasets of LSU, ITS, *tef1*-α, and *rpb2* gene regions. All type strains are indicated by a superscript T, and the new isolate is highlighted in bold red. The phylogenetic tree was rooted using Readeriellipsidaceae taxa (Abdollahzadeh et al. 2020) as the outgroup. Bootstrap support (BS) values that are equal to or exceed 70% and Bayesian posterior probabilities (PP) values that are equal to or exceed 0.9 are shown above the nodes. Type strains: superscript T and ET; new isolate: red bold; unavailable sequences are denoted by -.

Taxonomy

Conidiocarpus Woron., Ann. Mycol. 24: 250 (1927)

Type species. *Conidiocarpus caucasicus* Woron., Key to fungi (fungi imperfecti) 2: 743 (1917).

Notes. *Conidiocarpus* was recognized as the asexual morph of *Phragmocapnias* (Hughes 1976) and was treated as a synonym of *Phragmocapnias* (Hughes 1976; Chomnunti et al. 2011; Chomnunti et al. 2014; Hongsanan et al. 2015). Following the nomenclatural priority principle, Bose et al. (2014) adopted Hughes' (1976) viewpoint and transferred species from *Phragmocapnias* to *Conidiocarpus*. However, Abdollahzadeh et al. (2020) proposed that these two are separate, independent genera based on morphological and phylogenetic analyses. According to MycoBank (2026) (<https://www.mycobank.org>, January 2026), the genus *Conidiocarpus* includes fifteen validly recognized species, including *C. asiaticus*, *C. betel*, *C. callitris*, *C. caucasicus*, *C. fici-septicae*, *C. fuliginodes*, *C. guilanensis*, *C. heliconiae*, *C. imperspicuus*, *C. longicollus*, *C. penzigii*, *C. philippinensis*, *C. plumeriae*, *C. siamensis*, and *C. cinnamomi* (Abdollahzadeh et al. 2020; Khodaparast et al. 2020; Tennakoon et al. 2022; Pem et al. 2024; Yang et al. 2024; Yang et al. 2025). *Conidiocarpus* is distinguished from other genera by the following characteristics: the swollen part of the sporogenous structure is located in the middle–upper part of the conidioma, accompanied by a long stipe and an elongated base (Chomnunti et al. 2011).

Conidiocarpus nanshanense J.P. Sun & X.Y. Zeng, sp. nov.

Index Fungorum: IF904721

Fig. 2

Etymology. Refers to the type locality, “Nanshan Park.”

Holotype. IFRD 99050.

Description. *Epiphytic* on the leaf surface of *Schefflera macrostachya*, forming a sooty coating on adaxial surface (Fig. 2). **Thallus** composed of brown, septate, ellipsoidal, smooth-walled hyphae. **Asexual morph.** *Conidiomata* (342–493 (–513) × 27–53 µm, $\bar{x}$ = 423 × 39 µm, n = 15) long, pycnidial, elongate, superficial, stipitate, with a long stalk (118–280 µm, $\bar{x}$ = 226 µm, n = 15), black, rigid, a distinct neck (66–120 µm, $\bar{x}$ = 101 µm, n = 15) height, and a prominently ovoid-swollen conidiogenous region, the swollen area producing conidia inside. *Conidiomata apex* gradually light brown to hyaline, with a circular **ostiole** (7–14 (–17) µm, $\bar{x}$ = 11 µm, n = 15) diam, consisting of rectangular, compact cells, surrounded by hyaline hyphae. **Conidiogenous region**, located in the upper-middle part of the conidiomata, (25–47 µm, $\bar{x}$ = 39 µm, n = 15) wide, brown, composed of cylindrical, thin-walled cells. *Conidia* 4.1–6.4 × 2.4–3.3 µm ($\bar{x}$ = 5 × 2.7 µm, n = 30), hyaline, single-celled, ellipsoidal, smooth-walled, guttulate with 1–2 distinct refractive oil droplets, exuding in creamy masses from the ostiole. **Sexual morph.** Undetermined.

Culture characteristics. Colonies growing on PDA reaching 16 mm in diam. after 14 days at 26 °C in the dark, colony surface gradually erumpent, with hyphae growing downward and immersed in the medium, olivaceous. Aerial hyphae pale to dark brown, cylindrical, distinctly verrucose, branched, with thin, inconspicuous septa (not constricted), walls becoming thickened.




Figure 2. *Conidiocarpus nanshanense* (IFRD 99050). **a.** Black mycelium covering the leaf surface; **b, c.** Conidiomata on the host; **d.** Conidioma; **e.** Ostiole neck; **f.** A prominently ovoid-swollen conidiogenous region; **g.** Mycelial network; **h.** Conidia; **i, j.** Germinated conidia; **k.** Front view of the colony on the PDA; **l.** Reverse view of the colony on the PDA. Scale bars: 500 μm (**b**); 50 μm (**c, d, g, i, j**); 20 μm (**e, f**); 10 μm (**h**).

Material examined. CHINA • Guangdong Province, Shenzhen City, Nanshan Park, on living leaves of *Schefflera macrostachya*. (22°50'25"N, 113°91'26"E), 24 March 2025, Jipeng Sun, (IFRD 99050, holotype), ex-type living culture IFRDCC 25-0001.

Notes. In the phylogenetic tree, *C. nanshanense* (IFRDCC 25-0001) clusters with *C. fici-septicae* (MFLUCC 19-0072), with BS/PP support values of 96%/0.96, and forms a sister clade with the strains *C. cinnamomi* (SICAU 23-0100), *C. siamensis* (MFLUCC 10-0061), and *C. caucasicus* (GUMH 937). There are 99.25% (535/539, 4 gaps) and 99.77% (879/881, 1 gap) similarities in the ITS and LSU regions between *C. nanshanense* (IFRDCC 25-0001) and *C. fici-septicae* (MFLUCC 19-0072), respectively. Comparison of *tef1*- α and *rpb2* sequences could not be conducted due to the lack of sequence data for *C. fici-septicae* (MFLUCC 19-0072). Tian et al. (2024) reported the first Chinese record of *C. caucasicus* (UESTCC 23.0246) and supplemented the molecular data for the ex-type strain *C. caucasicus* (GUMH 937) with ITS, LSU, *tef1*- α , and *rpb2* sequences. Multi-gene alignment between *C. nanshanense* (IFRDCC 25-0001) and *C. caucasicus* (UESTCC 23.0246) revealed the following sequence similarities: ITS 99.80% (511/512, 0 gaps), LSU 99.63% (806/809, 0 gaps), *tef1*- α 97.90% (841/859, 0 gaps), and *rpb2* 99.29% (981/988, 0 gaps). Multi-gene alignment between *C. nanshanense* (IFRDCC 25-0001) and *C. cinnamomi* (SICAU 23-0100) showed the sequence similarities as follows: ITS 98.88% (528/534, 2 gaps) and LSU 99.10% (877/885, 1 gap). In terms of morphological characteristics, *C. nanshanense* (IFRDCC 25-0001) differs from *C. fici-septicae* (MFLUCC 19-0072) by possessing larger conidiomata (342–493 μm vs. 240–280 μm), a broader base (27–53 μm vs. 7–8 μm), a wider ostiole (7–17 μm vs. 6–9 μm), and larger conidia (5 $\times$ 2.7 vs. 4.5 $\times$ 1.5 μm) (Tennakoon et al. 2021). Compared to *C. caucasicus* (GUMH 937), *C. nanshanense* (IFRDCC 25-0001) has smaller conidiomata (342–493 μm vs. 420–930 μm), narrower ostioles (7–14 μm vs. 14–28 μm), and larger conidia (4.1–6.4 $\times$ 2.4–3.3 μm vs. 3.5–5 $\times$ 1.5–2 μm) (Khodaparast 2006). When compared with *C. cinnamomi* (SICAU 23-0100), *C. nanshanense* (IFRDCC 25-0001) exhibits smaller conidiomata (342–493 μm vs. 500–860 μm) and wider ostioles (7–17 μm vs. 4–8 μm) (Yang et al. 2025). However, a morphological comparison with *C. siamensis* (MFLUCC 10-0061), which was isolated from an unknown host, could not be conducted due to the lack of a detailed morphological description for this strain (Chomnunti et al. 2011; Abdollahzadeh et al. 2020). *Conidiocarpus caucasicus*, *C. fici-septicae*, and *C. cinnamomi* were isolated from *Citrus* sp. (Rutaceae, Sapindales), *Ficus septica* (Moraceae, Rosales), and *Cinnamomum japonicum* (Lauraceae, Laurales), respectively (Khodaparast 2006; Tennakoon et al. 2021; Yang et al. 2025), whereas *C. nanshanense* was discovered on *Schefflera macrostachya* (Araliaceae, Apiales) in this study. Accordingly, the species represented by this strain is described as a new species, *C. nanshanense*.

***Polychaeton* (Pers.) Lév., Dictionnaire universelle d'histoire naturelle 8: 493 (1846)**

Type species. *Polychaeton quercinum* (Pers.) Kuntze, Revisio generum plantarum 3 (3) (1891).

Notes. *Polychaeton* was originally established by Persoon (1822) as a sub-genus within *Fumago* and was later elevated to generic rank by L  veill   (1847). The taxonomic concept of *Polychaeton* has subsequently been re-evaluated by Hughes (1976) and Chomnunti et al. (2011). Hughes (1976) recognized *Polychaeton citri* and *P. quercinum* as representative species of the genus and formally designated *P. quercinum* as the lectotype. In contrast, Chomnunti et al. (2011) interpreted *Capnodium* as the sexual morph associated with *Polychaeton* and, in accordance with the “one fungus = one name” principle, adopted *Capnodium* as the accepted genus. According to MycoBank (2026) (<https://www.mycobank.org>, January 2026), the genus *Polychaeton* currently comprises 19 species, although molecular sequence data are available for only *Polychaeton citri*. *Polychaeton* is distinguished from other genera by the position of the swollen conidiogenous region in the middle to lower part of the conidiomata and by the absence of a stalk (Chomnunti et al. 2011).

***Polychaeton cengongense* J. P. Sun & X. Y. Zeng, sp. nov.**

Index Fungorum: IF904722

Fig. 3

Etymology. Refers to the type locality, “Cengong County.”

Holotype. IFRD 99051.

Description. *Epiphytic* on the leaf surface of *Dalbergia assamica*, forming a sooty coating on adaxial surface (Fig. 3). **Thallus** composed of brown, septate, ellipsoidal, smooth-walled hyphae. Asexual morph. **Conidiomata** (182–263 × 19–46 (–54) µm, $\bar{x}$ = 224 × 34 µm, n = 20) long, pycnidial, elongate, superficial, stipitate, lack stalk, with a distinct neck (62–125 µm, $\bar{x}$ = 95 µm, n = 20) height, a prominently ovoid-swollen conidiogenous region, the swollen area producing conidia inside. **Conidiomata apex** light brown to hyaline, with a circular **ostiole** (10–23 µm, $\bar{x}$ = 16 µm, n = 20) diam, consisting of rectangular, compact cells, surrounded by hyaline hyphae. **Conidiogenous region**, located in the lower-middle part of the conidiomata, (30–62 µm, $\bar{x}$ = 37 µm, n = 20) wide, brown, composed of cylindrical, thin-walled cells. **Conidia** 4.2–6.5 × 1.7–4 µm ($\bar{x}$ = 5.1 × 2.3 µm, n = 30), hyaline, single-celled, ellipsoidal, smooth-walled, guttulate with 1–2 distinct refractive oil droplets, exuding in creamy masses from the ostiole. **Sexual morph.** Undetermined.

Culture characteristics. Colony up to 15.6 mm in diam on PDA at 26 °C after 14 days in darkness. Colony superficial, velvety, with an entire margin, olivaceous-green. Aerial hyphae pale to dark brown, cylindrical, distinctly verrucose (with 1–2 µm tall warts), branched, with thin, inconspicuous septa (not constricted), walls becoming thickened.

Material examined. CHINA • Guizhou Province, Cengong County, on living leaves of *Dalbergia assamica* (27°44'20"N, 108°44'20"E), 14 July 2025, Jipeng Sun, (IFRD 99051, holotype), ex-type living culture IFRDCC 25-0002.

Notes. In the phylogenetic tree, *Polychaeton cengongense* (IFRDCC 25-0002) forms a sister clade to *Polychaeton citri* (CBS 116435), with BS/PP support values of 96%/1, respectively. *Polychaeton cengongense* (IFRDCC 25-0002) and *P. citri* (CBS 116435) have 99.06% (508/518, 3 gaps) and 93.91% (756/805, 0 gaps) similarity in the ITS and LSU regions, respectively, while comparison of their




Figure 3. *Polychaeton cengongense* (IFRD 99051). **a.** Black mycelium covering the leaf surface; **b, c.** Conidiomata on the host; **d.** Conidioma; **e.** Ostiole neck; **f.** A prominently ovoid-swollen conidiogenous region; **g.** Mycelial network; **h.** Conidia; **i.** Germinated conidia; **j.** Front view of the colony on the PDA; **k.** Reverse view of the colony on the PDA. Scale bars: 1000 µm (**b**); 100 µm (**c**); 50 µm (**d–g**); 10 µm (**h**); 25 µm (**i**).

tef1-α and *rpb2* regions reveals 93.91% (756/805, 0 gaps) and 89.67% (738/823, 0 gaps) sequence similarity, respectively. In morphological data, compared to *P. citri* (CBS 116435), *P. cengongense* (IFRDCC 25-0002) has smaller conidiomata (182–263 μm vs. 345–391 μm), narrower bases (19–46 μm vs. 36–40 μm), wider ostioles (10–23 μm vs. 13–15 μm), and smaller conidia (5.1 × 2.3 μm vs. 6.5 × 5 μm) (Chomnunti et al. 2011). *P. citri* was isolated from *Citrus aurantium* (Rutaceae, Sapindales), while *P. cengongense* in this study was obtained from *Dalbergia assamica* (Fabaceae, Fabales), representing a new host record for this species. The hosts of these two species exhibit remarkable differences at the higher taxonomic ranks of family and order with no overlap at all. Therefore, the species represented by this strain is described as a new species, *P. cengongense*.

Discussion

China, one of the world's most significant biodiversity hotspots, harbors a remarkably rich and underexplored diversity of sooty mould fungi (Capnodiaceae). Although the family is globally distributed across tropical and subtropical ecosystems, only a few Capnodiaceae species have been formally documented in China (Lu et al. 2022; Tennakoon et al. 2022; Thungdee et al. 2023; Tian et al. 2024; Yang et al. 2025), leaving a significant gap in our understanding of their ecological roles and evolutionary history. The discovery of *Conidiocarpus nanshanense* and *Polychaeton cengongense* highlights the vast, untapped fungal diversity yet to be uncovered within China's subtropical forests, especially in undersampled areas such as Guizhou and Guangdong Provinces. These findings not only expand the species inventory of Capnodiales in China but also highlight the urgent need for more systematic surveys to better understand the ecological interactions and biogeographic patterns of sooty mould fungi in this megadiverse country. Such efforts are crucial for establishing a solid taxonomic framework and guiding conservation strategies for China's unique fungal flora.

Our morphological and phylogenetic analyses of the isolated strains confirm the distinctiveness of *Conidiocarpus nanshanense* and *Polychaeton cengongense* as new species. Species delimitation was supported by an integrative taxonomic approach combining detailed morphological characterization and multi-gene phylogenetic analysis. Both new species form well-supported clades in the phylogenetic trees and exhibit consistent morphological and molecular differences from closely related taxa. These findings enrich the known species diversity of *Conidiocarpus* and *Polychaeton* in China and provide crucial morphological and molecular data to facilitate further taxonomic studies of Capnodiales.

In traditional understanding, sooty mould fungi are regarded as "harmful fungi" because they cover leaf surfaces and interfere with photosynthesis (Chomnunti et al. 2011; Thungdee et al. 2023). However, field observations in this study revealed that the colonization densities of *Conidiocarpus nanshanense* and *Polychaeton cengongense* show a positive correlation with the aphid population density on host plants, without causing host defoliation. This suggests that sooty mould fungi may be obligate decomposers of insect honeydew. By decomposing honeydew secreted by aphids, they prevent secondary fungal infections on leaves induced by honeydew accumulation and thus actually act as "buffers" in plant–insect interactions. Experimental evidence has confirmed that certain species of sooty mould fungi can produce secondary metabolites,

including those from three genera, *Capnodium*, *Leptoxylphium*, and *Trichomerium* (Herath et al. 2012; Haituk et al. 2022). These secondary metabolites inhibit the growth of common airborne fungi such as *Penicillium* and *Mucor*. Currently, the taxonomic status of species in the genus *Conidiocarpus* remains unclear. For some closely related species, such as *C. fici-septicae*, only ITS/LSU sequences are available, lacking markers with higher resolution (*tef1-α* and *rpb2*), which results in inaccurate species delimitation. To address the issue of insufficient molecular markers, priority should be given to supplementing *tef1-α*, *rpb2*, and other gene sequences for type strains of sooty mould fungi, thereby establishing a multi-gene reference database.

The morphological characteristics of sooty mould fungi (such as conidiomata and conidia dimensions) are prone to morphological convergence across taxa under environmental influences, while the same species may also exhibit morphological divergence due to varying environmental conditions (Abdollahzadeh et al. 2020; Thungdee et al. 2023; Tian et al. 2024). This not only leads to the misidentification of distinct taxa as the same species and to the erroneous division of a single species into multiple taxa, but also results in ambiguity in the taxonomic boundaries within *Conidiocarpus*. However, *C. nanshanense* and *C. fici-septicae* exhibit high morphological similarity; combined phylogenetic inference based on the multi-locus dataset (ITS+LSU+*tef1-α*+*rpb2*) reveals that they are not conspecific taxa. Instead, they cluster within the same clade of the genus *Conidiocarpus* as distinct divergent lineages, and this phylogenetic relationship is supported by strong statistical evidence (BS/PP = 96%/0.96). This finding suggests that morphological trait convergence does not imply complete congruence in phylogenetic relationships. This result supports the hypothesis that the genus *Conidiocarpus* is a monophyletic group (Wheeler et al. 2013; Williams et al. 2016) and revises the traditional morphological classification criterion that delimits *Conidiocarpus* species based on conidiomata size. In this study, the conidiomatal sizes and swollen regions of *C. nanshanense* and *C. caucasicus* overlap, whereas the ostiole width and spore length–width ratio differ. Prior to this study, most known species of the genus *Conidiocarpus* have been reported from Southeast Asia, Thailand, and West Asia, Iran (Chomnunti et al. 2011; Khodaparast et al. 2020). The discovery of the new *Conidiocarpus* species in China reveals the characteristics of both connection and isolation between the sooty mould fungal flora of Southwest China and that of Southeast Asia.

In the taxonomic studies of sooty mould fungi, Chomnunti et al. (2011) first clarified the taxonomic correspondence between *Polychaeton* and *Capnodium* by constructing a phylogenetic tree based on multiple specimens of polychaeton-like taxa and combining SSU and LSU gene sequences. For their part, Abdollahzadeh et al. (2020) retrieved *Polychaeton citri* CBS 116435 from the Westerdijk Fungal Biodiversity Institute (CBS), Utrecht, the Netherlands. This strain was isolated from the host plant *Citrus aurantium* in Iran (Chomnunti et al. 2011; Hyde et al. 2011). Through rejuvenation culture on oatmeal agar (OA) and synthetic low nutrient agar (SNA), they found that the morphology of its conidiomata exhibited medium-dependent differences: on SNA medium, the strain possessed swollen conidiogenous structures and a slender neck with a short stalk, whereas on OA medium, its neck was further elongated, while the swelling characteristic was absent. By contrast, the conidiomata of taxa in the genus *Conidiocarpus* maintained a stable morphology of a swollen main body, long

neck, and long stalk on both media. Despite the studies mentioned above confirming the taxonomic independence of *Polychaeton*, numerous controversies remain regarding the accurate delimitation of intra-generic species. Meanwhile, the new species *Polychaeton cengongense* described in this study represents the first record of *Polychaeton* on the host *Dalbergia assamica* in Guizhou Province, China, and adds to the literature, laying a critical foundation for clarifying the taxonomic boundaries of this genus and improving species identification.

The successful amplification of the *rbcL* gene from *Polychaeton cengongense* provides molecular confirmation of its host association with *Dalbergia assamica*. The complete *rbcL* gene sequence of the host, with a BLAST 100% (573/573, 0 gaps) similarity, is provided in Suppl. material 1. Combined with previous reports of *Polychaeton* species on *Citrus aurantium* (Rutaceae, Sapindales) (Chomnunti et al. 2011), this indicates ecological differentiation in host associations of this genus within woody plant taxa. For *Conidiocarpus nanshanense*, the inability to obtain host *rbcL* data underscores the challenge of working with preserved, field-collected samples, where prolonged storage can lead to DNA degradation. However, the morphological confirmation of *Schefflera macrostachya* as its host, together with the distinct geographical distribution (Guizhou/Guangdong Province vs. Taiwan/Sichuan Province for closely related species), further supports the ecological differentiation of *C. nanshanense* and its status as a novel species. These host associations also contribute to our understanding of the diversity and host specificity of sooty mould fungi in subtropical China.

The two new species described in this study are not only new records of sooty mould fungal diversity in China but also provide novel scientific perspectives for research on Capnodiaceae along two dimensions: taxonomic revision and re-evaluation of ecological functions. Future research could focus on the phylogeography of global sooty mould fungi and their contributions to honeydew nutrient cycling to deepen the understanding of their taxonomy, evolution, and ecological functions.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

No use of AI was reported.

Funding

This work was supported by the National Natural Science Foundation of China (No. 32400013) and the Guizhou Provincial Basic Research Program (Natural Science) ZK[2023] general 087.

Author contributions

Conceptualization: Jipeng Sun and Xiangyu Zeng; methodology: Jipeng Sun and Xiangyu Zeng; investigation: Jipeng Sun and Tao Chen; data curation: Tao Chen and Shengxi Chen; formal analysis: Jipeng Sun; software: Tao Chen; validation: Tao Chen

and Shengxi Chen; funding acquisition: Xiangyu Zeng; supervision: Xiangyu Zeng; writing—original draft: Jipeng Sun; writing—review and editing: Entaj Tarafder, Sinang Hongsan, Chada Norphanphoun, and Samantha C. Karunarathna.

Author ORCIDs

Jipeng Sun  <https://orcid.org/0009-0004-0065-6920>

Entaj Tarafder  <https://orcid.org/0000-0002-3680-3433>

Tao Chen  <https://orcid.org/0009-0004-7806-4503>

Sinang Hongsan  <https://orcid.org/0000-0003-0550-3152>

Shengxi Chen  <https://orcid.org/0009-0005-4945-5094>

Chada Norphanphoun  <https://orcid.org/0000-0002-5756-7206>

Samantha C. Karunarathna  <https://orcid.org/0000-0001-7080-0781>

Xiangyu Zeng  <https://orcid.org/0000-0003-1341-1004>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Abdollahzadeh J, Groenewald JZ, Coetzee MPA, Wingfield MJ, Crous PW (2020) Evolution of lifestyles in *Capnodiales*. *Studies in Mycology* 95: 381–414. <https://doi.org/10.1016/j.simyco.2020.02.004>
- Ariyawansa HA, Hyde KD, Jayasiri SC, Buyck B, Chethana KWT, Dai DQ, Dai YC, Daranagama DA, Jayawardena RS, Lücking R, Ghobad-Nejhad M, Niskanen T, Thambugala KM, Voigt K, Zhao RL, Li GJ, Doilom M, Boonmee S, Yang ZL, Cai Q, Cui YY, Bahkali AH, Chen J, Cui BK, Chen JJ, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, Hashimoto A, Hongsan S, Jones EBG, Larsson E, Li WJ, Li QR, Liu JK, Luo ZL, Maharachchikumbura SSN, Mapook A, McKenzie EHC, Norphanphoun C, Konta S, Pang KL, Perera RH, Phookamsak R, Phukhamsakda C, Pinruan U, Randrianjohany E, Singtripop C, Tanaka K, Tian CM, Tibpromma S, Abdel-Wahab MA, Wanasinghe DN, Wijayawardene NN, Zhang JF, Zhang H, Abdel-Aziz FA, Wedin M, Westberg M, Ammirati JF, Bulgakov TS, Lima DX, Callaghan TM, Callac P, Chang CH, Coca LF, Dal-Forno M, Dollhofer V, Fliegerová K, Greiner K, Griffith GW, Ho HM, Hofstetter V, Jeewon R, Kang JC, Wen TC, Kirk PM, Kytövuori I, Lawrey JD, Xing J, Li H, Liu ZY, Liu XZ, Liimatainen K, Lumbsch HT, Matsumura M, Moncada B, Nuankaew S, Parnmen S, de Azevedo Santiago ALCM, Sommai S, Song Y, de Souza CAF, de Souza-Motta CM, Su HY, Suetrong S, Wang Y, Wei SF, Wen TC, Yuan HS, Zhou LW, Réblová M, Fournier J, Camporesi E, Luangsa-ard JJ, Tسانathai K, Khonsanit A, Thanakitpipattana D, Somrithipol S, Diederich P, Millanes AM, Common RS, Stadler M, Yan JY, Li XH, Lee HW, Nguyen TTT, Lee HB, Battistin E, Marsico O, Vizzini A, Vila J, Ercole E, Eberhardt U, Simonini G, Wen HA, Chen XH, Miettinen O, Spirin V, Hernawati (2015) Fungal diversity notes 111–252-taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 75(1): 27–274. <https://doi.org/10.1007/s13225-015-0346-5>
- Bose T, Reynolds DR, Berbee ML (2014) Common, unsightly and until now undescribed: *Fumiglobus pieridicola* sp. nov., a sooty mold infesting *Pieris japonica* from western North America. *Mycologia* 106(4): 746–756. <https://doi.org/10.3852/13-288>
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25(15): 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>

- Cheewangkoon R, Groenewald JZ, Hyde KD, To-anun C, Crous PW (2012) Chocolate spot disease of Eucalyptus. *Mycological Progress* 11(1): 61–69. <https://doi.org/10.1007/s11557-010-0728-8>
- Chomnunti P, Schoch CL, Aguirre-Hudson B, Ko-Ko TW, Hongsanan S, Jones EBG, Kodsueb R, Phookamsak R, Chukeatirote E, Bahkali AH, Hyde KD (2011) Capnodiaceae. *Fungal Diversity* 51(1): 103–134. <https://doi.org/10.1007/s13225-011-0145-6>
- Chomnunti P, Hongsanan S, Aguirre-Hudson B, Tian Q, Peršoh D, Dhami MK, Alias AS, Xu JC, Liu XZ, Stadler M, Hyde KD (2014) The sooty moulds. *Fungal Diversity* 66(1): 1–36. <https://doi.org/10.1007/s13225-014-0278-5>
- Crous PW, Braun U, Groenewald JZ (2007) *Mycosphaerella* is polyphyletic. *Studies in Mycology* 58: 1–32. <https://doi.org/10.3114/sim.2007.58.01>
- Crous PW, Schoch CL, Hyde KD, Wood AR, Gueidan C, de Hoog GS, Groenewald JZ (2009) Phylogenetic lineages in the *Capnodiales*. *Studies in Mycology* 64: 17–47. <https://doi.org/10.3114/sim.2009.64.02>
- Crous PW, Groenewald JZ, Shivas RG, Edwards J, Seifert KA, Alfenas AC, Alfenas RF, Burgess TI, Carnegie AJ, Hardy GE, Hiscock N, Hüberli D, Jung T, Louis-Seize G, Okada G, Pereira OL, Stukely MJ, Wang W, White GP, Young AJ, McTaggart AR, Pascoe IG, Porter IJ, Quaedvlieg W (2011) Fungal Planet description sheets: 69–91. *Persoonia* 26(1): 108–156. <https://doi.org/10.3767/003158511X581723>
- Czachura P, Janik P, Piątek M (2025) *Chaetocapnodium magnum* and *Chaetocapnodium polonicum* from conifer resins disclose an unknown lifestyle in the *Capnodiales* (Dothideomycetes). *MycKeys* 119: 315–333. <https://doi.org/10.3897/mycokeys.119.159094>
- Haituk S, Withee P, Sangta J, Senwanna C, Khamsaw P, Karunarathna A, Hongsibsong S, Sringarm K, Prasad SK, Sommano SR, Cheewangkoon R (2022) Production of Non-Volatile Metabolites from Sooty Molds and Their Bio-Functionalities. *Processes* 10(2): 329. <https://doi.org/10.3390/pr10020329>
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Herath K, Jayasuriya H, Zink DL, Sigmund J, Vicente F, de la Cruz M, Basilio A, Bills GF, Polishook JD, Donald R, Phillips J, Goetz M, Singh SB (2012) Isolation, Structure Elucidation, and Antibacterial Activity of Methiosetin, a Tetramic Acid from a Tropical Sooty Mold (*Capnodium* sp.). *Journal of Natural Products* 75(3): 420–424. <https://doi.org/10.1021/np200857y>
- Höhnelt F (1910) Fragmenta zur Mykologie (Xi Mitteilung, Nr. 527 bis 573). *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien, Abt. I*, 119: 617–679. <https://www.biodiversitylibrary.org/part/234030>
- Hongsanan S, Tian Q, Hyde KD, Chomnunti P (2015) Two new species of sooty moulds, *Capnodium coffeicola* and *Conidiocarpus plumeriae* in Capnodiaceae. *Mycosphere* 6(6): 814–824. <https://doi.org/10.5943/mycosphere/6/6/14>
- Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie E, Sarma VV, Boonmee S, Jeewon R (2020) Refined families of Dothideomycetes: Dothideomycetidae and Pleosporomycetidae. *Mycosphere* 11(1): 1553–2107. <https://doi.org/10.5943/mycosphere/11/1/13>
- Hughes SJ (1976) Sooty Moulds. *Mycologia* 68(4): 693–820. <https://doi.org/10.1080/00275514.1976.12019958>
- Hyde KD, McKenzie E, Ko Ko T (2011) Towards incorporating anamorphic fungi in a natural classification-checklist and notes for 2010. *Mycosphere* 2: 1–88.

- Index Fungorum (2026) Index Fungorum (accessed 1 January 2026). <http://www.index-fungorum.org/Names/Names.asp>
- Jouraeva VA, Johnson DL, Hassett JP, Nowak DJ, Shipunova NA, Barbarossa D (2006) Role of sooty mold fungi in accumulation of fine-particle-associated PAHs and metals on deciduous leaves. *Environmental Research* 102(3): 272–282. <https://doi.org/10.1016/j.envres.2006.06.004>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Khodaparast SA (2006) A survey on citrus sooty mold fungi in Gilan Province, Iran. *Rostaniha* 7: 59–65. <https://civilica.com/doc/1542941> [Botanical Journal of Iran]
- Khodaparast SA, Pourmoghaddam MJ, Amirmijani A, Byrami F (2020) Phylogenetic structure of the Iranian *capnodiaceous* sooty mould fungi inferred from the sequences of rDNA regions and TEF1- α . *Mycological Progress* 19(2): 155–169. <https://doi.org/10.1007/s11557-019-01551-w>
- Kirk PM, Cannon PF, Minter DW, Stalpers JA (2008) Ainsworth and Bisby's Dictionary of the Fungi, 10th edn. CABI International, Wallingford, UK. <https://doi.org/10.1079/9780851998268.0000>
- Kress WJ, Erickson DL, Jones FA, Swenson NG, Perez R, Sanjur O, Bermingham E (2009) Plant DNA barcodes and a community phylogeny of a tropical forest dynamics plot in Panama. *Proceedings of the National Academy of Sciences of the United States of America* 106(44): 18621–18626. <https://doi.org/10.1073/pnas.0909820106>
- Léveillé JH (1847) Mycologie, mycétologie. Dictionnaire Universel d'Histoire naturelle. Victor Masson, Paris, 261–303.
- Li WJ, McKenzie EHC, Liu JK, Bhat DJ, Dai DQ, Camporesi E, Tian Q, Maharachchikumbura SSN, Luo ZL, Shang QJ, Zhang JF, Tangthirasun N, Karunarathna SC, Xu JC, Hyde KD (2020) Taxonomy and phylogeny of hyaline-spored coelomycetes. *Fungal Diversity* 100(1): 279–801. <https://doi.org/10.1007/s13225-020-00440-y>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Lu ZH, Wanasinghe DN, Madrid H, Maharachchikumbura SSN (2022) *Hyphocapnodia sichuanensis* gen. et sp. nov. (Capnodiaceae), a novel hyphomycete from Sichuan Province, China. *Phytotaxa* 564: 084–094. <https://doi.org/10.11646/phytotaxa.564.1.6>
- Lumbsch T, Huhndorf SM (2011) Myconet Vol. 14. Part One. Outline of Ascomycota-2009. Part Two. Notes on Ascomycete Systematics. Nos. 4751–5113. *Fieldiana. Life and Earth Sciences* 1: 1–64. <https://doi.org/10.3158/1557.1>
- MycoBank (2026) MycoBank. [accessed 1 January 2026] <https://www.mycobank.org>
- Pem D, Hyde KD, McKenzie EHC, Hongsanan S, Wanasinghe DN, Boonmee S, Darmostuk V, Bhat DJ, Tian Q, Htet Z, Senanayake IC, Niranjana M, Sarma VV, Doilom M, Dong W (2024) A comprehensive overview of genera in Dothideomycetes. *Mycosphere* 15(1): 2175–4568. <https://doi.org/10.5943/mycosphere/15/1/18>
- Persoon CH (1822) *Mycologia Europaea, seu Completa omnium fungorum in variis Europaea regionibus detectorum enumeratio, methodo naturali disposita. Sectio prima*. Erlangae: Impensis Ioannis Iacobi Palmii, 9–9.
- Reynolds DR, Gilbert GS (2005) Epifoliar fungi from Queensland, Australia. *Australian Systematic Botany* 18(3): 265–289. <https://doi.org/10.1071/SB04030>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian Phylogenetic

- Inference and Model Choice Across a Large Model Space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Schoch CL, Crous PW, Groenewald JZ, Boehm EWA, Burgess TI, de Gruyter J, de Hoog GS, Dixon LJ, Grube M, Gueidan C, Harada Y, Hatakeyama S, Hirayama K, Hosoya T, Huhndorf SM, Hyde KD, Jones EBG, Kohlmeyer J, Kruys Å, Lumbsch HT, Marvanová L, Mbatchou JS, McVay AH, Miller AN, Mugambi GK, Muggia L, Nelsen MP, Nelson P, Owensby CA, Phillips AJL, Phongpaichit S, Pointing SB, Pujade-Renaud V, Raja HA, Rivas Plata E, Robbertse B, Ruibal C, Sakayaroj J, Sano T, Selbmann L, Shearer CA, Shirouzu T, Slippers B, Suetrong S, Tanaka K, Volkmann-Kohlmeyer B, Wingfield MJ, Wood AR, Woudenberg JHC, Yonezawa H, Zhang Y, Spatafora JW (2009) A class-wide phylogenetic assessment of Dothideomycetes. *Studies in Mycology* 64: 1–15. <https://doi.org/10.3114/sim.2009.64.01>
- Spatafora JW, Sung G-H, Johnson D, Hesse C, O'Rourke B, Serdani M, Spotts R, Lutzoni F, Hibbett DS, Miadlikowska J, Reeb V, Gueidan C, Fraker E, Lumbsch HT, Schmitt I, Hosaka K, Aptroot A, Roux C, Schoch CL (2006) A five-gene phylogeny of *Pezizomycotina*. *Mycologia* 98(6): 1018–1028. <https://doi.org/10.1080/15572536.2006.11832630>
- Tennakoon DS, Kuo C-H, Maharachchikumbura SSN, Thambugala KM, Gentekaki E, Phillips AJL, Bhat DJ, Wanasinghe DN, de Silva NI, Promputtha I, Hyde KD (2021) Taxonomic and phylogenetic contributions to *Celtis formosana*, *Ficus ampelas*, *F. septica*, *Macaranga tanarius* and *Morus australis* leaf litter inhabiting microfungi. *Fungal Diversity* 108(1): 1–215. <https://doi.org/10.1007/s13225-021-00474-w>
- Tennakoon DS, Kuo C-H, Purahong W, Gentekaki E, Pumas C, Promputtha I, Hyde KD (2022) Fungal community succession on decomposing leaf litter across five phylogenetically related tree species in a subtropical forest. *Fungal Diversity* 115(1): 73–103. <https://doi.org/10.1007/s13225-022-00508-x>
- Theissen F (1915) *Mykologische Abhandlungen. Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 66: 231–234. <https://www.biodiversitylibrary.org/item/238853>
- Thungdee S, Haituk S, Withee P, Cheewangkoon R, Suwannarach N, Marasinghe D, Hongsan S (2023) Unraveling Capnodiaceae species in Northern Thailand. *Phytotaxa* 620(2): 143–156. <https://doi.org/10.11646/phytotaxa.620.2.2>
- Tian WH, Liu JW, Jin Y, Chen YP, Liu JK, Maharachchikumbura S, Wanasinghe D, Su PW (2024) Morphological and phylogenetic studies of *Ascomycota* from gymnosperms in Sichuan Province, China. *Mycosphere* 15(1): 1794–1900. <https://doi.org/10.5943/mycosphere/15/1/16>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wheeler Q, Assis L, Rieppel O (2013) Heed the father of cladistics. *Nature* 496(7445): 295–296. <https://doi.org/10.1038/496295a>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğan M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Hawksworth DL, Jones EBG, Karunarathna SC, Thines M (2022) Outline of Fungi and fungus-like taxa–2021. *Mycosphere* 13(1): 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>

- Williams D, Schmitt M, Wheeler Q [Eds] (2016) The Future of Phylogenetic Systematics: The Legacy of Willi Hennig. Cambridge University Press, Cambridge. <https://doi.org/10.1017/CBO9781316338797>
- Yang H, Ariyawansa HA, Wu HX, Hyde KD (2014) The genus *Leptoxyphium* (Capnodiaceae) from China. *Phytotaxa* 176(1): 174–183. <https://doi.org/10.11646/phytotaxa.176.1.17>
- Yang CL, Xu XL, Zeng Q, Liu LJ, Liu F, Deng Y, Wang FH, Sun QR, Jeewon R, Hyde KD, Jayawardena RS, McKenzie EHC, Wanasinghe DN, Liu YG, Xiao QG, Han S, Yang H, Li SJ, Liu L, Xie JL (2024) Bambusicolous mycopathogens in China with an update on taxonomic diversity, novel species, pathogenicity, and new insights. *Mycosphere* 15(1): 4788–4918. <https://doi.org/10.5943/mycosphere/15/1/21>
- Yang CL, Li XY, Xiang SS, Xu XL, Zeng Q, Sun QR, Liu JK, Maharachchikumbura SSN, Lv YC, Wang FH, Ji-Zuo JL, A-Niu XY, Liu YG (2025) Microfungi associated with plant diseases on horticultural vegetation in southwestern China. *Mycosphere* 16(1): 1861–2001. <https://doi.org/10.5943/mycosphere/16/1/11>
- Zeng XY, Tan TJ, Tian FH, Wang Y, Wen TC (2023) OFPT: A one-stop software for fungal phylogeny. *Mycosphere* 14(1): 1730–1741. <https://doi.org/10.5943/mycosphere/14/1/20>

Supplementary material 1

***Dalbergia assamica* rbcL complete sequence**

Authors: Jipeng Sun, Entaj Tarafder, Tao Chen, Sinang Hongsan, Shengxi Chen, Chada Norphanphoun, Samantha C. Karunarathna, Xiangyu Zeng

Data type: fasta

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.129.182290.suppl1>