ETHNOBIOLOGY AND CONSERVATION

RESEARCH ARTICLE

Ethnobiology and Conservation, 15:13 (10 June 2026)
doi:10.15451/ec2026-06-15-13-1-16
ISSN 2238-4782 ethnobioconservation.com

Future climatically stable habitat of a vulnerable medicinal tree, *Warburgia salutaris* (Pepper-bark tree), overlaps with some protected areas in southern Africa

Percy Jinga[1,*] and Tawanda Manyangadze[2]

ABSTRACT

Climate change will affect the distribution of medicinal plant species in sub-Saharan Africa, with potential negative consequences to especially rural and poor communities that depend on them for traditional medicines. The aim of the study was to determine future climatically stable habitat of *Warburgia salutaris*, a commonly used vulnerable medicinal plant species. An ensemble species distribution model (SDM) was built from eight algorithms and used to estimate the current and future distributions of the species. Six environmental variables and 146 occurrence records were used to calibrate individual SDMs before combining them into the ensemble SDM. The ensemble SDM was projected to 2041–2060 and 2081–2100 under two shared socioeconomic pathways (SSPs), namely SSP2-4.5 and SSP5-8.5. The projections were made under three general circulation models (GCMs), which were CanESM5, IPSL-CM6A-LR and MIROC6. Results show that the current suitable habitat of the species occurs in South Africa, Eswatini, Malawi, Mozambique and Zimbabwe. Under both time periods and SSPs, the suitable habitat of the species will markedly contract, especially under SSP5-8.5 in 2081–2100 (CanESM5 = -79%, IPSL-CM6A-LR = -87%, MIROC6 = -71%). Climate refuges that overlap with the future suitable habitat of *W. salutaris* were identified in protected areas (PAs), such as Malalotja Nature Reserve (Eswatini), Itala Nature Reserve (South Africa), and Nyanga National Park (Zimbabwe). The climate refuges in PAs may be priority areas to introduce the species.

Keywords: Climate change, Conservation, Protected area, Traditional medicine, *Warburgia salutaris*.

1 Biological Sciences Department, Bindura University of Science Education, Private Bag 1020, Bindura, Zimbabwe.
2 Geosciences Department, Bindura University of Science Education, Private Bag 1020, Bindura, Zimbabwe.
* Corresponding author ✉. E-mail address: PJ (pjinga@buse.ac.zw, TM (tmanyangadze@buse.ac.zw)

SIGNIFICANCE STATEMENT

We used ensemble modeling in the R package 'biomod2' to estimate distribution of the current suitable habitat of *W. salutaris*, a medicinally important and vulnerable tree species, and to predict shifts in the distribution in future. We used eight algorithms to build an ensemble model for the species. Our work is original and is not under consideration elsewhere. To our knowledge, this is the first study to use ensemble modeling to predict the impacts of climate change on the species across its entire natural range. Our study shows that the size of the suitable habitat of *W. salutaris* will contract under climate change. The contraction is likely to impact people who rely on the species for traditional medicine. Our results also show that there are climate refuges in protected areas (PAs) within the natural range of the species. The climate refuges in PAs may serve as conservation areas of the species.

INTRODUCTION

Warburgia salutaris (Bertol. f.) Chiov. (pepper-bark tree, Canellaceae) is an ethnomedicinally important tree to especially rural dwellers of southern Africa (Meddows-Taylor and Ramadwa 2025). It is naturally distributed in Swaziland, Lesotho, Malawi, Mozambique, South Africa and Zimbabwe (Maroyi 2013; Senkoro et al. 2020; Glennon et al. 2023). Habitat degradation and overharvesting have resulted in population declines of *W. salutaris* throughout its native range (Senkoro et al. 2019). The species is overharvested mainly because of its use as traditional medicine for a variety of ailments (Hollmann and Van der Schijff 1996; Rabe and van Staden 2000; Leonard and Viljoen 2015; Maroyi 2013). Leaf, bark and root extracts have been shown to have pharmacological effects, such as anti-inflammatory, antioxidant, antimycobacterial, antimalarial, and antifungal activities (Glennon et al. 2023, Khumalo et al. 2024). Several bioactive compounds from bark (Khumalo et al. 2019; Abdelfattah et al. 2022) and leaf (Steenkamp et al. 2013; Mashamba et al. 2022) have already been reported, and biomedical studies on the pharmacological efficacy of the compounds are ongoing (Meddows-Taylor and Ramadwa 2025). Adult individuals are severely debarked, which causes tree mortality and ultimately local population extinction (Dludlu et al. 2017). As a result, *W. salutaris* is classified as vulnerable on the IUCN Red List (IUCN 2024). Nationally, the species is classified as endangered in South Africa (van den Bosch et al. 2023) and Malawi (Msekandiana and Mlangeni 2002), critically endangered in Swaziland (Dlamini and Dlamini 2002), and vulnerable in Mozambique (Izidine and Bandeira 2002). In Zimbabwe, the status is more extreme as the species is classified as extinct in the wild (Maroyi 2008), signaling that plans for reintroduction of *W. salutaris* in the country are necessary.

Local ecological knowledge and cultural practices have historically been used to ameliorate some of the threats faced by *W. salutaris* in its native range (Senkoro et al. 2019). The cultural practices include rotational harvesting that allows regeneration, selective harvesting of only mature individuals (Terer et al. 2012; Senkoro et al. 2019), regulating harvesting time according to critical life phases of the species (Schmidt and Ticktin 2012), and controlled debarking that allows bark regeneration and avoids tree mortality (Ssenku et al. 2022). However, recently, the traditional conservation practices of *W. salutaris* are failing to curb overharvesting of especially the bark (Maroyi 2013). Human population growth, the comparatively high cost of modern pharmaceuticals, and the strengthening belief that traditional medicines have fewer side effects than modern drugs are some factors that have led to unsustainable harvesting of medicinally important indigenous tree species, such as *W. salutaris* (Ssenku et al. 2022). To regenerate populations of *W. salutaris*, reintroductions into the wild (Maroyi 2011), distribution of seedlings to traditional healers for regeneration (van den Bosch et al. 2023) and establishment of domesticated populations (Maroyi 2013) have been explored. Introductions, however, should be informed by maps showing where the species is likely to successfully thrive. Introduction in climatically stable protected areas (PAs) has also been suggested as a viable alternative for the conservation of *W. salutaris* since the PAs may provide a higher chance of successful growth under climate change (Hannweg et al. 2016).

In addition to habitat fragmentation and overharvesting, climate change has been identified as a severe threat to medicinal plant species, particularly in sub-Saharan Africa (Jinga et al. 2020; Lyam et al. 2022; Esser and Weldemariam 2023). Mean temperature in sub-Saharan Africa has increased by 0.7°C during the 20th century, and it is forecasted to increase by 0.2°C – 0.5°C per decade in the 21^{st} century (Hulme et al. 2001). While mean temperature is forecasted to increase across the African continent, precipitation will likely show regional variability, with a decrease in southern Africa (Almazroui et al. 2020; Dosio et al. 2021). Temperature and precipitation changes are likely to be major causes of plant biodiversity loss, especially in sub-Saharan Africa (Midgley and Bond 2015; Sintayehu 2018). Range contraction of several African medicinal plant species has already been forecasted under climate change, including *Pterocarpus angolensis* (De Cauwer et al. 2014), *Uapaca kirkiana* (Jinga et al. 2020), *Senegalia senegal* (Lyam et al. 2022), *Afzelia africana* (Balima et al. 2022) and *Boswellia* spp. (Esser and Weldemariam 2023). The forecasted future disappearance of the suitable habitat of medicinal plant species in Africa is especially disastrous to approximately 80% of the population that relies on traditional plant medicines (Ampomah et al. 2023). It is important, therefore, to investigate the impact of climate change on medicinal plant species in southern Africa, such as *W. salutaris*, and to suggest conservation options for long term viability and persistence of populations. The impact of climate change on *W. salutaris* has been investigated at the local level in Mozambique using a single modeling algorithm and fewer occurrence records (Senkoro et al. 2024). This study investigates the impact of climate change over the entire natural range of the species which is not constrained by political boundaries in southern Africa. This study also uses ensemble modeling with more occurrence records, making the results more robust and broad-based for conservation planning across the natural range.

The impact of climate change on the distribution and abundance of species is now commonly investigated by species distribution models (SDMs). SDMs correlate environmental variables to species occurrence records to predict the potential geographic distribution of the species (Sillero et al. 2021). SDMs are also used to forecast changes in the distribution of species under climate change (Santini et al. 2021). The proliferation of online public data portals hosting species occurrence records as well as potential environmental variables have increased the use of SDMs (Zurell et al. 2020). The premise in SDMs is that species are in pseudoequilibrium with the environment (Guisan and Thuiller 2005; Araujo and Guisan 2006; Sinclair et al. 2010). Thus, correlating occurrence records and environmental variables should predict suitable habitat of the species (Sinclair et al. 2010). However, the pseudoequilibrium assumption can be problematic, especially for species under severe pressure due to dispersal limitation and demographic stochasticity (Soley-Gaurdia et al. 2024). Other notable shortcomings of SDMs include failure to simulate dispersal and competition (Sinclair et al. 2010; Moonlight et al. 2020). Regardless of these limitations, SDMs have been successfully applied in conservation planning (Regos et al. 2021), identification of niches under climate change (Estrada et al. 2018), and delineating boundaries of PAs (Carvalho et al. 2011), among several fields. For long-term viability after introductions and or reintroductions of imperiled species, such as *W. salutaris*, SDMs may be used to identify target areas that will remain climatically stable.

The aim of the study was to determine the current geographic distribution and impact of future climate change on *W. salutaris*. Distribution maps are especially important to locate areas where the species may be introduced and to direct botanical explorations and plant surveys. Using an ensemble SDM, we first sought to (i) determine the current suitable habitat of *W. salutaris*, then (ii) project how the suitable habitat of *W. salutaris* changes under different climate change scenarios in 2041–2060 and 2081–2100, and finally (iii) we used the projections to identify climate refuges in national parks and other PAs that are priority targets for conservation and reintroduction efforts. Climate refuges in PAs are usually protected from illegal harvesting (Nutall et al. 2022), and, thus, they are ideal areas to introduce and or reintroduce species at risk from climate change, overharvesting and other anthropogenic threats.

MATERIAL AND METHODS

Occurrence records and environmental variables

Occurrence records were obtained from the Global Biodiversity Information Facility online data portal (www.gbif.org) (GBIF 2024). Automatic filters on the GBIF portal were used to screen occurrence records with potential errors, such as those with out of range coordinates, swapped coordinates, and invalid precision and accuracy coordinates. The advantage of including GBIF occurrence records is that they have a comprehensive coverage since some of them are collected by citizen science tools (Dyderski et al. 2018). Citizen science tools are especially valuable in sampling areas not covered by forest inventories and too large for reliable sampling by researchers. Occurrence records were also obtained from the RAINBIO online data portal (`https://gdauby.github.io/rainbio/download_page.html`) (Dauby et al. 2016). The RAINBIO database is a compilation of 13 datasets of georeferenced occurrence records of African vascular plant species located south of the Sahel. The Tropicos (`http://www.tropicos.org`) online data portal (Missouri Botanical Garden, St. Louis, MO, USA) was also a source of occurrence records. Additional occurrence records were obtained from Senkoro et al. (2020) and Dludlu et al. (2017). All occurrence records falling out of the known south-east African range of *W. salutaris* were removed. Occurrence records collected before 1970 were also discarded to suit the temporal range of bioclimatic variables.

After removing duplicate occurrences in Microsoft Excel using conditional formatting to highlight repeated coordinates, a total of 146 occurrence records were obtained. To have a relatively even distribution, the occurrence records were resampled in R (R Core Team 2024) using the 'spThin' version 0.2.0 R-package (Aiello-Lammens et al. 2015). SDMs built with occurrence records showing uneven sampling will fit a biased environmental signal that causes inflated model performances (Hijmans 2012). A minimum distance of 10 km between occurrence records was used in resampling to obtain a relatively uniform distribution and to ensure that at most one record per pixel was retained for use in SDMs (Figure 1).

Bioclimatic variables (Booth et al. 2014) and elevation obtained from the Worldclim version 2.1 online public data portal (www.worldclim.org) (Fick and Hijmans 2017) were used as environmental variables. Bioclimatic variables describe quarterly or yearly variations in temperature and precipitation and are more influential in determining plant distributions than monthly climate variables (Bede-Fazekas and Somodi 2020). Bioclimatic variables are ecologically and phys-




Figure 1. Location of thinned occurrence records of *Warburgia salutaris* (Pepper-bark tree) in southern Africa that were used to calibrate species distribution models.

iologically more relevant to vegetation establishment and growth (Title and Bemmels 2018), and they have been widely used in SDMs to delineate conservation areas (Varela et al. 2022), forecast impact of climate change (Esser and Weldemariam 2023), and to generate distribution maps of species (Hama and Khwarahm 2023), among several applications.

The bioclimatic variables and elevation were selected for calibration of individual SDMs after correlation analysis using the 'usdm' version 2.1-7 R-package (Naimi et al. 2014). Correlated variables, among several drawbacks, increase computation time, decrease model transferability, increase over-fitting, and decrease signal-to-noise ratio (De Cauwer et al. 2014; Jarnevich et al. 2015).Variables with a Spearman's correlation coefficient (r) of less than 0.70 (Dormann et al. 2013; Jarnevich et al. 2015) and a variance inflation factor (VIF) of less than 10 (Steen et al. 2017) were used in individual SDM calibrations. The selected variables, downloaded at a resolution of 2.5 min, were mean diurnal range (BIO2), isothermality (BIO3), mean temperature of wettest quarter (BIO8), precipitation of driest month (BIO14), precipitation of warmest quarter (BIO18), and elevation.

Calibration, evaluation and ensembling of SDMs

An R-based software, 'biomod2' version 4.2-4 (Thuiller 2003; Thuiller et al. 2009, 2023) was used to calibrate and ensemble individual SDMs. Up to 12 individual modeling algorithms can be calibrated simultaneously and ensembled in 'biomod2.' Eight algorithms were used in this study, and these were artificial neural network (ANN), classification tree analysis (CTA), flexible discriminant analysis (FDA), generalized boosted model (GBM), generalized linear model (GLM), multivariate adaptive regression splines (MARS), random forest (RF) and surface range envelope (SRE). To account for spatial autocorrelation and reduce inflation of model performance due to spatial sorting bias, we implemented spatial block cross-validation. Occurrence records were combined with randomly generated pseudo-absences (n = 10,000) sampled within the study extent (10°S – 38°S latitude and 10°E – 42°E longitude). A rela-

tively large pseudo-absence sample was used to stabilize model parameter estimation and ensure adequate representation of background environmental conditions. Environmental variables were extracted for all presence-pseudo-absence locations, and records containing missing environmental values were removed before modeling. Spatial partitioning was conducted using the "spatialBlock" function in the R-package "blockCV" version 3.2-0 (Valavi et al. 2019). Blocks were generated using a distance threshold of 400 km and partitioned into $k = 10$ folds. The 400 km block size was selected to ensure meaningful geographic separation between training and testing subsets while retaining sufficient sample size within each fold. Ten folds were used to maximize spatial independence across validation runs while maintaining adequate representation of presence and pseudo-absence records per fold. Spatial partitioning provides a more conservative and ecologically realistic estimate of predictive performance compared to random cross-validation, which can inflate evaluation metrics in spatially structured datasets (Roberts et al. 2017; Yates et al. 2023).

Blocks were randomly allocated to folds across 100 iterations to ensure balanced representation of presences and pseudoabsences. Folds containing only a single response class were excluded to prevent model fitting errors. A user-defined cross-validation table was constructed such that, for each run, models were trained on $k - 1$ spatial folds and evaluated on the withheld fold. This spatially explicit cross-validation strategy ensures that training and testing data were geographically separated, thereby providing a more realistic estimate of predictive performance under spatial extrapolation. The SDMs were evaluated using the area under the receiver operating characteristic curve (ROC) and true skill statistics (TSS).

To determine whether spatial structure remained in model residuals, we quantified spatial autocorrelation using Moran's *I*. Predicted probabilities were extracted for each algorithm and averaged across cross-validation runs to obtain a mean prediction per observation point. Spatial coordinates were projected to a metric coordinate reference system (UTM Zone 35S; EPSG: 32735) prior to distance-based neighbor construction. Moran's I statistic was computed from the models' residuals to test for significant spatial autocorrelation. Statistical significance was assessed using the analytical approximation implemented in the 'moran.test' function in the R-package "spdep' version 1.4-2 (Bivand 2022). This procedure allowed evaluation of whether spatial block cross-validation sufficiently reduced spatial dependence in model residuals. Significant positive Moran's I values would indicate remaining spatial autocorrelation and potential overestimation of model predictive ability.

The cross-validated individual SDMs were combined into an ensemble SDM using a weighted mean approach (Emwmean), retaining only individual models with ROC ≥ 0.80. A threshold of ROC ≥ 0.80 was applied to ensure that only models demonstrating at least fair discriminatory capacity contributed to the ensemble, thereby enhancing predictive robustness and minimizing the influence of low-performing algorithms (Kafle et al. 2023). Overall ensemble SDM performance has been shown to be highest when individual SDMs are weighted according to their evaluation score compared to other methods (Marmion et al. 2009; Hao et al. 2019). The ensemble SDM was used to determine the current suitable habitat of *W. salutaris* as well as to forecast under climate change. Distribution maps were converted into binary presence/absence maps using a ROC-derived threshold (ROC = 0.901) calculated during ensemble model evaluation in 'biomod2'. The binary maps were used to calculate range size changes under different climate change scenarios.

Identification of future climatically stable areas

The ensemble SDM was projected in southern Africa to 2041–2060 and 2081–2100 under two shared socio-economic pathways (SSPs) and three general circulation models (GCMs). SSPs are successors to representative concentration pathways (RCPs) and they take into consideration population growth, economic advancement, urbanization and technological development in determining future greenhouse gas emissions (Gidden et al. 2019; Riahi et al. 2017). In this study, the middle-of the-road SSP2-4.5 and the pessimistic SSP5-8.5 were used. The two SSPs were used in order to capture the full range of the impact of climate change, from the realistic to the most pessimistic. A map generated under SSP5-8.5 for 2081–2100 was used further to identify climate refuges that corresponded with PAs under climate change. Climate refuges identified under the pessimistic SSP5-8.5 scenario are most likely to be climatically conducive in future even under moderate and optimistic scenarios.

Variability in results of SDMs emanates from GCMs (Goberville et al. 2015). GCMs assess the response of climate to greenhouse gas emissions due to trends in land and energy use (Rogelj et al. 2012). The GCMs vary due to several factors, including use of different spatial resolutions and parameterizations of natural processes (Timbal 2004). To cater for inter-GCM variability, three GCMs were used, and these are the Canadian Earth System Model version 5 (CanESM5) (Swart et al. 2019), the Institut Pierre-Simon Laplace Climate Model version 6A Low Resolution (IPSL-CM6A-LR) (Boucher et al. 2020), and the Model for Interdisciplinary Research on Climate

version 6 (MIROC6) (Tatebe et al. 2019). The GCMs were selected due to their predictive accuracy of climatic variables over the African continent (Zebaze et al. 2025) and completeness of data over all SSPs. A single consensus distribution map was produced in ArcGIS software showing presence in all maps generated from the three GCMs for each SSP and time period.

The binary consensus distribution map of *W. salutaris* for 2081–2100 under SSP5-8.5 was overlaid on a map of PAs in southern Africa to identify climate refuges that overlap with or are within the areas under protection. The period 2081–2100 was used to identify climate refuges that will remain stable in the long-term. The map of PAs was obtained from the world database on PAs of the United Nations Environment Program-World Conservation Monitoring Centre (www.protectedplanet.net) (UNEP-WCMC and IUCN 2024). The climate refuges that overlap with the suitable habitat of *W. salutaris* may help conservationists with options on where the species may be introduced and thrive successfully in future.

RESULTS

Model evaluation and current distribution

Mean ROC scores for the individual SDMs ranged from 0.744 to 0.892, while TSS scores ranged from 0.489 to 0.663 (Additional File 1). Averaged across all the SDMs, the scores showed robust performance (ROC = 0.836, TSS = 0.530). The Moran's I value (I = 0.165, $P < 0.05$) indicated a weak positive but significant spatial autocorrelation, suggesting that spatial structure in the residuals was minimal and that the spatial blocking strategy sufficiently reduced spatial dependence among folds. The ensemble SDM of the current distribution of *W. salutaris* also showed robust performance (ROC = 0.901, TSS = 0.895). After three permutations, two variables, Elevation and precipitation of warmest quarter (BIO18), showed the highest mean variable importance scores (Additional File 2). The variable importance scores show the contribution of each variable to the predictive power of the ensemble SDM. Response curves showed that the optimum precipitation of warmest quarter (BIO18) for *W. salutaris* is around 600 mm while the optimum mean temperature of the wettest quarter (BIO8) ranges from 20°C to 25°C (Additional File 3). Response curves also showed that suitability of the habitat to *W. salutaris* increases with increasing precipitation of the driest month (BIO14) up to 40 mm, and decreases beyond an altitude of 2,000 m.

The current distribution map shows potential presence of *W. salutaris* in South Africa, Swaziland, Malawi, Mozambique and Zimbabwe (Figure 2). In South Africa, the species was predicted to occur predominantly in the north-eastern provinces of KwaZulu Natal, Mpumalanga and Limpopo. Eswatini, formerly Swaziland, was shown to be largely suitable for *W. salutaris*. The eastern highlands in Zimbabwe and the southern region of Malawi are also priority areas for botanical exploration in search of the species. Mozambique has the largest potential distribution, especially in the provinces of Manica, Maputo and Zambezia. The relatively restricted potential distribution of *W. salutaris* in southern Africa has implications for viability of populations, supply of the species' medicinal products and designing of conservation strategies.

Suitable habitat under climate change

Averaged across three GCMs, the suitable habitat of *W. salutaris* contracted under the two SSPs and time periods. Under SSP2-4.5, the suitable habitat contracted from approximately 647,000 km^2 to 421,000 km^2 (-35%) in 2041—2060, and to 304,000 km^2 (-53%) in 2081–2100. Under SSP5-8.5, the contractions were approximately to 381,000 km^2 (-41%) in 2041—2060, and to 135,000 km^2 (-79%) in 2081–2100. The three GCMs showed similar results of contraction under all the SSPs and time periods (Table 1).

The largest contraction was observed under the pessimistic SSP5-8.5, thus, highlighting the grim consequences of uncontrolled greenhouse gas emission to the occurrence of *W. salutaris*. Under all projections, contraction of the natural range was especially notable in Malawi, Mozambique and Zimbabwe (Figure 3).

There are several PAs that contain or overlap with predicted climate refuges for *W. salutaris* in southern Africa in 2081–2100 (Table 2).The PAs include national parks, nature reserves, forest reserves and state forests. In Eswatini, Malalotja Nature Reserve was identified to contain climate refuges for *W. salutaris*. Most PAs were identified in South Africa and they include Barberton, Coleford, Doreen Clark and Itala Nature Reserves. In Zimbabwe, where *W. salutaris* has been classified as extinct in the wild, the PAs that overlap with climate refuges include Chipinge Safari Area and Nyanga National Park. Some entire PAs are climate refuges, such as Qudeni Forest Reserve in South Africa and Vumba Botanical Garden in Zimbabwe. The larger PAs, such as Addo-Elephant National Park and Maloti Drakensberg Conservation and Development Area in South Africa, partially overlap with climate refuges of *W. salutaris*.




Figure 2. Geographic distribution of the current suitable habitat of *Warburgia salutaris* (Pepper-bark tree) in southern Africa determined by ensemble species distribution modeling.

Table 1. Percentage changes of size of the suitable habitat of *Warburgia salutaris* (Pepper-bark tree) estimated for two time periods by an ensemble species distribution model under three general circulation models (GCM) and two shared socio-economic pathways (SSPs).

	Shared socio-economic pathway (SSP)/ Year (%)			
GCM	**SSP 2-4.5 / 2060**	**SSP 5-8.5 / 2060**	**SSP 2-4.5 / 2100**	**SSP 5-8.5 / 2100**
CanESM5	-39.872	-44.573	-59.701	-78.921
IPSL-CM6A-LR	-29.424	-38.190	-51.869	-87.292
MIROC6	-34.824	-40.502	-46.618	-71.109
Mean change	-34.707	-41.088	-52.729	-79.107

DISCUSSION

The current distribution of *W. salutaris* has often been described as a list of countries where the species has been sampled (e.g., Maroyi et al. 2013; Glennon et al. 2023; van den Bosch et al. 2023) or as occurrence records where samples were collected (e.g., Dludlu et al. 2017; Senkoro et al. 2020). Although useful to some extent, country-level descriptions have the disadvantage of assuming presence even in areas where the species does not occur within the countries (Rathore and Sharma 2023). Similarly, maps of occurrence records only show locations that were accessed by researchers, while those that are inaccessible for sampling due to rough terrain and absence of roads are not represented (Fourcade et al. 2014). The map of the current suitable habitat established in this study represents a first attempt to show the spatial distribution of *W. salutaris* within countries and also in locations beyond the reach of traditional herbalists and researchers. The ensemble SDM map is an additional resource that may be used to determine forest management practices, guide botanical explorations and to suggest areas of introduction or preservation.

Warburgia salutaris is an evergreen species that occupies diverse habitats including coastal and riverine regions, dunes, mundane forests, open woodland, thickets and kloofs (Kotina et al. 2014; Leonard et al. 2023). The suitability of habitat remains high at a wide range of precipitation of driest months. The high habitat suitability under a wide precipitation range infers that the species has some evolutionary traits to respond to water extremes up to some point beyond which suitability declines. The evolutionary traits include stomatal density and thickness of cuticle (Kotina

et al. 2014). The leaves have less stomata and thicker cuticle on the adaxial compared to the abaxial leaf surface (Kotina et al. 2014). On the adaxial surface, the cuticle has a mean thickness of 5.0 μm while on the abaxial surface, the mean cuticle thickness is 2.4 μm (Kotina et al. 2014). The thicker cuticle and less stomata on the adaxial surface reduce water loss by evaporative cooling. The ability to reduce water loss likely enables *W. salutaris* to occupy habitats of different water availability levels. However, as temperature of wettest quarter exceeds 25 °C and precipitation of warmest quarter goes below 250 mm, the habitat suitability decreases. This study has shown that although there are some evolutionary traits to respond to water and temperature stress levels, the projected climate change conditions are beyond the tolerance limit of *W. salutaris*, likely rendering the evolutionary traits inadequate for adaptation.

Globally, climate change has been shown to reduce the natural range of medicinal plants. The range contractions are a result of several factors, such as intolerance to climate extremes of the medicinal plants (Applequist et al. 2019) and climate change-accelerated spread of invasive species and exotic pathogens (Pyšek et al. 2020; Guégan et al. 2023). Intolerance to climate extremes has been shown in *Draba litamo* (Rodríguez et al. 2018), *Tylophora hirsuta*, (Khanum et al. 2013), *Rosa arabica* (Abdelaal et al. 2019), *Pterocarpus angolensis* (De Cauwer et al. 2014) and *Boswellia* sp. (Lemenih et al. 2014; Bongers et al. 2019). This study has also demonstrated the intolerance of *W. salutaris* to climate change. Overharvesting and climate change will combine to exacerbate population contraction that will cause inbreeding and loss of genetic variation, making the small populations less resilient to pests and diseases. In addition to disruption of the primary healthcare of economically vulnerable communities, the unavailability of medicinal plants due to climate change in Africa would also be devastating to collectors and vendors who rely on selling traditional medicinal products.

The forecasted higher contraction under SSP5-8.5 shows the dire consequences of unmitigated greenhouse gas emissions and associated climatic disruption. In southern Africa, the predicted long-term increase in mean temperature (Hulme et al. 2001) and decrease in mean annual precipitation (Dosio et al. 2021) due to climate change will likely disrupt establishment and growth, especially of those plants adapted to mesic environments. To mitigate the effects of the forecasted contraction of the natural range in the immediate term, current cultivation efforts by communities living in areas found in this study to be suitable for the species should be encouraged and supported. Once populations of *W. salutaris* are established in climatically suitable habitat, the communities may also be educated on the importance of sustainable harvesting and the practice of other traditional ecological man-




Figure 3. Consensus maps generated from three General Circulation Models of the forecasted distribution of the suitable habitat of *Warburgia salutaris* (Pepper-bark tree) under SSP2-4.5 in 2041–2060 (a), SSP5-8.5 in 2041–2060 (b), SSP2-4.5 in 2081-2100 (c), and SSP5-8.5 in 2081-2100 overlaid with protected areas (d).

Table 2. Protected areas that overlap with or contain predicted climate refuges of *Warburgia salutaris* (Pepper-bark tree) in 2081–2100 in southern Africa.

Country	Name of protected area	Type of protected area
Eswatini	Malatotja	Nature Reserve
South Africa	Addo-Elephant	National Park
	Barberton	Nature Reserve
	Coleford	Nature Reserve
	Doreen Clark	Nature Reserve
	Entumeni	Nature Reserve
	Golden Gate Highlands	National Park
	Karkloof	Nature Reserve
	Maloti Drakensberg	Conservation Area
	Mount Currie	Nature Reserve
	Nkandla	Nature Reserve
	Qudeni	Nature Reserve
Zimbabwe	Chipinge	Safari Area
	Stapleford	State Forest
	Nyanga	National Park
	Erin	State Forest
	Nyangui	State Forest

agement practices.

The establishment of PAs has been hailed as a significant conservation strategy to prevent biodiversity loss (Elsen et al. 2020; Leberger et al. 2020). In recognizing their role in conservation, countries under the Convention on Biological Diversity set a target to extend PAs to cover at least 17% of terrestrial and inland water, and 10% of coastal and marine areas (CBD 2024). PAs often contain large contiguous patches of natural land cover that provide critical habitat for sensitive and threatened species. The contiguous natural land cover helps to maintain intact, functioning ecosystems (Langdon and Lawler 2015). Climate refuges within PAs may potentially support species that migrate into them by providing climatically stable habitat (Hole et al. 2009; Sweet et al. 2019). In southern Africa, the PAs shown to contain climate refuges for *W. salutaris* may play a crucial role in conservation of the vulnerable species.

The climate refuges within PAs identified in this study may be used to establish pilot plots of *W. salutaris*. Establishment of the plots should ideally be a joint activity among government agencies responsible for PAs, university researchers, NGOs, and other interested stakeholders. The pilot plots may be used to track ecological data on development and survival before full-scale introduction. The established populations in the PAs may serve as future sources of seeds or cuttings for afforestation where the species can be harvested. It is, therefore, recommended that the identified climate refuges in PAs be considered as potential areas where *W. salutaris* may be introduced and maintained for long term persistence.

CONCLUSION

The study applied ensemble species distribution modeling to more accurately map the current range of *W. salutaris*, a culturally and medicinally tree species threatened by climate change, habitat loss, and overharvesting. Climate projections indicate a marked contraction of its suitable habitat, highlighting the species' vulnerability to rising temperatures and shifting precipitation patterns. However, the identification of climate refuges within PAs located in the natural range of the species offers a strategic opportunity for conservation planning. By integrating ensemble mod-

eling with spatial conservation priorities, this research offers a strategic framework for safeguarding *W. salutaris* in the face of climate uncertainty. While the ensemble modeling results are robust and provide a valuable foundation for conservation planning, future studies would benefit from incorporating biotic interactions, such as competition and dispersal dynamics. These additions in modeling would enhance ecological realism and improve predictive accuracy, especially under rapidly changing environmental conditions.

ACKNOWLEDGMENTS

Appreciation goes to the Biological Sciences and Geosciences Departments at Bindura University of Science Education for supporting the study by providing computing resources and GIS software.

DATA AVAILABILITY

Occurrence records of *Warburgia salutaris* are publicly available at https://doi.org/10.15468/dl.f935qg, `https://gdauby.github.io/rainbio/download_page.html`, and `http://www.tropicos.org`. Ocurrence records are also available from Senkoro et al. (2020) and Dludlu et al. (2017). Environmental variables are publicly available at `www.worldclim.org`. The global map of protected areas is available at `http://protectedplanet.net`. The R code for model calibration, validation, ensembling and projection is available from the authors upon request.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

CONTRIBUTION STATEMENT

Conceived of the presented idea: PJ, TM Carried out the experiment: PJ
Carried out the data analysis: PJ
Wrote the first draft of the manuscript: PJ
Review and final write of the manuscript: PJ, TM
Supervision: PJ, TM

REFERENCES

Abdelaal M, Fois M, Fenu G, Bacchetta G (2019) **Using MaxEnt modeling to predict the potential distribution of the endemic plant *Rosa arabica* Crép. in Egypt**. *Ecological Informatics* 50: 68–75. doi: 10.1016/j.ecoinf.2019.01.003.

Abdelfattah MAO, Dmirieh M, Bakrim WB, Mouhtady O, Ghareeb MA, Wink M, Sobeh M (2022) **Antioxidant and anti-aging effects of *Warburgia salutaris* bark aqueous extract: Evidences from *in silico*, *in vitro* and *in vivo* studies.** *Journal of Ethnopharmacology* 292: 115187. doi: 10.1016/j.jep.2022.115187.

Aiello-Lammens EM, Boria AR, Radosavljevic A, Vilela B, Anderson PR (2015) **spThin: an R package for spatial thinning of species occurrence records for use in ecological niche models**. *Ecography* 38: 541–545. doi: 10.1111/ecog.01132.

Almazroui M, Saeed F, Saeed S, Islam MN, Ismail M, Klutse NAB, Siddiqui MH (2020) **Projected change in temperature and precipitation over Africa from CMIP6**. *Earth Systems and Environment* 4: 455–475. doi: 10.1007/s41748-020-00161-x.

Ampomah IG, Malau-Aduli BS, Seidu A-A, Malau-Aduli AEO, Emeto TI (2023) **Integrating traditional medicine into the Ghanaian health system: perceptions and experiences of traditional medicine practitioners in the Ashanti region**. *International Health* 15: 414–427. doi: 10.1093/inthealth/ihac059.

Applequist WL, Brinckmann JA, Cunningham AB, Hart RE, Heinrich M, Katerere DR, van Andel T (2019) **Scientists' warning on climate change and medicinal plants**. *Planta Medica* 86: 10–18. doi: 10.1055/a-1041-3406.

Araujo MB, Guisan A (2006) **Five (or so) challenges to species distribution modelling**. *Journal of Biogeography* 33: 1677–1688. doi: 10.1111/j.1365-2699.2006.01584.x.

Balima HL, Nacoulma BMI, Da SS, Ouédraogo A, Soro D, Thiombiano A (2022) **Impacts of climate change on the geographic distribution of African oak tree (*Afzelia africana* Sm.) in Burkina Faso, West Africa.** *Heliyon* 8: e08688. doi: 10.1016/j.heliyon.2021.e08688.

Bede-Fazekas A, Somodi I (2020) **The way bioclimatic variables are calculated has impact on potential distribution models**. *Methods in Ecology and Evolution* 11: 1559–1570. doi: 10.1111/2041-210X.13488.

Bivand R (2022) **R Packages for analyzing spatial data: A comparative case study with Areal data**. *Geographical Analysis* 54: 488–518. doi: 10.1111/gean.12319.

Bongers F, Groenendijk P, Bekele T, Birhane E, Damtew A, Decuyper M, Eshete A, Gezahgne A, Girma A, Khamis MA, Lemenih M, Mengistu T, Ogbazghi W, Sass-Klaassen U, Tadesse W, Teshome M, Tolera M, Sterck FJ, Zuidema PA (2019) **Frankincense in peril**. *Nature Sustainability* 2: 602–610.

doi: 10.1038/s41893-019-0322-2.

Booth TH, Nix HA, Busby JR, Hutchinson MF (2014) **BIOCLIM: The first species distribution modelling package, its early applications and relevance to most current MAXENT studies**. *Diversity and Distributions* 20: 1–9. doi: 10.1111/ddi.12144.

Boucher O, Servonnat J, Albright AL, Aumont O, Balkanski Y, Bastrikov V, Bekki S, Bonnet R, Bony S, Bopp L, Braconnot P, Brockmann P, Cadule P, Caubel A, Cheruy F, Codron F, Cozic A, Cugnet D, D'Andrea F, Davini P, de Lavergne C, Denvil S, Deshayes J, Devilliers M, Ducharne A, Dufresne J-L, Dupont E, Éthé C, Fairhead L, Falletti L, Flavoni S, Foujols M-A, Gardoll S, Gastineau G, Ghattas J, Grandpeix J-Y, Guenet B, Guez LE, Guilyardi E, Guimberteau M, Hauglustaine D, Hourdin F, Idelkadi A, Joussaume S, Kageyama M, Khodri M, Krinner G, Lebas N, Levavasseur G, Lévy C, Li L, Lott F, Lurton T, Luyssaert S, Madec G, Madeleine J-B, Maignan F, Marchand M, Marti O, Mellul L, Meurdesoif Y, Mignot J, Musat I, Ottlé C, Peylin P, Planton Y, Polcher J, Rio C, Rochetin N, Rousset C, Sepulchre P, Sima A, Swingedouw D, Thiéblemont R, Traore AK, Vancoppenolle M, Vial J, Vialard J, Viovy N, Vuichard N (2020) **Presentation and evaluation of the IPSL-CM6A-LR climate model**. *Journal of Advances in Modeling Earth Systems* 12: e2019MS002010. doi: 10.1029/2019MS002010.

Carvalho SB, Brito JC, Crespo EG, Watts ME, Possingham HP (2011) **Conservation planning under climate change: toward accounting for uncertainty in predicted species distributions to increase confidence in conservation investments in space and time**. *Biological Conservation* 144: 2020–2030. doi: 10.1016/j.biocon.2011.04.024.

CBD (2024) **Strategic plan for biodiversity 2011-2020, including Aichi biodiversity targets.** [https://www.cbd.int/sp/targets] Accessed 7 August 2024.

Dauby G, Zaiss R, Blach-Overgaard A, Catarino L, Damen T, Deblauwe V, Dessein S, Dransfield J, Droissart V, Duarte MC, Engledow H, Fadeur G, Figueira R, Gereau RE, Hardy OJ, Harris DJ, de Heij J, Janssens SB, Klomberg Y, Ley AC, Mackinder BA, Meerts P, van de Poel JL, Sonké B, Sosef MS, Stévart T, Stoffelen P, Svenning JC, Sepulchre P, van der Burgt X, Wieringa JJ, Couvreur TLP (2016) **RAINBIO: a mega-database of tropical African vascular plants distributions**. *PhytoKeys* 74: 1–18. doi: 10.3897/phytokeys.74.9723.

De Cauwer V, Muys B, Revermann R, Trabucco A (2014) **Potential, realized, future distribution and environmental suitability for *Pterocarpus angolensis* DC in southern Africa**. *Forest Ecology and Management* 315: 211–226. doi: 10.1016/j.foreco.2013.12.032.

Dlamini TS, Dlamini GM (2002) **Swaziland**. In: Golding JS (ed) *Southern African plant red data lists*. Southern African Botanical Diversity Network Report No. 14, SABONET, Pretoria, pp. 121–134.

Dludlu MN, Dlamini PS, Sibandze GS, Vilane VS, Dlamini CS (2017) **The distribution and conservation status of the endangered pepperbark tree** *Warburgia salutaris (Canallaceae) in Swaziland*. *Oryx* 51: 441–454.

Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B, Lafourcade B, Leitão PJ, Münkemüller T, McClean C, Osborne PE, Reineking B, Schröder B, Skidmore AK, Zurell D, Lautenbach S (2013) **Collinearity: A review of methods to deal with it and a simulation study evaluating their performance.** *Ecography* 36: 27–46. doi: 10.1111/j.1600-0587.2012.07348.x.

Dosio A, Jury MW, Almazroui M, Ashfaq M, Diallo I, Engelbrecht FA, Klutse NAB, Lennard C, Pinto I, Sylla MB, Tamoffo AT (2021) **Projected future daily characteristics of African precipitation based on global (CMIP5, CMIP6) and regional (CORDEX, CORDEX-CORE) climate models.** *Climate Dynamics* 57: 3135–3158. doi: 10.1007/s00382-021-05859-w.

Dyderski KM, Paz S, Frelich EL, Jagodzinski MA (2018) **How much does climate change threaten European forest tree species distributions?** *Global Change Biology* 24: 1150–1163. doi: 10.1111/gcb.13925.

Elsen PR, Monahan WB, Dougherty R, Merenlender AM (2020) **Keeping pace with climate change in global terrestrial protected areas**. *Science Advances* 6: eaay0814. doi:

Esser LF, Weldemariam EC (2023) **Climate changes could jeopardize a main source of livelihood in Africa's drylands.** *Journal of Nature Conservation* 71: 126319. doi: 10.1016/j.jnc.2022.126319.

Estrada A, Barbosa AM, Real R (2018) **Changes in potential mammal diversity in national parks and their implications for conservation**. *Current Zoology* 64: 671–679. doi: 10.1093/cz/zoy001.

Fick SE, Hijmans RJ (2017) **WorldClim 2: New 1 km spatial resolution climate surfaces for global land areas.** *International Journal of Climatology* 37: 4302–4315. doi: 10.1002/joc.5086.

Fourcade Y, Engler JO, Rodder D, Secondi J (2014)

Mapping species distributions with MAXENT using a geographically biased sample of presence data: A performance assessment of methods for correcting sampling bias. *PLoS ONE* 9: e97122. doi: 10.1371/journal.pone.0097122.

GBIF.org (2024) **GBIF Occurrence Download.** doi: 10.15468/dl.pduaes Accessed 15 July 2024.

Gidden MJ, Riahi K, Smith SJ, Fujimori S, Luderer G, Kriegler E, van Vuuren DP, van den Berg M, Feng L, Klein D, Calvin K, Doelman JC, Frank S, Fricko O, Harmsen M, Hasegawa T, Havlik P, Hilaire J, Hoesly R, Horing J, Popp A, Stehfest E, Takahashi K (2019) **Global emissions pathways under different socioeconomic scenarios for use in CMIP6: A dataset of harmonized emissions trajectories through the end of the century.** *Geoscientific Model Development* 12: 1443–1475. doi: 10.5194/gmd-12-1443-2019.

Glennon KL, Le Roux JJ, Thompson DI (2023) **Genetic insights into pepper bark tree (*Warburgia salutaris*) reproduction in South Africa**. *Conservation Genetics* 24: 883–891. doi: 10.1007/s10592-023-01531-4.

Goberville E, Beaugrand G, Hautekeete N.-C, Piquot Y, Luczak C (2015) **Uncertainties in the projection of species distributions related to general circulation models.** *Ecology and Evolution* 5: 1100–1116. doi: 10.1002/ ece3.1411.

Guégan J.-F, de Thoisy B, Gomez-Gallego M, Jactel H (2023) **World forests, global change, and emerging pests and pathogens.** *Current Opinion in Environmental Sustainability* 61: 101266. doi: 10.1016/j.cosust.2023.101266.

Guisan A, Thuiller W (2005) **Predicting species distribution: offring more tha simple habitat models.** *Ecology Letters* 8: 993–1009. doi: 10.1111/j.1461-0248.2005.00792.x.

Hama AA, Khwarahm NR (2023) **Predictive mapping of two endemic oak tree species under climate change scenarios in a semiarid region: Range overlap and implications for conservation.** *Ecological Informatics* 73: 101930. doi: 10.1016/j.ecoinf.2022.101930.

Hannweg K, Sippel A, Hofmeyr M, Swemmer L, Froneman W (2016) **Strategies for the conservation of** *Warburgia salutaris (family: Canellaceae), a red data list species - development of propagation methods. Acta Horticulturae* 1125: 33–40. doi: 10.17660/ActaHortic.2016.1125.4.

Hao T, Elith J, Guillera-Arroita G, Lahoz-Monfort JJ (2019) **A review of evidence about use and performance of species distribution modelling ensembles like BIOMOD.** *Diversity and Distributions* 25: 839–852. doi: 10.1111/ddi.12892.

Hijmans RJ (2012) **Cross-validation of species distribution models: removing spatial sorting bias and calibration with a null model.** *Ecology* 93: 679–688. doi: 10.1890/11-0826.1.

Hole DG, Willis SG, Pain DJ, Fishpool LD, Butchart SHM, Collingham YC, Rahbek C, Huntley B (2009) **Projected impacts of climate change on a continent-wide protected area network.** *Ecology Letters* 12: 420-431. doi: 10.1111/j.1461-0248.2009.01297.x.

Hollmann J, Van der Schijff M (1996) **Portrait of a medicinal tree: Has the publicity of being one of the trees of the year for 1996 helped conserve the pepper bark tree?** *Veld and Flora* 82: 115–116.

Hulme M, Doherty R, Ngara T, New M, Lister D (2001) **African climate change: 1900–2100.** *Climate Research* 17: 145–168. doi: 10.3354/cr017145.

IUCN (2024) **The IUCN Red List of Threatened Species. Version 2024-1.** [https://www.iucnredlist.org] Accessed 13 July 2024.

Izidine S, Bandeira SO (2002) **Mozambique.** In: Golding JS (ed) Southern African plant red data lists. Southern African Botanical Diversity Network Report Series No. 14. SABONET, Pretoria, pp. 43–53.

Jarnevich CS, Stohlgren TJ, Kumar S, Morisette JT, Holcombe TR (2015) **Caveats for correlative species distribution modeling.** *Ecological Informatics* 29: 6–15. doi: 10.1016/j.ecoinf.2015.06.007.

Jinga P, Palagi J, Chong JP, Bobo ED (2020) **Climate change reduces the natural range of African wild loquat (*Uapaca kirkiana* Müll. Arg., Phyllanthaceae) in south-central Africa**. *Regional Environmental Change* 20: 108. doi: 10.1007/s1011 3-020-01700-y.

Kafle S, Thapa D, Ghimire K, Gautam J (2023) **Ensemble modeling of *Rhododendron arboreum* distribution in Nepal: Assessing current patterns and projecting future changes**. *Species* 24: e81s1590. doi:

Khanum R, Mumtaz AS, Kumar S (2013) **Predicting impacts of climate change on medicinal asclepiads of Pakistan using Maxent modeling.** *Acta Oecologica* 49: 23–31. doi: 10.1016/j.actao.2013.02.007.

Khumalo GP, Loa-Kum-Cheung W, Van Wyk BE, Feng Y, Cock IE (2024) **Leaf extracts of eight selected southern African medicinal plants modulate pro-inflammatory cytokine secretion in LPS-stimulated RAW**

264.7 macrophages. *Inflammopharmacology* 32: 1607–1620. doi: 10.1007/s10787-023-01420-9.

Khumalo GP, Sadgrove NJ, Van Vuuren S, Van Wyk BE (2019) **Antimicrobial activity of volatile and nonvolatile isolated compounds and extracts from the bark and leaves of *Warburgia salutaris* (Canellaceae) against skin and respiratory pathogens**. *South African Journal of Botany* 122: 547–550. doi: 10.1016/j.sajb.2018.10.018.

Kotina EL, Van Wyk BE, Tilney PM (2014) **Anatomy of the leaf and bark of** *Warburgia salutaris (Canellaceae), an important medicinal plant from South Africa*. *South African Journal of Botany* 94: 177–181. doi: 10.1016/j.sajb.2014.06.008.

Langdon JGR, Lawler JJ (2015) **Assessing the impacts of projected climate change on biodiversity in the protected areas of western North America.** *Ecosphere* 6: 87. doi: 10.1890/ES14-00400.1.

Leberger R, Rosa IMD, Guerra CA, Wolf F, Pereira HM (2020) **Global patterns of forest loss across IUCN categories of protected areas.** *Biological Conservation* 241: 108299. doi: 10.1016/j.biocon.2019.108299.

Lemenih M, Arts B, Wiersum K, Bongers F (2014) **Modelling the future of *Boswellia papyrifera* population and its frankincense production**. *Journal of Arid Environments* 105: 33–40. doi: 10.1016/j.jaridenv.2014.02.006.

Leonard C, Chen W, Kamatou G (2023) ***Warburgia salutaris***. In: Viljoen A, Sandasi M, Fouche G, Combrinck S, Vermaak I (eds) *The South African Herbal Pharmacopoeia*. Academic Press, pp. 531–556. doi: 10.1016/B978-0-323-99794-2.00010-6.

Leonard CM, Viljoen AM (2015) **Warburgia: a comprehensive review of the botany, traditional uses and phytochemistry**. *Journal of Ethnopharmacology* 165: 260–285. doi: 10.1016/j.jep.2015.02.021.

Lyam PT, Duque-Lazo J, Hauenschild F, Schnitzler J, Muellner-Riehl AN, Greve M, Ndangalasi H, Myburgh A, Durka W (2022) **Climate change will disproportionally affect the most genetically diverse lineages of a widespread African tree species.** *Scientific Reports* 12: 7035. doi: 10.1038/s4159 8-022-11182-z.

Marmion M, Parviainen M, Luoto M, Heikkinen RK, Thuiller W (2009) **Evaluation of consensus methods in predictive species distribution modelling.** *Diversity and Distributions* 15: 59–69. doi: 10.1111/j.1472-4642.2008.00491.x.

Maroyi A (2008) **Ethnobotanical study of two threatened medicinal plants in Zimbabwe.** *International Journal of Biodiversity Science and Management* 4: 148–153. doi: 10.3843/Biodiv.4.3:2.

Maroyi A (2011) **Community attitudes towards the reintroduction programme for the endangered pepperbark tree *Warburgia salutaris*: implications for plant conservation in south-east Zimbabwe**. *Oryx* 46: 213–218. doi: 10.1017/S0030605311000767.

Maroyi A (2013) ***Warburgia salutaris* (Bertol. f.) Chiov.: A multi-use ethnomedicinal plant species**. *Journal of Medicinal Plants Research* 7: 53–60.

Mashamba TG, Adeosun IJ, Baloyi IT, Tshikalange ET, Cosa S (2022) **Quorum sensing modulation and inhibition in biofilm forming foot ulcer pathogens by selected medicinal plants**. *Heliyon* 8: e09303. doi: 10.1016/j.heliyon.2022.e09303.

Meddows-Taylor S, Ramadwa TM (2025) **A comprehensive review of the traditional uses, pharmacological activity and phytochemistry of *Warburgia salutaris* in southern Africa**. *South African Journal of Botany* 179: 134–146. doi: 10.1016/j.sajb.2025.02.016.

Midgley G, Bond W (2015) **Future of African terrestrial biodiversity and ecosystems under anthropogenic climate change.** *Nature Climate Change* 5: 823–829. doi: 10.1038/nclimate2753.

Moonlight PW, Silva de Miranda PL, Cardoso D, Dexter KG, Oliveira-Fihlo T, Pennington RT, Ramos G, Sarkinen TE (2020) **The strengths and weaknesses of species distribution models in biome delimitation.** *Global Ecology Biogeogreophy* 29: 1770–1784. doi: 10.1111/geb.13149.

Msekandiana G, Mlangeni E (2002) **Malawi.** In: Golding JS (ed) *Southern African plant red data lists.* Southern African Botanical Diversity Network Report No. 14. SABONET, Pretoria, pp. 135–156.

Naimi B, Hamm NAS, Groen TA, Skidmore AK, Toxopeus AG (2014) **Where is positional uncertainty a problem for species distribution modelling?** *Ecography* 37: 191–203. doi: 10.1111/j.1600-0587.2013.00205.x.

Nuttall MN, Griffin O, Fewster RM, McGowan PJK, Abernethy K, O'Kelly H, Nut M, Sot V, Bunnefeld N (2022) **Long-term monitoring of wildlife populations for protected area management in Southeast Asia**. *Conservation Science and Practice* 4: e614. doi: 10.1111/csp2.614.

Pyšek P, Hulme PE, Simberloff D, Bacher S, Black-

burn TM, Carlton JT, Dawson W, Essl F, Foxcroft LC, Genovesi P, Jeschke JM, Kühn I, Liebhold AM, Mandrak NE, Meyerson LA, Pauchard A, Pergl J, Roy HE, Seebens H, van Kleunen M, Vilà M, Wingfield MJ, Richardson DM (2020) **Scientists' warning on invasive alien species.** *Biological Reviews* 95: 1511-1534. doi: 10.1111/brv.12627.

R Core Team (2024) **R: A language and environment for statistical computing. R Foundation for Statistical Computing**, Vienna, Austria. https://www.R-project.org.

Rabe T, van Staden J (2000) **Isolation of an antibacterial sesquiterpenoid from *Warburgia salutaris*.** *Journal of Ethnopharmacology* 73: 171–174. doi: 10.1016/S0378-8741(00)00293-2.

Rathore MK, Sharma LK (2023) **Efficacy of species distribution models (SDMs) for ecological realms to ascertain biological conservation and practices**. *Biodiversity and Conservation* 32: 3053–3087. doi: 10.1007/s10531-023-02648-1.

Regos A, Arenas-Castro S, Tapia L, Domínguez J, Honrado JP (2021) **Using remotely sensed indicators of primary productivity to improve prioritization of conservation areas for top predators.** *Ecological Indicators* 125: 107503. doi: 10.1016/j.ecolind.2021.107503.

Riahi K, van Vuuren DP, Kriegler E, Edmonds J, O'Neill BC, Fujimori S, Bauer N, Calvin K, Dellink R, Fricko O, Lutz W, Popp A, Cuaresma JC, KC S, Leimbach M, Jiang L, Kram T, Rao S, Emmerling J, Ebi K, Hasegawa T, Havlik P, Humpenöder F, Da Silva LA, Smith S, Stehfest E, Bosetti V, Eom J, Gernaat D, Masui T, Rogelj J, Strefler J, Drouet L, Krey V, Luderer G, Harmsen M, Takahashi K, Baumstark L, Doelman JC, Kainuma M, Klimont Z, Marangoni G, Lotze-Campen H, Obersteiner M, Tabeau A, Tavoni M (2017) **The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: An overview.** *Global Environmental Change* 42: 153–168. doi: 10.1016/j.gloenvcha.2016.05.009.

Roberts DR, Bahn V, Ciuti S, Boyce MS, Elith J, Guillera-Arroita G, Hauenstein S, Lahoz-Monfort JJ, Schröder B, Thuiller W, Warton DI, Wintle BA, Hartig F, Dormann CF (2017) **Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure**. *Ecography* 40: 913–929. doi: 10.1111/ecog.02881.

Rodríguez MA, Angueyra A, Cleef AM, van Andel TR (2018) **Ethnobotany of the Sierra Nevada del Cocuy-Güicán: climate change and conservation strategies in the Colombian Andes.** *Journal of Ethnobiology and Ethnomedicine* 14: 34. doi: 10.1186/s13002-018-0227-6.

Rogelj J, Meinshausen M, Knutti R (2012) **Global warming under old and new scenarios using IPCC climate sensitivity range estimates.** *Nature Climate Change* 2: 248–253. doi: 10.1038/nclimate1385.

Santini L, Benítez-López A, Maiorano L, Čengić M, Huijbregts MAJ (2021) **Assessing the reliability of species distribution projections in climate change research.** *Diversity and Distributions* 27: 1035–1050. doi: 10.1111/ddi.13252.

Schmidt IB, Ticktin T (2012) **When lessons from population models and local ecological knowledge coincide—Effects of flower stalk harvesting in the Brazilian savanna.** *Biological Conservation* 152: 187–195. doi: 10.1016/j.biocon.2012.03.018.

Senkoro AM, Munt DD, Shackleton CM, Ribeiro-Barros AI, Voeks RA (2024) **The case of a threatened medicinal tree with optimistic prospects under climate change.** *Global Ecology and Conservation* 54: e03126. doi: 10.1016/j.gecco.2024.e03126.

Senkoro AM, Shackleton CM, Voeks RA, Ribeiro AI (2019) **Uses, knowledge, and management of the threatened pepper-bark tree (*Warburgia salutaris*) in southern Mozambique**. *Economic Botany* 73: 304–324. doi: 10.1007/s12231-019-09468-x.

Senkoro AM, Talhinhas P, Simoes F, Batista-Santos P, Shackleton CM, Voeks RA, Marques I, Ribeiro-Barros AI (2020) **The genetic legacy of fragmentation and overexploitation in the threatened medicinal African pepper bark tree, *Warburgia salutaris***. *Scientific Reports* 10: 19725. doi: 10.1038/s41598-020-76654-6.

Sillero N, Arenas-Castro S, Enriquez-Urzelai U, Vale CG, Sousa-Guedes D, Martinez-Freiria F, Real R, Barbosa AM (2021) **Want to model a species niche? A step-by-step guideline on correlative ecological niche modelling.** *Ecological Modelling* 456: 109671. doi: 10.1016/j.ecolmodel.2021.109671.

Sinclair SJ, White MD, Newell GR (2010) **How useful are species distribution models for managing biodiversity under future climates?** *Ecology and Society* 15: 8.

Sintayehu DW (2018) **Impact of climate change on biodiversity and associated key ecosystem services in Africa: a systematic review.** *Ecosystem Health and Sustainability* 4: 1530054. doi: 10.1080/20964129.2018.1530054.

Soley-Guardia M, Alvarado-Serrano DF, Anderson RP (2024) **Top ten hazards to avoid when model-**

ing species distributions: a didactic guide of assumptions, problems, and recommendations. *Ecography* 2024: e06852. doi: 10.1111/ecog.06852.

Ssenku JE, Okurut SA, Namuli A, Kudamba A, Tagume P, Matovu P, Wasige G, Kafeero HM, Walusansa A (2022) **Medicinal plant use, conservation, and the associated traditional knowledge in rural communities in Eastern Uganda.** *Tropical Medicine and Health* 50: 39. doi: 10.1186/s41182-022-00428-1.

Steen V, Sofaer HR, Skagen SK, Ray AJ, Noon BR (2017) **Projecting species' vulnerability to climate change: Which uncertainty sources matter most and extrapolate best?** *Ecology and Evolution* 7: 8841–8851. doi: 10.1002/ece3.3403.

Steenkamp V Nkwane O, Van Tonder J, Dinsmore A, Gulumian M (2013) **Evaluation of the phenolic and flavonoid contents and radical scavenging activity of three southern African medicinal plants**. *African Journal of Pharmacy and Pharmacology* 7: 703–709.

Swart NC, Cole JNS, Kharin VV, Lazare M, Scinocca JF, Gillett NP, Anstey J, Arora V, Christian JR, Hanna S, Jiao Y, Lee WG, Majaess F, Saenko OA, Seiler C, Seinen C, Shao A, Sigmond M, Solheim L, von Salzen K, Yang D, Winter B (2019) **The Canadian earth system model version 5 (CanESM5.0.3).** *Geoscientific Model Development* 12: 4823–4873. doi: 10.5194/gmd-12-4823-2019.

Sweet LC, Green T, Heintz JGC, Frakes N, Graver N, Rangitsch JS, Rodgers JE, Heacox S, Barrows CW (2019) **Congruence between future distribution models and empirical data for an iconic species at Joshua Tree National Park.** *Ecosphere* 10: e02763. doi: 10.1002/ecs2.2763.

Tatebe H, Ogura T, Nitta T, Komuro Y, Ogochi K, Takemura T, Sudo K, Sekiguchi M, Abe M, Saito F, Chikira M, Watanabe S, Mori M, Hirota N, Kawatani Y, Mochizuki T, Yoshimura K, Takata K, O'ishi R, Yamazaki D, Suzuki T, Kurogi M, Kataoka T, Watanabe M, Kimoto M (2019) **Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6.** *Geoscientific Model Development* 12: 2727–2765. doi: 10.5194/gmd-12-2727-2019.

Terer T, Muasya AM, Dahdouh-Guebas F, Ndiritu GG, Lugwig-Triestr L (2012) **Integrating local ecological knowledge and management practices of an isolated semi-arid papyrus swamp (Loboi, Kenya) into a wider conservation framework.** *Journal of Environmental Management* 93: 71–84. doi: 10.1016/j.jenvman.2011.08.005.

Thuiller W (2003) **BIOMOD—Optimizing predictions of species distributions and projecting potential future shifts under global change.** *Global Change Biology* 9: 1353–1362. doi: 10.1046/j.1365-2486.2003.00666.x.

Thuiller W, Georges D, Gueguen M, Engler R, Breiner F, Lafourcade B, Patin R (2023) **Biomod2: Ensemble platform for species distribution modeling. R Package Version 4.2-4.** `https://cran.r-project.org/package=biomod2`.

Thuiller W, Lafourcade B, Engler R, Araújo MB (2009) **BIOMOD—A platform for ensemble forecasting of species distributions.** *Ecography* 32: 369–373. doi: 10.1111/j.1600-0587.2008.05742.x.

Timbal B (2004) **Southwest Australia past and future rainfall trends.** *Climate Research* 26: 233–249. doi: 10.3354/cr026233.

Title PO, Bemmels JB (2018) **ENVIREM: An expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling**. *Ecography* 41: 291–307. doi: 10.1111/ecog.02880.

UNEP-WCMC, IUCN (2024) **Protected Planet: The World Database on Protected Areas (WDPA) and World Database on Other Effective Area-based Conservation Measures (WDOECM), Cambridge, UK, UNEP-WCMC and IUCN.** [`www.protectedplanet.net`] Accessed 10 July 2024.

Valavi R, Elith J, Lahoz-Monfort JJ, Guillera-Arroita G (2019) **blockCV: An R package for generating spatially or environmentally separated folds for k-fold cross-validation of species distribution models.** *Methods in Ecology and Evolution* 10: 225–232. doi: 10.1111/2041-210X.13107.

van der Bosch K, Witkowski ETF, Thompson DI, Cron GV (2023) **Reproductive ecology offers some answers to the pepperbark tree persistence puzzle in the Kruger National Park, South Africa.** *Global Ecology and Conservation* 41: e02330. doi: 10.1016/j.gecco.2022.e02330.

Varela D, Romeiras MM, Silva L (2022) **Implications of climate change on the distribution and conservation of Cabo Verde endemic trees**. *Global Ecology and Conservation* 34: e02025. doi: 10.1016/j.gecco.2022.e02025.

Yates LA, Aandahl Z, Richards SA, Brook BW (2023) **Cross validation for model selection: A review with examples from ecology**. *Ecological Monographs* 93: e1557. doi: 10.1002/ecm.1557.

Zebaze S, Anand A, Fotso-Nguemo TC, Taguela TN,

Ngavom Z, Mbienda AJK, Fotso-Kamga G, Choumbou PC, Vondou DA (2025) **Assessing the performance of the CMIP6 multi model mean in simulating precipitation and temperature across Africa**. *Modeling Earth Systems and Environment* 11: 374. doi: 10.1007/s40808-025-02560-3.

Zurell D, Franklin J, König C, Bouchet PJ, Dormann CF, Elith J, Fandos G, Feng X, Guillera-Arroita G, Guisan A, Lahoz-Monfort JJ, Leitão PJ, Park DS, Peterson AT, Rapacciuolo G, Schmatz DR, Schröder B, Serra-Diaz JM, Thuiller W, Yates KL, Zimmermann NE, Merow C (2020) **A standard protocol for reporting species distribution models.** *Ecography* 43: 1261–1277. doi: 10.1111/ecog.04960.

Received: 12 April 2025
Accepted: 02 June 2026
Published: 10 June 2026

Editor: Thiago Gonçalves-Souza

Additional Files


Add File 1. Mean evaluation scores of SDMs combined into an ensemble SDM used to predict the suitable habitat of Warburgia salutaris in southern Africa.

Algorithm	Validation TSS	Validation ROC
ANN	0.642	0.831
CTA	0.489	0.744
FDA	0.535	0.789
GBM	0.626	0.877
GLM	0.663	0.875
MARS	0.654	0.867
RF	0.009	0.892
SRE	0.624	0.812

Add File 2. Importance scores of variables used in an ensemble species distribution model to determine the current suitable habitat of Warburgia salutaris (Pepper-bark tree) in southern Africa.

Variable	**Permutation Score**			**Mean Score**
	1	2	3	
BIO2	0.122	0.118	0.127	0.122
BIO3	0.032	0.033	0.030	0.032
BIO8	0.085	0.083	0.085	0.084
BIO14	0.120	0.121	0.121	0.121
BIO18	0.155	0.152	0.153	0.153
Elevation	0.187	0.180	0.189	0.185

BIO2 = Mean diurnal range, BIO3 = Isothermality, BIO8 = Mean temperature of wettest quarter, BIO14 = Precipitation of driest month, BIO18 = Precipitation of warmest quarter



Add File 3. Response curves for variables used in an ensemble species distribution model to determine the current and future suitable habitat of Warburgia salutaris (Pepper-bark tree).

Bio_2 = Mean diurnal range, Bio_3 = Isothermality, Bio_8 = Mean temperature of wettest quarter, Bio_14 = $Precipitation of driest month$, $Bio_18 = Precipitation of warmest quarter$, $ele = Elevation$