

Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete chloroplast genome sequence of *Huperzia javanica* (sw.) C. Y. Yang in Lycopodiaceae

Hong-rui Zhang, Jong-Soo Kang, Ronald L. L. Viane & Xian-Chun Zhang

To cite this article: Hong-rui Zhang, Jong-Soo Kang, Ronald L. L. Viane & Xian-Chun Zhang (2017) The complete chloroplast genome sequence of *Huperzia javanica* (sw.) C. Y. Yang in Lycopodiaceae, Mitochondrial DNA Part B, 2:1, 216-218, DOI: [10.1080/23802359.2017.1310603](https://doi.org/10.1080/23802359.2017.1310603)

To link to this article: <https://doi.org/10.1080/23802359.2017.1310603>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 09 Apr 2017.



Submit your article to this journal [↗](#)



Article views: 734



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

MITOGENOME ANNOUNCEMENT



The complete chloroplast genome sequence of *Huperzia javanica* (sw.) C. Y. Yang in Lycopodiaceae

Hong-rui Zhang^{a,b*}, Jong-Soo Kang^{a,b*}, Ronald L. L. Viane^c and Xian-Chun Zhang^a

^aState Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing, China; ^bUniversity of Chinese Academy of Sciences, Beijing, China; ^cResearch Group Pteridology, Department of Biology, Ghent University, Gent, Belgium

ABSTRACT

Huperzia javanica (Sw.) C. Y. Yang is a valuable medical herb used for treating Alzheimer's disease. Here, we described the complete chloroplast genome of *H. javanica* using Illumina paired-end sequencing. The total genome length is 154,415 bp, containing 119 unique genes, with 86 protein-coding genes, 29 tRNA genes, and 4 rRNA genes. The gene content and their order are consistent with two previously reported *Huperzia* genomes. The overall GC content of the chloroplast genome of *H. javanica* is 36.4%. The topology of our maximum-likelihood tree is consistent with topologies found in previous studies, with *H. javanica* sister to a clade of *H. serrata* and *H. lucidula*. We support the recognition of *H. javanica* as an independent species. *Huperzia serrata* is more closely related to *H. lucidula* than to *H. javanica*.

ARTICLE HISTORY

Received 6 March 2017
Accepted 21 March 2017

KEYWORDS

Huperzia javanica; *Huperzia*; Lycopodiaceae; chloroplast genome

Huperzia javanica C. Y. Yang (Lycopodiaceae) is a perennial, evergreen, South East Asian herb found in South China, India, Japan, South Korea, and the Philippines (Sun 2015). It grows on shady places in humid forests and valleys at an elevation ranging from 300–1200 m (Shrestha & Zhang 2015; Sun 2015). *Huperzia* species are known as valuable medicinal herbs. Especially, *H. javanica* and *H. serrata* are used for the treatment of Alzheimer's disease (Tang 1996; Wang et al. 1998; Yang et al. 2008). Recently, *H. javanica* was distinguished as independent species from *H. serrata* complex through a previous study using morphological and climatic data (Shrestha & Zhang 2015), but we could not distinguish these species on the dried condition for making the oriental medicine based on only morphology. Chloroplast genomes are widely used in molecular phylogeny, to assess genetic diversity in conservation genetics, and in molecular identification such as DNA barcoding (Burke et al. 2012; Huang et al. 2014; Walker et al. 2014). In this study, we describe the chloroplast genome of *H. javanica* and reveal the phylogenetic relationship between *H. javanica* and *H. serrata*, thus contributing to the molecular identification and the conservation biology of *H. javanica*. The annotated sequence was deposited in GenBank under accession number KY609860.

Leaf material of *H. javanica* was collected in Shaoguan, Guangdong province in south China. The voucher specimen (*R. Wei CBL011*) has been deposited in the Herbarium of the Institute of Botany, Chinese Academy of Sciences (PE). Total DNA was extracted from silica gel dried leaves with a modified cetyltrimethylammonium bromide (CTAB) method

(Li et al. 2013). The NEBNext DNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA) was used for library construction. Paired-end reads of 2 × 150 bp then were generated using an Illumina HiSeq PE150 (Illumina, San Diego, CA). A total of 8,688,705 paired-end sequence reads of 150 bp were generated, of which 230,793 paired-ends reads belong to the chloroplast genome. The chloroplast genome data were extracted using *H. lucidula* (AY660566) as a reference and assembled *de novo* with Geneious v. R9.0.5 (Kearse et al. 2012). The initial annotation of *H. javanica* chloroplast genome was performed using Dual Organellar GenoMe Annotator (DOGMA; Wyman et al. 2004). After initial annotation, the putative starts, stops, and intron positions were determined by comparison with homologous genes in previously reported *Huperzia* chloroplast genomes. The tRNA genes were annotated using DOGMA and tRNAscan-SE (Schattner et al. 2005). The circular chloroplast genome map was drawn using the OGDRAW program (Lohse et al. 2013).

The complete size of the *H. javanica* chloroplast genome is 154,415 bp, which includes a pair of inverted repeat (IR) regions of 15,315 bp separated by a small single copy (SSC) region of 19,667 bp and a large single copy (LSC) region of 104,120 bp similar to the previously reported *Huperzia* chloroplast genomes (Wolf et al. 2005; Guo et al. 2016). The *H. javanica* chloroplast genome contains 119 genes, 11 of which are duplicated in the IR region, giving a total of 130 genes. The chloroplast genome of *H. javanica* contains 29 distinct tRNAs, four of which are duplicated in the IR region. 13 genes (*atpF*, *ndhA*, *ndhB*, *petB*, *petD*, *rps12*, *rps16*, *rpl2*,

CONTACT Xian-Chun Zhang ✉ zhangxc@ibcas.ac.cn State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing, China

*These authors have equally contributed to this study.

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Phylogenetic tree based on 72 protein-coding genes using the ML method. Taxon in bold is the new genome reported in this study. Bootstrap values are shown above the nodes.

rpl16, *rpoC1*, *trnA-UGC*, *trnL-UAA*, *trnV-UAC*) contain one intron, while two genes (*clpP*, *ycf3*) contain two introns. The *rps12* gene is divided in two independent transcription units. The overall GC content is 36.4%, in particular 34.5% in the LSC, 32.9% in the IR, and 45.0% in the SSC region.

Phylogenetic analysis based on 72 protein-coding genes was performed using extra species from Lycopodiaceae to Cyatheales in the PPG I system (PPG I 2016), and two Bryophyte species (*Anthoceros angustus* and *Marchantia polymorpha*) as the outgroup (Figure 1). A total of 55,034 bp was aligned using MAFFT (Kato et al. 2002). Maximum-likelihood (ML) analysis was performed using RAXML v. 7.4.2 with 1000 bootstrap replicates and the GTR+I+G model (Stamatakis 2006; Darriba et al. 2012). Our ML tree topology is consistent with topologies published in previous studies (Lu et al. 2015; PPG I 2016) (Figure 1). Although *H. javanica* is morphologically more similar to *H. serrata*, the phylogenetic relationship of *H. serrata* is closer to that of *H. lucidula* than to that of *H. javanica* (Figure 1). Moreover, this is confirmed by the sequence variation rates of the data matrix of 72 aligned protein-coding gene sequences of these three *Huperzia* species, which show that the sequence variation rate between *H. serrata* and *H. javanica* (0.411%) is slightly higher than that between *H. serrata* and *H. lucidula* (0.099%). These results are consistent with those of a previous study using morphological and climatic data, in which *H. javanica* was separated from *H. serrata* and treated as independent species (Shrestha & Zhang 2015).

Acknowledgments

The authors thank Ran Wei at the Institute of Botany, Chinese Academy of Sciences for help with collecting the sample.

Disclosure statement

The authors report no conflicts of interest, and are solely responsible for the content and writing of this paper.

Funding

This study was supported by the National Natural Science Foundation of China (31670205).

References

- Burke SV, Grennan CP, Duvall MR. 2012. Plastome sequences of two New World bamboos-*Arundinaria gigantea* and *Cryptochloa strictiflora* (Poaceae)-extend phylogenomic understanding of Bambusoideae. *Am J Bot.* 99:1951–1961.
- Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods.* 9:772.
- Guo ZY, Zhang HR, Shrestha N, Zhang XC. 2016. Complete chloroplast genome of a valuable medicinal plant, *Huperzia serrata* (Lycopodiaceae), and comparison with its congener. *Appl Plant Sci.* 4:1600071.
- Huang H, Shi C, Liu Y, Mao SY, Gao LZ. 2014. Thirteen *Camellia* chloroplast genome sequences determined by high-throughput sequencing: genome structure and phylogenetic relationships. *BMC Evol Biol.* 14:151.

- Katoh K, Misawa K, Kuma KI, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30:3059–3066.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics.* 28:1647–1649.
- Li J, Wang S, Yu J, Wang L, Zhou S. 2013. A modified CTAB protocol for plant DNA extraction. *Chin Bull Bot.* 48:72–78.
- Lohse M, Drechsel O, Kahlau S, Bock R. 2013. OrganellarGenomeDRAW—a suite of tools for generating physical maps of plastid and mitochondrial genomes and visualizing expression data sets. *Nucleic Acids Res.* 41:575–581.
- Lu JM, Zhang N, Du XY, Wen J, Li DZ. 2015. Chloroplast phylogenomics resolves key relationships in ferns. *J Syst Evol.* 53:448–457.
- PPG I. 2016. A community-derived classification for extant lycophytes and ferns. *J Syst Evol.* 54:563–603.
- Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res.* 33:W686–W689.
- Shrestha N, Zhang XC. 2015. Recircumscription of *Huperzia serrata* complex in China using morphological and climatic data. *J Syst Evol.* 53:88–103.
- Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics.* 22:2688–2690.
- Sun BY. 2015. Lycopodiaceae. In: Park CW, editors. *Flora of Korea*, vol. 1. Incheon, South Korea: National Institute of Biological Resources; p. 13–16.
- Tang XC. 1996. Huperzine A (shuangyiping): a promising drug for Alzheimer's disease. *Zhongguo Yao Li Xue Bao.* 17:481–484.
- Walker JF, Zanis MJ, Emery NC. 2014. Comparative analysis of complete chloroplast genome sequence and inversion variation in *Lasthenia burkei* (Madieae, Asteraceae). *Am J Bot.* 101:722–729.
- Wang L, Ji X, Wong Q, Yang J. 1998. Observations of 36 cases on Alzheimer's disease treatment using Huperzine A. *J Nantong Med Coll.* 18:486–488.
- Wolf PG, Karol KG, Mandoli DF, Kuehl J, Arumuganathan K, Ellis MW, Mishler BD, Kelch DG, Olmstead RG, Boore JL. 2005. The first complete chloroplast genome sequence of a lycophyte, *Huperzia lucidula* (Lycopodiaceae). *Gene.* 350:117–128.
- Wyman SK, Jansen RK, Boore JL. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics.* 20:3252–3255.
- Yang HJ, Huang H, Chen DJ, Zhang JC. 2008. Ultrasonic extraction and determination of Huperzine A from *H. Javanica* in Sanming region. *Appl Chem Ind.* 12:008.