



First Phytochemical Profiling and In-Vitro Antiprotozoal Activity of Essential Oil and Extract of *Plagiochila porelloides*

Anaïs Pannequin, Joëlle Quetin-Leclercq, Jean Costa, Aura Tintaru, Alain Muselli

► To cite this version:

Anaïs Pannequin, Joëlle Quetin-Leclercq, Jean Costa, Aura Tintaru, Alain Muselli. First Phytochemical Profiling and In-Vitro Antiprotozoal Activity of Essential Oil and Extract of *Plagiochila porelloides*. *Molecules*, 2023, 28 (2), pp.616. <10.3390/molecules28020616>. <hal-04524487>

HAL Id: hal-04524487

<https://hal.science/hal-04524487v1>

Submitted on 28 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

First phytochemical profiling and *in-vitro* antiprotozoal activity of essential oil and extract of *Plagiochila porelloides*

Anaïs Pannequin¹, Joëlle Quetin-Leclercq², Jean Costa¹, Aura Tintaru^{3*} and Alain Muselli^{1*}

¹ Université de Corse, UMR CNRS 6134, Laboratoire Chimie des Produits Naturels, BP 52, 20250 Corte, France; pannequin_a@univ-corse.fr (A.P.); costa_d@univ-corse.fr (J.C.); muselli_a@univ-corse.fr (A.M.)

² UCLouvain, Louvain Drug Research Institute, Pharmacognosy Group, Avenue E. Mounier 72, B-1200 Brussels, Belgium; joelle.leclercq@uclouvain.be (J.Q.-L.)

³ Aix Marseille Univ, CNRS, Centre Interdisciplinaire de Nanoscience de Marseille, UMR7325, 13288 Marseille, France; aura.tintaru@univ-amu.fr (A.T.)

* Correspondence: aura.tintaru@univ-amu.fr (A.T.); muselli_a@univ-corse.fr (A.M.)

Abstract: Volatiles metabolites from the liverwort *Plagiochila porelloides* harvested in Corsica were investigated by chromatographic and spectroscopic methods. In addition to already reported constituents, three new compounds were isolated by preparative chromatography and their structures were elucidated by mass spectrometry (MS) and NMR experiments. Hence, an atypical aliphatic compound, namely (3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol and two isomers, (E) and (Z), possessing an unusual humbertiane skeleton (called *p*-menth-1-en-3-[2-methylbut-1-enyl]-8-ol) are newly reported and fully characterized in this work.

The *in-vitro* antiprotozoal activity of essential oil and extract of *P. porelloides* against *Trypanosoma brucei brucei* and *Leishmania mexicana mexicana* and cytotoxicity were determined. Essential oil and Et₂O extract showed a moderate activity against *T. brucei* with IC₅₀ values: 2.03 and 5.18 µg/mL, respectively. It is noteworthy that only the essential oil showed a high selectivity (SI = 11.7). Diethyl oxide extract exhibited moderate anticancer (cancerous macrophage-like murine cells) activity and also cytotoxicity (human normal fibroblast) with IC₅₀ values: 1.25 and 2.96 µg/mL.

Keywords: Bryophytes, *Plagiochila porelloides*, humbertiane, farnesane, sesquiterpenes, antiprotozoal

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Molecules* **2022**, *27*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Lastname

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Bryophytes are largely located in various ecosystems characterized by humid climates, such as lakes, rivers, swamps, etc. where the water, an essential element for their development and sexual reproduction, is abundantly present. Considered as the oldest green plants, they are being the first vegetables adapted to terrestrial life 500 million years ago [1]. Furthermore, following their taxonomy, the bryophytes are classified among pteridophytes and algae, and could be sub-divided into three coordinate phyla: liverworts (Marchantiophyta or Hepaticae), mosses (Bryophyta) and hornworts (Anthocerotophyta). Nowadays, approximately 25 000 species of Bryophytes were identified and worldwide disseminated. Among those, 1 800 species are located in Europe and almost 75% were identified on the French territory [1].

In particular, Corsica has more than 500 species of Bryophytes spread over all the vegetation levels of the island [2]. Most of the Bryophytes live in fresh and humid places but they are also found in dry and open habitats. They are also present in running water, around streams and lakes, in marshes and bogs. In addition, these plants that contribute significantly to flora diversity and play an essential role in the functioning of many ecosystems (peat bogs, forests, etc...). Bryophytes constitute an important plant biomass available throughout the year, which is not economically exploited, yet. To our knowledge, the Bryophytes of Corsica have been the subject of only one phytochemical study carried out earlier by our group [3,4]. Nevertheless, even if Bryophytes are currently poorly investigated, based on the results reported by the few studies conducted on this topic [5–10], they are depicted as “the pharmacy of tomorrow”. For all these reasons, the characterization of the molecular constituents of essential oils, volatile fractions, and extracts produced from the Bryophytes, combined with their screening of active compounds, constitute an interesting scientific challenge to be performed at the scale of Corsican region.

Plagiochila species is the largest genus in the Marchantiophyta, with at least 1 600 varieties [11]. This genus is characterized not only by morphological diversity but also by a huge chemical variety. From the composition of solvent extracts of almost sixty species of *Plagiochila*, the author distinguished twelve types [11] as following I: 2,3-secoaromadendrane sesquiterpene-type (subdivided depending on the degree of oxidation and acetylation of 2,3-secoaromadendrane); II: bibenzyl-type; III: cuparane-herbertane sesquiterpene-type; IV: bibenzyl-cuparane-herbertane-type; V: gymnomitrane (barbatane)-bicyclogermacrane sesquiterpene-type; VI: bicyclogermacrane-spathulenol sesquiterpene-type; VII: pinguisane sesquiterpene-type; VIII: 2,3-secoaromadendrane-sesquiterpene lactone-type; IX: cyclic bis-bibenzyl-2,3-secoaromadendrane-type; X: sesquiterpene lactone-type; XI: epiverrucosane type and XII: fusicoccane-labdane type. According to two studies concerning compounds extracted by diethyl oxide, *P. porelloides* can be integrated to type I. The first report of *P. porelloides* compounds led to the isolation 3 α -acetoxybicyclogermacrene, bicyclogermacrene, plagiochilines A, -C, -D et -H as main constituents from Swiss species [[12]]. The second investigation reported on the isolation of three sesquiterpene esters from a German sample, derived from mono esterification of a 2,3-secoaromadendrane-type sesquiterpenoid by different fatty acids, in addition to spathulenol [13].

The volatiles constituents of the *Plagiochila* genus have been poorly studied. To our knowledge, only four studies have been carried out on the chemical composition of the essential oils prepared from six *Plagiochila* species: *P. biffaria* [14,15], *P. maderensis*, *P. retrorsa*, *P. stricta* [14], *P. asplenoides* [16] and *P. ovalifolia* [17] (see Table 1 and

Figure 1). The bibliographic data highlighted a chemical variability according to each studied species. However, the richness in hydrocarbon monoterpenes and oxygenated sesquiterpenes is the common point of all reported varieties. Moreover, all the investigated species constituted a source of new compounds. In *P. biffaria*, three eudesmane type sesquiterpenes, eudesm-4-en-6-one, eudesm-4(15)-en-6-one, 7-hydroxyeudesm-4-en-6-one were isolated and identified as new natural products [15]. In *P. asplenoides*, one aromadendrane sesquiterpene, aromadendra-1(10),3-diene, two aromatic sesquiterpene hydrocarbons, bisabola-1,3,5,7(14)-tetraene and bisabola-1,3,5,7-tetraene, three sesquiterpene oxides, muurolan-4,7-peroxide, plagiochilines W and X were described for the first time [16].




Figure 1. Structures of main components reported from the essential oils of *Plagiochila* species. (Numbers refers to table 1).

77

78

79

Table 1. Main compounds of *P. biffaria*, *P. maderensis*, *P. retrorsa*, *P. stricta*, *P. aspenoides* and *P. ovalifolia* [14],[15], [16],[17].

Species	Hydrocarbon terpenes	Oxygenated terpenes
<i>P. biffaria</i> [14]		(-)-5R,7R,10S-eudesm-4(15)-en-6-one (9-19%, 1)
		Methyl everminate (1-35%, 2)
		peculiaroxide (13-16%, 3)
<i>P. maderensis</i> [14]	terpinolene (34-60%, 4)	
<i>P. retrorsa</i> [14]	β -phellandrene (16-46%, 5)	peculiaroxide (9-12%, 3)
<i>P. retrorsa</i> [14]	<i>allo</i> -ocimene (15%, 6)	peculiaroxide (12%, 3)
	<i>neo-allo</i> -ocimene (10%, 7)	
	terpinolene (13%, 4)	
<i>P. stricta</i> [14]	<i>allo</i> -ocimene (7-19%, 7)	peculiaroxide (11-21%, 3)
	bicyclogermacrene (4-17%, 8)	Spathulenol (2-14%, 9)
	<i>neo-allo</i> -ocimene (4-11%, 7)	
<i>P. biffaria</i> [15]		ent-eudesm-4-en-6-one (10),
		ent-eudesm-4(15)-en-6-one (1)
		ent-7-hydroxyeudesm-4-en-6-one (11),
<i>P. aspenioides</i> [16]	(-)-selina-5,7(11)-diene (8%, 12)	plagio-4,7-peroxide (20%, 13)
		maalian-5-ol (19%, 14)
<i>P. ovalifolia</i> [17]		ent-4 β ,10 α 10 α -dihydroxyaromadendrane (15)
		Acetoxyisoplagiochilide (16)
		Maalian-5-ol (13)
		plagiochiline C (17)
		Plagiochiline N (18)

The aim of the present work was to investigate the volatile metabolites of *P. porelloides* prepared by hydro-distillation (essential oil and hydrosol), hexane and diethyl oxide cold-extractions and microwave assisted extractions using a combination of techniques involving liquid Column Chromatography, GC/FID (using retention indices), GC-MS (EI), HRMS and NMR spectroscopy (^1H -, ^{13}C - and 2D-NMR). As liverwort chemicals were generally very complex mixtures, our study deals firstly with the identification of components not recorded in MS-libraries; in addition, we are also reporting the isolation and structure determination of three unknown natural products showing a farnesane and humbertiane skeletons, respectively. Finally, the cytotoxicity and some anti-protozoal activities of the bulk extracts were investigated *in vitro*, to evaluate their potential pharmacological properties.

2. Results

In order to carry out a more exhaustive study of the volatile metabolites of *P. porelloides*, samples were prepared by multiple extraction procedures from dried plant material: essential oil and hydrosol were obtained by hydrodistillation, solvent extracts by cold maceration and assisted microwave extractions as well as, the volatiles were sampled using SPME. The identification of components involved a methodology first based on the comparison of RI and MS data with those contained in the in-house library or commercial libraries. After this preliminary

analysis, components matched by standards from the in-house library were considered as definitely identified while components matched only by commercial library database needed identification-confirmation. In the present work, several components remained unidentified. So, preparative liquid chromatography and additional NMR experiments were carried out to achieve an unambiguous compound identification, as well as the complete NMR assignment.

Our study allowed identifying 50 compounds representing 76.9 % of the essential oil (EO) and 52.6% of the hydrosol extract (HY), 82.6% and 77.9 % of hexane and diethyl oxide solvent extracts (EXT_H and EXT_O), 89.4 % of microwave extract (MW) and 90.3% volatile fraction (VF). Among them, the presence of three unknown sesquiterpenoids was revealed in the diethyl oxide extract and the essential oil of *P. porelloides*.

2.1. Resolution of ambiguous identifications of sesquiterpenes

Preparative liquid chromatography of *P. polleroiodes* essential oil was performed to obtained rich-sesquiterpene fractions. As GC-MS identification of sesquiterpenes in complex mixture can be a complex task [18], unambiguous identification of components **45**, **48**, **50**, **51**, **54** and **55** were definitively established using NMR-Extraction procedure [19]. Among these, the presence of the isomers spathulenol **48**, globulol **50** and viridiflorol **51** with close RI and mass spectra was confirmed by comparison of their ¹³C-NMR data with those described in the literature. The same procedure allowed the identification of 4-epi-maaliol **45**, rosifoliol **54**, maalian-5-ol **55** (Figure 2).




Figure 2. Structures of *P. porelloides* essential oil components identified by GC-MS and ensured by NMR. (Numbers refers to table 4).

The compounds **23**, **27** and **36** were concentrated in the hydrocarbon fraction of the essential oil and identified as aristolene **23**, β -barbatene **27** and bicyclogermacrene **36** (Figure 2).

2.2. Structural elucidations of new natural compounds

Column chromatography of *P. porelloides* EXT_O carried out using a gradient of polarity with hexane and diisopropyl oxide, was producing two fractions in which **58** (65%) was isolated from the polar fraction. EI mass spectra of **58** exhibited a base peak at m/z 107 and a signal at m/z 222, which could be attributed to the molecular ion. ESI (+)-HRMS measurements confirms the molecular formula C₁₅H₂₆O, (detected ion C₁₅H₂₆ONa⁺ (m/z)_{exp} 245.1879 and (m/z)_{th} 245.1876, error +1.2 ppm). The formula indicated a saturation degree of 3. The ¹H-NMR (CDCl₃, 300K) spectrum of **58** (Table 2), showed the presence of five methyl groups δ 1.3, 1.6, 1.7, 1.8 (s, H₁₃, H₁₅, H₁₂ and H₁₄, respectively) and δ 0.9 (t, J = 7.44 Hz, H₁) and four olefinic protons δ 5.1 (t, J = 6.0 Hz, H₁₀), δ 5.6 (d, J = 15.3 Hz, H₄), δ 5.9 (d, J = 10.8 Hz, H₆) and δ 6.5 (dd, J = 15.3, 10.8 Hz, H₅). The ¹³C-NMR spectrum contained 15 resonances: five methyls, three methylenes, four methines and three quaternary carbons. The occurrence of a quaternary carbon at 73.38 ppm suggested the presence of a tertiary alcohol. Spectral data were similar to those of nerolidol, which suggests a farnesane-type compound. The HSQC and HMBC correlations (Figure 3, (A)) and NOE correlations (Figure 3, (B)) indicated that the oxygenated carbon (C₃) was correlated with the methyl (C₁₃), the methylene group (C₂) and the methine (C₄). The signal between 5.5 and 6.6 ppm of ¹H-NMR spectrum showed coupling of methine protons (H₄, H₅ and H₆) (). This correlation indicated the presence of a conjugated ethylenic system. The signal of H₅ is observed as a doublet of doublets pattern due to its interaction with H₆ and H₄; the ³J

corresponding coupling constant values are 10.8 and 15.3 Hz, respectively. The coupling constant $^3J_{H5,H6}$, indicated the E-configuration of the diene fragment [20]. In addition, the coupling constant $^3J_{H4,H5}$ (15.6 Hz) corresponds to an E-configuration of the C₄-C₅ double bond, confirmed by NOE correlation among H₅ and H₁₃. The configuration of the C₆-C₇ double bond, was established from NOE correlation between H₆ → H₈ and H₅ → H₁₄, respectively, suggesting an E configuration. Finally, the structure of compound **58** was determined as (3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol.

Table 2. Full NMR data of (3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol **58** (500MHz, 300K and CDCl₃)

(3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol 58								
¹³ C NMR			¹ H NMR			¹ H- ¹ H and ¹ H- ¹³ C 2D correlations		
δ (ppm)	Type		δ (ppm)	Mult	J (Hz)	HMBC*	NOESY	COSY
1 8.37	CH ₃		0.9	t	7.4	H ₂	H ₂	-
2 35.43	CH ₂		1.6	dd	7.44 & 1.6	H ₁ H ₁₃	H ₁ H ₁₃	-
3 73.38	C-O		-	-	-	H ₁ H ₂ H ₄ H ₅ H ₁₃	-	-
4 137.80	CH=		5.6	d	15.3	H ₂ H ₆ H ₁₃	H ₆ H ₁₃	H ₆
5 123.99	CH=		6.5	dd	15.3 & 10.8	H ₆ H ₁₃	H ₁₃ H ₁₄	H ₁₄
6 124.13	CH=		5.9	d	10.8	H ₄ H ₅ H ₈ H ₁₄	H ₈ / H ₉	H ₈ / H ₉
7 138.75	C=		-	-	-	H ₅ H ₈ H ₁₄	-	-
8 39.97	CH ₂		2.1	m	-	H ₉ H ₁₄	H ₆	-
9 26.62	CH ₂		2.1	m	-	H ₈	H ₁₀ H ₁₄ H ₁₅	H ₁₄ H ₁₅
10 124.03	CH=		5.1	t	6.0	H ₈ H ₁₂ H ₁₅	H ₉ H ₁₂ H ₁₅	H ₁₂
11 131.71	C=		-	-	-	H ₁₂ H ₁₅	-	-
12 25.71	CH ₃		1.7	s	-	H ₁₀ H ₁₅	H ₁₀	-
13 27.58	CH ₃		1.3	s	-	H ₄	H ₁	-
14 16.75	CH ₃		1.8	s	-	H ₆	H ₅	-
15 17.70	CH ₃		1.6	s	-	H ₁₀ H ₁₂	H ₁₀	-

*HMBC correlation are reported as C->H




Figure 3. Long range ^{13}C - ^1H HMBC (A) and NOE (B) correlations observed in compound **58**




Figure 4. H-H coupling pattern of the ethylenic system of **58**

Compounds **42a** and **42b** were isolated from the fraction **F27** (12 mg) obtained from *P. porrellioides* essential oil by column chromatography using hexane and diisopropyl oxide (90:10). Both apolar and polar GC chromatograms of **F27** exhibited one major signal amounted for 70% of the FID-response. ^{13}C -NMR spectra of **F27** exhibited 29 signals including one carbon atom at δ_{c} 74.4 ppm with double relative intensity. These observations support the hypothesis of the occurrence of a mixture of diastereoisomers. While the molecular ion of **42** has not been observed, the EI-MS spectra showed a base peak at m/z 107; the other intense fragment ion, detected at m/z 59 suggests the presence of a 2-hydroxy isopropyl group, characteristic of tertiary alcohol [21]. ESI (+)-HRMS measurements allowed to determine the molecular formula $\text{C}_{15}\text{H}_{26}\text{O}$, (detected ion $\text{C}_{15}\text{H}_{26}\text{ONa}^+$ (m/z)_{exp} 245.1878 and (m/z)_{th} 245.1876, error +0.8 ppm). According to the δ_{c} intensities, two sets of resonances with a ratio (2:1) assigned to **42a** and **42b** respectively, could be extracted from the ^{13}C -NMR spectra of **F27**. Attached Proton Test (APT) experiment confirms the molecular formula $\text{C}_{15}\text{H}_{26}\text{O}$ for both isomers and the assignment of the 15 carbon signals was carried out as follow: 5 methyl groups, 3 methylenes, 4 methynes of which two ethylenics at δ_{c} 124.88 and 129.59 ppm and 3 quaternary carbons at δ_{c} 74.38, 137.34 and 134.16 ppm (Table 3). 2D HSQC, HMBC and COSY experiments confirmed the structure of p-menthane framework displaying a side chain composed of five carbon atoms, three of which have sp^3 hybridization and two of sp^2 type, forming a double bond (). The isopropyl alcohol group was confirmed by HMBC assignment, where appeared the correlations between protons of both methyl terminal groups (δ_{H} at 1.17 and 1.24 ppm, respectively) and the quaternary carbon atom at δ_{c} 74.38 ppm. Moreover, our ^1H and ^{13}C NMR data are in good agreement with those of α -terpineol, an alcohol monoterpene with p-menth-1-en-8-ol structure. Regarding the side chain assignment, the δ_{CH_2} and one δ_{CH_3} of one isomer exhibited excessive $\Delta\delta$ relative to the other (7.4 ppm and 6.3 ppm, respectively), suggesting steric γ -effects generated by double bond stereochemistry. As ^1H -coupling constants of the allylic system were not sufficiently to resolve the stereochemistry, NOESY experiments were acquired to elucidate the spatial proximities determined by the double bond for each isomer. Concerning **42a**, the sp^2 methine proton $\delta_{\text{H}11}$ at 5.15 ppm showed a strong NOE connectivity to H_{13} and H_{14} , while the H_3 had a strong NOE cross peak to the allylic proton of the methyl group C_{15} at δ c 16.21 ppm. This clearly indicated that the double bond in the side chain of **42a** have E configuration. These spectral features require a structure of p-menth-1-en-3-[2-(E)-methylbut-1-enyl]-8-ol for **42**. Contrary, the double

bond brought by **42b** side chain was found as Z stereochemistry; these was assigned according to correlations between $H_3 \rightarrow H_{13}$, and $H_{11} \rightarrow H_{15}$, respectively ().

Table 3. Full NMR data of *p*-menth-1-en-3-[2-methylbut-1-enyl]-8-ol isomers **42a** and **42b** (in $CDCl_3$, at 500MHz and 300K)

42a							42b						
¹³ C NMR			¹ H NMR		2D NMR		¹³ C NMR			¹ H NMR		2D NMR	
	δ (ppm)	DEPT	δ(ppm)	Mult (J, Hz)	HMBC*	NOESY	δ (ppm)	DEPT	δ(ppm)	Mult (J, Hz)	HMBC*	NOESY	
1	134.16	C	-	-	H ₇		134.29	C	-	-	H ₇		
2	124.88	CH	4.95	d (1.26)	H ₇	H ₃ H ₇ H ₂ H ₅	125.19	CH	4.95 s	d (1.26)	H ₇	H ₃ H ₇	
3	37.18	CH	3.08	m		H ₉ H ₁₁ H ₁₅	36.93	CH	3.08	m	H ₅		
4	50.48	CH	1.5	m	H ₁₁ H ₂ H ₁₀ H ₉	H ₅ H ₆	50.54	CH	1.34	m	H ₉ H ₁₀	H ₅ H ₆	
5	25.37	CH ₂	1.83/1.31	m	H ₃ H ₄ H ₆		25.29	CH ₂	1.83/1.31	m			
6	30.7	CH ₂	2.03	m	H ₂ H ₇		30.65	CH ₂	1.96	m	H ₅ H ₇	H ₅	
7	23.31	CH ₃	1.67	s	H ₂	H ₆	23.33	CH ₃	1.64	s		H ₇	
8	74.38	C-OH	-	-	OH H ₁₀ H ₉		74.38	C-OH	-	-	H ₉ H ₁₀		
9	25.78	CH ₃	1.17	s	H ₁₀	H ₃	25.71	CH ₃	1.17	s	H ₁₀	H ₃	
10	28.93	CH ₃	1.24	s	OH H ₉	H ₄ H ₅ H ₄ H ₆	28.99	CH ₃	1.23	s	H ₈ H ₉	H ₄ H ₅	
11	129.59	CH	5.15	dd (10.1 and 1.26)	H ₁₃ H ₁₅	H ₁₃ H ₁₄ OH	130.69	CH	5.11	d(10.1)	H ₁₃ H ₁₅	H ₁₅	
12	137.34	C	-	-	H ₁₃ H ₁₅ H ₁₄		137.4	C	-	-	H ₁₃ H ₁₄ H ₁₅		
13	32.5	CH ₂	2.03	q (7.32)	H ₁₁ H ₁₄ H ₁₅	H ₁₄ H ₁₅	25.11	CH ₂	2.19/2.09	m	H ₁₄ H ₁₅	H ₃ H ₁₄	
14	12.55	CH ₃	1.02	t (7.32)	H ₁₃	H ₁₃	12.85	CH ₃	1.06	t (7.23)	H ₁₃	H ₁₃	
15	16.21	CH ₃	1.73	d (0.88)	H ₁₁ H ₁₃	H ₃ H ₉ H ₁₃ H ₁₄	22.98	CH ₃	1.74	d (1.47)	H ₁₁ H ₁₃	H ₃ H ₉ H ₁₃ H ₁₄	
		OH	2.6	OH				OH	2.6	OH			

* HMBC correlations are reported as C->H




Figure 5. Structure of **42a** and schematic representation of HMBC correlations




Figure 6. Zoom of NOESY spectrum showing specific correlations used to resolve stereochemistry of the acyclic double bond in **42a**, **b**

Both alcohol isomers **42a** and **42b** possess a humbertiane skeleton, a relatively rare sesquiterpene pattern. To our knowledge, only four isomeric isohumbertiols, structurally related to the alcohols **42a** and **42b**, were identified from the wood of *Humbertia madagascariensis* Lam [22]. In addition, two analogous structures were isolated after fungal transformation of α -farnesene while the NMR data (^1H and ^{13}C) reported in the literature [23], differed significantly from the experimental data described here for both new sesquiterpene alcohols.

2.3. *P. porelloides* volatile components: chemical compositions of specific plant extracts

The chemical compositions of essential oil (EO), hydrosol extract (HY), both hexane and diethyle oxide extracts (EXT_H and EXT_O), volatile fraction (VF) as well as microwave extract (MW) were investigated by GC/RI, GC-MS and ^{13}C -NMR (Table 4).

Table 4. Chemical compositions of *P. porelloides* samples from Corsica

217

		Samples									
No ¹	Compounds	L RI ^{A 2}	RI ^{A 3}	RI ^{P 4}	EO	HY	EXT		MW	VF	
							EXT _H	EXT _O			
1	hexanal	770	771	819	t	-	t	t	-	t	IR, MS
2	heptanal	876	876	1079	t	-	t	t	-	t	IR, MS
3	α -pinene	931	931	1032	t	-	t	t	-	1.0	IR, MS
4	camphene	943	944	1066	t	-	t	t	-	0.1	IR, MS
5	6-methylhept-5-en-2-one	963	954	1343	0.1	-	t	-	-	0.1	IR, MS
6	oct-1-en-3-ol	959	962	1453	0.4	1.9	-	t	-	t	IR, MS
7	octen-3-one	963	980	1260	t	-	-	t	-	0.1	IR, MS
8	octan-3-ol	982	982	1401	t	-	-	t	-	-	IR, MS
9	phenylaldehyde	1013	1010	1616	t	-	-	-	-	t	IR, MS
10	β -phellandrene	1021	1021	1219	t	-	-	0.1	-	0.4	IR, MS
11	nonanal	1083	1083	1396	0.1	-	t	-	-	-	IR, MS
12	octen-3-yl acetate	1094	1094	1377	0.1	-	t	0.6	0.2	2.4	RI, MS, Ref
13	decanal	1186	1185	1496	t	-	-	-	-	-	RI, MS, Ref
14	bicycloelemene	1334	1334	1552	0.3	-	0.9	1.1	0.4	2.0	RI, MS, Ref
15	maali-1,3-diene	1347	1346	1532	0.8	-	0.5	0.6	-	1.5	RI, MS, Ref
16	anastreprene	1373	1369	-	0.2	-	0.4	1.2	0.8	1.4	RI, MS, Ref
17	isodene	1372	1373	-	0.1	-	0.1	0.2	-	0.2	IR, MS
18	α -copaene	1379	1376	1498	0.2	-	0.4	0.4	-	1.7	IR, MS
19	β -elemene	1388	1388	1592	0.2	-	0.4	0.6	-	1.4	IR, MS
20	african-3-ene	1391	1390	-	-	-	0.0	0.3	-	1.1	RI, MS, Ref
21	α -barbatene	1414	1409	1565	1.8	-	1.3	2.5	0.4	4.7	RI, MS, Ref
22	tritomarene	1416	1413	1410	0.2	-	-	0.5	-	1.1	RI, MS, Ref
23	aristolene	1420	1419	1581	2.6	-	5.0	1.5	4.1	7.3	RI, MS, NMR
24	γ -maaliene	1428	1425	1613	1.2	-	2.9	0.2	3.0	0.5	RI, MS, Ref
25	calarene	1439	1438	1603	0.6	-	0.5	0.6	0.5	1.0	IR, MS
26	α -maaliene	1440	1440	-	0.5	-	t	0.6	0.8	-	RI, MS, Ref
27	β-barbatene	1445	1445	1663	28.7	-	23.1	19.5	32.6	39.5	RI, MS, NMR
28	aromadendrene	1447	1458	1601	0.3	-	0.4	0.5	-	0.6	RI, MS, Ref
29	α -acoradiene	1464	1460	-	0.6	-	0.2	0.3	-	0.9	RI, MS, Ref
30	β -acoradiene	1465	1462	-	0.1	-	0.4	0.2	-	0.4	RI, MS, Ref
31	γ -curcumene	1475	1471	-	0.1	-	0.7	0.5	-	0.4	RI, MS, Ref
32	β -chamigrene	1478	1473	-	1.3	-	0.5	0.8	-	1.5	RI, MS, Ref

33	α -curcumene	1473	1481	1738	0.1	-	0.1	0.4	-	0.7	RI, MS, Ref
34	β -selinene	1483	1485	-	1.1	-	0.0	0.1	-	-	RI, MS, Ref
35	β -maaliene	1480	1481	-	0.1	-	0.1	0.3	-	-	RI, MS, Ref
36	bicyclogermacrene	1494	1494	1744	8.2	-	16.4	14.1	17.8	14.2	RI, MS, NMR
37	204[M] ⁺ ; 107(100); 93(90)	1498	1497	-	1.5	-	0.4	0.9	-	1.0	
38	ledene	1494	1498	1706	t	-	-	0.9	-	0.4	RI, MS
39	α -chamigrene	1503	1500	-	t	-	-	0.4	-	0.4	RI, MS, Ref
40	γ -cadinene	1507	1514	1775	t	-	-	0.6	-	0.7	RI, MS
41	α -alaskene	1512	1513	-	t	-	-	0.5	-	0.4	RI, MS, Ref
42a	<i>p</i>-menth-1-en-3-[2-methyl-1E-butenyl]-8-ol	-	1536	1917	1.2	3.4	0.3	1.2	-	-	RI, MS, NMR
42b	<i>p</i>-menth-1-en-3-[2-methyl-1Z-butenyl]-8-ol	-	1536	1917	3.8	-	0.6	-	-	-	RI, MS, NMR
43	tamariscol	1535	1536	1917	1.2	-	0.6	1.1	-	-	RI, MS
44	204[M] ⁺ ; 161(100); 91(56)	-	1532	1917	3.7	-	-	-	-	-	
45	4-epi-maaliol	-	1544	-	0.7	-	0.6	0.3	-	-	RI, MS, NMR
46	222[M] ⁺ ; 107(100); 135(51)	-	1545	1832	0.9	4.7	-	-	-	-	-
47	pallustrol	1567	1558	1923	t	-	-	t	-	-	RI, MS, Ref
48	spathulenol	1557	1561	2090	0.7	9.3	1.1	0.6	3.5	-	RI, MS, NMR
49	204[M] ⁺ ; 107(100); 135(56)	-	1567	-	0.9	-	-	-	-	-	-
50	globulol	1571	1578	2077	4.4	17.2	1.9	1.3	0.2	0.9	RI, MS, NMR
51	viridiflorol	1591	1585	2085	5.8	4.2	1.9	-	-	0.3	RI, MS, NMR
52	222[M] ⁺ ; 107(100)	-	1588	-	1.2	-	-	-	-	-	-
53	238[M] ⁺ ; 149(100)	-	1591	1920	5.21	-	-	-	-	-	-
54	rosifoliol	1599	1587	2108	3.8	7.3	5.9	6.1	0.4	-	RI, MS, NMR
55	maalian-5-ol	1607	1595	2051	5.4	9.4	2.2	1.9	7.6	1.2	RI, MS, NMR
56	204[M] ⁺ ; 107(100); 135(76)	-	1611	1938	1.4	-	-	-	-	-	-
57	222; 107(100); 105(75)	-	1613	-	1.3	4.2	-	-	-	-	-
58	(3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol	-	1616	2136	-	-	13.3	15.7	17.2	-	RI, MS, NMR
Total identified					76.9	52.6	82.6	77.9	89.4	90.3	
Classes of compounds (%)											
Hydrocarbon compounds					49.3	-	54.4	49.2	60.4	85.4	
Oxygenated compounds					27.5	72.3	28.1	28.7	29.1	4.9	
Monoterpene hydrocarbons					-	-	-	0.1	-	1.5	
Monoterpene oxygenated					-	-	-	-	-	-	
Sesquiterpene hydrocarbons					49.3	-	54.4	49.1	60.4	83.8	
Sesquiterpene oxygenated					26.9	70.5	28.1	28.1	28.8	2.4	
Other					0.6	1.9	-	0.6	0.2	2.5	

¹ Order to elution is given on apolar column (Rtx-1) ² LRI_A: Literature retention indices on apolar column reported from literature [24]. ³ RI_A: Retention indices on Rtx-1 (apolar) column ⁴ RI_r: Retention indices on Rtx-Wax (polar) column ⁵ Percentages of individual components on Rtx-1 except those with the same RI_A; percentages given on Rtx-Wax column. ⁶ RI: Retention Indices; MS: Mass spectrometry; NMR: Nuclear Magnetic Resonance. Compounds in italic type were identified from commercial libraries [24]

EO : Essential oil obtained by Hydrodistillation; HY : LLE extract obtained from hydrosol; EXT_H and EXT_O: hexane and diethyl oxide extracts (cold maceration); MW : Assisted microwave extract; VF : Volatiles sampled by SPME.

P. porelloides EO was dominated by hydrocarbon compounds (49.4 %), among them two sesquiterpenes were predominated: β -barbatene **27** (28.7%) and bicyclogermacrene **36** (8.2 %). Oxygenated compounds were represented by 11 sesquiterpene alcohols (27.7%) and 10 non-terpenic compounds (0.7 %). The other main components were globulol **50** (4.4 %), viridiflorol **51** (5.8 %) and maalian-5-ol **55** (5.4%). Hydrosol extract (HY) was dominated by globulol **50** (17.2 %), maalian-5-ol **55** (9.4%), spathulenol **48** (9.3%) and rosifoliol **54** (7.3%). Relative to the essential oil, none hydrocarbon compounds has been detected in the LLE extract obtained from hydrosol.

Unlike the essential oil, both hexane and diethyl oxide extracts (EXT_H and EXT_O) were dominated by β -barbatene **27** (23.1 and 19.5 %, respectively) and (3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol **58** (13.3 and 15.7 %, respectively) which was not detected in the essential oil. bicyclogermacrene **36** (16.4 and 14.1 %, respectively), rosifoliol **54** (5.9 and 6.1 %, respectively) and aristolene **23** (5.0 and 1.5 %, respectively) were presents in remarkable amounts. The assisted microwave extract (MW) exhibited close chemical composition with β -barbatene **27** (32.6%), bicyclogermacrene **36** (17.8%) and (3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol **58** (17.2%), as main components. Finally, the volatiles emitted by the plant and sampled by SPME were β -barbatene **27** (39.5 %), aristolene **23** (7.32%), bicyclogermacrene **36** (14.2%) and maalian-5-ol **55** (1.2 %).

The volatile metabolites of *P. porelloides* were very atypical but the most surprising was the exclusive presence of an oxygenated linear sesquiterpene **58** in the solvent extracts. It is likely that hydrodistillation conditions cause the degradation of **58** into several compounds. However, the presence of **58** in the microwave extract proves that the temperature is not the only experimental parameter responsible for the degradation process. It is very likely that the protic character of water plays an important role. The absence of **58** in the volatile fraction emitted by the plant material can be explained by the conditions used for SPME; particularly the selectivity of the adsorbent phase seems to be implicated. The SPME experiment carried out on the plant extract support this hypothesis, allowing to detect the entire components previously listed except **58** [25].

It is difficult to accurately distinguish between the real essential oil constituents and those issued by the compound degradation. As the plant resources are limited, hemisynthesis trials becomes too complicated, especially as bicyclogermacrene is known to degrade into viridiflorol and spathulenol [26,27]. But in our work, these compounds are present in sample of cold and hot extractions, we may think that these compounds are naturally present in *P. porelloides*.

2.4. Evaluation of biological activity: antitrypanosomal, antileishmanial and cytotoxic activities

In the in vitro screening phase, a molecule is considered to have strong antiparasitic activity if its median inhibitory concentration (IC₅₀) is below 1 $\mu\text{g}\cdot\text{mL}^{-1}$. For complex mixtures such as essential oils, the activity becomes interesting when its IC₅₀ is less than 12.5 $\mu\text{g}\cdot\text{mL}^{-1}$. When the selectivity index (SI) is greater than 5, then the sample is considered a hit and is therefore likely to proceed to the in vivo stage [28].

It could be noted that *P. porelloides* EO and EXT_O could be considered as having a good activity with IC₅₀ values $\leq 20 \mu\text{g}/\text{mL}$ of *Leishmania mexicana mexicana* and a best activity ($< 6 \mu\text{g}/\text{mL}$) on *Trypanosoma brucei brucei*. Concerning the selectivity which was assessed here on the human non cancer fibroblast cell line WI38, only the essential oil had a sufficient selectivity (SI: 11.7) to be a good candidate for bioguided analysis.

Concerning cytotoxic activities, we observed that diethyl oxide extract of *P. porelloides* could be considered as having a good potential with IC₅₀ value $\leq 5 \mu\text{g}/\text{mL}$ of WI38 and J774 (murine cancer macrophages) cells but we did not find a clear selectivity on the cancer cell line. Nevertheless, this cytotoxicity may be interesting in the search for anticancer agents (Table 5).

It has to be mentioned that earlier studies focused on different liverworts reported also an antiprotozoal activity, notably for alpha-eudesmol [29] and marchantin A [30]

Table 5. The cytotoxicity (W138 and J774) and *in-vitro* activity of *P. porelloides* EO and EXT_o against *Leishmania mexicana mexicana* (Lmm) and *Trypanosoma brucei brucei* (Tbb).

Sample	Cytotoxicity		Antiprotozoal assay		Selectivity indices		
	IC ₅₀ ± SD in µg/mL (µM for Pure Compound)				SI = IC ₅₀ (WI38)/IC ₅₀ (J774 or parasite)		
	WI38	J774	Lmm	Tbb	J774	Lmm	Tbb
EO	23.85±4.39	28.81±0.45	15.99±0.85	2.03±0.12	0.8	1.5	11.7
EXT _o	2.96±0.17	1.25±0.08	17.73±1.14	5.18±0.81	2,4	0.2	0.6
Camptothecin	0.031±0.002	0.01±0.001					
Pentamidine			0.07±0.004				
Suramine				0.03±0.004			

W138: non cancer human fibroblasts; J774: cancerous macrophage-like murine cells; Tbb: *Trypanosoma brucei brucei* (bloodstream forms); Lmm: *Leishmania mexicana mexicana* promastigotes; Selectivity index calculated for antiparasitic activities compared to W138 cytotoxicity.

3. Materials and Method

3.1. Plant material

Fresh *P. porelloides* were harvested in 2019 in one location of Corsica (France). Botanical determination was performed according to the determination keys summarized in Bryophyte Flora [31] and voucher specimens were deposited in the herbarium of University of Corsica, Corte (France).

3.2. Essential oil and hydrosol isolation

After 15 day of drying, plant material (500 g) was subjected to hydrodistillation (HD) for 5 h using a Clevenger-type apparatus according to the method recommended in the *European pharmacopoeia* [32]. Hydrosol (300 mL) obtained by HD was submitted to Liquid/liquid Extraction (LLE) in order to obtain an liquid extract (HY, 57.3 mg). LLE was performed three times using 50 mL of diethyle oxide.

The essential-oil yield (0.2 %) were expressed in % (w/dw) based on the weight of the dried plant material.

3.3. SPME experiments

Volatile fraction (VF) emitted by the plant sample were extracted with HS-SPME method. Bryophytes samples (1 g) were crushed and disposed into 20 mL headspace vials. The vial was sealed with a silicon septum placed in 70°C dry bath, and equilibrated for 30 min. A 75 µm DVB/CAR/PDMS solid-phase fiber (Supelco, Bellefonte, PA, USA) was then plugged into the headspace of the vial for 60 min. Later, the volatiles sampled by the solid-phase fiber was analysed after desorption into the gas chromatography-mass spectrometry (GC-MS) injection port (5 min) in splitless mode. Each sample was done in triplicates.

3.4. Solvent extractions

Dried plant materials (200 g) were mechanically powdered and extracted with hexane and diethyl oxide at room temperature for 24 hours each in order to give after filtration and concentration in vacuum, both extract called EXT_H (400 mg) and EXT_O (640 mg), respectively.

3.5. Microwave-assisted extractions

Dried plant materials were mechanically powdered and extracted using Multiwave 3000 (Anton Paar, Gratz, Austria) apparatus provided with 16 ceramic vessels. For each vessel 5 g of dried bryophyte were introduced with 40 mL of hexane and extraction was realized at 180 °C during 20 min. Solvent was then filtered on activated carbon and concentrated under vacuum. Resulting extract was next taken up in absolute ethanol and centrifuged (20 min at 6000 rpm) and the supernatant was collected and concentrated to finally obtain the MAE extract.

3.6. Essential oil (EO) and diethyl oxide extract (EXT_O) fractionations

Essential oil (EO, 800 mg) and diethyle oxide extract (EXT_O, 640 mg) of *P. Porelloides* was submitted to column chromatography (CC) on a silica-gel column (200-500 µm, 12 g, Clarisep® Bonna Agela Technologies, Willington, USA) with Combi Flash apparatus (Teledyne ISCO, Lincoln, USA) equipped with a fraction collector monitored by an UV detector. Using gradients of (v/v) hexane/diisopropyl oxide (HEX/DIPO), forty-seven fractions (2 hydrocarbon fractions and 45 oxygenated fractions) were eluted from EO and two (one hydrocarbon fraction and one oxygenated fraction) were eluted from EXT_O.

3.7. GC-FID Conditions

Analyses were carried out using a *Perkin-Elmer Clarus 600* Gas Chromatography (GC) apparatus (Walton, MA, USA) equipped with a single injector and two flame ionization detectors (FIDs) for simultaneous sampling to two fused-silica capillary columns (60 m × 0.22 mm i.d., film thickness 0.25 µm; Restek, Bellefonte, PA, USA) with stationary phases of different polarity, *i.e.* a nonpolar *Rtx-1* (polydimethylsiloxane) and a polar *Rtx-Wax* (polyethylene glycol). The oven temperature was programmed rising from 60 to 230 °C at 2 °C min⁻¹ and held isothermal at 230 °C for 30 min. The injector temperature was maintained at 280 °C and detector temperature at 280 °C, the carrier gas was H₂ (0.7 mL.min⁻¹) and the samples were injected (0.1 µL of pure oil) in the split mode (1:80). Retention indices (RIs) of the compounds were determined relative to the retention times (*t_R*) of a series of n-alkanes (C₅–C₃₀; commercial solution obtained from *Restek*, Bellefonte, PA, USA) using the Van den Dool and Kraetz equation [33].

3.8. GC-MS Analysis

Essential oils, extracts and fractions obtained by CC were investigated using a *Perkin Elmer Turbo Mass* quadrupole detector directly coupled to a *Perkin Elmer Autosystem XL* (Walton, MA, USA), equipped with the two same fused-silica capillary columns as described above. Both columns were used with the same MS detector. The analyses were consecutively carried out on the nonpolar and on the polar column. Hence, for each sample, two reconstructed ionic chromatograms (*RIC*) were provided, which were investigated consecutively. The GC conditions were the same as described above and the MS parameters as follows: Ion-source temperature, 150 °C,

ionization energy, 70 eV; electron ionization mass spectra acquired over a mass range of 35–350 amu during a scan time 1 s. The injection volumes for the essential oil and the fractions were 0.1 μL , and 0.2 μL , respectively.

3.9. High resolution mass spectrometry experiments

High resolution mass spectrometry experiments were performed with a Synapt G2 HDMS quadrupole/time-of-flight (Manchester, UK) equipped with an electrospray source operating in positive mode. Samples were introduced at 10 $\mu\text{L}\cdot\text{min}^{-1}$ flow rate (capillary voltage +2.8 kV, sampling cone voltage: varied between +20 V and +30 V) under a curtain gas (N_2) flow of 100 $\text{L}\cdot\text{h}^{-1}$ heated at 35 $^{\circ}\text{C}$. Accurate mass experiments were performed using reference ions from CH_3COONa internal standard. The samples were dissolved and further diluted in methanol (Sigma-Aldrich, St-Louis - MO, USA) doped with formic acid (1% v/v) prior to analysis. Data analyses were conducted using MassLynx 4.1 programs provided by Waters.

3.10. NMR Conditions

Nuclear Magnetic Resonance (NMR) spectra were recorded on the CC-fraction **F27** obtained from the EO and the polar CC-fraction obtained from the EXT₀. NMR experiments were acquired in CDCl_3 (EuroIsotop, Saint Aubin, France), at 300K using a Bruker Avance DRX 500 NMR spectrometer (Karlsruhe, Germany) operating at 500.13 MHz for ^1H and 125.77 MHz for ^{13}C Larmor frequency with a double resonance broadband fluorine observe (BBFO) 5 mm probe head. ^{13}C -NMR experiments were recorded using one-pulse excitation pulse sequence (90° excitation pulse) with ^1H decoupling during signal acquisition (performed with WALTZ-16); the relaxation delay has been set at 2 s. For each analyzed sample, depending on the compound concentration, 3k up to 5k free induction decays (FID) 64k complex data points were collected using a spectral width of 30000 Hz (240 ppm). Chemical shifts (δ in ppm) were reported relative to residual signal of CDCl_3 (δ_{C} 77.04 ppm). Complete ^1H and ^{13}C assignments of the new compound were obtained using 2D gradient-selected NMR experiments, ^1H - ^1H COSY (CORrelation Spectroscopy), ^1H - ^{13}C HSQC (Heteronuclear Single Quantum Correlation), ^1H - ^{13}C HMBC (Heteronuclear Multiple Bond Coherence) and ^1H - ^1H NOESY (Nuclear Overhauser Effect Spectroscopy), for which conventional acquisition parameters were used, as described in the literature [34].

3.11. Identification of components

Identification of individual components in essential oil, extracts or CC-fractions was based on a methodology involving integrated techniques, such as GC retention indices, GC-MS (EI) and NMR. Identification of volatiles sampled by SPME were carried only by GC retention indices and GC-MS (EI). The identification of individual components was based i) on the comparison of the retention indices (*RI*s) determined on the polar and nonpolar columns with those of authentic compounds or literature data [24,35] ii) on computer matching of the mass spectra with commercial MS-libraries and the mass spectra with those listed in our homemade MS-library built of mass spectra of authentic compounds or literature data [36,37] iii) comparing the ^{13}C -NMR chemical shifts of CC-fraction components with those of reference spectra reported in the literature iv) NMR assignments using 1D and 2D data.

(3,7,11)-trimethyl-dodec-4E,6E,10-atrien-3-ol **58** : Yellow oil (20 mg); RI apolar 1616, RI polar 2136; for ^1H and ^{13}C NMR data (see Table 2); MS (EI; 70 eV) *m/z* (rel. Int) : 222 [M^+] (4), 107 (100), 41 (80), 135 (73), 69 (62), 91 (53), 93

(53), 105 (49), 43 (46), 79 (40), 204 (30), 119 (30), 161 (29), 77 (28), 55 (26). HRMS: detected ion m/z 245.1879 [MNa⁺] (calc. for C₁₅H₂₆ONa⁺, error: +1.2ppm).

p-menth-1-en-3-[2-methylbut-1-enyl]-8-ol **42a,b**: Colorless oil (7mg); RI apolar 1528, RI polar 1917; for ¹H and ¹³C NMR data (see Table 3); MS (EI; 70 eV) m/z (rel. Int): 222 [M⁺] (1), 107 (100), 59 (31), 161 (31), 91 (31), 41 (28), 175 (26), 105 (25), 135 (23), 79 (23), 119 (22), 93 (21), 108 (19), 43 (16), 77 (16). HRMS: detected ion m/z 245.1878 [MNa⁺] (calc. for C₁₅H₂₆ONa⁺, error: +0.8ppm).

3.12. Component Quantification

The quantification of components was performed using the methodology reported by Bicchi [38] and adapted in our laboratory [39]. Briefly, the compound quantification was carried out using peak normalization, including FID response factors relative to tridecane (0.7 g/100 g) used as internal standard, and expressed as normalized contents (% abundances).

3.13. Parasites, cells and media

Trypanosoma brucei brucei (strain427) bloodstream forms were cultured in vitro in HMI9 medium containing 10% heat-inactivated foetal bovine serum [40]. *Leishmania mexicana mexicana* promastigotes (MHOM/BZ/84/BEL46) were cultivated in vitro in a semi-defined medium (SDM-79) [41] supplemented with 15% heat-inactivated foetal bovine serum. The human normal fibroblast cell line, WI-38, was cultivated in vitro in DMEM medium containing 4 mM L-glutamine, 1 mM sodium pyruvate supplemented with 10% heat-inactivated foetal bovine serum and penicillin–streptomycin (100 UI/ml to 100 µg/ml). All cells were incubated in a humidified atmosphere with 5% CO₂ at 37 °C except the *Leishmania promastigotes* which were incubated at 28 °C.

3.14. In vitro test for antitrypanosomal and antileishmanial activity

The *in vitro* test was performed as described previously [42] Pentamidine isethionate salt (a commercial antileishmanial drug) and suramine sodium salt (a commercial antitrypanosomal drug) were used as positive controls in all experiments with an initial concentration of 10 µg/mL. First stock solutions of crude extracts and compounds were prepared in DMSO at 20 mg/mL (and 2 mg/mL for positive controls). The solutions were further diluted in medium in medium to give 0.2 mg/ml stock solutions. Essential oil and extracts were tested in eight serial three-fold dilutions (final concentration range: 100–0.05 µg/mL) in 96-well microtiter plates. All tests were performed in duplicate.

3.15. Cytotoxicity assay

The cytotoxicity of the essential oil and extracts on WI-38 and J774 cells was evaluated as previously described from the same stock solutions [43].

Supplementary Materials:

Author Contributions: Conceptualization, A.P., A.T. and A.M.; methodology, A.P., A.T. and A.M.; formal analysis, A.P.; investigation, A.P. and J.Q.L.; resources, A.P.; writing—original draft preparation, A.P., A.T. and A.M.;

writing—review and editing, A.P., A.T. J.C. and A.M; project administration, A.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments:

Conflicts of Interest: The authors declare no conflict of interest.

References

- Manneville, O. *LES BRYOPHYTES: MOUSSES, SPHAIGNES, HEPATIQUES Particularités et Cycles Biologiques, Divers Groupes, Écologie*; Station Alpine de Joseph Fourier: Grenoble, 2011;
- Sotiaux, A.; Pioli, A.; Royaud, A.; Schumacker, R.; Vanderpoorten, A. A Checklist of the Bryophytes of Corsica (France): New Records and a Review of the Literature. *Journal of Bryology* **2007**, *29*, 41–53, doi:10.1179/174328207X171872.
- Pannequin, A. Caractérisation chimique des bryophytes de Corse et propriétés biologiques. phdthesis, Université Pascal Paoli, 2019.
- Pannequin, A.; Tintaru, A.; Desjobert, J.-M.; Costa, J.; Muselli, A. New Advances in the Volatile Metabolites of *Frullania Tamarisci*. *Flavour Fragr J* **2017**, 1–10, doi:10.1002/ffj.3407.
- Asakawa, Y. Chemical Constituents of the Hepaticae. In *Fortschritte der Chemie organischer Naturstoffe / Progress in the Chemistry of Organic Natural Products*; Fortschritte der Chemie organischer Naturstoffe / Progress in the Chemistry of Organic Natural Products; Springer, Vienna, 1982; pp. 1–285 ISBN 978-3-7091-8679-4.
- Asakawa, Y. Biologically Active Compounds from Bryophytes. *Pure and Applied Chemistry* **2007**, *79*, 557–580, doi:10.1351/pac200779040557.
- Asakawa, Y.; Ludwiczuk, D.A.; Nagashima, D.F. Biologically Active Compounds of the Marchantiophyta and Bryophyta. In *Chemical Constituents of Bryophytes*; Progress in the Chemistry of Organic Natural Products; Springer Vienna, 2013; pp. 619–638 ISBN 978-3-7091-1083-6.
- Asakawa, Y.; Ludwiczuk, A.; Nagashima, F.; Editors. *Chemical Constituents of Bryophytes: Bio- and Chemical Diversity, Biological Activity, and Chemosystematics. (Progress in the Chemistry of Organic Natural Products, Volume 95).*; Springer, 2013;
- Zinsmeister, H.D.; Becker, H.; Eicher, T. Bryophytes, a Source of Biologically Active, Naturally Occurring Material? *Angewandte Chemie International Edition in English* **1991**, *30*, 130–147, doi:10.1002/anie.199101301.
- Asakawa, Y. Chemical Constituents of the Bryophytes. In *Progress in the Chemistry of Organic Natural Products*; Fortschritte der Chemie organischer Naturstoffe; Springer Vienna, 1995; pp. 1–562 ISBN 978-3-7091-7427-2.
- Asakawa, Y. Chemosystematics of the Hepaticae. *Phytochemistry* **2004**, *65*, 623–669, doi:10.1016/j.phytochem.2004.01.003.
- Asakawa, Y.; Inoue, H.; Toyota, M.; Takemoto, T. Sesquiterpenoids of Fourteen Plagiochila Species. *Phytochemistry* **1980**, *19*, 2623–2626, doi:10.1016/S0031-9422(00)83931-X.

13. Toyota, M.; Nakamura, I.; Huneck, S.; Asakawa, Y. Sesquiterpene Esters from the Liverwort *Plagiochila Porelloides*. *Phytochemistry* **1994**, *37*, 1091–1093, doi:10.1016/S0031-9422(00)89535-7. 435
436
14. Figueiredo, A.C.; Sim-Sim, M.; Costa, M.M.; Barroso, J.G.; Pedro, L.G.; Esquivel, M.G.; Gutierrez, F.; Lobo, C.; Fontinha, S. Comparison of the Essential Oil Composition of Four *Plagiochila* Species: *P. Bifaria*, *P. Maderensis*, *P. Retrorsa* and *P. Stricta*. *Flavour Fragr. J.* **2005**, *20*, 703–709, doi:10.1002/ffj.1627. 437
438
439
15. Hackl, T.; König, W.A.; Muhle, H. Three Ent-Eudesmenones from the Liverwort *Plagiochila Bifaria*. *Phytochemistry* **2006**, *67*, 778–783, doi:10.1016/j.phytochem.2006.01.001. 440
441
16. Adio, A.M.; König, W.A. Sesquiterpene Constituents from the Essential Oil of the Liverwort *Plagiochila Asplenioides*. *Phytochemistry* **2005**, *66*, 599–609, doi:10.1016/j.phytochem.2005.01.015. 442
443
17. Nagashima, F.; Tanaka, H.; Toyota, M.; Hashimoto, T.; Okamoto, Y.; Tori, M.; Asakawa, Y. 2,3-Secoaromadendrane-, Aromadendrane- and Maaliane-Type Sesquiterpenoids from Liverwort. *Journal of Essential Oil Research* **1995**, *7*, 343–345, doi:10.1080/10412905.1995.9698536. 444
445
446
18. Merfort, I. Review of the Analytical Techniques for Sesquiterpenes and Sesquiterpene Lactones. *Journal of Chromatography A* **2002**, *967*, 115–130, doi:10.1016/S0021-9673(01)01560-6. 447
448
19. Gerhard, U.; Thomas, S.; Russell, M.F. Accelerated Metabolite Identification by “Extraction-NMR.” *Journal of Pharmaceutical and Biomedical Analysis* **2003**, *32*, 531–538, doi:10.1016/S0731-7085(03)00218-8. 449
450
20. Bothner-By, A.A.; Harris, R.K. Nuclear Magnetic Resonance Studies of 1,3-Butadienes. II. The Relation of (H,H) Coupling Constants to Conformation1. *J. Am. Chem. Soc.* **1965**, *87*, 3451–3455, doi:10.1021/ja01093a029. 451
452
21. Silverstein, R.; Webster, F.; Kiemle, D. *Identification spectrométrique de composés organiques*; De Boeck, 2016; ISBN 978-2-8073-0293-8. 453
454
22. Weyerstahl, P.; Christiansen, C.; Marschall, H. New Sesquiterpene Ethers and Other Constituents Isolated from Brazilian Vassoura Oil. *Liebigs Annalen* **1995**, *1995*, 1039–1043, doi:10.1002/jlac.1995199506147. 455
456
23. Krings, U.; Hardebusch, B.; Albert, D.; Berger, R.G.; Maróstica, Mario; Pastore, G.M. Odor-Active Alcohols from the Fungal Transformation of α -Farnesene. *J. Agric. Food Chem.* **2006**, *54*, 9079–9084, doi:10.1021/jf062089j. 457
458
24. König, W.A.; Joulain, D. *Terpenoids and Related Constituents of Essential Oils, Library of Mass Finder 2.1*; Institute of organic chemistry: Hamburg, 2001; 459
460
25. Asfaw, N.; Storesund, H.J.; Skattebøl, L.; Aasen, A.J. Coexistence of Chrysanthenone, Filifolone and (Z)-Isogeranic Acid in Hydrodistillates. Artefacts! *Phytochemistry* **2001**, *58*, 489–492, doi:10.1016/S0031-9422(01)00254-0. 461
462
26. Njoroge, S.M.; Ukeda, H.; Sawamura, M. Changes in the Volatile Composition of Yuzu (Citrus Junos Tanaka) Cold-Pressed Oil during Storage. *J. Agric. Food Chem.* **1996**, *44*, 550–556, doi:10.1021/jf950284k. 463
464
27. Toyota, M.; Koyama, H.; Mizutani, M.; Asakawa, Y. (–)-Ent-Spathulenol Isolated from Liverworts Is an Artefact. *Phytochemistry* **1996**, *41*, 1347–1350, doi:10.1016/0031-9422(95)00798-9. 465
466
28. Monzote, L.; Piñón, A.; Setzer, W.N. Antileishmanial Potential of Tropical Rainforest Plant Extracts. *Medicines* **2014**, *1*, 32–55, doi:10.3390/medicines1010032. 467
468
29. Otoguro, K.; Iwatsuki, M.; Ishiyama, A.; Namatame, M.; Nishihara-Tukashima, A.; Kiyohara, H.; Hashimoto, T.; Asakawa, Y.; Ōmura, S.; Yamada, H. In Vitro Antitrypanosomal Activity of Plant Terpenes against Trypanosoma Brucei. *Phytochemistry* **2011**, *72*, 2024–2030, doi:10.1016/j.phytochem.2011.07.015. 469
470
471
30. Otoguro, K.; Ishiyama, A.; Iwatsuki, M.; Namatame, M.; Nishihara-Tukashima, A.; Kiyohara, H.; Hashimoto, T.; Asakawa, Y.; Ōmura, S.; Yamada, H. In Vitro Antitrypanosomal Activity of Bis(Bibenzyl)s and Bibenzyls from Liverworts against Trypanosoma Brucei. *J Nat Med* **2012**, *66*, 377–382, doi:10.1007/s11418-011-0587-x. 472
473
474
31. Smith, A. *The Moss Flora of Britain & Ireland*; Cambridge University Press: Cambridge, 1980; 475
32. Council of Europe *European Pharmacopoeia*; 3rd ed.; Council of Europe: Strasbourg, 1997; ISBN 978-92-871-2991-8. 476

-
33. Van Den Dool, H.; Dec. Kratz, P. A Generalization of the Retention Index System Including Linear Temperature Programmed Gas–Liquid Partition Chromatography. *Journal of Chromatography A* **1963**, *11*, 463–471, doi:10.1016/S0021-9673(01)80947-X.
34. Siegmara Braun; Hans-Otto Kalinowski; Berger Stefan *150 and More Basic NMR Experiments. A Practical Course.*; Wiley-VCH: Weinheim, 1998; Vol. 37; ISBN 3 527 29512 7.
35. Adams, R.P. *Identification of Essential Oil Components By Gas Chromatography/Mass Spectrometry*; Allured Publishing Corporation: Carol Stream, 2007;
36. NIST (National Institute of Standards and Technology) *Spectral Database for Organic Compounds.*; NIST, 2008;
37. National Institute of Standards and Technology (NIST) *PC Version of the NIST/EPA/NIH Mass Spectral Library*; Gaithersburg, MD, 2008;
38. Bicchi, C.; Liberto, E.; Matteodo, M.; Sgorbini, B.; Mondello, L.; Zellner, B. d'Acampora; Costa, R.; Rubiolo, P. Quantitative Analysis of Essential Oils: A Complex Task. *Flavour Fragr. J.* **2008**, *23*, 382–391, doi:10.1002/ffj.1905.
39. Andreani, S.; Barboni, T.; Desjobert, J.-M.; Paolini, J.; Costa, J.; Muselli, A. Essential Oil Composition and Chemical Variability of *Xanthium Italicum* Moretti from Corsica. *Flavour Fragr. J.* **2012**, *27*, 227–236, doi:10.1002/ffj.3084.
40. Hirumi, H.; Hirumi, K. Axenic Culture of African Trypanosome Bloodstream Forms. *Parasitol Today* **1994**, *10*, 80–84, doi:10.1016/0169-4758(94)90402-2.
41. Brun, R.; Lun, Z.-R. Drug Sensitivity of Chinese Trypanosoma Evansi and Trypanosoma Equiperdum Isolates. *Veterinary Parasitology* **1994**, *52*, 37–46, doi:10.1016/0304-4017(94)90033-7.
42. Catteau, L.; Schioppa, L.; Beaufay, C.; Girardi, C.; Hérent, M.-F.; Frederich, M.; Quetin-Leclercq, J. Antiprotozoal Activities of Triterpenic Acids and Ester Derivatives Isolated from the Leaves of Vitellaria Paradoxa. *Planta Medica* **2020**, doi:10.1055/a-1286-1879.
43. Abdul Khaliq, H.; Ortiz, S.; Alhouayek, M.; Muccioli, G.G.; Quetin-Leclercq, J. Dereplication and Quantification of Major Compounds of Convolvulus Arvensis L. Extracts and Assessment of Their Effect on LPS-Activated J774 Macrophages. *Molecules* **2022**, *27*, 963, doi:10.3390/molecules27030963.