

Mycorrhizal symbiosis and phosphorus supply determine interactions among plants with contrasting nutrient-acquisition strategies

Article

Accepted Version

Standish, R. J. ORCID: <https://orcid.org/0000-0001-8118-1904>, Albornoz, F. E. ORCID: <https://orcid.org/0000-0001-9526-0945>, Morald, T. K., Hobbs, R. J. ORCID: <https://orcid.org/0000-0003-4047-3147> and Tibbett, M. ORCID: <https://orcid.org/0000-0003-0143-2190> (2021) Mycorrhizal symbiosis and phosphorus supply determine interactions among plants with contrasting nutrient-acquisition strategies. *Journal of Ecology*, 109 (11). pp. 3892-3902. ISSN 1365-2745 doi: 10.1111/1365-2745.13766 Available at <https://centaur.reading.ac.uk/100378/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/1365-2745.13766>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

**Phosphorus supply affects seedling growth of mycorrhizal but not cluster-root forming
jarrah-forest species**

Rachel J. Standish^{1*}, Matthew I. Daws², Tim K. Morald³, Jane Speijers⁴, John M. Koch⁵,
Richard J. Hobbs³ and Mark Tibbett^{2,3}

¹Harry Butler Institute, Murdoch University, 90 South Street, Murdoch Western Australia
6150, Australia

²Department of Sustainable Land Management and Soil Research Centre, School of
Agriculture Policy and Development, University of Reading, Berkshire, RG6 6AR, United
Kingdom

³School of Biological Sciences, The University of Western Australia, Crawley, Western
Australia 6009, Australia

⁴Department of Agriculture and Food, 3 Baron-Hay Court, South Perth Western Australia
6151, Australia

⁵Mining Environmental Department, Alcoa of Australia Ltd, PO Box 172, Pinjarra Western
Australia 6208, Australia

*Corresponding author:

Mark Tibbett

Address as above

Email: m.tibbett@reading.ac.uk

ORCID: Rachel J. Standish 0000-0001-8118-1904, Matthew I. Daws 0000-0002-9120-
4724, Richard J. Hobbs 0000-0003-4047-3147, Mark Tibbett 0000-0003-0143-2190.

Abstract

Aims: Fertiliser is often used to kick-start ecological restoration despite growing evidence of the potentially negative impacts on plant diversity. Jarrah (*Eucalyptus marginata*) forest species growing on nutrient (especially phosphorus) impoverished soils in southwestern Australia have a suite of adaptations for phosphorus (P) acquisition, including the formation of cluster roots, and associations with mycorrhizal fungi. Here we investigated how escalating P supply, along with a stoichiometric adjustment of nitrogen (N) supply, impacted the growth and nutrition of a wide range of jarrah forest seedlings.

Methods: In a pot experiment, we measured seedling biomass and nutritional responses of 12 jarrah forest species to a gradient of P supply in relation to N supply, and for the mycorrhizal species, inoculation with arbuscular mycorrhizal fungi.

Results: Three cluster-root forming species did not respond to increasing P, probably because they were reliant on seed P. Generally, mycorrhizal species showed a positive biomass response to increasing P when N was available. Mycorrhizas benefited seedling growth at low P (9 mg P added per kg of jarrah forest soil) when N was also available, and were parasitic to seedling growth at high P (243 mg P/ kg soil) without additional N.

Conclusions: These results highlight importance of P and N supply in determining the nature of the symbiosis between plants and mycorrhizal fungi. Since P supply has the potential to reduce plant growth, for a range of species, our results suggest careful consideration of fertiliser amounts for ecological restoration of ecosystems adapted to nutrient poor soils.

Key words Arbuscular mycorrhizal fungi, ecological restoration, mycorrhizal growth dependency, N and P co-limitation, plant-soil interactions, pot experiment.

53 **Declarations**

54 **Funding**

55 This research was funded by the Australian Research Council, Alcoa of Australia Ltd and BHP
56 Billiton Worsley Alumina Pty Ltd (ARC Linkage Project 668931). The completion of this
57 work was supported by the Building Outstanding Impact Support Programme H&F38:
58 *Restoring biodiversity to phosphorus sensitive forests* and Research England Grant: *Policy*
59 *change to halt biodiversity loss and restore sustainable ecosystems after mining.*

60 **Conflicts of interest/competing interests**

61 The authors have no conflicts of interest or competing interests to declare.

62 **Availability of data and material**

63 The raw data are available on request to the corresponding author.

64 **Code availability**

65 Not applicable

66 **Authors' contributions**

67 RJS, TKM, JMK, RJH and MT designed the experiment; RJS and TKM established and
68 maintained the experiment; MID, JS and RJS analysed the data; RJS and MID drafted the
69 manuscript, and all the authors contributed to the final draft.

Introduction

Ecological restoration is required to meet global biodiversity conservation goals (United Nations Sustainable Development Goals 2015). Ecosystems degraded by human activity pose a significant challenge for ecological restoration especially where degraded soils impact establishment of native vegetation (Bradshaw 1997). Fertiliser is prescribed for ecological restoration where there is insufficient quantity or availability of soil nutrients (Whisenant 1999). Consequently, the application of fertiliser is considered necessary, to replace nutrients lost in vegetation and soils, and best practice, to kickstart restoration of native plants and ecosystems after surface mining in Brazil, Canada, and Australia (Bizuti et al. 2020; Pinno et al. 2012; Tibbett 2010). Yet, in other restoration contexts the removal of soil nutrients is often required to kickstart restoration (e.g. abandoned farmlands, sites exposed to high atmospheric nitrogen deposition). Here, nutrient enrichment can favour some species over others and alter species composition compared with historical reference plant communities (e.g. Smits et al. 2008; Wassen et al. 2005). These latter findings suggest caution in the use of fertilisers for mine restoration.

Yet evidence from mine restoration is mixed. Fertiliser application can positively impact plant growth, especially trees, and reduce the risk of soil erosion by increasing plant cover (e.g. Ward et al. 1990). However, it can also negatively impact some plant species by increasing competition with highly nutrient-responsive species, such as non-native weeds (e.g. Daws et al. 2013; Norman et al. 2006; Nussbaumer et al. 2016). Further, in regions with highly weathered, nutrient-deficient soils, fertiliser application, particularly P, can negatively affect establishment of some native species (Lambers et al. 2008). For example, the addition of 26 kg P ha⁻¹ for restoration of South African fynbos resulted in increased mortality of native proteaceous species three years after its application (Holmes 2001). These cluster-root forming species are sensitive to P-toxicity when supplied with P concentrations above those that they experience in nature (de Campos et al. 2013; Handreck 1991; Lambers et al. 2002; Pang et al.

2010; Shane et al. 2004a; Williams et al. 2019), potentially due to the loss of low affinity transporter systems (Huang et al. 2011). In south western Australia, P-fertiliser addition also decreased the abundance of slow-growing resprouter species in jarrah forest restored after mining (Daws et al. 2013; 2019 a, b).

Physical, chemical, and biological components of soil, and their interactions, play important roles in ecosystem development on mined lands (Rowland et al. 2009). This is particularly true in weathered, nutrient-poor soils because of the reliance of many plants on mycorrhizal fungi to assist with nutrient acquisition, particularly P (Bowen 1981). As for cluster-root forming species, a range of mycorrhizal plant species adapted to grow on nutrient impoverished soils have a limited ability to regulate P-uptake when the P-supply is increased (Shane et al. 2004a). For example, the dominant tree species in the jarrah forest (jarrah, *Eucalyptus marginata*) exhibited visible symptoms of toxicity in the glasshouse after two pulses of P; shoot P concentrations were between 1.8 and 5.5 mg P g⁻¹ DW (Kariman et al. 2014). However, arbuscular mycorrhizal (AM) fungal associations can reduce the severity of P-toxicity in jarrah by reducing P-uptake (Kariman et al. 2014). Conversely, fertilisation with phosphate can diminish the benefits of AM associations to plants and in the extreme case, generate parasitic associations (Johnson et al. 1997; Standish et al. 2021). These studies suggest a greater understanding of fertiliser impacts on plant-soil dynamics is required for restoration of P-impoverished ecosystems.

Here, we add to growing body of research to inform best practice use of fertiliser in the ecological restoration of jarrah forest after bauxite mining in southwestern Australia. Until recently, P-fertiliser had been generously applied (up to 80 kg P ha⁻¹) to kickstart restoration (Koch 2007) despite the inherently nutrient-impoverished soils characteristic of the reference ecosystem (Handreck 1997). The rationale for the high application rates has been to replace nutrients that are lost during the mining process and to promote the establishment of nitrogen-fixing legumes, assuming these species would facilitate the return of other plant species (Grant

2006; Koch 2007). While P-fertilisation has usually resulted in the restoration of jarrah forest similar in diversity to that cleared for mining (Koch and Hobbs 2007), in some cases legumes can dominate the early successional forest and so alter the ecosystem trajectory (Grant 2006; Norman et al. 2006). In addition, a range of slow-growing, long-lived resprouter species tend to occur at reduced abundances following the application of P (Daws et al. 2019a). This negative effect of P fertiliser on the abundance of slow growing species can potentially be explained by either competition with vigorous P-responsive species, such as legumes (Dell et al. 1987; Stoneman et al. 1995; Daws et al. 2019 a, b), or direct negative effects of elevated P on growth rates and survival. Even a single initial application of fertiliser to restored sites, including the jarrah forest, can result in elevated plant-available soil phosphorus concentrations after 20 or more years (Addison et al. 2019; Banning et al. 2008; Smits et al. 2008; Spain et al. 2015). Therefore, any potential negative effects of P-addition on competitive dynamics may persist in the long-term.

In this large glasshouse study, we tested seedling responses of a range of jarrah forest species, including both AM species and non-mycorrhizal cluster-root forming species, to P amendment, with and without additional N supply, where AM species were grown in treatments with and without AM inoculation. Individual seedlings of each species were grown in pots in the absence of competition. Specifically, we tested the hypotheses that: (1) plant responses to P are dependent on species' nutrient acquisition strategy; (2) seedlings will exhibit P-sensitivity to high external P concentrations in the absence of mycorrhiza; (3) Mycorrhizal symbiosis will limit growth suppression at high external P concentrations; (4) N addition will effect species-specific P responses. That is, the growth responses to P will be dependent on the addition of N due to: the potential for N and P co-limitation, differential AM fungal responses to N, differences in plant species nutrient stoichiometry, or some combination of these factors. Results are discussed in relation to plant mineral nutrition and growth and their implications for ecological restoration.

Materials and methods

Experimental design

We used a factorial design to test the effect of increasing P supply on seedling growth responses of 12 jarrah forest species (Table 1). Five P-levels were used: 0, 9, 27, 81 and 243 mg P kg⁻¹ of dry jarrah forest soil; these were selected to span current and previous rates of P-fertiliser application for jarrah forest restoration after bauxite mining (i.e., 20–80 kg P ha⁻¹; Standish et al. 2015). Jarrah forest soil is low in plant-available P and N (Standish et al. 2008) and readily adsorbs phosphate (Bolan et al. 1983); the highest P-level was included to try to elicit P-toxicity. We added AM inoculum to half the pots containing plant species that form mycorrhizal associations; AM inoculum was not added to pots containing non-mycorrhizal species. The design also included N addition (+N)/no N addition (–N) and there were three replicates of each treatment: in total there were 630 pots [i.e., (5 P-levels × 9 mycorrhizal species × 2 AM-treatments × 2 N-treatments × 3 replicates) + (5 P-levels × 3 non-mycorrhizal species × 2 N-treatments × 3 replicates)]. Pots, each containing one native plant, were arranged in a completely randomised block design within a glasshouse.

We used topsoil (0–10 cm depth) collected from under the jarrah forest at Huntly, Western Australia (32°35'06" S, 116°06'44" E). These soils tend to be low in available phosphorus, and nitrogen, and be slightly acidic (Table S1). Sampled soil was passed through an 8 mm sieve, steamed twice at 80°C for three hours, and dried at 100 °C to eradicate naturally occurring mycorrhizal inocula. Then, for the AM treatment, bulk soil portions were inoculated (1 inoculum: 9 parts soil) and 1.3 kg portions were poured into pots measuring 8 cm × 8 cm × 18 cm (depth). The AM inoculum consisted of *Rhizophagus irregularis* (Błaszk., Wubet, Renker and Buscot) C. Walker and A. Schüßler 2010 (formerly *Glomus intraradices*), hyphae and colonised leek (*Allium porrum* L.) roots, grown in sterilised river sand. *Rhizophagus* species are common in jarrah forest soils (Brundrett and Abbott 1994) and we

selected this species for its ease of culture and ability to rapidly colonise roots of Western Australian native plants. For the treatment without AM, we added one-part sterilised river sand to nine parts soil. Species belonging to the Fabaceae usually form associations with soil micro-organisms that fix nitrogen, but the use of steamed soil meant that only a few seedlings formed root nodules.

Phosphorus was added to the pots at the time of planting, as KH_2PO_4 (Chem-Supply Pty Ltd, Gillman SA, Australia) because it is readily available to plants. To ensure a constant ionic background and balanced potassium levels, potassium chloride (KCl) was added in inverse proportions to KH_2PO_4 amendments. The KH_2PO_4 and the KCl were mixed into the dry soil. Nitrogen, 30 mg N kg^{-1} soil, was added as NH_4NO_3 dissolved in a modified (minus P) Long Ashton solution containing micronutrients (Cavagnaro et al. 2001), twice during the experiment.

Seedlings were grown from seeds collected within a 20 km radius of Huntly or Boddington (32°48'13.61"S 116°28'25.04"E), Western Australia. Seeds were either immersed in boiling water for 30 seconds, treated with aerosol smoke or soaked in smoke-water to stimulate germination and surface sterilised by immersing in 0.5 % (v/v) sodium hypochlorite solution for 10 mins. Seeds were placed on moistened filter paper in Petri dishes, and these were placed in a dark 15 °C constant-temperature room. The germinants were planted into the pots between 31 July and 15 August 2007. Soils were watered to field capacity (18.7 % DI water by weight) every four days. Alkathene polyethylene beads (Qenos Pty Ltd, Altona, Victoria, Australia) were added to the soil surface within each pot to minimise evaporation and to help prevent mycorrhizal contamination of non-mycorrhizal treatments. Germinants that did not emerge, and dead seedlings, were replaced within six weeks after 31 July 2007.

Response variables

We harvested species in size order from largest to smallest; we started harvesting *Acacia*
celastrifolia on 21st November 2007 and finishing harvesting *Xanthorrhoea gracilis* on 6th
February 2008. The temperature in the glasshouse ranged from 8–39 °C during the growth
period and seedlings were aged between 14 and 27 weeks at harvest. Plant shoots (i.e.
aboveground material) were oven-dried at 70 °C and weighed. Shoot P was determined by
digesting ground plant material in a mixture of 70% (v/v) nitric acid and concentrated
perchloric acid and measured by ICP-OES (Zarcinus 1984). We determined seed P using the
same methodology, on seeds leftover from our original collection, and seed N was determined
using a Vario Macro combustion analyser (Elemental Analysis GmbH, Hanau, Germany).

The roots were washed, dried with paper towel, and weighed; the cluster roots and the
nodules were counted. Then, ~0.1 g (fresh weight) of the fine roots were removed and stored in
50% (v/v) ethanol pending assessment of mycorrhizal colonisation. The remaining roots were
oven-dried at 70 °C and weighed; the root biomass includes an estimate of the dry weight of
the fine-roots sampled for mycorrhizal colonisation. The biomass of the cluster roots was
estimated by subtracting the loss-on-ignition (at 650 °C for 16 hrs) of organic matter from soil
only (~2%; Table S1) with that from cluster roots with soil attached. Raw data were visualised
in biomass response plots to allow comparison to other P response studies. Subsequently, we
modelled P response using linear mixed models and linear and curvilinear relationships were
fitted where appropriate (see below).

The fine-root samples were cleared in 10 % (w/v) KOH at room temperature for 5 days,
and then stained in a 5 % (v/v) black ink vinegar solution for 1 hr before being transferred to a
solution of lactoglycerol (Walker 2005). The dark roots of *A. fraseriana*, *E. marginata* and *X.*
gracilis were bleached in a 0.5 % (v/v) sodium hypochlorite solution prior to staining. The line
intercept method at 40× magnification was used to assess the percentage root length colonised
by *Rhizopagus irregularis* (Giovanetti and Mosse 1980).

Mycorrhizal growth dependency is defined as the ratio of the biomass of plants inoculated with AM to that of uninoculated plants; a mycorrhizal dependency >0 means that plant growth was enhanced by AM inoculation (Gerdemann 1975). The mycorrhizal growth dependency of each plant species was determined for each combination of P and N treatments using the equation from Plenchette et al. (1983):

$$MD = \frac{\text{Biomass mycorrhizal plant} - \text{biomass non-mycorrhizal plant}}{\text{Biomass mycorrhizal plant}} \times 100\%$$

We used the average biomass values of mycorrhizal and non-mycorrhizal seedlings for each of the treatment combinations.

Statistical analyses

The changes in seedling biomass in response to the P gradient were analysed using linear mixed models. These models are used to analyse data with more than one variance component, and which include both fixed and random effects (Galwey 2006). They are also appropriate for the analysis of unbalanced designs such as this one (i.e. three non-mycorrhizal species were not inoculated with AM). We fitted up to four regression lines for each species, corresponding to the factorial combinations of N and AM treatments. The y (biomass of shoot + root + clusters) and x variables (P-levels) were natural-log transformed prior to analysis. The biomass response to P (hereafter P-response) was modelled by fitting linear and quadratic functions to the transformed data sets. N-treatment, species nested within nutrient-acquisition strategy (mycorrhizas or cluster roots), and AM nested within mycorrhizal species were included as fixed effects; the replicates, second and higher order interactions were included as random effects. The models were fitted using the residual maximum likelihood (REML) procedure in GenStat (Payne et al. 2008). The Wald statistic was estimated to test for the significance of the terms and their interactions (Elston 1998). The highest order interactions were dropped from the final model.

We used a two-factor ANOVA to compare mycorrhizal growth dependency (MD) among N and P treatment combinations followed by a Fisher's Least Significant Difference test to determine the pairwise differences between the P-treatment means. The MD values of plants grown at 243 mg P kg⁻¹ soil were highly variable, so we excluded these from the analyses so that the remaining dataset conformed to the assumption of homoscedasticity. Data were reflected and/or log_e transformed prior to analysis. We also used a two-factor ANOVA to compare shoot P concentrations; the factors were N-treatment and nutrient acquisition strategy. We excluded non-mycorrhizal seedlings from these analyses, so the sample sizes were equal. Data were log_e transformed prior to analysis where necessary to meet the assumptions of ANOVA. These and the preceding ANOVAs were conducted using GenStat Version 11.1 (Payne et al. 2008).

Results

Mycorrhizal colonisation and cluster root formation

For AM-inoculated seedlings, the percentage length of fine roots colonised by *Rhizophagus irregularis* varied among species: *Xanthorrhoea gracilis* 0–1 %, *Eucalyptus marginata* 0–6 %, *Acacia celastriifolia* 0–37 %, *Acacia pulchella* 0–47 %, *Allocasuarina fraseri* 0–48 %, *Bossiaea ornata* 0–51 %, *Bossiaea pulchella* 0–51 %, *Bossiaea aquifolium* 0–68 %, *Phyllanthus calycinus* 0–90 %. Percentage AM colonisation varied widely within and among P/N treatment combinations for all species except *X. gracilis* (i.e., AM colonisation was consistently low), and except for percentage AM colonisation at the highest P supply which was consistently low for all species except *P. calycinus* (Figure 1).

There were no statistically significant effects of N addition and increasing P on the biomass of cluster roots, expressed as a proportion of the total root biomass, for the three cluster-root-forming species (data not shown).

Plant growth

Plant growth was strongly affected by N addition in all but one species, *X. gracilis* (Figure 2). Eight of the nine mycorrhizal species showed positive (linear or curvilinear) biomass responses to increasing P addition for at least one of the N/AM treatment combinations (Figure 3). For these species, the effect of P was constrained by the availability of N, and the biomass response to N addition was greater than the biomass response to inoculation with the AM fungus (Figure 2). The remaining four species—the three proteaceous species and *X. gracilis*—did not show any biomass response to increasing P with or without additional N (Figure 4). For the mycorrhizal species (incl. *X. gracilis*), seedlings grown in soil containing AM inoculum and additional N attained the highest biomass compared with the other N/AM treatment combinations. Without additional N, the biomass of mycorrhizal seedlings declined with increasing soil P for six of the nine mycorrhizal species (Figure 3, Table 2 and next section). Most of the interaction terms in the model were statistically significant (Table S2).

The biomass of the seedlings reflected their seed masses and seed nutrient reserves; in particular, *B. grandis* attained greater biomass than seedlings of the other species (Figure 2, Table 1). The two species with the smallest seeds had the highest mean root: shoot ratios ($\pm$ SE): 3.09 ± 0.85 and 1.86 ± 0.19 for non-mycorrhizal *P. calycinus* and *Bo. ornata* respectively, but then, other small-seeded species such as *Bo. aquifolium*, had comparatively low mean root: shoot ratios (0.34 ± 0.02 for non-mycorrhizal seedlings).

Mycorrhizal growth dependency

Mycorrhizal growth dependency (MD) values were generally higher when seedlings had access to N, although the reverse was true for seedlings grown at 0P, and seedlings grown at 9 mg P kg⁻¹ benefitted from AM more than seedlings grown at the other P concentrations (Table 2; MS = 2.58, $F_{1,64} = 5.93$, $P = 0.02$ (N-treatment); $F_{3,64} = 6.41$, $P = 0.001$ (P-treatment); $F_{3,64} = 2.92$, $P = 0.04$ (N-treatment $\times$ P-treatment)). The mean difference in MD values ($\pm$ 95% CIs) is

presented in Table 2. Growth benefits at 9 mg P kg⁻¹ were particularly evident for small-seeded species (Table 2). In most cases, AM were parasitic on seedlings grown at 243 mg P kg⁻¹ without additional N (Table 2).

Shoot P concentration

For the 12 study species, shoot P concentrations increased with increasing external P concentrations reaching a maximum of ca. 1-3 mg P g⁻¹ DW of shoot tissue (Figure 3). Across the five external P concentrations, shoot P concentration was significantly higher both for the AM species compared with the cluster root species (Figure 5; $F_{1,20} = 4.82$, $P = 0.04$) and when seedlings were grown without additional N (Figure 5; $F_{1,20} = 4.91$, $P = 0.04$).

Discussion

Responses to P depend on nutrient acquisition strategy

Seedling responses of the 12 jarrah forest species to P could be separated into three broad groups (Figure 6). Group 1 included the mycorrhizal species *Allocasuarina fraseri*, *Acacia celastriifolia*, *Ac. pulchella*, *Bossiaea aquifolium*, *Bo. ornata*, *Bo. pulchella*, *Eucalyptus marginata* and *Phyllanthus calycinus*. These species showed a positive growth response to increasing soil P when N was available and a generally negative growth response to P in the absence of additional N (Figure 6). The second seedling response evident among the 12 jarrah forest species was represented by *Xanthorrhoea gracilis*. It did not respond like the other mycorrhizal species. This species fits the classical description of plants from nutrient-poor soils: low growth rate and no nutritional response to increasing P supply (Chapin 1980). Its low growth rate relative to the other species we tested, constrained its ability to respond to increased soil P and N, although it benefited from its association with *Rhizophagus irregularis*, irrespective of P supply (Figure 6).

The third group was defined by the proteaceous species *Banksia grandis*, *Hakea prostrata* and *H. undulata*. These species exhibited no growth response to increasing soil P (Figure 6). Instead, seedling growth probably relied on seed nutrient reserves (Barrow 1977; Pate et al. 1990; Stock et al. 1990) which may also explain the limited response of shoot P concentration to increasing P. This seedling strategy is also evident among proteaceous fynbos shrubs growing on similarly nutrient-poor soils (Allsopp and Stock 1995). Using published data for root [P] (Pate et al. 1990), we estimated that the seed P reserves of *H. undulata* were almost depleted when the seedlings were harvested, whereas the larger seed P reserves of *B. grandis* were probably only half consumed at harvest. Soil P will become more important as the seed nutrient reserves are depleted (Mitchell and Allsopp 1984) and the seedlings become reliant on cluster roots for nutrient acquisition. For these species, an increasing P supply did not suppress the formation of cluster roots as might be expected (e.g. Lamont 1972b; Shane and Lambers 2005, 2006). This result is potentially explained by the strongly P-fixing properties of jarrah forest soil (Bolan et al. 1983; Handreck 1997). The phosphorus retention index of jarrah forest soils can be as high as 90 (Tibbett et al. 2020).

P-sensitivity to high external P concentrations in the absence of mycorrhiza

P-toxicity symptoms were not observed for any of the study species even at the highest amendment rate of 243 mg P kg⁻¹. The shoot P concentrations we recorded for the study species were consistently below those at which we would expect visible symptoms of P-toxicity to develop, i.e., <10 mg g⁻¹ DM (Lambers et al. 2002; Shane et al. 2004b). While there were no visible symptoms of P-toxicity for the study species, we observed a decline in growth increasing P supply for at least half of the species, most commonly in the absence of exogenous N supply. For some species e.g. *Allocasuarina fraserii*, total biomass was reduced at 243 mg P kg⁻¹ compared with 81 mg P kg⁻¹, and this occurred irrespective of AM inoculation.

These responses may be partly due to the high P-fixing properties of jarrah forest soil, causing the amended P to become partially or wholly unavailable to plants. While highly weathered and P-deplete, these soils contain large amounts of reactive iron and aluminium, which adsorbs available P, and this capacity can be assessed by the P retention index (Allen and Jeffery 1990). A P retention index of 90 for jarrah forest soils (Tibbett et al. 2020) is high by global standards and even compared to other Western Australian soils (Bolland and Russell 2010; Bolland and Windsor 2007). This high P retention index may explain the modest responses to amended P, which would only be partially available to plants. The precise quotient is impossible to estimate with any accuracy.

The reduction in growth at high P amendment has some similarities to the work of Pang et al. (2010), who showed growth reductions for several species of perennial legumes, including native Australian species. These authors used higher P amendments (up to 384 mg kg⁻¹) and grew the plants in a washed river sand, in which nearly all amended P would be readily available. Some species show growth declines from as low as 24 mg kg⁻¹ for Australian natives such as *Kennedia* spp. and perhaps we would have observed physiological toxicities and growth reduction at high P amendment if the plants were grown in sand culture. Importantly however, we show growth reductions are possible even when seedlings were grown in native high P-fixing jarrah forest soils, which is more relevant than sand for informing fertiliser application for jarrah forest restoration.

Mycorrhizas limit growth suppression at high external P concentrations

While AM fungal inoculation has been shown to reduce the effect of elevated P on visible symptoms of toxicity in *Eucalyptus marginata* (Kariman et al. 2014), our study showed no evidence of AM inoculation moderating its decline in biomass at 81 and 243 mg P kg⁻¹. Across species, the shoot P concentration at which biomass declined was approx. 3 mg g⁻¹ DM, which is similar to the concentrations reported by Williams et al. (2019) at which negative effects on

growth were observed for woody species from the Great Western Woodlands of Western Australia. However, shoot P concentrations closer to 10 mg g⁻¹ DM were required to trigger growth reductions in perennial legumes.

There were three consistent effects of increasing P supply on mycorrhizal growth dependency among the jarrah forest species: (i) a decline in percentage colonisation of fine roots by the AM fungus at high P supply (243 mg P kg⁻¹ soil); (ii) the AM fungus was more beneficial to plant growth at low P (9 mg P kg⁻¹ soil) than at the other P-levels (0, 27, 81 and 243 mg P kg⁻¹ soil) and; (iii) seedling dry mass was less at high P supply compared with lower external P concentrations, which we interpret evidence for the AM fungus being parasitic on seedlings (*sensu* Johnson et al. 1997). Although the levels of mycorrhizal colonisation that we observed were low, they were consistent with those observed in other studies of jarrah forest species (Kariman et al. 2014; Nazeri et al. 2014). The decline in colonisation with increasing P confirms a common observation for both natural and agricultural systems (Smith and Read 2008 but see Jasper et al. 1989). The transition from mutualism to parasitism with increasing P supply supports other studies that have reported the importance of soil nutrient availability for determining the nature of the relationship between plants and mycorrhizal fungi (Johnson et al. 2010; Johnson 2010; Neuenkamp et al. 2019). The trade balance model predicts mutualism in soils with low P and N and conversely, parasitism in soils with high soil P and N (Johnson 2010). We confirmed this prediction for one fungus, a species considered ‘ruderal’ in Grime’s CSR framework (Chagnon et al. 2013); the next step would be to determine the nature of the symbiosis for other jarrah forest fungi.

N addition will effect species specific P responses

The treatment that gave the most consistent response across all species was N addition to the P treatments. This treatment caused a large variation in the amended N:P ratio, ~ 8-fold difference compared with pots without applied N, with the difference being maintained

throughout the experiment because pots were watered to weight (i.e. nutrients did not leach). Positive seedling growth responses to P were generally observed with N addition and did not occur without N addition. While shoot P concentrations were consistently higher without N addition, they were still lower than the range at which reductions in growth have been observed in tree seedlings (e.g. Williams et al. 2019). This suggests two important aspects of nutrition at play. Firstly, in unamended soils (-N, and -P) both N and P may be growth limiting, and the addition of N ensures only P is growth-limiting and causes biomass to increase broadly in line with increasing P application rate in Group 1 (mycorrhizal) species, with some growth decline at the highest amendment rate. Secondly, in the absence of N addition, Group 1 (mycorrhizal) species showed a decline in response broadly in line with increasing P rate and concomitant with tissue P concentration. We postulate that this is due to a poor capacity to regulate P uptake, due to the loss of high affinity transporters (Huang et al. 2011), in these Gondwanan trees without the ability to dilute acquired P in greater biomass allowed by N addition. This finding also demonstrates the importance of N:P ratio in forest ecosystem development (e.g. Amazonas et al. 2011), which is often overlooked in the assessment of ecosystem nutrition for restoration (Tibbett et al. 2019).

Further studies are required to determine the mechanism behind the reduction in growth when P, but not N, were added. Nonetheless, this growth reduction has the potential to be ecologically significant in a field setting. Current practice in large areas of post-mining restoration in the jarrah forest is to apply single superphosphate without additional N (e.g. Worsley Alumina; George et al. 2006). Thus, while for each individual species in Figure 2, the area encompassed by the suite of lines likely represents the potential range of growth responses to applied N and P, the actual seedling growth response in the field may be closer to the P only treatments, i.e. a reduction in growth across a range of species, at least for the species that do not fix nitrogen.

Our findings are consistent with field-based studies of restored jarrah forest where mycorrhizal species generally have access to arbuscular mycorrhizal fungi and N-fixing species have access to rhizobia and *Frankia* (Jasper 2007). For example, Tibbett et al. (2020) found that in one-year old restored sites, species richness was higher when N was included in the applied fertiliser mix as opposed to applying just single superphosphate. Indeed, our current data suggests this observation may result from the reduced growth of some species when just P is applied. Daws et al. (2013) reported that after 2.5 years the proteaceous species *B. grandis* exhibited no growth response to a single, initial application of up to 40 kg P ha⁻¹, while the AM species jarrah doubled in size as P increased from 0 to 40 kg ha⁻¹ P, and the two N-fixing AM species *Acacia drummondii* and *A. lateriticola* exhibited an almost six-fold increase in plant size as applied P increased from 0 to 40 kg P ha⁻¹. Indeed, P-addition can result in legumes dominating the understorey in restored sites because of their ability to fix N and exploit P (e.g. Grant 2006). Tibbett et al. (2020) reported that restored mine sites which received a single application of superphosphate were dominated by the fast-growing legume *Ac. celastrifolia* in just one year, with negative effects on the abundance of slower growing resprouter species. In contrast, in the zero P control, *Ac. celastrifolia* growth was minimal and the abundance of resprouter species. Indeed, from the perspective of plant community composition, a single initial application of P can reduce the species richness and density of long-lived resprouter species, including species such as *X. gracilis* and the cluster root forming Proteaceae, for 15 or more years after the initiation of restoration (Daws et al. 2019 a, b).

Conclusions

The conundrum for practitioners working to restore native plant communities that have evolved in ecosystems with naturally nutrient-poor soils, such as those under jarrah forest, is to apply P-fertiliser in amounts that will facilitate rapid plant establishment without favouring some species more than others. Our results imply that higher rates of P-fertiliser (27–243 kg P ha⁻¹)

are likely to promote the dominance of P-responsive jarrah forest species, especially those that fix N. Our data also suggest that AM fungi will benefit seedling establishment in soils where N and P are available, and AM fungi will not benefit seedling establishment where N and P are either not available or conversely, where P is high.

Seedlings of jarrah forest species exhibited a range of responses to applied N and P, which potentially explain field observations in restored jarrah forest. In particular, the unresponsiveness to fertiliser addition of the cluster-root forming species and slow growing species such as *X. gracilis* suggests that, where fertiliser is applied to restored sites, seedlings of these species will be susceptible to competition from more P-responsive species. In addition, the reduction in growth of a range of the AM species when P was applied, without additional N, merits further investigation as it may negatively impact seedling establishment. Consequently, our results imply that applying little or no P-fertiliser may be optimal for the return of species belonging to these groups and their mycorrhizas. Increased knowledge of belowground communities and their linkages to aboveground processes, particularly during the critical phase of seedling establishment, may lead to more sophisticated tools for ecosystem restoration (Harris 2009).

Acknowledgements

This research was funded by the Australian Research Council, Alcoa of Australia Ltd and BHP Billiton Worsley Alumina Pty Ltd (ARC Linkage Project 668931). The completion of this work was supported by the Building Outstanding Impact Support Programme H&F38: *Restoring biodiversity to phosphorus sensitive forests* and Research England Grant: *Policy change to halt biodiversity loss and restore sustainable ecosystems after mining*. We are grateful to TD Lardner, B Sadeghzadeh, TL Edmonds-Tibbett, JM Greatrex, EL Walker, BA Stokes, EL Cromer and TL Luitingh for assistance with this research.

References

- Addison SL, Smaill SJ, Garrett LG, Wakelin SA (2019) Effects of forest harvest and fertiliser amendment on soil biodiversity and function can persist for decades. *Soil Biology and Biochemistry* 135:194–205. <https://doi.org/10.1016/j.soilbio.2019.05.006>
- Allen DG, Jeffery RC (1990) Methods of analysis of phosphorus in Western Australian soils. Report of Investigation No. 37, Chemistry Centre, East Perth, Australia.
- Allsopp N, Stock WD (1995) Relationships between seed reserves, seedling growth and mycorrhizal responses in 14 related shrubs (Rosidae) from a low-nutrient environment. *Functional Ecology* 9:248–254. <https://doi.org/10.2307/2390571>
- Amazonas NT, Martinelli LA, de Cássia Piccolo M, Rodrigues RR (2011) Nitrogen dynamics during ecosystem development in tropical forest restoration. *Forest Ecology and Management* 262:1551–1557. <https://doi.org/10.1016/j.foreco.2011.07.003>
- Banning NC, Grant CD, Jones DL, Murphy DV (2008) Recovery of soil organic matter, organic matter turnover and nitrogen cycling in a post-mining forest rehabilitation chronosequence. *Soil Biology and Biochemistry* 40:2021–2031. <https://doi.org/10.1016/j.soilbio.2008.04.010>
- Barrow NJ (1977) Phosphorus uptake and utilization by tree seedlings. *Australian Journal of Botany* 25:571–584. <https://doi.org/10.1071/BT9770571>
- Bell DT, Plummer JA, Taylor SK (1993) Seed germination ecology in southwestern Western Australia. *Botanical Review* 59:24–73. <https://doi.org/10.1007/BF02856612>
- Bolan NS, Robson AD, Barrow NJ (1983) Plant and soil factors including mycorrhizal infection causing sigmoidal response of plants to applied phosphorus. *Plant and Soil* 73:187–201. <https://doi.org/10.1007/BF02197715>
- Bolland MDA, Russell WK (2010) Changes in chemical properties of 48 intensively grazed, rain-fed dairy paddocks on sandy soils over 11 years of liming in south-western Australia. *Soil Research* 48:682–692. <https://doi.org/10.1071/SR09199>

508 Bolland MDA, Windsor DP (2007) Converting reactive iron, reactive aluminium, and
 509 phosphorus retention index (PRI) to the phosphorus buffering index (PBI) for sandy soils
 510 of south-western Australia. *Soil Research* 45:262–265. <https://doi.org/10.1071/SR07026>
 511 Bowen GD (1981) Coping with low nutrients. In: Pate JS, McComb AJ (eds) *The Biology of*
 512 *Australian Plants*. The University of Western Australia Press, Nedlands, pp 33–64.
 513 Bradshaw AD (1997) The importance of soil ecology in restoration science. In: Urbanska
 514 KM, Webb NR, Edwards PJ (eds) *Restoration ecology and sustainable development*.
 515 Cambridge University Press, Cambridge, pp 33–64.
 516 Brundrett MC, Abbott LK (1991) Roots of jarrah forest plants. I. Mycorrhizal associations of
 517 shrubs and herbaceous plants. *Australian Journal of Botany* 39:445–457.
 518 <https://doi.org/10.1071/BT9910445>
 519 Brundrett MC, Abbott LK (1994) Mycorrhizal fungus propagules in the jarrah forest. 1.
 520 Seasonal study of inoculum levels. *New Phytologist* 127:539–546.
 521 <https://doi.org/10.1111/j.1469-8137.1994.tb03972.x>
 522 Cavagnaro TR, Smith FA, Lorimer MF, Haskard KA, Ayling SM, Smith SE (2001)
 523 Quantitative development of Paris-type arbuscular mycorrhizas formed between
 524 *Asphodelus fistulosus* and *Glomus coronatum*. *New Phytologist* 149:105–113.
 525 <https://doi.org/10.1046/j.1469-8137.2001.00001.x>
 526 Chagnon PL, Bradley RL, Maherali H, Klironomos JN (2013) A trait-based framework to
 527 understand life history of mycorrhizal fungi. *Trends in Plant Science* 18:484–491.
 528 <http://dx.doi.org/10.1016/j.tplants.2013.05.001>
 529 Chapin III FS (1980) The mineral nutrition of wild plants. *Annual Review of Ecology and*
 530 *Systematics* 11:233–260. <https://doi.org/10.1146/annurev.es.11.110180.001313>
 531 Daws MI, Richardson C (2015) An operational scale assessment of the effects of a reduction in
 532 fertiliser application rates on plant species responses in jarrah forest restored after bauxite
 533 mining. Alcoa of Australia Ltd, Research Note No. 38.

534 Daws MI, Grigg AH, Tibbett M Standish RJ (2019a) Enduring effects of large legumes and
 535 phosphorus fertiliser on jarrah forest restoration 15 years after bauxite mining. *Forest*
 536 *Ecology and Management* 438:204-214. <https://doi.org/10.1016/j.foreco.2019.02.029>
 537 Daws MI, Grigg AH, Standish, R.J, Tibbett M (2019b) Applied phosphorus has long-term
 538 impacts on vegetation responses in restored jarrah forest, In: Fourie AB, Tibbett M (eds)
 539 *Proceedings of the 13th international conference on mine closure*. Australian Centre for
 540 Geomechanics, Perth, Australia pp 693–704.
 541 Daws MI, Standish RJ, Koch JM, Morald TK (2013) Nitrogen and phosphorus fertiliser
 542 regime affect jarrah forest restoration after bauxite mining in Western Australia. *Applied*
 543 *Vegetation Science* 16:610–618. <https://doi.org/10.1111/avsc.12046>
 544 de Campos MCR, Pearse SJ, Oliveira RS, Lambers H (2013) Downregulation of net
 545 phosphorus-uptake capacity is inversely related to leaf phosphorus-resorption proficiency
 546 in four species from a phosphorus-impooverished environment. *Annals of Botany* 111:445–
 547 454. <https://doi.org/10.1093/aob/mcs299>
 548 Dell B, Jones S, Wilson SA (1987) Phosphorus nutrition of jarrah (*Eucalyptus marginata*)
 549 seedlings. *Plant and Soil* 97:369–379. <https://doi.org/10.1007/BF02383227>
 550 Elston DA (1998) Estimation of denominator degrees of freedom of F-distributions for
 551 assessing Wald statistics for fixed-effect factors in unbalanced mixed models. *Biometrics*
 552 54:1085–1096. <https://doi.org/10.2307/2533859>
 553 George SJ, Tibbett M, Braimbridge MF, Davis SG, Vlahos S, Ryan M (2006) Phosphorus
 554 fertiliser placement and seedling success in Australian jarrah forest. In: Fourie A, Tibbett
 555 M (eds) *Proceedings of the first international seminar on mine closure*, Australian Centre
 556 for Geomechanics, Perth Australia, pp 341–349.
 557 Gerdemann JW (1975) Vesicular-arbuscular mycorrhizae. In: Torrey JG, Clarkson DT (eds)
 558 *The development and function of roots*. Academic Press, New York, pp 575–591.

559 Giovanetti M, Mosse B (1980) An evaluation of techniques for measuring VAM infection in
 560 roots. *New Phytologist* 84:489–500. <https://www.jstor.org/stable/2432123>
 561 Grant CD (2006) State-and-transition successional model for bauxite mining rehabilitation in
 562 the jarrah forest of Western Australia. *Restoration Ecology* 14:28–37.
 563 <https://doi.org/10.1111/j.1526-100X.2006.00102.x>
 564 Handreck KA (1997) Phosphorus requirements of Australian native plants. *Australian Journal*
 565 *of Soil Research* 35:241–289. <https://doi.org/10.1071/S96060>
 566 Handreck KA (1991) Interactions between iron and phosphorus in the nutrition of *Banksia*
 567 *ericifolia* L. f. var. *ericifolia* (Proteaceae) in soil-less potting media. *Australian Journal of*
 568 *Botany* 39:373–384. <https://doi.org/10.1071/BT9910373>
 569 Harris JA (2009) Soil microbial communities and restoration ecology: facilitators or followers?
 570 *Science* 325:573–574. <https://doi.org/10.1126/science.1172975>
 571 Holmes PM (2001) Shrubland restoration following woody alien invasion and mining: effects
 572 of topsoil depth, seed source, and fertilizer addition. *Restoration Ecology* 9:71–84.
 573 <https://doi.org/10.1046/j.1526-100x.2001.009001071.x>
 574 Huang CY, Shirley N, Genc Y, Shi S, Langridge P (2011) Phosphate utilization efficiency
 575 correlates with expression of low-affinity phosphate transporters and noncoding RNA,
 576 IPS1, in barley. *Plant Physiology* 156:1217–1229. <https://doi.org/10.1104/pp.111.178459>
 577 Jasper DA (2007) Beneficial soil microorganisms of the jarrah forest and their recovery in
 578 bauxite mine restoration in southwestern Australia. *Restoration Ecology* 15:S74–S84.
 579 <https://doi.org/10.1111/j.1526-100X.2007.00295.x>
 580 Jasper DA, Abbott LK, Robson AD (1989) Acacias respond to additions of phosphorus and to
 581 inoculation with VA mycorrhizal fungi in soils stockpiled during mineral sand mining.
 582 *Plant and Soil* 115:99–108. <https://doi.org/10.1007/BF02220699>

583 Johnson NC (2010) Resource stoichiometry elucidates the structure and function of arbuscular
584 mycorrhizas across scales. *New Phytologist* 185:631–647. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8137.2009.03110.x)
585 [8137.2009.03110.x](https://doi.org/10.1111/j.1469-8137.2009.03110.x)

586 Johnson NC, Graham JH, Smith FA (1997) Functioning of mycorrhizal associations along the
587 mutualism–parasitism continuum. *New Phytologist* 135:575–585.
588 <https://doi.org/10.1046/j.1469-8137.1997.00729.x>

589 Johnson NC, Wilson GWT, Bowker MA, Wilson JA, Miller RM (2010) Resource limitation is
590 a driver of local adaptation in mycorrhizal symbioses. *Proceedings of the National*
591 *Academy of Sciences* 107:2093–2098. <https://doi.org/10.1073/pnas.0906710107>

592 Kariman K, Barker SJ, Finnegan PM, Tibbett M (2014) Ecto- and arbuscular mycorrhizal
593 symbiosis can induce tolerance to toxic pulses of phosphorus in jarrah (*Eucalyptus*
594 *marginata*) seedlings. *Mycorrhiza* 24:501–509. [https://doi.org/10.1007/s00572-014-0567-](https://doi.org/10.1007/s00572-014-0567-6)
595 [6](https://doi.org/10.1007/s00572-014-0567-6)

596 Koch JM (2007) Alcoa’s mining and restoration process in south Western Australia.
597 *Restoration Ecology* 15:S11–S16. <https://doi.org/10.1111/j.1526-100X.2007.00288.x>

598 Koch JM, Hobbs RJ (2007) Synthesis: is Alcoa successfully restoring a jarrah forest ecosystem
599 after bauxite mining in Western Australia? *Restoration Ecology* 15:S137–S144.
600 <https://doi.org/10.1111/j.1526-100X.2007.00301.x>

601 Lambers H, Juniper D, Cawthray GR, Veneklaas EJ, Martínez-Ferri E (2002) The pattern of
602 carboxylate exudation in *Banksia grandis* (Proteaceae) is affected by the form of
603 phosphate added to the soil. *Plant and Soil* 238:111–122.
604 <https://doi.org/10.1023/A:1014289121672>

605 Lambers H, Raven JA, Shaver GR, Smith SE (2008) Plant nutrient-acquisition strategies
606 change with soil age. *Trends in Ecology and Evolution* 23:95–103.
607 <https://doi.org/10.1016/j.tree.2007.10.008>

608 Lamont BB (1972a) The effect of soil nutrients on the production of proteoid roots by *Hakea*
609 species. Australian Journal of Botany 20:27–40. <https://doi.org/10.1071/BT9720027>

610 Lamont BB (1972b) The morphology and anatomy of proteoid roots in the genus *Hakea*.
611 Australian Journal of Botany 80:155–174. <https://doi.org/10.1071/BT9720155>

612 Malajczuk N, Bowen GD (1974) Proteoid roots are microbially induced. Nature 251:316–317.
613 <https://doi.org/10.1038/251316a0>

614 Mitchell DT, Allsopp N (1984) Changes in the phosphorus composition of seeds of *Hakea*
615 *sericea* (Proteaceae) during germination under low phosphorus conditions. New
616 Phytologist 96:239–247. <https://doi.org/10.1111/j.1469-8137.1984.tb03560.x>

617 Nazeri, NK, Lambers H, Tibbett M, Ryan MH (2014) Moderating mycorrhizas: arbuscular
618 mycorrhizas modify rhizosphere chemistry and maintain plant phosphorus status within
619 narrow boundaries. Plant, Cell and Environment 37:911–921.
620 <https://doi.org/10.1111/pce.12207>

621 Neuenkamp L, Prober SM, Price JN, Zobel M, Standish RJ (2019) Benefits of mycorrhizal
622 inoculation to ecological restoration depend on plant functional type, restoration context
623 and time. Fungal Ecology 40:140–149. <https://doi.org/10.1016/j.funeco.2018.05.004>

624 Norman MA, Koch JM, Grant CD, Morald TK, Ward SC (2006) Vegetation succession after
625 bauxite mining in Western Australia. Restoration Ecology 14:278–288.
626 <https://doi.org/10.1111/j.1526-100X.2006.00130.x>

627 Nussbaumer Y, Cole MA, Offler CE, Patrick JW (2016) Identifying and ameliorating nutrient
628 limitations to reconstructing a forest ecosystem on mined land. Restoration Ecology
629 24:202–211. <https://doi.org/10.1111/rec.12294>

630 Pang JY, Tibbett M, Denton MD, Lambers H, Siddique KHM, Bolland MDA, Revell CK,
631 Ryan MH (2010) Variation in seedling growth of 11 perennial legumes in response to
632 phosphorus supply. Plant and Soil 328:133–143. [https://doi.org/10.1007/s11104-009-](https://doi.org/10.1007/s11104-009-0088-9)
633 0088-9

634 Pate JS, Froend RH, Bowen BJ, Hansen A, Kuo J (1990) Seedling growth and storage
635 characteristics of seeder and resprouter species of Mediterranean-type ecosystems of SW
636 Australia. *Annals of Botany* 65:585–601.
637 <https://doi.org/10.1093/oxfordjournals.aob.a087976>

638 Payne RW, Harding SA, Murray DA, Soutar DM, Baird DB et al. (2008) GenStat Release 11.1
639 (PC/ Windows) Copyright. VSN International Ltd, Hetfordshire.

640 Pinno BD, Landhäusser SM, MacKenzie MD, Quideau SA, Chow PS (2012) Trembling aspen
641 seedling establishment, growth and response to fertilization on contrasting soils used in oil
642 sands reclamation. *Canadian Journal of Soil Science* 92:143–151.
643 <https://doi.org/10.4141/cjss2011-004>

644 Plenchette C, Fortin JA, Furlan V (1983) Growth response of several plant species to
645 mycorrhizae in a soil of moderate P-fertility I. Mycorrhizal dependency under field
646 conditions. *Plant and Soil* 70:199–209. <https://doi.org/10.1007/BF02374780>

647 Rowland SM, Prescott CE, Grayston SJ, Quideau SA, Bradfield GE (2009) Recreating a
648 functioning forest soil in reclaimed oil sands in northern Alberta: an approach for
649 measuring success in ecological restoration. *Journal of Environmental Quality* 38:1580–
650 1590. <https://doi.org/10.2134/jeq2008.0317>

651 Shane MW, Lambers H (2005) Cluster roots: a curiosity in context. *Plant and Soil* 274:101–
652 125. <https://doi.org/10.1007/s11104-004-2725-7>

653 Shane MW, Szota C, Lambers H (2004a) A root trait accounting for the extreme phosphorus
654 sensitivity of *Hakea prostrata* (Proteaceae). *Journal of Experimental Botany* 27:991–1004.
655 <https://doi.org/10.1111/j.1365-3040.2004.01204.x>

656 Shane MW, McCully ME, Lambers H (2004b) Tissue and cellular phosphorus storage during
657 development of phosphorus toxicity in *Hakea prostrata* (Proteaceae). *Journal of*
658 *Experimental Botany* 55:1033–1044. <https://doi.org/10.1093/jxb/erh111>

659 Shane MW, Lambers H (2006) Systematic suppression of cluster-root formation and net P-
660 uptake rates in *Grevillea crithmifolia* at elevated P supply: a proteacean with resistance for
661 developing symptoms of ‘P toxicity’. *Journal of Experimental Botany* 57:413–423.
662 <https://doi.org/10.1093/jxb/erj004>

663 Smits NAC, Willems JH, Bobbink R (2008) Long-term after-effects of fertilisation on the
664 restoration of calcareous grasslands. *Applied Vegetation Science* 11:279–286.
665 <https://doi.org/10.3170/2008-7-18417>

666 Smith SE, Read DJ (2008) Mycorrhizal symbiosis, 3rd edn. Academic Press, London.

667 Spain AV, Tibbett M, Hinz DA, Ludwig JA, Tongway DJ (2015) The mining-restoration
668 system and ecosystem development following bauxite mining in a biodiverse environment
669 of the seasonally dry tropics, Northern Territory, Australia. In: Tibbett M (ed) *Mining in*
670 *Ecologically Sensitive Landscapes*. CRC Press, Netherlands, pp 159–227.

671 Standish RJ, Morald TK, Koch JM, Hobbs RJ, Tibbett M (2008) Restoring jarrah forest after
672 bauxite mining in Western Australia—the effect of fertilizer on floristic diversity and
673 composition. In: Fourie AB, Tibbett M, Weiersbye IM, Dye PJ (eds) *Mine Closure 2008*.
674 Australian Centre for Geomechanics, Perth, Australia, pp 717–725.

675 Standish RJ, Daws MI, Gove AD, Didham RK, Grigg AH, Koch JM, Hobbs RJ (2015) Long-
676 term data suggest jarrah-forest establishment at restored mine sites is resistant to climate
677 variability. *Journal of Ecology* 103:78–89. <https://doi.org/10.1111/1365-2745.12301>

678 Standish RJ, Albornoz F, Morald TK, Hobbs RJ, Tibbett M (2021) Mycorrhizal symbiosis and
679 phosphorus supply determine interactions among plants with contrasting nutrient
680 acquisition strategies. *Journal of Ecology*, accepted 9 August 2021.

681 Standish RJ, Cramer VA, Hobbs RJ, Kobryn HT (2006) Legacy of land-use evident in soils of
682 Western Australia’s wheatbelt. *Plant and Soil* 280:189–207.
683 <https://doi.org/10.1007/s11104-005-2855-6>

684 Stock WD, Pate JS, Delfs J (1990) Influence of seed size and quality on seedling development
685 under low nutrient conditions in five Australian and South African members of the
686 Proteaceae. *Journal of Ecology* 78:1005–1020. <https://doi.org/10.2307/2260949>

687 Stoneman GL, Dell B, Turner NC (1995) Growth of *Eucalyptus marginata* (jarrah) seedlings
688 in Mediterranean-climate forest in south-west Australia in response to overstorey, site and
689 fertiliser application. *Forest Ecology and Management* 79:173–184.
690 [https://doi.org/10.1016/0378-1127\(95\)03608-3](https://doi.org/10.1016/0378-1127(95)03608-3)

691 Tibbett M (2010) Large-scale mine site restoration of Australian eucalypt forests after bauxite
692 mining: soil management and ecosystem development. In: Batty LC, Hallberg K (eds)
693 Ecology of industrial pollution. Cambridge University Press, Cambridge, pp 309–326.

694 Tibbett M, Daws MI, George SJ, Ryan MH (2020) The where, when and what of phosphorus
695 fertilisation for seedling establishment in a biodiverse jarrah forest restoration after bauxite
696 mining in Western Australia. *Ecological Engineering* 153:105907.
697 <https://doi.org/10.1016/j.ecoleng.2020.105907>

698 Tibbett M, O'Connor R, Daws MI (2019) Too much of a good thing: phosphorus over-
699 fertilisation in rehabilitated landscapes of high biodiversity value. In: Fourie AB, Tibbett
700 M (eds) Proceedings of the 13th International Conference on Mine Closure. Australian
701 Centre for Geomechanics, Perth, Australia, pp 651–666.

702 United Nations Sustainable Development Goals (2015) Transforming our world: the 2030
703 Agenda for Sustainable Development. Agenda adopted by the 193 Member States of the
704 United Nations in New York on 25 September 2015. Available at
705 <https://sustainabledevelopment.un.org/post2015/transformingourworld/publication>
706 (accessed 20 April 2021).

707 Walker C (2005) A simple blue staining technique for arbuscular mycorrhizal and other root-
708 inhabiting fungi. *Inoculum* 56:68–69.

- Ward SC, Koch JM, Nichols OG (1990) Bauxite mine rehabilitation in the Darling Range,
Western Australia. *Proceedings of the Ecological Society of Australia* 16:557–565.
- Wassen MJ, Venterink HO, Lapshina ED, Tanneberger F (2005) Endangered plants persist
under phosphorus limitation. *Nature* 437:547–550. <https://doi.org/10.1038/s41559-020-01323-w>
- Western Australian Herbarium. 1998–. FloraBase — The Western Australian Flora.
Department of Environment and Conservation, Perth, Western Australia. Available at
<http://florabase.dec.wa.gov.au/> (Accessed 31 March 2010).
- Whisenant SG (1999) Repairing damaged wildlands: a process-orientated, landscape-scale
approach. Cambridge University Press, Cambridge.
- Williams A, George S, Birt H, Daws MI, Tibbett M (2019) Sensitivity of seedling growth to
phosphorus supply in six tree species of the Australian Great Western Woodlands.
Australian Journal of Botany 67:390–396. <https://doi.org/10.1071/BT18247>
- Zarcinas BA (1984) Analysis of soil and plant material by inductively coupled plasma-optical
spectrometry: comparison of digestion procedures for major and trace constituents in plant
material. CSIRO Division of Soils Divisional Report No. 70, Canberra, Australia.

Table 1. Traits of 7 of the 12 jarrah forest plants included in this pot study. Species nomenclature follows that of the Western Australian Herbarium (1998–); life forms after Bell et al. (1993). ‡ Mycorrhizal associations recorded by Brundrett and Abbott (1991). †Cluster roots recorded by Lamont (1972a) and Malajczuk and Bowen (1974). Mass (mean $\pm$ SE; $n=10$) and nutrient contents per seed estimated from bulk samples. Plants belonging to the Fabaceae and Casuarinaceae form associations with nitrogen-fixing microorganisms. Data for *Acacia celastrifolia*, *Bossiaea aquifolium*, *Eucalyptus marginata*, *Hakea undulata* and *Phyllanthus calycinus* is included in Standish et al. (2021) except seed N, which is 0.97 mg, 0.48 mg, 0.81 mg, 2.05 mg and 0.06 mg, respectively.

Species	Family	Life form	P acquisition strategy	Seed mass (mg/seed)	Seed P (mg/seed)	Seed N (mg/seed)
<i>Allocasuarina fraseriana</i>	Casuarinaceae	Tree	AM/ECM‡	5 $\pm$ 0.5	0.03	0.19
<i>Acacia pulchella</i>	Fabaceae	Shrub	AM/ECM‡	11 $\pm$ 0.8	0.03	0.55
<i>Bossiaea ornata</i>	Fabaceae	Shrub	AM‡	3 $\pm$ 0.2	0.01	0.18
<i>Bossiaea pulchella</i>	Fabaceae	Shrub	AM	18 $\pm$ 0.9	0.05	0.70
<i>Banksia grandis</i>	Proteaceae	Small tree	Cluster roots†	78 $\pm$ 3.7	0.81	9.99
<i>Hakea prostrata</i>	Proteaceae	Shrub	Cluster roots†	44 $\pm$ 3.2	0.45	4.39
<i>Xanthorrhoea gracilis</i>	Xanthorrhoeaceae	Sub-shrub	AM‡	14 $\pm$ 1.2	0.07	0.57

Table 2. Mycorrhizal dependency (MD) values for jarrah forest plants grown with and without additional nitrogen (+N and –N respectively) at five P-levels. Extreme values are indicated in bold: both beneficial (MD >50) and parasitic (MD <–50) associations. ND = not determined as seedlings in the –N +AM treatment died.

Species (–N)	0P	9P	27P	81P	243P
<i>Allocasuarina fraseriana</i>	22.0	77.4	34.4	44.9	ND
<i>Acacia celastriifolia</i>	–7	–42.2	–33.7	–16.2	–13.7
<i>Acacia pulchella</i>	33.5	20.5	6.5	–5.8	–159.3
<i>Eucalyptus marginata</i>	4.6	50.7	39.5	17.7	–5.5
<i>Phyllanthus calycinus</i>	–34.0	–17.9	42.3	–98.5	–198.6
<i>Bossiaea aquifolium</i>	43.9	59.1	57.9	20.0	–7.6
<i>Bossiaea ornata</i>	48.6	39.2	–1.2	–50	ND
<i>Bossiaea pulchella</i>	61.3	48.9	61.7	20.2	–36.5
<i>Xanthorrhoea gracilis</i>	20.0	39.2	20.1	9.5	33.1
Means ± SE	21.4 ± 10.0	30.5 ± 12.7	25.3 ± 10.2	–6.5 ± 14.6	–55.4 ± 33.0
Species (+N)	0P	9P	27P	81P	243P
<i>Allocasuarina fraseriana</i>	14.4	61.7	28.9	32.5	17.3
<i>Acacia celastriifolia</i>	24.6	25.7	28.7	10.6	–9.0
<i>Acacia pulchella</i>	–32.6	59.7	29.1	54.2	32.9
<i>Eucalyptus marginata</i>	–25.8	20.9	69.2	33.2	40.7
<i>Phyllanthus calycinus</i>	–73.6	96.6	–42.1	22.9	52.8
<i>Bossiaea aquifolium</i>	–2.3	77.8	70.0	44.0	71.1
<i>Bossiaea ornata</i>	30.7	93.1	66.7	45.9	69.8
<i>Bossiaea pulchella</i>	51.1	85.2	55.0	19.0	–259.4
<i>Xanthorrhoea gracilis</i>	23.2	26.6	47.0	36.2	21.3
Means ± SE	1.1 ± 12.9	60.8 ± 10.0	39.2 ± 11.7	33.2 ± 4.6	1.2 ± 34.0




Figure 1. Percentage colonisation of fine roots by arbuscular mycorrhizal fungi at 0, 9, 27, 81 and 243 mg added P per kg dry soil, with and without additional nitrogen (30 mg per kg of soil), 14–27 weeks after seedlings were inoculated with *Rhizophagus irregularis*; ○ = *P. calycinus* and ● = data for eight other mycorrhizal species; $n = 3$ per species per N/P treatment combination.




Figure 2. Raw biomass response of jarrah forest seedlings to increasing soil P supply, with and without additional nitrogen (30 mg per kg of soil), and without and without inoculation with AM fungi. Values are means per treatment combination ($n = 3$) where ● = +N+AM; ○ = +N-AM; ■ = -N+AM and □ = -N-AM. Data for 0P are offset for clarity.




Figure 3. Modelled biomass response of jarrah forest seedlings to increasing [P]: 0, 9, 27, 81 and 243 mg added P per kg dry jarrah forest soil, with and without additional nitrogen (30 mg per kg of soil), and without and without inoculation with AM fungi. Values are means per treatment combination ($n = 3$) where: ● = +N +AM; ○ = +N -AM; ■ = -N +AM and □ = -N -AM. Linear and curvilinear relationships were fitted where appropriate ($P < 0.02$ for all; Table S2).




Figure 4. Shoot P concentrations of jarrah forest seedlings at 5 levels of soil P supply, with and without additional nitrogen (30 mg per kg of soil) and without and without inoculation with AM fungi. Values are means per treatment combination ($n = 3$) where $\bullet = +N+AM$; $\circ = +N-AM$; $\blacksquare = -N+AM$ and $\square = -N-AM$.




Figure 5. Shoot P concentration for 12 jarrah forest species, with and without additional nitrogen (30 mg per kg of soil), and without and without inoculation with AM fungi. AM = mycorrhizal seedlings (filled symbols); R = roots alone and CR = cluster roots (open symbols). Values are means $\pm$ SE of five external P concentrations, except for *Al. fraseri* R and *Bo. ornata* R at 243P as these seedlings died. Dashed lines are means for AM, R and CR seedlings, respectively.




Figure 6. Schematic representation of the three distinct nutritional responses of species' suggested by our data. Mycorrhizal species are represented by circles and N-fixing mycorrhizal species are represented by squares; with the sizes of each symbol relative to species' mycorrhizal dependency at 9 kg phosphorus (P) per kg of soil and with additional nitrogen (N). Cluster-root forming species are represented by triangles. Group 1: Positive growth response to P when N available; Group 2 No response to P with or without N; Group 3: No response to P with or without N. BA = *Bossiaea aquifolium*, BO = *Bo. ornata*, AF = *Allocasuarina fraseriana*, AP = *Acacia pulchella*, BP = *Bo. pulchella*, PC = *Phyllanthus calycinus*, AC = *Ac. celastriifolia*, *Eucalyptus marginata* is the unlabelled amber coloured circle in Group 1. *Xanthorrhoea gracilis* is unlabelled blue circle in Group 2. BG = *Banksia grandis*, HU = *Hakea undulata* and HP = *H. prostrata*.

Supplementary information

Table S1. Properties of jarrah forest soils (0-10 cm) depth at Huntly, southwestern Australia. Data are means $\pm$ SE ($n = 12$). Properties determined by CSBP Soil and Plant Laboratories (Bibra Lake, Perth) using methods described in Standish et al. (2006).

NH ₄ -N	3.42 $\pm$ 0.63 mg/kg
NO ₃ -N	< 1 mg/kg (below limit of detection)
Plant available phosphorus (Colwell)	1.70 $\pm$ 0.51 mg/kg
Potassium	35.33 $\pm$ 3.01 mg/kg
Sulphur	5.16 $\pm$ 0.70 mg/kg
Organic Carbon	2.00 $\pm$ 0.07 g/100g
Conductivity	0.02 $\pm$ 0.00 dS/m
pH (in CaCl ₂)	5.39 $\pm$ 0.05

Table S2. Results of linear mixed model of the effect of increasing [P] on the biomass of 12 jarrah forest species grown in pots with and without additional nitrogen (Ntt) and arbuscular mycorrhizal fungi (Mtt). Strategy refers to P-acquisition assisted by mycorrhizas or cluster roots.

Fixed term	Wald statistic (n.d.f.)	F statistic (d.d.f.)	P-value
Strategy	1277.15 (1)	1277.15 (83.1)	< 0.001
Strategy × species	735.9 (10)	73.6 (82.3)	< 0.001
Strategy × species × Mtt	54.4 (9)	6.04 (82.6)	< 0.001
Log _e (Ptt + 1) (linear effect)	45.5 (1)	45.5 (81.4)	< 0.001
C × [Log _e (Ptt + 1)] ² (quadratic effect)	4.34 (1)	4.34 (83.1)	0.04
Ntt	1745.4 (1)	1745.4 (83)	< 0.001
Strategy × Log _e (Ptt + 1)	4.17 (1)	4.17 (82.2)	0.04
Strategy × C × [Log _e (Ptt + 1)] ²	2.33 (1)	2.33 (82.3)	0.13
Strategy × Ntt	89.8 (1)	89.8 (84.7)	< 0.001
Ntt × Log _e (Ptt + 1)	237.2 (1)	237.2 (81.9)	< 0.001
Ntt × C × [Log _e (Ptt + 1)] ²	21.5 (1)	21.5 (84.1)	< 0.001
Strategy × species × Log _e (Ptt + 1)	50.7 (10)	5.07 (81.8)	< 0.001
Strategy × species × C × [Log _e (Ptt + 1)] ²	17.1 (10)	1.71 (83.4)	0.09
Strategy × species × Ntt	245.1 (10)	24.51 (83.5)	< 0.001
Strategy × species × Mtt × Log _e (Ptt + 1)	9.61 (9)	1.07 (81.8)	0.40
Strategy × species × Mtt × C × [Log _e (Ptt + 1)] ²	24.4 (9)	2.71 (83.9)	0.008
Strategy × species × Mtt × Ntt	28.3 (9)	3.14 (83.9)	0.003
Strategy × Ntt × Log _e (Ptt + 1)	28.6 (1)	28.6 (83.6)	< 0.001
Strategy × Ntt × C × [Log _e (Ptt + 1)] ²	0.89 (1)	0.89 (83.8)	0.35
Strategy × species × Ntt × Log _e (Ptt + 1)	69.79 (10)	6.98 (82.3)	< 0.001
Strategy × species × Ntt × C × [Log _e (Ptt + 1)] ²	32.02 (10)	3.20 (84.4)	0.002
Strategy × species × Mtt × Ntt × Log _e (Ptt + 1)	16.48 (9)	1.83 (82.1)	0.075
Strategy × species × Mtt × Ntt × C × [Log _e (Ptt + 1)] ²	5.34 (9)	0.59 (85.1)	0.80