



Are the invasive *Acacia melanoxylon* and *Eucalyptus globulus* drivers of other species invasion? Testing their allelochemical effects on germination

Sheila F. Riveiro¹ · Óscar Cruz² · Otilia Reyes¹

Received: 9 February 2022 / Accepted: 11 September 2023 / Published online: 16 September 2023
© The Author(s) 2023

Abstract

Many invasive alien species (IAS) produce secondary metabolites that affect how other plants function (allelopathic compounds) and can drive other species invasion, as proposed by the invasional meltdown hypothesis. *Acacia melanoxylon* and *Eucalyptus globulus* are two of such species. In this study, we analyzed the germination response of seven IAS (*Acacia dealbata*, *Acacia mearnsii*, *Acacia melanoxylon*, *Acacia longifolia*, *Eucalyptus globulus*, *Paraserianthes lophantha*, *Phytolacca americana*) and a native biotest species (*Lactuca sativa*) to the application of two different aqueous extracts at two different concentrations of donor species *A. melanoxylon* and *E. globulus*. Extract compounds were identified by UHPLC-ESI-QTOF-MS. *Eucalyptus* aqueous extracts significantly reduced germination in three species (*A. dealbata*, *E. globulus*, *P. americana*). The germination of all the species tested was reduced with acacia aqueous extracts. Our results support the postulates of the Biochemical Recognition Hypothesis in that seeds gauge establishment potential based on phytochemical release of other plants. Furthermore, *A. melanoxylon* and *E. globulus* lowered their own germination, suggesting that these species exhibit intraspecific biochemical recognition. We also found support for the Novel Weapons Hypothesis in the case of *L. sativa* as a native species. Our research shows that phytochemicals are a component of plant-plant interactions, including the invasion process.

Keywords Allelopathy · Biochemical recognition hypothesis · Invasional meltdown hypothesis · Novel weapons hypothesis · Invasive alien species · Seed germination

✉ Sheila F. Riveiro
sheila.riveiro@usc.es

¹ BIOAPLIC group. Área de Ecología, Dpto. de Biología Funcional, Facultade de Biología, Universidade de Santiago de Compostela, Santiago de Compostela 15782, Spain

² BIOAPLIC group. Área de Ecología, Dpto. de Biología Funcional, Universidade de Santiago de Compostela, EPSE Campus Terra, Lugo 27002, Spain

Introduction

Invasive alien species (IAS) are a threat to global biodiversity, because they outcompete native species, change the physico-chemical characteristics of soils (e.g., modification of the soil microbial activity—Qu et al. 2021; allelochemical release that reduces native plant establishment—Zhang et al. 2021), alter the rate of nutrient cycling (Vilà et al. 2011) and modify ecosystem fire regimes (D’Antonio and Vitousek 1992; Gaertner et al. 2014). Plant invaders reduce the richness and abundance of native species (Pyšek et al. 2012; Wohlgemuth et al. 2022) by modulating seed germination, inhibit seedling establishment and growth (Hussain et al. 2011a), modification of plant-pollinator interactions, and pose a significant impact at species, ecosystem and community levels and alter ecosystem services (Vilà et al. 2011). Additionally, IAS cause economic loss in forestry and agriculture (Pimentel et al. 2005) and are accelerated by anthropogenic perturbations (Young and Larson 2011). Coastal areas with temperate climates are a primary pathway for the introduction of IAS to Europe and the level of invasion in these areas is usually high (Chytrý et al. 2009) due to its favorable climate for plant establishment and growth (Stohlgren et al. 2003, 2006). The northwestern Iberian Peninsula constitutes a pathway of entry of IAS to Europe, and exhibits some early stages of IAS invasion (Capdevila-Argüelles et al. 2012; GISD 2022). This coastal region also has the highest forest fire activity in Europe (San-Miguel-Ayanz et al. 2013) and is one of the most affected regions by fire in the world (Archibald et al. 2013). Some of the IAS are tree species that cause major ecosystem problems, including the propagation of forest fires and making areas more fire prone (Brooks et al. 2004; Underwood et al. 2019; Aslan and Dickson 2020). In the northwest of the Iberian Peninsula several species stand out, including widely distributed species of *Acacia* and *Eucalyptus* genera (Sanz-Elorza et al. 2004), as well as others in early stages of invasion, such as the genera *Paraserianthes* (Dana et al. 2004) and *Phytolacca* (Valdés et al. 2011).

The invasional meltdown hypothesis (Simberloff and von Holle 1999) states that the establishment of one IAS facilitates invasion of other IAS (Kumar Rai and Singh 2020). Testing this hypothesis is the first step to know the repercussion of the initial invasions on subsequent ones and will be useful for exposing possible key species in the invasion process. *Acacia melanoxylon* R. Br. (Australian Blackwood) is one of the most damaging species for agriculture (Souto et al. 1994; Hussain et al. 2020) and *Eucalyptus globulus* Labill. (blue gum) for forestry ecosystems (Arán et al. 2013; Calviño-Cancela and Rubido-Bará 2013). Testing the invasional meltdown hypothesis for these IAS will allow us to discover if their presence helps the establishment of other IAS, namely other acacias, *Paraserianthes lophantha* (Willd.) I.C.Nielsen and *Phytolacca americana* L. This hypothesis postulates an adverse IAS diversity synergism on native species. Invasive species can also drive their own regeneration or produce negative feedback that limit it.

Seeds can recognize phytochemicals of conspecific seeds and establishing plants (biochemical recognition hypothesis, BRH; *sensu* Renne et al. 2004, 2014). Based on their dose-dependent assessment, they may decide to accelerate (e.g., avoid priority effects or enjoy conspecific facilitation; Dyer 2004; Orrock and Christopher 2010; Yamowo and Mukai 2017; Yannelli et al. 2020; Ohsaki et al. 2020) or delay germination (e.g., avoid competition-induced post-emergence mortality; Renne et al. 2014; Houseman and Mahoney 2015) to maximize establishment potential. In either case, this competition avoidance mechanism is predicted to maximize establishment potential (Renne et al. 2014) and thus, inva-

sion success. This process also applies to heterospecific recognition (Tielbörger and Prasse 2009; Yannelli et al. 2020; Fenesi et al. 2020). Some invasive plants may succeed because they bring novel phytotoxins to natural plant communities, as proposed by Callaway and Aschehoug (2000) in the novel weapons hypothesis (NWH). This would be a non-mutually exclusive mechanism to the BRH, for their own population's regeneration. Following this statement, native species would be more vulnerable to allelochemicals that they had never encountered. This is a chemically mediated mechanism that has been demonstrated in several studies (Callaway and Ridenour 2004; Becerra et al. 2018; Puig et al. 2018), and affects other species germination or seedling growth. This knowledge will be very useful for policymaking on control and management of these invasive alien species due to its wide distribution around the world (GISD 2022) since their interactions and synergies could also take place in other locations.

Many introduced species of *Acacia* and *Eucalyptus* produce allelopathic compounds, which reduce germination of some species (Souto et al. 2001; Hussain et al. 2011b; Lorenzo et al. 2012). Allelopathy is the ecological process in which biotic interference occurs through bioactive molecules (Singh et al. 1999). Allelochemicals (i.e., secondary metabolites, mostly phenolics and terpenoids) usually reduce native species and facilitate for IAS colonization in new habitats (Novoa et al. 2012; Kalisz et al. 2021; Zhang et al. 2021). The identification of phenolics, flavonoids and terpenoids can be difficult because they contain several structures but HPLS-MS is a useful tool to identify natural compounds in vegetal extracts (Quirantes-Piné et al. 2013; Jia et al. 2016). Allelochemicals could drive the invasion success of IAS, and it is important to know which compounds or doses affect their own germination and that of other species. Since germination is the most critical stage in the life cycle of many plants (Reyes et al. 1997) it is necessary to know how IAS seeds respond to their own phytochemicals and how the allelochemicals affect other IAS germination in already colonized areas.

In this study, we addressed the following objectives: (i) to determine the effect of different doses of allelopathic compounds from aqueous extracts of *A. melanoxylon* phyllodes and *E. globulus* leaves on their own germination and on other IAS germination, and (ii) to identify the possible compounds responsible for germination modification.

Materials and methods

Study species

We studied seven invasive alien species present in the northwest of the Iberian Peninsula and found in many other areas of the world: *Acacia dealbata* Link, *Acacia mearnsii* De Wild., *Acacia melanoxylon* R.Br., *Acacia longifolia* (Andrews) Willd., *Eucalyptus globulus* Labill., *Paraserianthes lophantha* (Willd.) I.C.Nielsen and *Phytolacca americana* L. The seeds of test species were collected in naturalized IAS populations in Galicia, Spain (northwestern Iberian Peninsula, Table S1). Additionally, we used *Lactuca sativa* L (cv. "Batavia") commercial seeds as a biotest species due to their fast germination and common use in allelopathic bioassays; seeds were purchased from Wamestrada S.L.L. (A Estrada, Spain).

Allelochemicals water extraction

Acacia melanoxylon and *E. globulus* were chosen as donor species of allelopathic compounds, due to their invasive character, their wide distribution and the density of their populations in the study area. *Acacia melanoxylon* is one of the IAS that has greatly increased in abundance and distribution in recent years (Martínez-Fernández et al. 2012); *E. globulus* is one of the most abundant species in the northwestern Iberian Peninsula, occupying 20% of the forested areas according to the IFN4 (IFN4 2011). Four aqueous extracts constituted the test treatments of IAS germination: two extracts at different concentrations from each of the two donor species. For the acacia extracts, we used *A. melanoxylon* abscised dry phyllodes and for the eucalyptus extracts we used *E. globulus* abscised dry leaves. Phyllodes and leaves were collected in naturalized adult populations in the Monte Pedroso area (Galicia, Spain). After collection, phyllodes and leaves were slashed in 2 cm² pieces (~1×2 cm). Extraction was performed in beakers covered with plastic film by soaking slashed phyllodes or leaves in distilled water at 200 g/L (hereafter, 200 g-acacia and 200 g-eucalyptus) and 100 g/L (100 g-acacia and 100 g-eucalyptus) for 72 h at room temperature. Then, leaves and phyllodes were separated from water by filtration and discarded, leading to the aqueous extracts. Following a similar methodology to Teerarak et al. (2010) and Nurjanah et al. (2020), the chosen values corresponded to the phyllodes or leaves maximum amount that is possible to soak in a known amount of water (200 g of phyllodes or leaves slashed pieces in 1 L of distilled water) and then reducing this amount to a half (100 g in 1 L).

Germination bioassays

For each of the seven species studied, in addition to four aqueous extract treatments, we used distilled water as a control to simulate natural conditions (control treatment). Hard-coated seeds were mechanically scarified before the beginning of the test to simulate germination. Scarification was performed with a scalpel, transversally cutting off a small part of the seed's distal end. For each treatment, five replicates of twenty-five seeds each were made. Each replicate was placed on a 9 cm diameter Petri dish, using two cellulose filter papers as substrate.

At the beginning of the test, 4 ml of the corresponding aqueous extract or distilled water was added to each replicate (Salgado et al. 2017). Subsequently, seed germination was checked every other day during a month. On those days, more water was added to keep the seeds moist, and seeds that had germinated (visible radicle) were removed from the Petri dish. Seed incubation was performed in a germination chamber (Climas AGP890) where the thermo-photoperiod was 16 h of light at 24 °C and 8 h of darkness at 16 °C, simulating favorable conditions for germination in the northwest of the Iberian Peninsula (Cruz et al. 2019; Riveiro et al. 2020).

Compound identification

To identify the *A. melanoxylon* and *E. globulus* chemical constituents present in the aqueous extracts, we conducted an ultrahigh-performance liquid chromatography coupled with quadrupole time-of-flight tandem mass spectrometry (UHPLC-ESI-QTOF-MS), a tool for characterizing complex natural products (Li et al., 2017). The compounds were identified

by comparing their retention times and MS/MS spectra provided by QTOF–MS with those of authentic standards whenever available. The remaining compounds were identified by interpreting their MS and MS/MS spectra obtained by QTOF–MS combined with the data provided in the literature. The score is a measure of identification confidence, based on the parameters of exact mass, retention time, mSigma and qualifier ions. Five score levels were obtained based on this: high (++++/+++), medium (++), tentative (+) and null (-); null level means that the qualifier ion was not found.

Data analysis

The data obtained were used to calculate the average germination percentage, the germination over time and the germination speed as T_{50} (which measures germination speed as the time required by seeds to reach 50% of final germination). T_{50} was calculated according to Cruz et al. (2022).

General linear models (GLMs) at a significance level of 0.05, with binomial error distribution, were carried out to test the effects of acacia and eucalyptus extracts on *germination percentage* and T_{50} (see supplementary material, Table S2). Independent analyses for each target species were carried out in a factorial analysis with the fixed factors *extract source* and *concentration of the extract*. Interaction between extract source and concentration of the extract was tested but it was not significant in any of the analyses, so we did not include it in the model to have a better model fit. We tested whether there were significant differences in *extract source* and *concentration of the extract* through ANOVA using the package *car* (Fox and Weisberg 2019); followed by a post-hoc HSD Tukey test using the package *agricolae* (de Mendiburu and Yassen 2020). GLMs for target species *E. globulus* and *L. sativa* regarding the response variable T_{50} did not include the predictor variable *concentration of the extract* for acacia extracts owing to the lack of data (there was no germination under these treatments). All the analyses were performed using R Software (R Core Team 2023).

Results

Germination percentage

Final germination varied by species and treatment (Table 1). Germination of all test species were significantly reduced by at least one treatment, and no treatment promoted germination. The germination percentage of *A. dealbata* and *P. lophantha* decreased in all treatments with respect to control and was especially low with 200 g-acacia treatment. Germination of *A. dealbata* decreased by 33% with 100 g-eucalyptus, 200 g-eucalyptus and 100 g-acacia and 75% with 200 g-acacia. *Paraserianthes lophantha* germination decreased 40% from control treatment with 200 g-acacia treatment ($p < 0.013$) and none of the other treatments modified its germination percentage. Germination of *A. mearnsii* decreased 36% with 200 g-acacia treatment ($p < 0.04$) with respect to control. *Acacia longifolia* and *A. melanoxylon* seed germination percentage was significantly reduced with both 100 g-acacia (61%, 87% respectively) and 200 g-acacia (71%, 68% respectively) treatments. Other treatments did not affect germination.

Table 1 Average final germination percentage ($\bar{x} \pm \text{SD}$) and T_{50} in days ($\bar{x} \pm \text{SD}$) for *A. dealbata*, *A. mearnsii*, *A. melanoxylon*, *A. longifolia*, *E. globulus*, *P. lophantha*, *P. americana* and *L. sativa* in control and 100 g-eucalyptus, 200 g-eucalyptus, 100 g-acacia and 200 g-acacia treatments. The first row for each species corresponds to germination percentage (G) and the second row to T_{50} . Different labels indicate significant differences for each species in germination (a, b, c, d) and in T_{50} (A, B, C) between the control and aqueous extracts in the Tukey contrasts performed. WD indicates that there were not enough replicates to perform the test

Treatment/Species		control	100 g-eucalyptus	200 g-eucalyptus	100 g-acacia	200 g-acacia
<i>A. dealbata</i>	G	96.0 $\pm$ 4.9 ^a	63.2 $\pm$ 25.8 ^b	57.6 $\pm$ 8.3 ^b	64.0 $\pm$ 10.2 ^b	24.0 $\pm$ 11.7 ^c
	T_{50}	3.0 $\pm$ 0.0 ^A	5.8 $\pm$ 1.8 ^B	6.2 $\pm$ 2.7 ^B	5.0 $\pm$ 0.0 ^B	5.0 $\pm$ 0.0 ^B
<i>A. mearnsii</i>	G	96.0 $\pm$ 4.0 ^a	79.2 $\pm$ 14.5 ^{ab}	77.6 $\pm$ 10.0 ^{ab}	88.0 $\pm$ 20.6 ^{ab}	60.0 $\pm$ 25.5 ^b
	T_{50}	5.0 $\pm$ 0.0 ^A	5.8 $\pm$ 1.8 ^A	5.8 $\pm$ 1.8 ^A	5.0 $\pm$ 0.0 ^A	6.0 $\pm$ 2.2 ^A
<i>A. melanoxylon</i>	G	50.4 $\pm$ 20.1 ^a	28.0 $\pm$ 17.7 ^{ab}	27.2 $\pm$ 21.4 ^{ab}	6.4 $\pm$ 5.4 ^b	16.0 $\pm$ 10.6 ^b
	T_{50}	5.0 $\pm$ 0.0 ^A	6.6 $\pm$ 2.2 ^A	9.0 $\pm$ 2.4 ^A	6.5 $\pm$ 3.0 ^A	5.8 $\pm$ 1.8 ^A
<i>A. longifolia</i>	G	78.4 $\pm$ 7.8 ^a	61.6 $\pm$ 37.3 ^{ab}	59.2 $\pm$ 18.0 ^{ab}	30.4 $\pm$ 22.0 ^b	22.4 $\pm$ 14.3 ^b
	T_{50}	8.2 $\pm$ 1.8 ^A	9.0 $\pm$ 0.0 ^A	9.0 $\pm$ 0.0 ^A	7.4 $\pm$ 2.2 ^A	9.0 $\pm$ 0.0 ^A
<i>E. globulus</i>	G	52.0 $\pm$ 22.8 ^{ab}	53.6 $\pm$ 12.8 ^a	15.2 $\pm$ 11.8 ^c	24.0 $\pm$ 17.4 ^c	0.0 $\pm$ 0.0 ^c
	T_{50}	5.0 $\pm$ 0.0 ^A	8.2 $\pm$ 1.8 ^A	11.8 $\pm$ 2.2 ^A	8.6 $\pm$ 2.2 ^A	WD
<i>P. lophantha</i>	G	76.0 $\pm$ 9.8 ^a	61.6 $\pm$ 12.5 ^{ab}	55.2 $\pm$ 13.1 ^{ab}	59.2 $\pm$ 12.8 ^{ab}	46.4 $\pm$ 10.4 ^b
	T_{50}	5.0 $\pm$ 0.0 ^A	5.0 $\pm$ 0.0 ^A	5.0 $\pm$ 0.0 ^A	5.0 $\pm$ 0.0 ^A	5.8 $\pm$ 1.8 ^A
<i>P. americana</i>	G	93.6 $\pm$ 3.6 ^a	87.2 $\pm$ 5.9 ^{ab}	76.0 $\pm$ 7.5 ^b	71.2 $\pm$ 10.0 ^{bc}	52.8 $\pm$ 11.1 ^d
	T_{50}	5.0 $\pm$ 0.0 ^A	8.2 $\pm$ 1.8 ^A	11.0 $\pm$ 2.0 ^A	7.6 $\pm$ 4.0 ^A	8.4 $\pm$ 5.6 ^A
<i>L. sativa</i>	G	96.0 $\pm$ 2.8 ^a	92.8 $\pm$ 5.2 ^a	77.0 $\pm$ 17.4 ^a	42.4 $\pm$ 25.2 ^{ab}	0.0 $\pm$ 0.0 ^c
	T_{50}	2.0 $\pm$ 0.0 ^A	5.0 $\pm$ 0.0 ^B	5.0 $\pm$ 0.0 ^B	5.0 $\pm$ 0.0 ^B	WD

In *E. globulus*, the 200 g-acacia treatment inhibited germination. 200 g-eucalyptus and 100 g-acacia reduced it by 70% and 53% with respect to control. Almost all treatments applied to *P. americana* significantly reduced germination ($p < 0.001$); 200 g-acacia reduced it by 43%, and 100 g-acacia and 200 g-eucalyptus produced smaller reductions (24%, 19% respectively). 100 g-eucalyptus did not affect germination of these two species. Biotest species *L. sativa* showed a marked decrease in germination percentage with 200 g-acacia ($p < 0.001$) from 89.6% in control treatment to 0%; other treatments did not affect its germination.

Acacia melanoxylon reduced target species germination with low (100 g-acacia) and high (200 g-acacia) concentration extracts, including its extracts to its own seeds. Regarding *E. globulus*, low concentration extract (100 g-eucalyptus) did not modify its own germination, but high concentrations (200 g-eucalyptus) reduced it. This effect was similar to that produced by *E. globulus* extracts in the other species studied, in which high concentration extract reduced germination more than low concentration extracts.

The most affected species by the applied treatments were *A. dealbata*, *E. globulus* and *P. americana*, whose germination was significantly reduced by 3 of the 4 treatments tested. In contrast, relative to the control, extracts generally did not affect the germination of *A. mearnsii*, *P. lophantha* and *L. sativa*, since their germination was reduced by only 1 out of 4 treatments. Eucalyptus aqueous extracts inhibited germination in 4 out of the 16 tests, and acacia aqueous extracts were inhibitory in 13 out of the 16 tests performed. Overall, the treatment which had the strongest effect was 200 g-acacia, which significantly reduced the germination percentage of seeds in all the species studied. 100 g-eucalyptus had the weakest effect, only modifying the germination of 1 out of 8 species.

T₅₀ and temporal distribution

For most of the species studied, T₅₀ increased slightly with some of the treatments, but the statistical analyses were not significant. The treatments applied only lead to a significant delay in the T₅₀ for *A. dealbata* and *L. sativa* ($p < 0.004$ and $p < 0.001$, respectively). The control germination of *A. dealbata* started on day 3 (Fig. 1) and ended on day 18, with a T₅₀ of 3 days. All treatments significantly increased the T₅₀ of this species (Table 1), reaching 5.0 to 6.2 days. Treated seeds had a delayed germination start on day 5. Control T₅₀ of *L. sativa* biotest species was 2.0 days, which significantly increased to 5.0 days ($p < 0.05$) for the seeds treated with 100 g-eucalyptus, 200 g-eucalyptus and 100 g-acacia. The control germination of *L. sativa* started on day 3, while the seeds' germination treated with 100 g-acacia started on day 5, and with 100 g-eucalyptus and 200 g-eucalyptus, germination started on day 9.

Acacia longifolia had the longest T₅₀ at 8 days, while all the other species reached T₅₀ at 5 days. In *A. mearnsii* control germination started on day 3 (Fig. 1) and reached a T₅₀ of 5.0 days. The control germination of *A. melanoxylon*, *E. globulus* and *P. americana* started on day 5, and reached T₅₀ on day 5 (Table 1). Treated seeds of *A. melanoxylon* started germination on day 9, delaying it with respect to control. *Eucalyptus globulus* seeds treated with 100 g-acacia started germination on the same day as control (Fig. 1) while seeds treated with 100 g-eucalyptus and 200 g-eucalyptus delayed their germination. Neither *A. melanoxylon* nor *E. globulus* modified their own T₅₀.

Germination of *A. longifolia* started on day 5 in control and treated seeds, with a T_{50} of 8.2 days and none of the treatments modified it (Table 1). Control seeds of *P. lophantha* and *P. americana* began their germination on day 3 while treated seeds started germination on day 5 (Fig. 1). The T_{50} of *P. lophantha* was very consistent, with no variation between control and treatments, being 5.0 days on average (Table 1).

To sum up, the T_{50} of most species remained unchanged after the application of treatments compared to control, except for *A. dealbata* and *L. sativa* in which case it was delayed (Table 1). Eucalyptus aqueous extracts delayed T_{50} in 4 out of 16 treatments, and acacia aqueous extracts delayed it in 3 out of 14 (Table 1). The treatments also delayed the start of germination, ranging from 2 to 6 days in all species except *A. longifolia*, whose treated seeds started germinating the same day as the control seeds (Fig. 1).

Identification and chemical analysis of compounds in aqueous extracts

The eucalyptus extracts contained 47 compounds, while acacia extracts had 20 compounds (Table S3, supplementary material). The maximum confidence level obtained in acacia extracts was medium (++), in 8 compounds and none of the compounds reached the highest level. From these compounds, the most abundant were trihydroxyoctadecenoic acid (17.5%), tyrosol glucoside (3.6%), gentistic acid (2.5%) and scopoletin (2.0%). Eleven compounds of eucalyptus extract obtained medium confidence level (++) and 4 compounds obtained a high level (++++). The high confidence level compounds were gallic acid, epicatechin, ferulic acid and ellagic acid, which represents 39.4% of the total eucalyptus sample analyzed. Thus, gallic acid and ellagic acid were the two main compounds in Eucalyptus extracts, and two of the most certain in the sample.

Eucalyptus and Acacia extracts shared 7 out of 16 high confidence level compounds. However, gallic acid and ellagic acid, the two most abundant compounds in eucalyptus extracts, were not present in the acacia extracts. Only 1 of the 16 high confidence compounds, tyrosol glucoside, was present in acacia extracts but not in eucalyptus extracts.

Discussion

Germination and ecological implications

Acacia and eucalyptus phytochemical compounds affected germination of the eight study species, and aqueous extract dose and origin were fundamental to this. Extracts of *A. melanoxylon* lowered germination in all species treated with 200 g-acacia, *E. globulus* extracts decreased germination in *A. dealbata*, *E. globulus* and *P. americana*, and none of the extracts fostered germination in any species. The germination reduction in all the studied species by *A. melanoxylon* extracts, at least at the highest concentration (200 g-acacia), matches the results of Hussain et al. (2011), who concluded that *Acacia* allelopathy effects depend on extract concentration. Of the five species whose germination was inhibited with the two concentrations tested, the inhibition effect increased with incrementing extract concentration only in *A. dealbata* and *P. americana*. The lack of germination promotion in any target species, does not support the invasional meltdown hypothesis (Simberloff and von Holle 1999). For example, *A. melanoxylon* and *E. globulus* did not facilitate the invasion of *A.*




Fig. 1 Temporal distribution of germination for control and treated seeds of *A. dealbata*, *A. mearnsii*, *A. melanoxylon*, *A. longifolia*, *E. globulus*, *P. lophantha*, *P. americana* and *L. sativa* in treatments without total suppression of germination. Values on day 30 constitute the final germination percentages

dealbata, *A. mearnsii*, *P. lophantha* and *P. americana* through allelopathy compounds in areas already colonized by *A. melanoxylon* or *E. globulus*. This is consistent with the fact that the majority of interactions between invaders are either neutral or negative (81% of interactions) and it is very rare the facilitation among them (Kuebbing and Nuñez 2015).

Regarding their native range, all target acacias are sympatric—as well as congeneric—so it is possible they have developed species-specific or phylogenetic-specific BR (Renne et al. 2014). The lowered acacia germination with *A. melanoxylon* extracts do not support the NWH, because the phytotoxins would not be new to the target acacias and it is unlikely that sharing habitat both in their native range and in their invaded range to not have developed some sort of mechanism to avoid the phytotoxic effects of sympatric, closely-related species. Importantly, *A. melanoxylon* extracts (100 g-acacia and 200 g-acacia) reduced germination of *A. melanoxylon* seeds, and 200 g-eucalyptus reduced *E. globulus* germination. This was found in other Fabaceae species (Khan and Shaukat 2006) treated with its own fruit extracts and in other woody species (Bran et al. 1990; Romane et al. 1992; Houseman and Mahoney 2015). The fact that the germination reduction is stronger on donor species than on the other IAS clearly supports the BRH. Donor species would be able to recognize high population density through phytochemicals released into the environment (intraspecific-BR) and react by limiting their germination to avoid strong intraspecific competition in the future (Renne et al. 2014; Hierro and Callaway 2021). This effect of suppression of its own germination is not detected in natural populations, probably due to the potential seed bank that those species presents; in *A. melanoxylon* the soil seed bank is large and persistent (Arán et al. 2017) and *globulus* has an aerial seed bank in which seeds can remain viable during years (Reyes and Casal 1998). Specifically, *A. melanoxylon*, in which we found the strongest germination reduction, tends to form dense monospecific stands which allow them to be permanently in contact with their conspecific phytochemicals, favoring BR development (Renne et al. 2014). The regulation of populational growth via BR would be ideal conditions for seed germination, with adult individuals releasing phytochemicals that postpone germination until the environment is conducive for establishment. This behavior would benefit adult plants of *A. melanoxylon* and *E. globulus*, as described by Aguilera et al. (2017) for *A. dealbata* in Chile.

Germination of biotest *L. sativa* was reduced with acacia extracts and delayed by all applied treatments. Similar results were obtained by Souto et al. 2001, who found reduced germination of *L. sativa* seeds from *A. melanoxylon* stands. Also, Hussain et al. (2011a, 2020) reported inhibition of germination of *L. sativa* seeds treated with flower and phyllodes extracts of *A. melanoxylon*; they also detected native species inhibition in Atlantic forests, such as *Dactylis glomerata* L., *Lolium perenne* L. and *Rumex acetosella* L. Meanwhile, *E. globulus* extracts did not affect *L. sativa* germination, neither in our study nor in Souto et al. (2001). Given that *L. sativa* is a native species, our results regarding acacia extracts support the NWH (Callaway and Aschehoug 2000). Nelson et al. (2021) did not find inhibitory effects of *E. globulus* on the germination of understory Californian native species either. By contrast Morsi and Abdelmigid (2016) found inhibitory effects on *Hordeum vulgare* L. germination, a native species of Saudi Arabian forests, using a high concentration of *E. globulus* leaf extracts. Additionally, despite its common use as biotest species in allelopathy studies, we are aware of the limitations of using *L. sativa* as a native species, due to its commercial use and its artificial selection for agriculture. For these reasons, it is necessary to perform more tests on a wider range of native species to extract solid conclusions about the NWH for the two donor species.

Different extracts and doses barely modified speed and temporal distribution of germination. In *A. dealbata*, treated seeds took longer to reach T_{50} when compared to control treatment. This also occurred in *L. sativa*, whose treated seeds also started germination later than control treatment. Other authors found that reduced germination is not linked to changes in T_{50} (Cruz et al. 2020). Start of germination was also delayed in some target species between 2 and 6 days; a delay in germination onset could be due to BR, since we only applied the aqueous extracts at the

start of the experiment, and subsequently applied distilled water. It is possible that phytochemical concentration in the seed environment was reduced, leading to more conducive conditions for germination. Comparing the eight studied species, the most sensitive ones were *A. dealbata*, *E. globulus* and *P. americana*, whose germination was lowered by three out of four tested treatments. Furthermore, the only species whose germination was totally inhibited were *L. sativa* and *E. globulus*, which were treated with 200 g-acacia. These two species were the only ones studied that did not present hard-coated seeds and had the smallest seed size. Nevertheless, all hard-coated seeds used in this study were scarified, so we cannot attribute this effect to the protective coat. In normal conditions, with the hard coat unaltered, these species take 1 to 3 months to germinate and their germination is low (Arán et al. 2017; Cruz et al. 2017; García-Duro et al. 2019; Riveiro et al. 2020).

Aqueous extract composition

Some of the main compounds found in the *A. melanoxylon* chemical analysis were reported in previous allelopathy studies. *p*-Hydroxybenzoic acid (Reigosa et al. 1999; Hussain et al. 2011b) was even reported to cause germination suppression and radicle growth inhibition at different concentrations in several species. In our study, *A. melanoxylon* extracts did not contain gallic acid, ferulic acid (Souto et al. 1994; Hussain et al. 2011b) and ellagic acid (Souto et al. 1994) despite being tested for them. Instead, we found that the most abundant compounds were trihydroxyoctadecenoic acid and tyrosol glucoside, which were not previously reported in *A. melanoxylon* extracts. In *E. globulus* extracts, the principal compounds gallic acid, epicatechin, ferulic acid and ellagic acid were found in other studies (Souto et al. 1994; Reigosa et al. 2000; Hussain et al. 2011b) and in other species of the *Eucalyptus* genus (Suresh and Vinaya Rai 1987; Li et al. 2010). Gallic acid and chlorogenic acid were also found by Puig et al. (2018) who concluded that the inhibitory effects observed could be attributed *a priori* to the phytotoxins present in the plant's aqueous extracts. Some of the compounds identified in our study also cause growth suppression in seedlings (Puig et al. 2018); Hussain et al. (2010) found a reduction on shoot, leaf and root length and a drop in fresh biomass of *L. sativa* using ferulic acid. This effect on the seedling growth would support the NWH if we use *L. sativa* to depict native species that have never been exposed to IAS allelochemicals of tested donor species, and consequently would experience negative effects after the exposure.

It would be interesting to clarify if the compounds found in our aqueous extracts cause inhibitory effects on germination when isolated from the mixtures. Other authors (Chon et al. 2003; Reigosa and Pazos-Malvido 2007) tested the efficacy of these compounds as germination suppressors, and they concluded that the compound's mixtures were more phytotoxic than individual compounds. As we did not find phytotoxic effects on IAS but biochemical recognition among them, we wonder if the BR would be stronger in the compound mixture than in the isolated compounds, or in contrast, whether there would be some compounds responsible for the BR. Furthermore, the allelochemical effect could be a temporary germination inhibition, reversible when the allelopathic compound concentration decreases, and the environment is more conducive for establishment. Lastly, given that seven of sixteen compounds are shared between acacia and eucalyptus test species (Table 2), it is possible that seeds use these phytochemical cues to indicate a competitive environment (Renne et al. 2014), regardless of whether an established acacia or eucalyptus individual is in close proximity.

Conclusion

From the seven studied IAS, only three were affected by adding eucalyptus aqueous extracts. In contrast, all of them were affected by acacia aqueous extracts. Regarding the treatments, eucalyptus aqueous extracts lowered germination in 4 out of 16 tests, and acacia aqueous extracts were inhibitory in 13 out of 16 tests performed. In all cases, the treatment that most reduced germination was 200 g-acacia (acacia aqueous extract at higher concentration). We also detected germination reduction in the two donor species treated with their own phytochemicals, suggesting intraspecific BR, and found support for the NWH depicting *L. sativa* as a native species. None of the donor species promoted germination in the tested species, discarding the meltdown hypothesis for *A. melanoxylon* and *E. globulus* aqueous extracts. In conclusion, our findings show that phytochemicals mediate plant-plant interactions by altering germination schedules, which likely scale up to population and community regulation.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11056-023-10001-1>.

Acknowledgements The authors thank the financial support of the Spanish Ministry of Science, Innovation and Universities, the Castilla y León Regional Government, the Galicia Regional Government and the European Regional Development Fund (ERDF) through FIRESEVES (AGL2017-86075-C2-2-R) and WUI-FIRECYL (LE005P20) projects, the Competitive Reference Group BIOAPLIC (ED431C2019/07) and the Strategic Researcher Cluster BioReDeS (ED431E 2018/09). S. F. Riveiro is financially supported by a PhD Fellowship-Contract (ED481A-2020/088) from Xunta de Galicia (Axudas de apoio á etapa predoutoral 2020 - Modalidade B). Ó. Cruz is financially supported by a Collaboration agreement between Xunta de Galicia and University of Santiago de Compostela which regulates the “Campus Terra” Specialization Campus. Authors would like to thank the use of RIAIDT-USC analytical facilities. The authors would like to thank the time dedicated by Mercedes Uscola (Associate Editor) and Ian J. Renne (Reviewer) to carefully review the first version of the manuscript, their comments were very valuable and contributed to considerably improve the final version of the manuscript.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Data Availability All data are available under request at Zenodo under the DOI: <https://doi.org/10.5281/zenodo.5910035>.

Declarations

Conflicts of interest The authors declare they have no conflicts of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Reference

- Aguilera N, Guedes LM, Becerra J, González L (2017) Is autotoxicity responsible for inhibition growth of new conspecific seedlings under the canopy of the invasive *Acacia dealbata* link? *Gayana Botánica* 74:1–14. <https://doi.org/10.4067/S0717-66432017005000101>
- Arán D, García-Duro J, Reyes O, Casal M (2013) Fire and invasive species: modifications in the germination potential of *Acacia melanoxylon*, *Conyza canadensis* and *Eucalyptus globulus*. *For Ecol Manage* 302:713. <https://doi.org/10.1016/j.foreco.2013.02.030>
- Arán D, García-Duro J, Cruz Ó et al (2017) Understanding biological characteristics of *Acacia melanoxylon* in relation to fire to implement control measurements. *Ann for Sci* 74:61. <https://doi.org/10.1007/s13595-017-0661-y>
- Archibald S, Lehmann CER, Gómez-Dans JL, Bradstock RA (2013) Defining pyromes and global syndromes of fire regimes. *Proc Natl Acad Sci* 110:6442–6447. <https://doi.org/10.1073/pnas.1211466110>
- Aslan CE, Dickson BG (2020) Non-native plants exert strong but under-studied influence on fire dynamics. *NeoBiota* 61:47–64. <https://doi.org/10.3897/NEOBIOTA.61.51141>
- Becerra PI, Catford JA, Inderjit et al (2018) Inhibitory effects of *Eucalyptus globulus* on understorey plant growth and species richness are greater in non-native regions. *Glob Ecol Biogeogr* 27:68–76. <https://doi.org/10.1111/GEB.12676>
- Bran D, Lobréaux O, Maistre M et al (1990) Germination of *Quercus ilex* and *Q. pubescens* in a *Q. ilex* coppice - long-term consequences. *Vegetatio* 87:45–50. <https://doi.org/10.1007/BF00045654>
- Brooks ML, D'Antonio CM, Richardson DM et al (2004) Effects of invasive alien plants on Fire Regimes | *BioScience* | Oxford Academic. *Bioscience* 54:677–688. [https://doi.org/10.1641/0006-3568\(2004\)054\[0677:EOIAP0\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0677:EOIAP0]2.0.CO;2)
- Callaway RM, Aschehoug ET (2000) Invasive plants versus their new and old neighbors: a mechanism for exotic invasion. *Sci* (80-) 290:521–523. <https://doi.org/10.1126/SCIENCE.290.5491.521/ASSET/C45238A7-1E34-42FC-ADAF-9A1315822366/ASSETS/GRAPHIC/SE4008916004.JPEG>
- Callaway RM, Ridenour WM (2004) Novel weapons: invasive success and the evolution of increased competitive ability - callaway - 2004 - *frontiers in Ecology and the Environment* - Wiley Online Library. *Front Ecol Environ* 2:436–443. [https://doi.org/10.1890/1540-9295\(2004\)002\[0436:NWISAT\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2004)002[0436:NWISAT]2.0.CO;2)
- Calviño-Cancela M, Rubido-Bará M (2013) Invasive potential of *Eucalyptus globulus*: seed dispersal, seedling recruitment and survival in habitats surrounding plantations. *For Ecol Manage* 305:129–137. <https://doi.org/10.1016/J.FORECO.2013.05.037>
- Capdevila-Argüelles L, Zilletti B, Suárez Álvarez V (2012) Plan Estratéxico Galego de xestión das Especies Exóticas Invasoras e para o desenvolvemento dun sistema estandarizado de. *Análise de Risco para as especies exóticas en Galicia*
- Chon S-U, Kim Y-M, Lee J-C (2003) Herbicidal potential and quantification of causative allelochemicals from several Compositae weeds. *Weed Res* 43:444–450. <https://doi.org/10.1046/j.0043-1737.2003.00361.x>
- Chytrý M, Pyšek P, Wild J et al (2009) European map of alien plant invasions based on the quantitative assessment across habitats. *Divers Distrib* 15:98–107. <https://doi.org/10.1111/j.1472-4642.2008.00515.x>
- R Core Team (2023) R: A language and environment for statistical computing
- Cruz Ó, García-Duro J, Casal M, Reyes O (2017) Can the mother plant age of *Acacia melanoxylon* (Leguminosae) modulate the germinative response to fire? *Aust J Bot* 65:593. <https://doi.org/10.1071/BT17083>
- Cruz Ó, García-Duro J, Casal M, Reyes O (2019) Role of serotiny on *Pinus pinaster* Aiton germination and its relation to mother plant age and fire severity. *iForest* 12:491–497. <https://doi.org/10.3832/ifor2968-012>
- Cruz Ó, García-Duro J, Riveiro SF et al (2020) Fire severity drives the natural regeneration of *Cytisus scoparius* L. (link) and *Salix atrocinerea* brot. *Communities and the germinative behaviour of these species. Forests* 11:124. <https://doi.org/10.3390/f11020124>
- Cruz Ó, Riveiro SF, Casal M, Reyes O (2022) Effect of fire factors (smoke, ash, charcoal and heat) on seeds of plant species. *MethodsX* 9:101679. <https://doi.org/10.1016/J.MEX.2022.101679>
- D'Antonio CM, Vitousek PM (1992) Biological Invasions by exotic grasses, the Grass/Fire cycle, and global change. *Annu Rev Ecol Syst* 23:63–87. <https://doi.org/10.1146/annurev.es.23.110192.000431>
- Dana ED, Sobrino E, Sanz-Elorza M (2004) Plantas invasoras en España: un nuevo problema en las estrategias de conservación. In: Bañares Á, Blanca G, Gümes J et al (eds) *Atlas y libro rojo de la flora vascular amenazada. Dirección General de Conservación de la Naturaleza*, Madrid, pp 1010–1029
- de Mendiburu F, Yassen M (2020) agricolae: Statistical Procedures for Agricultural Research
- Dyer AR (2004) Maternal and sibling factors induce dormancy in dimorphic seed pairs of *Aegilops triuncialis*. *Plant Ecol* 172:211–218. <https://doi.org/10.1023/B:VEGE.0000026339.61069.33/METRICS>
- Fenesi A, Kelemen K, Sándor D et al (2020) Influential neighbours: seeds of dominant species affect the germination of common grassland species. *J Veg Sci* 31:1028–1038. <https://doi.org/10.1111/JVS.12892>
- Fox J, Weisberg S (2019) *An R companion to Applied Regression*, 3rd edn. Sage, Thousand Oaks (CA)

- Gaertner M, Biggs R, Te Beest M et al (2014) Invasive plants as drivers of regime shifts: identifying high-priority invaders that alter feedback relationships. *Divers Distrib* 20:733–744. <https://doi.org/10.1111/ddi.12182>
- García-Duro J, Cruz Ó, Casal M, Reyes O (2019) Fire as driver of the expansion of *Paraserianthes lophantha* (Willd.) I. C. Nielsen in SW Europe. *Biol Invasions* 21:1427–1438. <https://doi.org/10.1007/s10530-018-01910-w>
- GISD (2022) Global Invasive Species Database (GISD). <https://www.cabi.org/isc/abstract/20067200496>. Accessed 14 Dec 2020
- Herrero-Borgoñón J (2007) Dos Mimosoideas (Leguminosae) nuevas para la flora castellanense. *Flora Montiberica* 28:26–28
- Hierro JL, Callaway RM (2021) The ecological importance of Allelopathy. *Annu Rev* 52:25–45. <https://doi.org/10.1146/ANNUREV-ECOLSYS-051120-030619>
- Houseman GR, Mahoney AK (2015) Intraspecific seed interactions alter seedling emergence of *Lespedeza cuneata* under field conditions. *Popul Ecol* 57:539–544. <https://doi.org/10.1007/S10144-015-0495-0>
- Hussain MI, González L, Reigosa MJ (2010) Phytotoxic effects of allelochemicals and herbicides on photosynthesis, growth and carbon isotope discrimination in *Lactuca sativa*. *Allelopathy J* 26:157–174
- Hussain MI, González L, Reigosa MJ (2011a) Allelopathic potential of *Acacia melanoxylon* on the germination and root growth of native species. *Weed Biol Manag* 11:18–28. <https://doi.org/10.1111/j.1445-6664.2011.00401.x>
- Hussain MI, González L, Souto XC, Reigosa MJ (2011b) Ecophysiological responses of three native herbs to phytotoxic potential of invasive *Acacia melanoxylon* R. Br. *Agrofor Syst* 83:149–166. <https://doi.org/10.1007/s10457-011-9433-0>
- Hussain MI, El-Sheikh MA, Reigosa MJ (2020) Allelopathic potential of aqueous extract from *acacia melanoxylon* r. br. On *lactuca sativa*. *Plants* 9:1–13. <https://doi.org/10.3390/plants9091228>
- IFN4 (2011) Cuarto Inventario Forestal Nacional, Galicia
- Jia M, Li Y, Zhai X et al (2016) Qualitative analysis and quality evaluation of *Cnidium monnieri* using UHPLC-ESI-Q-TOF/MS. *Chin Herb Med* 8:323–330. [https://doi.org/10.1016/s1674-6384\(16\)60058-8](https://doi.org/10.1016/s1674-6384(16)60058-8)
- Kalisz S, Kivlin SN, Bialic-Murphy L (2021) Allelopathy is pervasive in invasive plants. *Biol Invasions* 23:367–371. <https://doi.org/10.1007/S10530-020-02383-6/FIGURES/1>
- Khan D, Shaikat S (2006) Phytotoxic effects of *Prosopis juliflora* Swartz. DC. Against some of its field associates and the cultivated species. *Int J Biol Biotechnol* 3:353–366
- Kuebbing SE, Nuñez MA (2015) Negative, neutral, and positive interactions among nonnative plants: patterns, processes, and management implications. *Glob Chang Biol* 21:926–934. <https://doi.org/10.1111/GCB.12711>
- Kumar Rai P, Singh JS (2020) Invasive alien plant species: their impact on environment, ecosystem services and human health. *Ecol Indic* 111:106020. <https://doi.org/10.1016/j.ecolind.2019.106020>
- Li ZH, Wang Q, Ruan X et al (2010) Phenolics and Plant Allelopathy. *Mol* 2010, Vol 15, Pages 8933–8952 15:8933–8952. <https://doi.org/10.3390/MOLECULES15128933>
- Li H, Yao W, Liu Q et al (2017a) Application of UHPLC-ESI-Q-TOF-MS to identify multiple constituents in processed products of the herbal medicine *ligustri lucidi fructus*. *Molecules* 22. <https://doi.org/10.3390/molecules22050689>
- Li N, Yang W, Fang S et al (2017b) Dispersal of invasive *Phytolacca americana* seeds by birds in an urban garden in China. *Integr Zool* 12:26–31. <https://doi.org/10.1111/1749-4877.12214>
- Lorenzo P, Pazos-Malvido E, Rubido-Bará M et al (2012) Invasion by the leguminous tree *Acacia dealbata* (Mimosaceae) reduces the native understorey plant species in different communities. *Aust J Bot* 60:669. <https://doi.org/10.1071/BT12036>
- Martínez-Fernández J, Hernández L, de la Vázquez A, Cañellas I (2012) (PDF) Distribución y dinámica del género *Acacia* en Galicia: un análisis a partir del Tercer y Cuarto Inventario Forestal Nacional. In: EEI 2012 Notas Científicas 4º Congreso Nacional sobre Especies Exóticas Invasoras. pp 128–132
- Morsi MM, Abdelmigid HM (2016) Allelopathic activity of *Eucalyptus globulus* leaf aqueous extract on *Hordeum vulgare* growth and cytogenetic behaviour. *AJCS* 10:1835–2707. <https://doi.org/10.21475/ajcs.2016.10.11.PNE122>
- Nelson KM, Bisbing S, Grossenbacher DL et al (2021) Testing an invasion mechanism for *Eucalyptus globulus*: is there evidence of allelopathy? *Am J Bot* 108:607–615. <https://doi.org/10.1002/AJB2.1635>
- Novoa A, González L, Moravcová L, Pyšek P (2012) Effects of Soil characteristics, Allelopathy and Fruivory on Establishment of the Invasive Plant *Carpobrotus edulis* and a Co-Occurring native, *Malcolmia littorea*. *PLoS ONE* 7:e53166. <https://doi.org/10.1371/JOURNAL.PONE.0053166>
- Nurjanah U, Setyowati N, Simarmata M (2020) Allelopathic potential of aqueous extract of *Archidendron jiringa* (Jering) pods for weed control in Swamp paddy field. *Int J Agric Technol* 16:1153–1164

- Ohsaki H, Mukai H, Yamowo A (2020) Biochemical recognition in seeds: germination of *Rumex obtusifolius* is promoted by leaves of facilitative adult conspecifics. *Plant Species Biol* 35:233–242. <https://doi.org/10.1111/1442-1984.12275>
- Orrock JL, Christopher CC (2010) Density of intraspecific competitors determines the occurrence and benefits of accelerated germination. *Am J Bot* 97:694–699. <https://doi.org/10.3732/AJB.0900051>
- Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol Econ* 52:273–288. <https://doi.org/10.1016/j.ecolecon.2004.10.002>
- Puig CG, Reigosa MJ, Valentão P et al (2018) Unravelling the bioherbicide potential of *Eucalyptus globulus* Labill: Biochemistry and effects of its aqueous extract. *PLoS ONE* 13:e0192872. <https://doi.org/10.1371/JOURNAL.PONE.0192872>
- Pyšek P, Jarošík V, Hulme PE et al (2012) A global assessment of invasive plant impacts on resident species, communities and ecosystems: the interaction of impact measures, invading species' traits and environment. *Glob Chang Biol* 18
- Qu T, Du X, Peng Y et al (2021) Invasive species allelopathy decreases plant growth and soil microbial activity. *PLoS ONE* 16:e0246685. <https://doi.org/10.1371/JOURNAL.PONE.0246685>
- Quirantes-Piné R, Lozano-Sánchez J, Herrero M et al (2013) HPLC-ESI-QTOF-MS as a powerful Analytical Tool for Characterising Phenolic Compounds in Olive-leaf extracts. *Phytochem Anal* 24:213–223. <https://doi.org/10.1002/pca.2401>
- Reigosa MJ, Pazos-Malvido E (2007) Phytotoxic Effects of 21 plant secondary metabolites on *Arabidopsis thaliana* Germination and Root Growth. *J Chem Ecol* 33:1456–1466. <https://doi.org/10.1007/S10886-007-9318-X>
- Reigosa MJ, Souto XC, González L (1999) Effect of phenolic compounds on the germination of six weeds species. *Plant Growth Regul* 28:83–88. <https://doi.org/10.1023/A:1006269716762>
- Reigosa MJ, González L, Souto XC, Pastoriza JE (2000) Allelopathy in forest ecosystems. In: *Allelopathy in Ecological Agriculture and Forestry*. Springer Netherlands, pp 183–193
- Renne JJ, Rios BG, Fehmi JS, Tracy BF (2004) Low allelopathic potential of an invasive forage grass on native grassland plants: a cause for encouragement? *Basic Appl Ecol* 5:261–269. <https://doi.org/10.1016/J.BAAE.2003.11.001>
- Renne JJ, Sinn BT, Shook GW et al (2014) Eavesdropping in plants: delayed germination via biochemical recognition. *J Ecol* 102:86–94. <https://doi.org/10.1111/1365-2745.12189>
- Reyes O, Casal M (1998) Germination of *Pinus pinaster*, *P. radiata* and *Eucalyptus globulus* in relation to the amount of ash produced in forest fires. *Ann des Sci for* 55:837–845. <https://doi.org/10.1051/forest:19980707>
- Reyes O, Casal M, Traubaud L (1997) The influence of population, fire and time of dissemination on the germination of *Betula pendula* seeds. *Plant Ecol* 133:201–208. <https://doi.org/10.1023/A:1009751513547>
- Riveiro SF, Cruz Ó, Casal M, Reyes O (2020) Fire and seed maturity drive the viability, dormancy, and germination of two invasive species: *Acacia longifolia* (Andrews) Willd. And *Acacia mearnsii* De Wild. *Ann For Sci* 77:60. <https://doi.org/10.1007/s13595-020-00965-x>
- Romane F, Bacillieri R, Bran D, Bouchet MA (1992) Natural Degenerate Mediterranean Forests: Which Future ? The Examples of the Holm Oak (*Quercus ilex* L.) and Chestnut (*Castanea sativa* Mill.) Coppice Stands. In: *Responses of Forest Ecosystems to Environmental Changes*. Springer Netherlands, pp 374–380
- Romero Buján MI (2007) Flora exótica de Galicia (noroeste ibérico). *Bot Complut* 31:113–125
- Salgado J, Parajó JJ, Teijeira T et al (2017) New insight into the environmental impact of two imidazolium ionic liquids. Effects on seed germination and soil microbial activity. *Chemosphere* 185:665–672. <https://doi.org/10.1016/j.chemosphere.2017.07.065>
- San-Miguel-Ayán J, Moreno JM, Camia A (2013) Analysis of large fires in European Mediterranean landscapes: Lessons learned and perspectives. *For Ecol Manage* 294:11–22. <https://doi.org/10.1016/J.FORECO.2012.10.050>
- Sanz-Elorza M, Dana ED, Sobrino-Vesperinas E (2004) Atlas de las plantas alóctonas invasoras de España. Dirección General para la Biodiversidad, Madrid
- Simberloff D, von Holle B (1999) Positive Interactions of Nonindigenous Species: Invasional Meltdown? *Biol Invasions* 1999 11 1:21–32. <https://doi.org/10.1023/A:1010086329619>
- Singh HP, Batish DR, Kohli RK (1999) Autotoxicity: Concept, organisms, and ecological significance. Taylor & Francis Group
- Souto XC, González L, Reigosa MJ (1994) Comparative analysis of allelopathic effects produced by four forestry species during decomposition process in their soils in Galicia (NW Spain). *J Chem Ecol* 20:3005–3015. <https://doi.org/10.1007/BF02098405>

- Souto XC, Bolaño JC, González L, Reigosa MJ (2001) Allelopathic effects of tree species on some soil microbial populations and herbaceous plants. *Biol Plant* 44:269–275. <https://doi.org/10.1023/A:1010259627812>
- Stohlgren TJ, Barnett DT, Kartesz JT (2003) The rich get richer: patterns of plant invasions in the United States. *Front Ecol Environ*. [https://doi.org/10.1890/1540-9295\(2003\)001\[0011:TRGRPO\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2003)001[0011:TRGRPO]2.0.CO;2) 1:
- Stohlgren TJ, Jarnevich C, Chong GW, Evangelista PH (2006) Scale and plant invasions: a theory of biotic acceptance. *Měřítko studia a rostlinné invaze: teorie biotické akceptance*. *Preslia* 78:405–426
- Suresh KK, Vinaya Rai RS (1987) Studies on the allelopathic effects of some agroforestry tree crops. *Int Tree Crop J* 4:109–115. <https://doi.org/10.1080/01435698.1987.9752816>
- Teerarak M, Laosinwattana C, Charoenying P (2010) Evaluation of allelopathic, decomposition and cytogenetic activities of *Jasminum officinale* L. f. var. *Grandiflorum* (L.) Kob. On bioassay plants. *Bioresour Technol* 101:5677–5684. <https://doi.org/10.1016/J.BIORTECH.2010.02.038>
- Tielbörger K, Prasse R (2009) Do seeds sense each other? Testing for density-dependent germination in desert perennial plants. *Oikos* 118:792–800. <https://doi.org/10.1111/J.1600-0706.2008.17175.X>
- Underwood EC, Klinger RC, Brooks ML (2019) Effects of invasive plants on fire regimes and postfire vegetation diversity in an arid ecosystem. *Ecol Evol* 9:12421–12435. <https://doi.org/10.1002/ECE3.5650>
- Valdés B, Melero D, Girón V (2011) Plantas americanas naturalizadas en el territorio de Doñana (SO de la Península Ibérica). *Lagascalia* 31:7–20
- Vilà M, Espinar JL, Hejda M et al (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol Lett* 14:702–708
- Wohlgemuth T, Gossner MM, Campagnaro T et al (2022) Impact of non-native tree species in Europe on soil properties and biodiversity: a review. *NeoBiota* 78:45–69. <https://doi.org/10.3897/NEOBIOTA.78.87022>
- Yamowo A, Mukai H (2017) Seeds integrate biological information about conspecific and allospecific neighbours. *Proc R Soc B Biol Sci* 284:20170800. <https://doi.org/10.1098/rspb.2017.0800>
- Yannelli FA, Novoa A, Lorenzo P et al (2020) No evidence for novel weapons: biochemical recognition modulates early ontogenetic processes in native species and invasive acacias. *Biol Invasions* 22:549–562. <https://doi.org/10.1007/S10530-019-02110-W/FIGURES/3>
- Young AM, Larson BMH (2011) Clarifying debates in invasion biology: a survey of invasion biologists. *Environ Res* 111:893–898. <https://doi.org/10.1016/j.envres.2011.06.006>
- Zhang Z, Liu Y, Yuan L et al (2021) Effect of allelopathy on plant performance: a meta-analysis. *Ecol Lett* 24:348–362. <https://doi.org/10.1111/ELE.13627>