

LETTER • OPEN ACCESS

Contrasting shrub species respond to early summer temperatures leading to correspondence of shrub growth patterns

To cite this article: Stef Weijers *et al* 2018 *Environ. Res. Lett.* **13** 034005

View the [article online](#) for updates and enhancements.

You may also like

- [Greater effect of increasing shrub height on winter versus summer soil temperature](#)
Mélicca Paradis, Esther Lévesque and Stéphane Boudreau
- [Shrub growth in the Alps diverges from air temperature since the 1990s](#)
Loïc Francon, Christophe Corona, Irène Till-Bottraud et al.
- [Are low altitude alpine tundra ecosystems under threat? A case study from the Parc National de la Gaspésie, Québec](#)
Catherine Dumais, Pascale Ropars, Marie-Pier Denis et al.

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED

15 November 2017

REVISED

29 December 2017

ACCEPTED FOR PUBLICATION

8 January 2018

PUBLISHED

19 February 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Contrasting shrub species respond to early summer temperatures leading to correspondence of shrub growth patterns

Stef Weijers^{1,3} , Roland Pape¹, Jörg Löffler¹ and Isla H Myers-Smith²¹ Department of Geography, University of Bonn, Meckenheimer Allee 166, D-53115 Bonn, Germany² School of GeoSciences, King's Buildings, University of Edinburgh, West Mains Road, Edinburgh, EH9 3FF, United Kingdom³ Author to whom any correspondence should be addressed.E-mail: weijers@uni-bonn.de**Keywords:** alpine, browning, greening, NDVI, productivity, shrubs, tundraSupplementary material for this article is available [online](#)

Abstract

The Arctic-alpine biome is warming rapidly, resulting in a gradual replacement of low statured species by taller woody species in many tundra ecosystems. In northwest North America, the remotely sensed normalized difference vegetation index (NDVI), suggests an increase in productivity of the Arctic and alpine tundra and a decrease in productivity of boreal forests. However, the responses of contrasting shrub species growing at the same sites to climate drivers remain largely unexplored.

Here, we test growth, climate, and NDVI relationships of two contrasting species: the expanding tall deciduous shrub *Salix pulchra* and the circumarctic evergreen dwarf shrub *Cassiope tetragona* from an alpine tundra site in the Pika valley in the Kluane Region, southwest Yukon Territories, Canada.

We found that annual growth variability of both species at this site is strongly driven by early summer temperatures, despite their contrasting traits and habitats. Shrub growth chronologies for both species were correlated with the regional climate signal and showed spatial correspondence with interannual variation in NDVI in surrounding alpine and Arctic regions. Our results suggest that early summer warming represents a common driver of vegetation change for contrasting shrub species growing in different habitats in the same alpine environments.

1. Introduction

Arctic-alpine ecosystems are sensitive to climate change (Settele *et al* 2014) and the rate of warming increases with latitude and elevation (Pepin *et al* 2015). Consequently, the Arctic-alpine biome in the northern hemisphere is expected to shift north- and upward (Settele *et al* 2014). Evidence from experimental warming (Walker *et al* 2006, Elmendorf *et al* 2012a), repeated vegetation surveys (Elmendorf *et al* 2012b), repeated photography (Sturm *et al* 2001, Tape *et al* 2006), and dendrochronology (Myers-Smith *et al* 2011, 2015a), suggests that shrub growth is sensitive to summer warming and that shrub cover has increased in response to climate warming throughout the tundra biome. Pollen records indicate greater shrub dominance in tundra during past warm episodes in Late

Quaternary North America (Hu *et al* 2002, Higuera *et al* 2008) and Russia (Velichko *et al* 1997). In addition, a growing number of studies indicate shrubline ecotone advance up- and north-ward into tall-shrub free tundra (Myers-Smith and Hik 2018). However, growth responses of contrasting shrub species with different traits and habitats, occupying separate niches, are rarely compared in the same plant communities. Thus, we do not yet know whether all shrub functional groups are responding similarly to the rapidly warming tundra climate.

In general, taller deciduous shrubs are expected to replace lower statured species, and positive effects of ambient and experimental warming were predominantly observed on the abundance of taller deciduous shrubs (Elmendorf *et al* 2012a, 2012b). Dwarf shrub cover was found to decline in a

tundra biome-wide synthesis study on experimental warming (Elmendorf *et al* 2012a), especially in warmer areas; perhaps as a result of increased competition for light. Evergreen dwarf shrub species have, however, been shown to be climate sensitive (Bär *et al* 2008, Buizer *et al* 2012, Weijers *et al* 2012, 2017) and have been observed to increase their leaf size and height in response to experimental warming (Hudson *et al* 2011) and cover in response to ambient warming (Hudson and Henry 2009) at some High Arctic sites.

Satellite observations of vegetation productivity, as measured by the normalized difference vegetation index (NDVI), have shown a greening of large parts of the tundra biome in the northern hemisphere (Jia *et al* 2003, Goetz *et al* 2005 2011, Guay *et al* 2014). Increases in NDVI over northern (Forbes *et al* 2010, Macias-Fauria *et al* 2012), northeastern Siberia (Blok *et al* 2011), and northwest North America (Tape *et al* 2012) have been related to increases in growth of deciduous shrubs and summer temperatures. Such links have yet to be explored for evergreen dwarf shrub species. Using aerial photography and satellite imagery, an increase in NDVI over the Low Arctic Tuktoyaktuk Coastal Plain, Northern Territories, western Canada, was shown to be related with an increase in shrub canopy cover (Fraser *et al* 2014). If remote sensing data are documenting tundra shrub expansion, then these data indicate that multiple species are responding synchronously to the changing climate conditions.

Here, we explore the links between NDVI, climate, and annual shrub growth of the tall deciduous shrub *Salix pulchra* and the evergreen dwarf shrub *Cassiope tetragona* co-occurring at an alpine tundra site in the Pika valley in the Kluane Region, southwest Yukon Territories, Canada. In this region *Salix* species have recently been shown to be expanding upward through new recruitment along the shrubline in response to winter warming and their growth has been shown to be sensitive to summer temperatures (Myers-Smith and Hik 2018). Annual growth of *C. tetragona* has been shown to be driven by summer temperatures at many High Arctic sites (Callaghan *et al* 1989, Havström *et al* 1995, Johnstone and Henry 1997, Rayback and Henry 2006, Rozema *et al* 2009, Weijers *et al* 2010, 2012, 2017), but may be more sensitive to fluctuations in nutrient availability at warmer sites in the Subarctic (Havström *et al* 1993). However, climate sensitivity of contrasting shrub species including a tall deciduous willow and a dwarf evergreen *Cassiope* species have yet to be directly compared.

Our research questions were:

1. What are the main climate drivers of annual growth variability of the co-occurring evergreen dwarf shrub *Cassiope tetragona* and the deciduous tall shrub *Salix pulchra* in the Pika valley of the Kluane Region, Yukon Territory, Canada?
2. Does the climate signal in the growth chronologies of the two species represent the regional climate?
3. How does the interannual variation in growth correspond with interannual variation in productivity as observed through the NDVI?

2. Materials and methods

2.1. Study site and sampling

Our research site is located on the Pika valley slopes in the Kluane Region, Yukon Territory, northwest Canada (61.22 °N, 138.28 °W; figure 1(a)).

On August 16, 2010, 16 samples of *Cassiope tetragona* were collected from the east-northeast-facing slope of the valley (figure 1(b)) at elevations of 1640–1673 m above sea level (a.s.l.). Several meters distances were kept between sampling spots to prevent repeated sampling of the same genet. *Cassiope tetragona* (L.) D. Don. (Ericaceae), or Arctic bell heather, is a multi-branched, clonal, hemi-prostrate, evergreen dwarf shrub with a circumpolar distribution (Eidesen *et al* 2007, Weijers *et al* 2017). At this site, *C. tetragona* is dominant in depressions with long-lasting snow cover and grows up to 30 cm in height.

Samples of the tall deciduous shrub *Salix pulchra* were collected between 19 and 21 August, 2007 from both valley slopes (figure 1(b)). Samples were taken at the shrubline (~1812 and 1970 m a.s.l. on the east-northeast and west-southwest-facing slope, respectively) and at more downslope positions with a *Salix* species cover of approx. 50% (~1715 m a.s.l.). In total, 17 *S. pulchra* specimens were sampled: four and three at the shrubline and three and seven downslope on the east-northeast and west-southwest-facing slopes, respectively. A 3–5 cm long disc was taken just above the stem-root boundary of the largest stem of each individual for growth ring analysis. Only distinct *Salix* patches were sampled, likely representing distinct genets. *Salix pulchra* Cham. (Salicaceae) is a canopy-forming deciduous shrub found in the Siberian and northwest North American tundra (CYSIP 2017). It is the most dominant willow species in the Kluane Region east of Kluane Lake (Myers-Smith and Hik 2018). The species reaches heights between 20 and 80 cm at our site.

2.2. Climate

We calculated monthly precipitation sums (mm), and mean, mean minimum, and mean maximum monthly temperatures (°C) as valid, i.e. lapse rate adjusted, for our research site for each month over the period 1901–2012 in ClimateWNA v5.40 (Wang *et al* 2017). In this program monthly normal data from climate mapping systems are downscaled to scale-free point data, which are then used as a baseline, together with monthly anomaly data from the Climate Research Unit Time Series (CRU TS3.23; Harris *et al* 2014),





Figure 1. Location of the Kluane Region in the Yukon Territory northwest Canada indicated with a yellow square on the inset map and the location of the Pika valley (yellow circle) northwest of Kluane Lake (a). Satellite image map of the Pika valley and adjacent areas (b); sampling spots of *Cassiope tetragona* are marked with red dots; sampling locations of *Salix pulchra* are indicated with green dots.

for the calculation of historical climate variables for specific locations and elevations (Wang *et al* 2016).

The mean annual temperature at our study site is -4.35°C and mean annual precipitation sum is 502 mm (figure 2(a)). July is both the warmest and the wettest month with a mean temperature of 9.28°C and mean precipitation sum of 69 mm. January is the coldest month (mean temperature -17.25°C). We defined November–March as the winter months,

April–May as spring, June–August as summer, and September–October as autumn, and calculated seasonal means and trends therein over two periods: 1950–2012 (figures 2(b) and (c)) and 1902–2012 (figure S1), as trends over these periods may have been relevant for shrub growth (cf. figure S2). Seasonal mean temperatures have risen significantly in winter ($r = 0.29$, $p < 0.05$), spring ($r = 0.38$, $p < 0.01$), and summer ($r = 0.33$, $p < 0.01$) over the period 1950–2012.



Over the period 1902–2012 mean temperatures have risen significantly in spring ($r = 0.26$, $p < 0.01$) and summer ($r = 0.38$, $p < 0.001$; figure S1). Trends in seasonal precipitation sums were not significant.

2.3. Climate-growth analyses: linear mixed models

We measured annual growth of *C. tetragona* as shoot length increments and that of *S. pulchra* as ring width increments. See supplementary information available at stacks.iop.org/ERL/13/034005/mmedia for details on annual growth measurement. We used linear mixed model analyses to test the influence of monthly and seasonal climatic parameters on shrub growth. Climate-growth models were compared over the periods 1902–2009 and 1949–2009 (*C. tetragona*), or 1949–2006 (*S. pulchra*). The R-package nlme (Pinheiro *et al* 2017) was used for the mixed model analyses, with maximum likelihood estimation for model comparison and restricted maximum likelihood estimation for slope estimates (Crawley 2007). Before the analyses, the climate and shrub shoot length and ring-width chronologies were normalized at the individual level through subtraction of the mean, followed by a division by the standard deviation. Annual shoot lengths or ring widths of individual shrubs were included in the models as the response variable, and climate variables were included as fixed effects. A random intercept for year and an autocorrelation structure (AR1, autoregressive process of order one) were additionally included in the models. Conditional pseudo- R^2 values were calculated for each model with the `r.squaredGLMM` function of the MuMIn package (Nakagawa and Schielzeth 2013).

We tested 94 climate-growth models including temperature means, mean maximums, mean minimums, and precipitation sums from 17 individual months (previous June to current October), the four seasons, and early (June–July) and late (July–August) summer as fixed effects, besides a null model for both

species. In a first step, climate models that performed better than the null model were selected based on the Akaike Information Criterion ($\Delta AIC > 2$). Subsequently, Akaike weights for the selected models were calculated for model comparison. Akaike weights are a relative weight of evidence for each model and can be interpreted as the probability a certain model is the best model, given the selected set of models, for the observed data (Johnson and Omland 2004). High numbers of models in a comparison analysis increases the chances that a model has a lower AIC-value than the accompanying null model. By including 94 models for each species, models might thus come up by chance having a ΔAIC -value greater than 2. However, it is unlikely that such models will have an AIC value much lower than the related null model or have a high Akaike weight.

2.4. Climate-growth analyses: site chronologies

We tested climate-growth relationships between monthly climate parameters and standardized site chronologies. See supplementary information for details on standardization, and chronology construction and statistics (table S1 and figure S2).

We calculated Pearson's correlation coefficients and response function coefficients between each standardized site chronology and monthly mean temperatures as well as monthly precipitation sums, with monthly parameters from previous June to current October. This was done over the period 1907–2009 for *C. tetragona* and 1971–2006 for *S. pulchra*. Parts of the chronologies with $n < 5$, were excluded from the analyses, as were the most recent growth years. The latter were excluded due to possible incomplete growth at the time of harvest. In addition, Pearson's correlation coefficients and response coefficients were calculated over 25 year moving windows over the period 1907–2009, with a two-year window offset with the same monthly parameters, separately for

mean monthly temperatures and monthly precipitation sums. This was only done for *C. tetragona* (1907–2009; $n \geq 5$), due to the relatively shortness of the *S. pulchra* chronology (1971–2006; $n \geq 5$). Correlation and response coefficient significance was determined through a 1000 bootstrapped iterations. Response function analysis takes multicollinearity between climate parameters into account, through the regression of growth chronologies against principal components of climate parameters (Zang and Biondi 2015). The correlation and response analyses were performed with the R-package treeclim (Zang and Biondi 2015) in R version 3.4.1 (R Core Team 2017).

Spatial relationships in the area between 50°–75°N and 170°–120°W over the period 1981–2006 were calculated between the standardized site chronologies, June–July NDVI (0.22° spatial resolution), and June–July mean maximum temperatures from the 0.5° × 0.5° monthly gridded meteorological dataset CRU TS4.00 (Harris *et al* 2014). Furthermore, the spatial relationships in the same area and over the same period between local mean June–July maximum temperatures (ClimateWNA v5.40), mean June–July NDVI, and mean June–July maximum temperatures (CRU TS4.00) were calculated. These calculations were carried out in KNMI Climate Explorer (Trouet and van Oldenborgh 2013), which uses a Monte Carlo approach for the determination of confidence intervals. The months June and July were chosen for the spatial analyses, because of their influence on growth of both species (see below). NDVI data for the period 1981–2006 were obtained from the GIMMS dataset (Tucker *et al* 2005) via KNMI Climate Explorer.

3. Results

3.1. Climate-growth analyses

Variation in annual growth of both species was best explained by early summer temperature models (tables 1 and S2), with evidence for the mean June–July temperature model and the mean maximum June–July temperature as best model for *C. tetragona* and the mean maximum June–July temperature model as best model for *S. pulchra*. In addition, we found some support for mean (maximum) summer temperature-growth models for *C. tetragona* and mean (maximum) July temperature-growth models for *S. pulchra* (table 1). Besides summer climate models some current spring as well previous year late summer and early autumn temperature models were among the selected models for *C. tetragona*.

A similar picture arises from the correlation and response function analyses, with positive correlations between mean monthly temperatures of April–August of the growing season, as well as previous August and September and *C. tetragona* growth (figure 3(a)). Mean June, July, August temperatures of the current year, and previous August remain significant predictors of

C. tetragona shoot length growth, taken collinearity between the monthly climate parameters into account (figure 3(b)). Mean June and July temperatures were again identified as the main predictors of *S. pulchra* radial growth (figures 3(c) and (d)).

The influence of mean June and July temperatures has been stable throughout the growth record of *C. tetragona* (figure 4). There has been a shift, however, in the influence of August temperatures from the current year, which have influenced growth until the late 1950s. Thereafter, August temperatures started to influence *C. tetragona* growth in the next year (figure 4(a)). The negative moving correlations found between *C. tetragona* growth and monthly precipitation sums, mainly of summer months (figure 4(b)), are likely a result of collinearity between the monthly climate variables (cf. figure 4(d)). The influence of June and July temperatures may have been shifting back and forth between these two months (figure 4(c)).

3.2. Spatial analyses

Both standardized chronologies as well as the Pika valley mean June–July maximum temperatures are strongly correlated to mean June–July maximum temperatures over a large area in northwest North America (figures 5(a)–(c)). Furthermore, the *C. tetragona* chronology reflects mean June–July NDVI-values from a large area, mainly from the tundra and boreal treeline vegetation at higher elevations and/or latitudes such as at the Stikine plateau, Mackenzie mountains, Coast mountains, Babine range, Muskwa ranges, and in the area north of the Brooks ranges (figure 5(d)). The standardized *S. pulchra* chronology shares the spatial correlations with June–July NDVI over the alpine tundra southeast to our research site, as found for *C. tetragona*, but lacks those to the north and northwest (figure 5(e)). Neither chronology is related to the June–July NDVI-values and temperatures of the boreal forests of the Alaskan interior. The spatial relationship between the lapse rate corrected mean June–July maximum temperatures as valid for the Pika valley and NDVI (figure 5(f)) is similar as that for the *C. tetragona* chronology, but it lacks the relationships with June–July NDVI north of the Brooks Range.

4. Discussion

We have shown that *Cassiope tetragona* and *Salix pulchra* growth at the Pika valley in the Kluane Region, Yukon Territory, Canada is driven by early (June–July) growing season temperatures. Despite contrasting plant traits of dwarf evergreen versus tall deciduous shrubs and different habitats related to snow distribution, consistent climate sensitivity to early growing season temperatures was observed. Early summer temperatures, which coincide with the time of year with maximum insolation, are warming across northwest North America and shrub growth chronologies for


Table 1. Results of the mixed model analyses with annual shoot lengths of *Cassiope tetragona* or annual ring widths of *Salix pulchra* included in the models as the response variable and climate variables as fixed effects, calculated over the period 1949–2009 (*C. tetragona*) or 1949–2006 (*S. pulchra*). Selected models are models with AIC values of at least two lower than the corresponding null model. R^2 -values are conditional pseudo- R^2 values. Akaike weights are the relative weight of evidence for each model. T: mean temperature; $T_{\max}$: mean maximum temperature; $T_{\min}$: mean minimum temperature; P: precipitation sum.

Species	Selected Models	Δ AIC	Slope $\pm$ SE	R^2	Akaike weight
<i>Cassiope tetragona</i>	June–July T	29.78	0.37 ± 0.06	0.29	0.53
	June–July $T_{\max}$	29.02	0.36 ± 0.06	0.29	0.36
	Summer T	25.25	0.34 ± 0.06	0.29	0.05
	Summer $T_{\max}$	25.13	0.34 ± 0.06	0.29	0.05
	July T	17.05	0.30 ± 0.06	0.29	<0.01
	June–July $T_{\min}$	15.75	0.29 ± 0.06	0.29	<0.01
	June T	14.61	0.28 ± 0.06	0.29	<0.01
	June $T_{\max}$	14.49	0.28 ± 0.06	0.29	<0.01
	Previous August T	12.71	0.27 ± 0.07	0.29	<0.01
	Summer $T_{\min}$	12.35	0.26 ± 0.07	0.29	<0.01
	July–August T	11.93	0.26 ± 0.07	0.29	<0.01
	July $T_{\min}$	11.85	0.26 ± 0.07	0.29	<0.01
	July $T_{\max}$	10.27	0.24 ± 0.07	0.29	<0.01
	Previous August $T_{\min}$	10.22	0.25 ± 0.07	0.29	<0.01
	July–August $T_{\max}$	9.57	0.24 ± 0.07	0.29	<0.01
	Previous August $T_{\max}$	8.61	0.23 ± 0.07	0.29	<0.01
	June $T_{\min}$	8.46	0.23 ± 0.07	0.29	<0.01
	July–August $T_{\min}$	7.03	0.21 ± 0.07	0.29	<0.01
	Previous July $T_{\min}$	4.96	0.19 ± 0.07	0.29	<0.01
	April $T_{\max}$	4.37	0.18 ± 0.07	0.29	<0.01
	May $T_{\min}$	4.15	0.18 ± 0.07	0.29	<0.01
	Spring $T_{\max}$	3.57	0.17 ± 0.07	0.29	<0.01
	August $T_{\max}$	3.48	0.17 ± 0.07	0.29	<0.01
	Spring T	3.41	0.17 ± 0.07	0.29	<0.01
	August T	2.90	0.16 ± 0.07	0.29	<0.01
	Summer P	2.75	-0.16 ± 0.07	0.29	<0.01
	April T	2.28	0.15 ± 0.07	0.29	<0.01
	Previous September T	2.13	0.15 ± 0.07	0.29	<0.01
	May T	2.01	0.15 ± 0.07	0.29	<0.01
<i>Salix pulchra</i>	June–July $T_{\max}$	18.66	0.28 ± 0.06	0.16	0.80
	July T	14.74	0.26 ± 0.06	0.16	0.11
	July $T_{\max}$	12.99	0.25 ± 0.06	0.17	0.05
	June–July T	11.78	0.24 ± 0.06	0.16	0.03
	Summer $T_{\max}$	9.63	0.23 ± 0.06	0.16	<0.01
	June $T_{\max}$	6.97	0.19 ± 0.06	0.15	<0.01
	Summer T	6.33	0.20 ± 0.07	0.16	<0.01
	July $T_{\min}$	5.26	0.19 ± 0.07	0.17	<0.01
	July–August T	4.06	0.17 ± 0.07	0.16	<0.01
	July–August $T_{\max}$	4.06	0.17 ± 0.07	0.16	<0.01
	June T	3.26	0.15 ± 0.06	0.15	<0.01
	Summer P	2.99	-0.16 ± 0.07	0.15	<0.01
	June–July P	2.65	-0.14 ± 0.06	0.15	<0.01
	June–July $T_{\min}$	2.02	0.14 ± 0.07	0.16	<0.01

both species were correlated with the regional climate and showed spatial correspondence with interannual variation in NDVI in surrounding alpine and Arctic regions. Our results suggest a common driver of vegetation change for contrasting shrub species growing in different habitats in the same alpine environments.

In contrast to *S. pulchra*, we found some influence of late summer and early autumn temperatures of the previous year on growth of *C. tetragona*, besides the influence of early summer temperature, as was earlier observed for this species on Svalbard (Weijers *et al* 2010). This is likely a result of the evergreen nature of *C. tetragona* leaves and the formation of its primordial leaves at the end of the growing season. Photosynthesis in *C. tetragona* leaves starts before snowmelt is completed (Starr and Oberbauer 2003) and shoot elongation may thus benefit from warm early summer temperatures. Annual growth of *C. tetragona* has previously been shown to be driven by summer

temperatures at several High Arctic sites, through experimental warming in Ellesmere Island, Canada (Hudson *et al* 2011), Greenland (Campioli *et al* 2013) and Svalbard (Havström *et al* 1993, Rozema *et al* 2009, Weijers *et al* 2012), and dendrochronological analyses in Ellesmere Island, Canada (Havström *et al* 1995, Johnstone and Henry 1997, Rayback and Henry 2006), North Greenland (Weijers *et al* 2017), and Svalbard (Callaghan *et al* 1989, Aanes *et al* 2002, Weijers *et al* 2010, 2012, 2013b). However, at warmer Subarctic alpine tundra sites, as our site, environmental growth controls of *C. tetragona* may be less uniform. Growth in Subarctic Abisko, North Sweden, for example, showed a greater response to nitrogen addition than to experimental warming, and may be more nitrogen- than temperature-limited (Havström *et al* 1993). Still, after longer timespans (22 years) neither fertilization nor warming was found to affect *C. tetragona* growth at the same experiment (Campioli *et al* 2012).



The *C. tetragona* shrubs in our study were from downslope depressions with late snowmelt. Winter snow depth at Arctic-alpine localities is, unlike at flat lowland sites, relatively independent of the winter precipitation sum, as excess snow is removed by wind and redistributed according to topography (Erickson *et al* 2005). Snowmelt date and growing season length at these *C. tetragona* localities are thus likely largely determined by temperature instead of precipitation, which may explain the strong relationship found between early summer temperatures and growth. In addition, mean growth rate at our site ($5.21 \text{ mm year}^{-1}$) is close to the $5.05 \text{ mm year}^{-1}$ reported for High Arctic Svalbard (Weijers *et al* 2012), despite the warmer conditions at our site (mean July temperatures of 9.28°C versus 6.43°C). This suggests relatively harsh conditions at the downslope snowbeds on the east-northeast-facing slope at our site, which may explain the dominance there of *C. tetragona*, which is generally a more High Arctic species. Relatively harsh conditions may persist at these places due to a shortened growing season in snowbeds, cooling of soils by (upslope) meltwater, and low amounts of direct sunlight due to the east-northeast slope aspect. During mornings, sunlight is blocked by the opposing mountain.

Early growing season temperatures have earlier been identified as the most important factor influencing *S. pulchra* growth in the northeastern Siberian tundra (Blok *et al* 2011) and may stimulate its growth

as leaf flush in *S. pulchra* takes place within days after snowmelt (Borner *et al* 2008). Moreover, *Salix spp.* growth was found to be driven predominantly by summer temperatures, and their recruitment by winter temperatures, throughout the Kluane Region (Myers-Smith and Hik 2018). In contrast, secondary growth of *S. pulchra* showed little response to fertilization and warming treatments when in competition with *Betula nana* (Bret-Harte *et al* 2002) and its abundance declined under both treatments (Bret-Harte *et al* 2001) in the moist tussock tundra at Toolik Lake in the northern foothills of the Brooks Range, Alaska. However, at a nearby site, June temperatures were found to be important for radial growth of *S. pulchra* (Ackerman *et al* 2017). Despite some variation among studies, June and July temperatures appear to be consistent drivers of variation in *S. pulchra* shrub growth.

We found some indication of a negative, but no positive, influence of summer precipitation on growth of both species. This corresponds with the lack of a *C. tetragona* growth response to experimentally doubled summer precipitation in Svalbard (Weijers *et al* 2013a) and the probably high growing season soil moisture content at the *C. tetragona* localities resulting from snowmelt. *S. pulchra* growth may be limited by precipitation at some sites, as it was found to increase its shoot length growth in response to experimentally increased precipitation (Keuper *et al* 2012).




Figure 4. Moving correlation coefficients (a) and (b) and response function coefficients (c) and (d), calculated over 25 year windows with 2 year offsets over the period 1907–2009, between the standardized *Cassiope tetragona* shoot length chronology and mean monthly temperatures as well as monthly precipitation sums, as modelled for the Pika valley, Kluane Region, Yukon Territory, Canada. Asterisks (*) indicate significant coefficients. Abbreviations of months from the year of growth are in lowercase letters and those of the previous year are given in capitals.

However, the low-Arctic northeast Siberian tundra site at which that experiment was conducted receives less than half of the precipitation annually falling in our study area, which may explain the lack of a positive influence of precipitation at our site. Still, a positive influence of summer precipitation was reported for *Salix spp.* at some other sites in the Kluane Region (Myers-Smith and Hik 2018).

Negative links between summer precipitation and *C. tetragona* growth were earlier reported by Callaghan *et al* (1989) for sites in Svalbard and Swedish Lapland. *C. tetragona* may suffer from soil subsidence due to thaw of deeper ice-rich permafrost layers under wetter conditions, as observed in a snow accumulation experiment in Svalbard (Blok *et al* 2015). However, the observed negative correlations between precipitation sums during summer months and growth at our site likely result from the negative relationship between summer precipitation and temperature at our site. This hypothesis is further supported by the low Akaike weights for summer precipitation models for both species and the near absence of significant response function coefficients between growth and monthly summer precipitation.

Annual *C. tetragona* and *S. pulchra* growth at our site correspond with mean maximum June–July temperatures for a large part of northwest North America.

C. tetragona shoot length growth furthermore tracked June–July NDVI-values of the vegetation at higher elevations and latitudes in this region and radial growth of *S. pulchra* corresponded with mean maximum NDVI values of a large area southeast of the study site. This suggests that annual growth variability of *C. tetragona* and *S. pulchra* respond to climate drivers with wide spatial extents and that productivity in these alpine regions may correspond across the whole region. Our findings correspond with NDVI observations of a greening Alaskan and Yukon tundra since 1982 (Verbyla 2008, Beck and Goetz 2011, Guay *et al* 2014, Ju and Masek 2016). Similar inter-correlations between annual shrub growth, summer temperatures, and NDVI have earlier been found for the deciduous shrubs *S. lanata* and *Alnus fruticosa* in northern Siberia (Forbes *et al* 2010, Macias-Fauria *et al* 2012) and for *S. pulchra* in northeastern Siberia (Blok *et al* 2011).

The climate signal in *S. pulchra* is somewhat less strong than that in *C. tetragona* shrubs and its growth corresponded with NDVI variability over a smaller area than *C. tetragona* growth. This might be because the evergreen dwarf shrub *C. tetragona* is not grazed upon, in contrast to many deciduous Arctic-alpine shrub species (Mallik *et al* 2011). At our site, the deciduous *S. pulchra* shrubs may have been prone to ptarmigan browsing new buds in spring, non-cyclic




Figure 5. Pearson correlation coefficients calculated over the period 1981–2006 between the standardized *Cassiope tetragona* chronology, the standardized *Salix pulchra* chronology, Pika valley mean maximum June–July temperatures, and gridded mean June–July maximum temperatures from CRU TS4.00 (a)–(c), and mean June–July NDVI from GIMMS (d)–(f). Only significant correlations are shown on the maps ($p < 0.05$). Black diamonds indicate the location of our research site at Pika valley in the Kluane Region, Yukon Territory, Canada.

insect herbivores, stem herbivory by small mammals including marmots and rare browsing by moose or other large herbivores (Myers-Smith and Hik 2018). Expansion of deciduous shrubs has earlier been shown to be suppressed by reindeer grazing in the Scandinavian alpine tundra, in contrast to that of evergreen shrubs (Vowles *et al* 2017).

Although Arctic tundra greening has mainly been attributed to an expansion of erect and tall deciduous shrub cover (Elmendorf *et al* 2012a, 2012b, Fraser *et al* 2014), we found that annual growth of the hemi-prostrate evergreen dwarf shrub *C. tetragona* largely corresponds with NDVI as a proxy for productivity over large parts of the northwest North American tundra. *C. tetragona* might be able to expand its cover through a densification of existing shrub patches in

the warmer parts of the tundra biome in certain environmental niches with long lasting snow cover, where relatively harsh conditions may persist during the growing season. *C. tetragona* is capable of forming dense mats, which might inhibit the recruitment of other taller shrubs in such locations.

5. Conclusions

We found a strong positive growth response of the co-occurring evergreen dwarf shrub *Cassiope tetragona* and the deciduous tall shrub *Salix pulchra* to early summer warming in the Pika valley of the Kluane Region, Yukon Territory, Canada. Despite differences in plant species traits and habitats, the two contrasting

species demonstrated surprisingly consistent growth responses to climate drivers. Moreover, our findings show that the annual growth variability of these species are likely representative of the annual variability in tundra vegetation productivity of large parts of north-west North America. Early summer warming has likely enhanced growth rates of entire shrub communities in this region, resulting in the densification of shrub stands and a greening of the Arctic-alpine tundra of Alaska and the Yukon, as reported in the literature.

Acknowledgments

We thank the Kluane First Nation for the opportunity to do research on their traditional lands. Field and laboratory assistance was provided by M M Grabowski, T L Elliot, A Briggs, C Henry, J Pongracz, and C McDonald. This research was partly funded by the Canon National Parks Science Scholars Program, Government of Canada International Polar Year Program (PPS Arctic Canada Project), NSERC (Discovery, Northern Research Supplement, and International Polar Year Programs), the Arctic Institute of North America, the Canada Foundation for Innovation, the Northern Research Institute at Yukon College, Alberta Ingenuity, the Northern Scientific Training Program, and the Circumpolar/Boreal Alberta Research Grant Program, Canadian Circumpolar Institute, and Air North, Yukon's Airline (flight discounts), and the German Research Foundation (DFG), grant WE 5472/1-1. We thank both anonymous referees for their helpful comments on the manuscript.

ORCID iDs

Stef Weijers  <https://orcid.org/0000-0003-3386-5417>

References

- Aanes R, Sæther B-E, Smith F M, Cooper E J, Wookey P A and Øritsland N A 2002 The Arctic Oscillation predicts effects of climate change in two trophic levels in a high-arctic ecosystem *Ecol. Lett.* **5** 445–53
- Ackerman D, Griffin D, Hobbie S E and Finlay J C 2017 Arctic shrub growth trajectories differ across soil moisture levels *Glob. Change Biol.* **23** 4294–302
- Bär A, Pape R, Bräuning A and Löffler J 2008 Growth-ring variations of dwarf shrubs reflect regional climate signals in alpine environments rather than topodimatic differences *J. Biogeogr.* **35** 625–36
- Beck P S A and Goetz S J 2011 Satellite observations of high northern latitude vegetation productivity changes between 1982 and 2008: Ecological variability and regional differences *Environ. Res. Lett.* **6** 45501
- Blok D, Sass-Klaassen U, Schaepman-Strub G, Heijmans M M P D, Sauren P and Berendse F 2011 What are the main climate drivers for shrub growth in Northeastern Siberian tundra? *Biogeosciences* **8** 1169–79
- Blok D, Weijers S, Welker J M, Cooper E J, Michelsen A, Löffler J and Elberling B 2015 Deepened winter snow increases stem growth and alters stem $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in evergreen dwarf shrub *Cassiope tetragona* in high-arctic Svalbard tundra *Environ. Res. Lett.* **10** 44008
- Borner A P, Kielland K and Walker M D 2008 Effects of simulated climate change on plant phenology and nitrogen mineralization in Alaskan Arctic tundra *Arctic Antarct. Alpine Res.* **40** 27–38
- Bret-Harte M S, Shaver G R and Chapin F S 2002 Primary and secondary stem growth in arctic shrubs: implications for community response to environmental change *J. Ecol.* **90** 251–67
- Bret-Harte M S, Shaver G R, Zoerner J P, Johnstone J F, Wagner J L, Chavez A S, Gunkelman R F, Lippert S C and Laundre J A 2001 Developmental plasticity allows *Betula nana* to dominate tundra subjected to an altered environment *Ecology* **82** 18–32
- Buizer B, Weijers S, van Bodegom P eter M, Alsos I G, Eidesen P B, van Breda J, de Korte M, van Rijkkevorsel J and Rozema J 2012 Range shifts and global warming: ecological responses of *Empetrum nigrum* L. to experimental warming at its northern (high Arctic) and southern (Atlantic) geographical range margin *Environ. Res. Lett.* **7** 25501
- Callaghan T V, Carlsson B A and Tyler N J C 1989 Historical records of climate-related growth in *Cassiope tetragona* from the Arctic *J. Ecol.* **77** 823–37
- Campioli M, Leblans N and Michelsen A 2012 Twenty-two years of warming, fertilisation and shading of subarctic heath shrubs promote secondary growth and plasticity but not primary growth *PLoS ONE* **7** e34842
- Campioli M, Schmidt N M, Albert K R, Leblans N, Ro-Poulsen H and Michelsen A 2013 Does warming affect growth rate and biomass production of shrubs in the High Arctic? *Plant Ecol.* **214** 1049–58
- Crawley M J 2007 *The R Book* (Chichester: John Wiley)
- CYSIP 2017 Central Yukon Species Inventory Project (CYSIP) (www.flora.dempstercountry.org) (Accessed: 17 August 2017)
- Eidesen P B, Carlsen T, Molau U and Brochmann C 2007 Repeatedly out of Beringia: *Cassiope tetragona* embraces the Arctic *J. Biogeogr.* **34** 1559–74
- Elmendorf S C *et al* 2012a Global assessment of experimental climate warming on tundra vegetation: heterogeneity over space and time *Ecol. Lett.* **15** 164–75
- Elmendorf S C *et al* 2012b Plot-scale evidence of tundra vegetation change and links to recent summer warming *Nat. Clim. Change* **2** 453–7
- Erickson T A, Williams M W and Winstral A 2005 Persistence of topographic controls on the spatial distribution of snow in rugged mountain terrain, Colorado, United States *Water Resour. Res.* **41** W04014
- Forbes B C, Macias-Fauria M and Zetterberg P 2010 Russian Arctic warming and 'greening' are closely tracked by tundra shrub willows *Glob. Change Biol.* **16** 1542–54
- Fraser R H, Lantz T C, Olthof I, Kokelj S V and Sims R A 2014 Warming-induced shrub expansion and lichen decline in the western Canadian Arctic *Ecosystems* **17** 1151–68
- Goetz S J *et al* 2011 Recent changes in Arctic vegetation: satellite observations and simulation model predictions *Eurasian Arctic Land Cover and Land Use in a Changing Climate* ed G Gutman and A Reissell (Dordrecht: Springer) pp 9–36
- Goetz S J, Bunn A G, Fiske G J and Houghton R A 2005 Satellite-observed photosynthetic trends across boreal North America associated with climate and fire disturbance *Proc. Natl Acad. Sci. USA* **102** 13521–5
- Guay K C, Beck P S A, Berner L T, Goetz S J, Baccini A and Buermann W 2014 Vegetation productivity patterns at high northern latitudes: a multi-sensor satellite data assessment *Glob. Change Biol.* **20** 3147–58
- Harris I, Jones P D, Osborn T J and Lister D H 2014 Updated high-resolution grids of monthly climatic observations—the CRU TS3.10 Dataset *Int. J. Climatol.* **34** 623–42
- Havström M, Callaghan T V and Jonasson S 1993 Differential growth responses of *cassiope tetragona*, an Arctic dwarf-shrub, to environmental perturbations among three contrasting high- and subarctic sites *Oikos* **66** 389–402
- Havström M, Callaghan T V, Jonasson S and Svoboda J 1995 Little ice age temperature estimated by growth and flowering differences between subfossil and extant shoots of *Cassiope tetragona*, an Arctic heather *Funct. Ecol.* **9** 650–4

- Higuera P E, Brubaker L B, Anderson P M, Brown T A, Kennedy A T and Hu F S 2008 Frequent fires in ancient shrub tundra: Implications of paleorecords for Arctic environmental change *PLoS ONE* **3** e0001744
- Hu F S, Lee B Y, Kaufman D S, Yoneji S, Nelson D M and Henne P D 2002 Response of tundra ecosystem in southwestern Alaska to younger-dryas climatic oscillation *Glob. Change Biol.* **8** 1156–63
- Hudson J M G and Henry G H R 2009 Increased plant biomass in a high Arctic heath community from 1981–2008 *Ecology* **90** 2657–63
- Hudson J M G, Henry G H R and Cornwell W K 2011 Taller and larger: shifts in Arctic tundra leaf traits after 16 years of experimental warming *Glob. Change Biol.* **17** 1013–21
- Jia G J, Epstein H E and Walker D A 2003 Greening of arctic Alaska, 1981–2001 *Geophys. Res. Lett.* **30** 2067
- Johnson J B and Omland K S 2004 Model selection in ecology and evolution *Trends Ecol. Evol.* **19** 101–8
- Johnstone J F and Henry G H R 1997 Retrospective analysis of growth and reproduction in *Cassiope tetragona* and relations to climate in the Canadian high Arctic *Arctic Alpine Res.* **29** 459–69
- Ju J and Masek J G 2016 The vegetation greenness trend in Canada and US Alaska from 1984–2012 Landsat data *Remote Sens. Environ.* **176** 1–16
- Keuper F, Parmentier F-J W, Blok D, van Bodegom P M, Dorrepaal E, van Hal J R, van Logtestijn R S P and Aerts R 2012 Tundra in the rain: differential vegetation responses to three years of experimentally doubled summer precipitation in Siberian shrub and Swedish bog tundra *Ambio* **41** 269–80
- Macías-Fauria M, Forbes B C, Zetterberg P and Kumpula T 2012 Eurasian Arctic greening reveals teleconnections and the potential for structurally novel ecosystems *Nat. Clim. Change* **2** 613–8
- Mallik A U, Wdowiak J V and Cooper E J 2011 Growth and reproductive responses of *Cassiope tetragona*, a circumpolar evergreen shrub, to experimentally delayed snowmelt *Arctic Antarct. Alpine Res.* **43** 404–9
- Myers-Smith I H *et al* 2011 Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities *Environ. Res. Lett.* **6** 45509
- Myers-Smith I H *et al* 2015 Climate sensitivity of shrub growth across the tundra biome *Nat. Clim. Change* **5** 887–91
- Myers-Smith I H and Hik D S 2018 Climate warming as a driver of tundra shrubline advance *J. Ecol.* in preparation (<https://doi.org/10.1111/1365-2745.12817>)
- Nakagawa S and Schielzeth H 2013 A general and simple method for obtaining R^2 from generalized linear mixed-effects models *Methods Ecol. Evol.* **4** 133–42
- Pepin N *et al* 2015 Elevation-dependent warming in mountain regions of the world *Nat. Clim. Change* **5** 424–30
- Pinheiro J, Bates D, DebRoy S and Sarkar D 2017 *NLME: linear and nonlinear mixed effects models* (R package version 3.1–131)
- R Core Team 2017 *R: A language and environment for statistical computing* (Vienna: R Foundation for Statistical Computing)
- Rayback S A and Henry G H R 2006 Reconstruction of summer temperature for a Canadian high Arctic site from retrospective analysis of the dwarf shrub, *Cassiope tetragona* *Arctic Antarct. Alpine Res.* **38** 228–38
- Rozema J, Weijers S, Broekman R, Blokker P, Buizer B, Werleman C, El Yaqine H, Hoogedoorn H, Fuertes M M and Cooper E J 2009 Annual growth of *Cassiope tetragona* as a proxy for Arctic climate: developing correlative and experimental transfer functions to reconstruct past summer temperature on a millennial time scale *Glob. Change Biol.* **15** 1703–15
- Settele J *et al* 2014 Terrestrial and inland water systems *Climate Change 2014: Impacts, Adaptation, and Vulnerability*. ed C B Field *et al* (Cambridge: Cambridge University Press) pp 271–359
- Starr G and Oberbauer S F 2003 Photosynthesis of Arctic evergreens under snow: implications for tundra ecosystem carbon balance *Ecology* **84** 1415–20
- Sturm M, Racine C and Tape K 2001 Climate change: Increasing shrub abundance in the Arctic *Nature* **411** 546–7
- Tape K, Sturm M and Racine C 2006 The evidence for shrub expansion in northern Alaska and the pan-Arctic *Glob. Change Biol.* **12** 686–702
- Tape K D, Hallinger M, Welker J M and Ruess R W 2012 Landscape heterogeneity of shrub expansion in Arctic Alaska *Ecosystems* **15** 711–24
- Trouet V and van Oldenborgh G J 2013 KNMI climate explorer: a web-based research tool for high-resolution paleoclimatology *Tree Ring Res.* **69** 3–13
- Tucker C J, Pinzon J E, Brown M E, Slayback D A, Pak E W, Mahoney R, Vermote E F and El Saleous N 2005 An extended AVHRR 8-km NDVI dataset compatible with MODIS and SPOT vegetation NDVI data *Int. J. Remote Sens.* **26** 4485–98
- Velichko A A, Andreev A A and Klimanov V A 1997 Climate and vegetation dynamics in the tundra and forest zone during the late glacial and holocene *Quatern. Int.* **41** 71–96
- Verbyla D 2008 The greening and browning of Alaska based on 1982–2003 satellite data *Glob. Ecol. Biogeogr.* **17** 547–55
- Vowles T, Gunnarsson B, Molau U, Hickler T, Klemmedtsson L and Björk R G 2017 Expansion of deciduous tall shrubs but not evergreen dwarf shrubs inhibited by reindeer in scandes mountain range *J. Ecol.* **105** 1547–61
- Walker M D *et al* 2006 Plant community responses to experimental warming across the tundra biome *Proc. Natl Acad. Sci. USA* **103** 1342–6
- Wang T, Hamann A and Spittlehouse D 2017 ClimateWNA v5: A program to generate climate normal, annual, seasonal and monthly data for historical and future periods in western North America
- Wang T, Hamann A, Spittlehouse D and Carroll C 2016 Locally downscaled and spatially customizable climate data for historical and future periods for North America *PLoS ONE* **11** e0156720
- Weijers S, Broekman R and Rozema J 2010 Dendrochronology in the high Arctic: July air temperatures reconstructed from annual shoot length growth of the circumpolar dwarf shrub *Cassiope tetragona* *Quatern. Sci. Rev.* **29** 3831–42
- Weijers S, Buchwal A, Blok D, Löffler J and Elberling B 2017 High Arctic summer warming tracked by increased *Cassiope tetragona* growth in the world's northernmost polar desert *Glob. Change Biol.* **23** 5006–20
- Weijers S, Greve Alsos I, Bronken Eidesen P, Broekman R, Loonen M J J E and Rozema J 2012 No divergence in *Cassiope tetragona*: persistence of growth response along a latitudinal temperature gradient and under multi-year experimental warming *Ann. Bot.* **110** 653–65
- Weijers S, Auliahierliaty L, van Logtestijn R S P and Rozema J 2013a Effects of manipulated precipitation and shading on *Cassiope tetragona* growth and carbon isotope discrimination: a high Arctic field study *Arctic Antarct. Alpine Res.* **45** 132–42
- Weijers S, Wagner-Cremer F, Sass-Klaassen U, Broekman R and Rozema J 2013b Reconstructing High Arctic growing season intensity from shoot length growth of a dwarf shrub *Holocene* **23** 721–31
- Zang C and Biondi F 2015 treeclim: an R package for the numerical calibration of proxy-climate relationships *Ecography* **38** 431–6