

Analysis of global Napier grass (*Cenchrus purpureus*) collections reveals high genetic diversity among genotypes with some redundancy between collections

Meki Muktar

International Livestock Research Institute

Tadelech Bizuneh

Ethiopian Institute of Agricultural Research, Holeta agricultural research centre

William Anderson

USDA-ARS

Yilikal Assefa

International Livestock Research Institute

Alemayehu T. Negawo

International Livestock Research Institute

Abel Teshome

International Livestock Research Institute

Ermias Habte

International Livestock Research Institute

Alice Muchugi

International Livestock Research Institute

Tileye Feyissa

Addis Ababa University

Chris S. Jones (✉ c.s.jones@cgiar.org)

International Livestock Research Institute

Article

Keywords:

Posted Date: May 30th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2972072/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on September 4th, 2023. See the published version at <https://doi.org/10.1038/s41598-023-41583-7>.

Abstract

Genetic diversity amongst genotypes of several Napier grass collections was analyzed and compared with the diversity in a set of open pollinated progeny plants. A total of 114,886 SNP and 46,293 SilicoDArT genome-wide markers were generated on 574 Napier grass genotypes. Of these, 89% of the SNP and 76% of the SilicoDArT markers were mapped onto the fourteen chromosomes of the Napier grass genome. For genetic diversity analysis, a subset of highly polymorphic and informative SNP markers, filtered using a maximum of 10% missing values, a minimum minor allele frequency of 5%, and a maximum linkage-disequilibrium value of 0.5, from those mapped across the Napier grass genome was used. Extensive genetic variability, with an average Nei's genetic distance value of 0.23, was identified in the material. The genotypes clustered into three major and eleven sub-clusters with high levels of genetic variation contained both within (54%) and between (46%) clusters. However, we found that there was low to moderate genetic differentiation among the collections and some overlap and redundancy occurred between collections. The progeny plants were genetically diverse and diverged from the germplasm collections, with an average F_{ST} value of 0.08. The findings of this study offer useful information for Napier grass breeding strategies, enhancement of genetic diversity, and provide a guide for the management and conservation of the collections.

Introduction

Napier grass (*Cenchrus purpureus* (Schumach.) Morrone syn. *Pennisetum purpureum* Schumach.), also known as elephant grass, is a monocotyledonous perennial C4 flowering plant belonging to the grass family Poaceae and the genus *Cenchrus*, one of the richest genera, consisting of about 140 species including important cultivated species such as Napier grass, pearl millet and kikuyu grass^{1,2,3}. Napier grass is native to Sub-Saharan Africa and has been distributed to other tropical and subtropical regions around the world, adapting to a wide range of agro-ecologies, from sea level to 2500 meters above sea level (m.a.s.l.), and has been acclimatized in areas of Central and South America, tropical parts of Asia, Australia, the Middle East and the Pacific islands^{1,4}.

Napier grass is mainly cultivated as a forage crop for animal feed due to its high palatability and feed quality, high dry matter production (up to 78 tons of dry matter/ ha/ year), and year-round availability under supplementary irrigation^{1,5,6,7}. In addition, Napier grass has the potential to produce biofuels such as alcohol, ethanol and butanol, and methane since it has high cellulose content that can be used as a carbon (energy) source^{8,9,10,11}. It has also many more beneficial characteristics, which include resistance to most pests and diseases, ease of establishment and fast regrowth capacity withstanding repeated cuttings, resistance to high temperatures, and low water and nutrient requirements^{4,12}. However, its productivity is primarily checked by the unpredictable weather arising from changes in climatic conditions^{13,14} and biotic stressors, such as the head smut and Napier grass stunt diseases, particularly in Central and Eastern African countries^{15,16,17}.

The availability of genetic diversity is a prerequisite for breeding and genetic improvement programs to identify novel alleles for agronomically important traits, such as disease and pest resistance, drought tolerance, and to withstand the current challenges associated with climate change^{1,18}. Napier grass is a cross-pollinating allotetraploid species with a chromosome number of $2n = 4x = 28$ (genome A'A'BB)^{19,20}. The Napier grass A'A' genome has been reported to be homologous to the AA genome of pearl millet (*Cenchrus americanus* Morrone syn. *Pennisetum glaucum*; $2n = 2x = 14$; AA)²¹. and the two species are cross-compatible producing triploid (AA'B genome) and sterile hybrids that can only be propagated by vegetative means, which ensures a true-to-type variety^{1,22,23}. The hybrids are favored mainly as a forage as they combine the superior forage quality of pearl millet with the high yielding ability of Napier grass¹. A number of important feed quality traits, including nutritional quality, have been introgressed into the genome of Napier grass from pearl millet through conventional breeding^{1,4,24,25} and hybrids have become a crucial part of the improved forage crop value chain in Africa, Asia and South America^{1,4}. Several national and international genebanks have conserved a substantial number of Napier grass genotypes collected from various geographical regions, for example: the International Livestock Research Institute (ILRI); the Brazilian Agricultural Research Corporation, Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA), United States Department of Agriculture (USDA-ARS), International Crops Research Institute for the Semi-Arid Tropics (ICRISAT) and the Millennium Seed Bank Kew (<https://www.genesys-pgr.org>). Although Napier grass exhibits high genetic variation due to its strict out crossing and self-incompatibility nature, it is expected that the global diversity is low due to the vegetative propagation nature and global use of cuttings as planting material, which could have limited the exchange of genes between genotypes. However, limited information exists about the overall available genetic diversity and germplasm exchange among the global collections.

The self-incompatibility and outcrossing nature of Napier grass^{20,26} results in a mixed lot of seeds from a genotype, thus progeny plants are heterogeneous and unpredictable in terms of performance¹. Additionally, reports of poor seed-setting, seed shattering, low seed germination, and weak seedlings indicated that seed propagation of this grass would be difficult⁴. However, in our experience the majority of the ILRI Napier grass genotypes growing in the Bishoftu and Ziway conservation sites can produce viable seeds²⁷, and their progeny plants have been grown with varying degrees of performance, indicating their potential for breeding and improvement of Napier grass. Progeny selection from open pollination and targeted crossing have been the breeding approaches in Napier grass and have resulted in the production of many improved cultivars¹. High biomass producing tall cultivar 'Merkeron' and the high yielding, nutritious dwarf leafy cultivar 'Mott', which is more suitable for grazing, are among the well-known cultivars produced from targeted crossing^{1,28,29,30}.

Wider genetic diversity is an essential source of novel and useful alleles for improvement of a species by breeding³¹ and will have an immediate practical impact on genetic improvement and productivity¹⁸. In this study, we analyzed and compared the between and within genetic diversity in Napier grass worldwide collections and included progeny plants raised from seeds using high density genome-wide markers, with

the aim of assessing the overall genetic diversity in the global collections and comparing it with the diversity in progeny plants to generate information useful for designing breeding strategies in Napier grass.

Results

Napier grass populations used in the study

With the aim of enhancing the genetic diversity and increasing the population size of the Napier grass collection held in the ILRI forage genebank, we analysed 32 new genotypes from ICRISAT, 93 from KALRO, 86 from EMBRAPA, and 23 from the USDA-ARS collections. The genotypes from the USDA-ARS were selected, based on a previous genetic diversity study on the collection²⁸, to represent as much of the diversity as possible held in the USDA-ARS's collection. In addition, we raised more than 219 progeny plants from 13 seed setting Napier grass genotypes of the ILRI collection (7 to 22 progeny plants per seed parent) (Supplementary Table S2), with the aim of releasing the genetic diversity held in the collection by dissecting the previously observed long haplotype blocks¹⁸. An additional 21 genotypes, composed of two commercial varieties, Maralfalfa and Super-Napier (*Cenchrus purpureus* x *Cenchrus americanus*), and other genotypes with unknown or different names, were included. We also included the 60 ILRI and 47 EMBRAPA genotypes, which have been studied previously^{18,32} to assess and visualize the overall diversity and identify new and unique genotypes. Out of these initially selected 581 Napier grass genotypes, individual genotypes with high missing values for the genome wide SNP markers were removed and a final set of 574 (Supplementary Table S1) genotypes were studied.

High density genome-wide DArTseq markers and their polymorphism

Genotyping generated 114,881 SNP and 46,293 SilicoDArT markers, distributed over the fourteen chromosomes of the Napier grass genome. Approximately 86% (98,535) of the SNP and 66% (30,745) of the SilicoDArT markers were aligned on to the fourteen chromosomes and about 2% (2,653 SNPs and 918 SilicoDArTs) of the markers were mapped onto different contigs of the Napier grass genome¹⁹. The highest number of markers were mapped onto chromosome CpA01 (8,057 SNPs and 2,947 SilicoDArTs) while the lowest number mapped to CpB07 (4,411 SNPs and 1,245 SilicoDArTs).

The percentages of missing values of the markers ranged from 0 to 22% for the SilicoDArT markers and 0 to 62% for the SNP markers. Heterozygosity (He) values for the SNP markers ranged from 0 to 0.5 with an average value of 0.15 while polymorphic information content (PIC) values ranged from 0 to 0.38 with an average value of 0.13 (Fig. 1). For the SilicoDArT markers, in which there is no difference between PIC and He, the values ranged from 0 to 0.38 with an average value of 0.15. More than 37% of the SilicoDArT markers and 30% of the SNP markers had values above 0.25 for PIC.

In order to assess the genetic diversity of the collections, a subset of 2,186 robust SNP markers were selected based on the marker's minor allele frequency ($MAF \geq 5\%$), missing values ($\leq 10\%$), independence from each other (Linkage disequilibrium- $LD \leq 0.5$), and their distribution across the Napier grass genome. The PIC and H_e values of the selected markers ranged from 0.08 to 0.38 and 0.09 to 0.50 with an average value of 0.26 and 0.32, respectively, and about 60% of the markers had values above 0.25 for PIC and H_e (Fig. 2). All the selected markers were able to be mapped onto the Napier grass genome¹⁹.

The overall genetic diversity among Napier grass genotypes from different collections

Genetic diversity and population structure in all genotypes from the different collections were analysed using the Bayesian algorithm implemented in STRUCTURE software, hierarchical clustering, and PCA analysis. In the STRUCTURE analysis, the delta K showed the presence of three major and up to sixteen subclusters (Fig. 3A and B). At a 60% membership probability threshold of $k = 3$, 158 genotypes were assigned to Cluster I, 246 genotypes to Cluster II, and 170 genotypes to Cluster III (Supplementary Table S3). Cluster I was composed mainly of genotypes from the ILRI collection (25 out of 59, 42%), progeny plants (83 out of 219, 38%) and the ICRISAT collection (10 out of 26, 38%). In addition, about 20% of genotypes from KALRO (19 out of 93), 11% (15 out of 133) from EMBRAPA, and 9% (2 out of 23) from USDA-ARS collections were clustered in this group. Cluster II contains the majority of the EMBRAPA (106 out of 133, 86%), KALRO (74 out of 93, 80%), and ILRI (31 out of 59, 53%) collections and about half of the genotypes from the ICRISAT and USDA-ARS collections. Only three progeny plants from NS-12 (Napier grass seedling number 12) were clustered in this group. Cluster III was mostly represented by progeny plants (133 out of 219, 61%) followed by USDA-ARS genotypes (10 out of 23, 43%) and three genotypes from ICRISAT, twelve from EMBRAPA, and three from the ILRI collection. Each major cluster was further partitioned into two to five subclusters that made eleven optimum number of subclusters and three admixture groups (Supplementary Table S3). A total of 242 genotypes, composed of 69% of the ICRISAT collection, 49% of the progeny, 47%, 43%, 37%, and 15% of the EMBRAPA, USDA-ARS, ILRI and KALRO collections, respectively, were under three different admixture groups. The three major clusters and the eleven subclusters were also detected by hierarchical clustering, in which the genotypes assignment in each major and the subclusters were mostly consistent with the clusters detected by the STRUCTURE analysis (Fig. 3C). A similar result was observed by the PCA, in which the first and second principal components explained about 16% of the molecular variance (Fig. 3D).

The pairwise F_{ST} values ranged from 0.14 to 0.15 in the three major clusters and from 0.17 (between sub-clusters I and II) to 0.47 (between sub-clusters III and VI) in the eleven sub-clusters (Table 1). AMOVA showed significant ($P < 0.01$) genetic variation among and within the major as well as the sub-clusters (Table 2).

Table 1. The N_e 's genetic distance (above the diagonal) and estimated fixation index (F_{ST}) (below the diagonal) between the eleven sub-clusters.

SC	SC-I	SC-II	SC-III	SC-IV	SC-V	SC-VI	SC-VII	SC-VIII	SC-IX	SC-X	SC-XI
SC-I	0.00	0.09	0.09	0.10	0.16	0.22	0.15	0.14	0.20	0.15	0.16
SC-II	0.17	0.00	0.10	0.10	0.12	0.19	0.12	0.11	0.17	0.13	0.13
SC-III	0.20	0.18	0.00	0.12	0.17	0.23	0.17	0.15	0.21	0.17	0.17
SC-IV	0.21	0.19	0.25	0.00	0.11	0.17	0.11	0.12	0.16	0.08	0.13
SC-V	0.33	0.25	0.36	0.26	0.00	0.11	0.13	0.14	0.17	0.16	0.17
SC-VI	0.45	0.35	0.47	0.37	0.33	0.00	0.11	0.16	0.12	0.15	0.18
SC-VII	0.28	0.22	0.30	0.22	0.27	0.25	0.00	0.09	0.10	0.08	0.11
SC-VIII	0.26	0.20	0.29	0.23	0.29	0.34	0.20	0.00	0.15	0.12	0.14
SC-IX	0.42	0.32	0.44	0.35	0.40	0.37	0.23	0.33	0.00	0.15	0.18
SC-X	0.30	0.25	0.33	0.20	0.33	0.34	0.19	0.26	0.33	0.00	0.10
SC-XI	0.30	0.23	0.32	0.25	0.34	0.40	0.22	0.26	0.38	0.23	0.00

SC = sub-cluster

Table 2. Analysis of molecular variance (AMOVA) showing the genetic variance among and within the three major clusters and the eleven sub-clusters.

Clusters	Source	df	SS	MS	P	Variance (%)
Major clusters	Between clusters	2	55931.66	27965.83	0.001	23.97
	Within clusters	404	268629.39	664.92	0.001	76.03
	Total	406	324561.05	799.41		100.00
Sub clusters	Between sub-clusters	10	114326.6	11432.66	0.001	46.17
	Within sub-clusters	321	141540.68	440.94	0.001	53.83
	Total	331	255867.28	773.01		100.00
Df = degrees of freedom; SS = sum of squares; MS = mean squares						

Table 3. Nei's genetic distance (above the diagonal) and estimated fixation index (F_{ST}) (below the diagonal) between the five Napier grass collections and progeny plants.

Collection	Progeny	EMBRAPA	USDA-ARS	ICRISAT	ILRI	KALRO
Progeny	0.000	0.040	0.040	0.041	0.035	0.050
EMBRAPA	0.079	0.000	0.021	0.032	0.018	0.029
USDA-ARS	0.071	0.031	0.000	0.037	0.022	0.039
ICRISAT	0.071	0.050	0.048	0.000	0.029	0.040
ILRI	0.067	0.032	0.024	0.029	0.000	0.018
KALRO	0.101	0.058	0.072	0.072	0.034	0.000

Genetic diversity between and within the collections and progeny plants

The genetic diversity between and within the five collections (ILRI, EMBRAPA, USDA-ARS, ICRISAT, KALRO) and progeny plants were compared using Nei's genetic distance³³ and fixation index (F_{ST})³⁴. The pairwise Nei's genetic distance ranged from 0.018, between ILRI and KALRO/ EMBRAPA, to 0.05, between the KALRO collection and progeny plants. Similarly, the estimated F_{ST} values ranged from 0.029, between ILRI and ICRISAT collections, to 0.101, between the KALRO collection and progeny plants (Table 3) which reflects moderate genetic variation between the collections. However, the variation among the five collections was low, the moderate variation observed was mainly between the progeny plants and the five collections (Table 3; Fig. 4A), indicating that there was unique genetic makeup in the progeny plants. Variance components calculated by AMOVA were also significant ($P < 0.001$) both between and within collections, however, higher variation was detected within (90%) than between (10%) collections (Table 4). The range and average of Nei's genetic distance within each collection and the progeny plants is shown in Fig. 4B. Higher within genetic diversity was observed in the ICRISAT collection (0.29 average Nei's genetic distance), followed by the ILRI collection (0.22), and the progeny plants (0.22).

Table 4. Analysis of molecular variance (AMOVA) showing the genetic variance between and within the five Napier grass collections and progeny plants.

Source	df	SS	MS	<i>P</i>	Variance (%)
Between Collections	5	45225.229	9045.046	0.001	10
Within Collections	1142	669442.872	586.202	0.001	90
Total	1147	714668.100			100
Df = degrees of freedom; SS = sum of squares; MS = mean square					

Table 5. Number of duplicated genotypes between collections.

Collection	EMBRAPA	KALRO	ILRI	ICRISAT	USDA-ARS	Progeny
EMBRAPA	0					
KALRO	9	0				
ILRI	7	45	0			
ICRISAT	0	0	0	0		
USDA-ARS	3	4	3	0	0	
Progeny	0	0	0	0	0	0

Potential duplicate genotypes in each collection were identified using Nei's genetic distance at a threshold of 0.025, which was equivalent to the identity-by-descent (IBD) kinship threshold ($k \geq 0.45$) for self, or duplicate, identification. A greater number of potential duplicates were detected in the KALRO collection, while all genotypes from ICRISAT were distinct (Fig. 5). Across collections, the largest overlap was between the ILRI and KALRO collections, followed by the KALRO and EMBRAPA, and the ILRI and EMBRAPA collections (Table 5). From the overall collection of 574 genotypes composed of the six collections, a total of 432 distinct (unique) genotypes that could be used in downstream genetic studies were selected (Supplementary Table S4).

Parentage analysis in the progeny plants

We estimated the parentage of the 219 progeny plants, raised from 13 ILRI open pollinated seed-setting genotypes, using identity-by-descent (IBD)³⁵ and Mendelian-inconsistent errors^{36,37} analyses. First, we examined the genetic relationship between the recorded seed parents and progeny plants by the IBD

method of the maximum likelihood (ML) relatedness estimator³⁵, and these were corrected based on the IBD result whenever there was a discrepancy between the recorded seed parent and the IBD result.

The most likely pollen donor of the progeny plants was analyzed by counting the Mendelian-inconsistent errors, considering a low percentage of errors as an indication of a putative parent-offspring relationship^{36,37}. The Mendelian error counts ranged from 2 to 316 for PC1 (parent-child relationship for potential seed parent), 1 to 401 for PC2 (parent-child relationship for potential pollen parent), and 37 to 937 for PPC (parent-parent-child relationship for possible crosses or as potential complete parents), respectively. In general, we identified or confirmed seed parents, potential pollen parents, and possible crosses as complete parents for 189 (86%) of the progeny plants (Supplementary Table S5). In addition, we classified the estimated relationships into strong-evidence vs weak-evidence based on the Mendelian error counts of the progeny plants for which the recorded seed parents were confirmed by the IBD analysis and determined a threshold for each relationship. A distinct gap in Mendelian error counts was observed at about 52 for PC1, at about 99 for PC2, and at about 94 for PPC, however, the gap for PC1 and PC2 was less clear compared to the gap for PPC (Fig. 6). Therefore, the Mendelian error counts for 53 (28%) of the PPC relationships that showed less than 94 Mendelian error counts were considered as real parents with strong evidence (Supplementary Table S5). We identified eleven ILRI genotypes as potential seed and pollen parents, however, none of the progeny plants showed parent-child relationships with two of the recorded ILRI parent genotypes (ILRI_P2_16789 and ILRI_P4_16783). ILRI_P1_1026 was the dominant pollen donor, followed by ILRI_P9_16821, ILRI_P8_16803, and ILRI_P7_16837; while ILRI_P11_16810 was the most infrequent pollen donor.

Discussion

It is thanks to the diversity of plant genetic resources which serves as a source of new alleles for the development of new cultivars with desirable qualities, including breeder' and farmers' preferred traits such as high biomass yield, high nutritional quality, larger seeds, and pest and disease resistance^{1,4,31}. In order to assess the level of global genetic variation in Napier grass, diverse set of genotypes acquired from five international institution collections were used in this study. Each collection had a different number of genotypes, ranging from 26 in USDA-ARS to 133 in EMBRAPA. In addition, progeny plants raised from 13 open pollinated seed-setting ILRI genotypes were evaluated for genetic diversity and compared to the diversity held in the collections with the aim of enhancing the genetic diversity and generating information useful for designing breeding strategies for Napier grass. Each collection represented a different region of the world; for example, the ILRI and KALRO collections were mostly from Africa^{18,38}, while the EMBRAPA materials were mainly from Brazil^{39,40}. Similarly, most of the genotypes from ICRISAT's collection had an Asian origin⁴¹. The USDA-ARS materials were from Latin-America (mainly from Puerto Rico), Africa (mainly from Kenya), and Tifton (genotypes derived from the cultivar "Merkeron" in a breeding program)^{28,42}.

We employed the GBS method of the DArTseq technology⁴³, an efficient and cost-effective sequencing technology that concurrently generates the dominant SilicoDArT and the co-dominant SNP markers (<https://www.diversityarrays.com/services/dartseq/>) with a high density and genome coverage. Similarly to our previous reports^{5,18,44}, we obtained a large number, 114,881 SNP and 46,293 SilicoDArT, of markers distributed over the fourteen chromosomes of the Napier grass genome, and more than 80% of the markers (from both marker types) were able to be mapped on to the genome, approximately equally on to the A' and B sub genomes¹⁹. We applied strong SNP filtering criteria by setting a maximum of 10% missing values, a minimum of 5% minor allele frequency, a maximum of 0.5 Linkage-disequilibrium (LD) value to ensure independent markers and, markers that are able to be mapped across the genome. The filtered polymorphic markers, with average PIC and He values of 0.26 and 0.32, respectively, were used in the genetic diversity analysis.

Genetic diversity in worldwide Napier grass collections

The Napier grass genotypes from the worldwide collections and progeny plants examined in this study demonstrated substantial genetic variability with an average value of 0.23 (0 to 0.41 range) Nei's genetic diversity coefficients³³. These genetic distance values were greater than the range that Wanjala et al. (2013)⁴⁵ reported for 281 Napier grass genotypes. The high genetic variability detected could be attributed to the high levels of heterozygosity that exists in Napier grass due to its self-incompatibility and obligate outcrossing nature²⁶. Napier grass is a highly heterozygous tetraploid species with a wide parental diversity and rich gene pool^{23,40}.

Furthermore, we detected three major clusters and eleven sub-clusters that were significantly different from each other, based on the admixture model in STRUCTURE, hierarchical clustering, PCA, and AMOVA analysis. The F_{ST} values ranged from 0.14 to 0.15 between the three clusters and 0.17 to 0.47 between the 11 sub-clusters. This substantial genetic variation was detected both within and between clusters, while within-cluster variation was slightly larger (54%) than between-cluster variation (46%). Generally, outcrossing species show higher levels of genetic variation among genotypes and less population divergence, compared to self-pollinating species⁴⁶. Similar results were reported in other outcrossing species such as Rhodes grass (*Chloris gayana*)⁴⁷ and perennial rye grass (*Lolium perenne*)⁴⁸. Nevertheless, about 46% of the genetic variation was detected between the eleven subclusters. This higher divergence could be explained by the clonal propagation of Napier grass through stem cuttings, which reduces genetic drift and gene flow among genotypes within a population^{45,46,49}. Genotype adaptation to specific environmental conditions might have also contributed to the high genetic diversity among the clusters, given that the genotypes were collected from different parts of the world.

With an F_{ST} value of about 0.47, cluster III (which consisted of progeny plants, mainly from NS-11) and cluster VI (mainly from the ILRI and KALRO collections) had the highest genetic distance, followed by between cluster I (Progeny plants, mainly from two seed parents, NS-6 and NS-9) and cluster VI, and cluster III and IX (comprised mainly of genotypes from EMBRAPA, ILRI, and KALRO). The results indicated

a divergence between the collections and progenies, showing the potential of the progeny plants, that were produced from open pollination, for enhancement of genetic diversity in Napier grass collections and potential for future use in breeding programs, such as heterosis breeding. Heterotic contributions to increased biomass production in the F1 hybrids of Napier grass have previously been reported^{50,51}.

The ILRI genotypes were distributed across sub-clusters, mainly in sub-clusters II, VI, VII, IX, and the admixture groups, and their clustering was mostly consistent with our previous report¹⁸. The genotypes in cluster II are low to medium biomass yielders, whereas the genotypes in clusters VI and VII are medium to high biomass yielders⁵. From the EMBRAPA collection, the majority of the elite lines (indicated by the acronym 'CNPGL' CNPGL in Supplemental Table S3) and the purple genotypes were in the admixture-2 group, while most of the genebank materials (indicated by the acronym 'BAGCE') were in Cluster VIII, which was mostly consistent with our previous report¹⁸ and reflected the breeding history of the collection. BAGCE_30 from cluster VIII and CNPGL-92-66-3 from the admixture group were reported to be high biomass yielding and drought tolerant genotypes under Ethiopian conditions⁵. Cluster V contained genotypes mainly from the KALRO collection, including Kakamega 1 and 2 genotypes, which have been reported to be tolerant to Napier grass head smut disease⁵²). This cluster included two important ILRI genotypes (ILRI_16791 and ILRI_16802), which have been identified for their superior biomass yields, drought tolerance, and comparatively higher protein content⁵.

Cluster XI was primarily comprised of USDA-ARS genotypes, with a few EMBRAPA genotypes. In line with the report by Harris et al., 2009²⁸, genotypes derived from the cultivar 'Merkeron' were clustered together in this group. 'Merkeron' is a well-known cultivar in the USA, released by a breeding program in Tifton, Georgia⁵⁰ and was derived from an intraspecific cross between a high yielding clone and a dwarf leafy clone, with improved yield and disease resistance²⁹. Among these, the well-known variety "Mott" (Tift N75), a dwarf leafy type selected from progeny of selfed Merkeron³⁰, is included in this cluster. The variety "Mott" was introduced to ILRI and has been preserved in the genebank with accession number ILRI_15743 and has been distributed to farmers and national research institutes in Ethiopia. However, none of the samples collected from farmers' fields and national research institutes, as well as the genotype labeled as ILRI_15743, clustered in this group. Similar findings were reported by Harris et al. (2009)²⁸, who found that none of the "Merkeron" samples taken from the various countries where "Merkeron" was distributed were genetically identical, which possibly happened due to genetic mixing up through outcrossing.

Cluster VII consisted of most *C. purpureus* × *C. americanus* hybrids including the commercial variety 'Maralfalfa'⁵³, 'Super-Napier', and the hybrids from the ILRI collection ILRI_16840, ILRI_16837, ILRI_15357¹⁸. The ILRI_15357 showed very high genetic similarity to 'Maralfalfa', with more than 0.47 IBD value. Maralfalfa has been reported for its adaptability to higher altitudes (3000 m asl) and for high biomass yield (60 tn/ha) with higher protein content⁵³. Many genotypes from Kiambu under the KALRO collection were also found in this cluster, perhaps suggesting that there are also hybrids held in this collection.

The progeny plants were almost exclusively distributed in clusters I, III, IV, and X, which were significantly different from each other with a range of F_{ST} values of 0.20 to 0.33 with an average value of 0.25. The progenies in cluster I and III were particularly distinct from the other clusters with an average F_{ST} value of 0.32, showing their divergence from genotypes and other progenies. The Napier grass's high heterozygosity within genotypes, general self-incompatibility and obligate outcrossing nature⁵⁴, which can increase the likelihood of developing new genetic make-up through recombination and produce distinct genotypes, may be responsible for this high level of diversity and population stratification in progeny plants.

The collections displayed less genetic divergence and different levels of genotype redundancy

The collections we studied generally exhibited low to moderate genetic differentiation with an F_{ST} range of 0.024 between the ILRI and USDA-ARS collections to 0.072 between the KALRO and ICRISAT collections. The genetic variation between collections explained only about 10% of the variation as compared to 90% of the variation explained by genotypes within collections, indicating a degree of overlap and redundancy that may have resulted from sharing and exchanging materials among the genebanks. Although Napier grass genotypes are primarily maintained via vegetative multiplication using cuttings, there is still some degree of hybridization existed in nature and also controlled crossing manner in some collections^{1,26}. Such redundancy of genotypes between genebanks could also be arisen because each genebank or breeding program gives its own different accession numbers, as there is no standardized labelling and registration procedure. Facilitated by the International Treaty on Plant Genetic Resources for Food and Agriculture, the ILRI forage genebank has assigned a digital object identifier (DOI) to each genotype held in its genebank^{18,47}, which could be a promising solution to the issue but should then be adopted by national and international genebanks and research institutions.

The presence of genotype overlaps and redundancy among the collections was further confirmed by Nei's genetic distance and identity-by-descent (IBD) analysis; the largest overlap was between the ILRI and KALRO collections, followed by between the KALRO and EMBRAPA, and the ILRI and EMBRAPA collections. The substantial overlap between the ILRI and KALRO collections could be due to the close proximity of the two institutes and historical sharing of genotypes, as indicated in the genotypes name, a large number of KALRO's genotypes were sourced from the ILRI collection. This suggests that the introduction and incorporation of new genotypes to a collection should be carried out strategically to minimize duplications and capture only new genetic diversity that is distinct from the existing one, as conservation is a costly business³².

Within collection genotype redundancy analysis, also showed a greater number of potential duplicates in the KALRO collection whereas all genotypes from ICRISAT were distinct. While genotype redundancies and overlaps between collections from various institutions, especially those from different continents like Africa, South America, Asia, and North America, can be taken into consideration and should be encouraged to preserve genotypes with specific adaptations to a wide range of farming systems and

agroecological conditions, overlaps within a collection should be reduced as it imposes a conservation cost. However, it is crucial to carefully remove overlaps using molecular marker analysis supported by field phenotype and passport data whenever possible in order to preserve unique alleles in closely related genotypes.

Greater genetic variation was detected in progeny plants

When compared to the variation among the five collections, the progeny plants were genetically separated, with F_{ST} values ranging from 0.067 between the progenies and ILRI collection to 0.101 between the progenies and KALRO collection. This finding further supports the potential of progeny plants to increase genetic diversity with the potential to be used in breeding programs in the future. Open-pollinated progeny selection has been one of the breeding approaches in Napier grass and has resulted in the generation of improved cultivars¹. Progeny plants from an open-pollinated population of cultivars 'A146', 'A148' and 'A149' were developed in Taiwan, and the progenies produced 20% more dry matter than the seed parents, had very few hairs on the leaves, flowered late and grew more erect^{1,55}. However, the use of open pollination breeding schemes allows the control of only one-half of the genome of the resulting progeny material because contributions from pollen parents are not controlled, and the transfer of undesirable traits from unimproved lines can limit or slow the successful release of new cultivars. Therefore, Napier grass breeding programs should focus on targeted crossing that have the potential to yield higher rates of success⁵⁰ in rapidly generating desirable genotypes and eventually culminating in hybrid breeding wherein a superior individual can be maintained by vegetative propagation after being identified.

In this study, we were able to identify potential seed and pollen parents for more than 80% of the progeny plants using identity-by-descent (IBD) analysis based on the maximum likelihood (ML) relatedness estimator and Mendelian-inconsistent errors^{36,37}. The progenies in clusters I and III were produced primarily from the ILRI_P9_16821 and ILRI_P6_16835 genomes, while the progenies in clusters IV and X were dominantly from ILRI_P7_16837 and ILRI_P13_16790 and ILRI_P1_1026 and ILRI_P8_16803 genomes, respectively. These results could offer a stronger basis for selecting superior progeny plants for further selection and improvement of Napier grass.

Conclusions and recommendations

1. This study represents the first comprehensive genetic diversity study in global Napier grass collections and revealed the presence of extensive genetic variation among Napier grass genotypes. The analysis detected major clusters and subclusters that were significantly different from each other with higher within cluster/sub-cluster genetic variation. The study provides a global picture of the available genetic diversity of this important tropical perennial crop and the findings could be used to guide targeted use and conservation of the diversity by breeders and genebanks. Additionally, the information can be used to determine heterotic groups for future Napier grass breeding and improvement.

2. The genetic variation amongst the different genebank collections studied here was generally low and various levels of genotype redundancy were detected both within and across collections. Genotype overlaps between genebanks, particularly those from different continents, should be encouraged in order to preserve genotypes with specific adaptations to a wide range of agroecological conditions. To reduce the cost of conservation, overlaps within a genebank should be minimized. However, overlaps within a genebank must be carefully removed to maintain unique alleles in closely related genotypes.
3. Higher genetic variation and divergence was detected in progeny plants produced from open pollination, which creates new genetic makeup and diversity. Incorporating some of the distinctive progeny plants into the genebank collections will be advantageous to enhance the genetic diversity and population size of the collections. However, a few genotypes predominated as a pollen source, as suggested by the parentage analysis, which invites further research on the mechanisms of pollination in Napier grass genotypes maintained in the ILRI forage genebank.

Methods

Plant Materials

A total of 581 Napier grass genotypes, comprising 219 progeny plants that were raised from open pollinated seeds and 341 genotypes from worldwide collections maintained by five institutes, were included in the study. These include 60 genotypes from the ILRI genebank collection, 133 genotypes from the EMBRAPA collection, 93 genotypes from the Kenya Agricultural and Livestock Research Organization (KALRO), 23 genotypes from the USDA-ARS and, 32 genotypes from the ICRISAT genebank. Two commercial varieties, Maralfalfa⁵³ and Super-Napier (Pakchong 1) - both reported as hybrids of Napier grass and pearl millet with high biomass yield and protein content, were included. An additional 19 genotypes, collected from different parts, with the name 'Mott' or with unknown names were also included (Supplementary Table S1).

About 40 to 50 Napier grass seeds, collected from 13 seed-bearing ILRI genotypes and maintained in the ILRI forage genebank, were obtained from the genebank and pre-germinated on agar medium containing potassium nitrate. The germinated seeds were transplanted into pots filled with soil and maintained in a screen house until they produced three to four leaves, at which point plant samples for DNA extraction were taken. On average, about 17 progeny plants from each of the 13 genotypes were sampled to assess the level of genetic diversity within and between the groups of progenies, as well as to assess the genetic diversity in progeny plants compared to the collections²⁷.

The ILRI forage genebank, EMBRAPA, KALRO, ICRISAT, or USDA-ARS provided all Napier grass samples, including seeds, leaves, or DNA samples. The studies were carried out in accordance with the guidelines and legislation outlined in the IUCN Policy Statement on Research Involving Species at Risk of Extinction and the Convention on the Trade in Endangered Species of Wild Fauna and Flora. Napier grass is a

popular forage species which is cultivated across the tropics and not at risk of extinction or a wild species.

DNA extraction and genotyping

Young leaf samples from each individual plant were collected into 2 ml Eppendorf tubes, stored in an icebox containing frozen ice, and transferred to a -80 freezer as quickly as possible. The samples were freeze-dried for about 48 hours and ground into fine powder using a tissue lyser. Genomic DNA was extracted from the leaf powder using a DNeasy® Plant Mini Kit (250) (Qiagen Inc., Valencia, CA) following the manufacturer's procedures. The concentration and quality of the extracted DNA was assessed using a Nano-drop spectrometer (DeNovix DS-11 FX spectrophotometer) and 0.8% agarose gel electrophoresis. The concentration of the DNA was adjusted to 50 to 100 ng/μl in 96 well semi-skirted plates and sent to SEQART-AFRICA (<https://www.seqart.net/>), Nairobi, Kenya for genotyping.

The genotyping was conducted by using the GBS (Genotyping By Sequencing) method of the DArTseq platform⁴³, which provides combined marker identification and genotyping for SNP and SilicoDArT (presence/absence of DNA fragments in genomic representations) markers, combining the DArT complexity reduction with sequencing on the Next Generation Sequencing (NGS) platforms^{18,43}. The markers were aligned with the Napier grass genome¹⁹ using the short sequence fragments associated with each marker to generate information on map position of the sequences and markers across the genome. A heatmap showing the density and distribution of markers across the fourteen Napier grass chromosomes was generated using a free online tool at <https://www.bioinformatics.com.cn/en>. For genetic diversity analysis, polymorphic markers were filtered based on the marker's minor allele frequency (MAF), missing values, and independence from one other (based on pairwise linkage disequilibrium, LD) using the dartR⁵⁶ and SNPRelate⁵⁷ R-packages in R statistical software (R Core Team, 2022).

Genetic diversity and population structure analysis

Population structure and sub clustering among the Napier grass genotypes was assessed based on the Bayesian algorithm implemented in STRUCTURE (ver. 2.3.4) software⁵⁸ using the filtered polymorphic markers. The burn-in time and number of iterations were both set to 100,000 with 10 repetitions, assuming hypothetical subpopulations (K) ranging from 1 to 20 in an admixture model with correlated allele frequencies. The results of the run were uploaded to the software "Structure Harvester" (<http://taylor0.biology.ucla.edu/structureHarvester/>) and the most likely number of subpopulations was determined by the Evanno method⁵⁹. Genotypes with less than 60% membership probability were considered admixed. Population structure was further examined using hierarchical clustering and principal component analysis (PCA) using Poppr⁶⁰ and adegenet⁶¹ R-packages. Genetic diversity was estimated using pairwise Nei's³³ genetic distance.

Pairwise estimates of population differentiation and divergence between clusters, subclusters, and collections were analysed by Analysis of MOlecular VAriance (AMOVA)⁶² and fixation index (F_{ST})³⁴ in the R-packages Poppr⁶⁰, adegenet⁶¹, and APE⁶³. AMOVA was used for partitioning the total variance in allele frequencies within and among clusters, subclusters, and collections and to assess the significance level of the variations. F_{ST} , which ranges from 0 to 1, assesses the degree of genetic differentiation between clusters, subclusters, and collections. A value of 0 indicates no differentiation while a value of 1 indicates complete differentiation. In general, an F_{ST} value of 0:00 to 0:05 indicates low differentiation, 0:05 to 0:15 indicates moderate differentiation, 0:15 to 0:25 high differentiation, while $F_{ST} > 0.25$ are considered to represent very high levels of differentiation⁶⁴.

Parentage analysis

Parentage and genetic relationship analyses were conducted using identity-by-descent (IBD) analysis based on the maximum likelihood (ML) relatedness estimator³⁵ using the R package SNPRelate⁵⁷ and by counting the Mendelian-inconsistent errors using a custom R script reported in Vanderzande et al. (2018)³⁶. The IBD analysis was mainly used to confirm the genetic relationship between the recorded seed parents and progeny plants, whereas Mendelian-inconsistent errors count was used to determine any putative set of parent-child (PC) and parent-parent-child (PPC) relationships. In the Mendelian-inconsistent errors count, a low percentage of errors was used as an indication of putative parent-offspring relationships while higher percentages were indicative of more distant relationships^{36,65}.

Declarations

Data Availability

Most of the datasets generated for the current study are found in the supplemental information, and further data can be obtained from the the corresponding author upon reasonable request. The ILRI accessions are freely available to researchers who accept the terms and conditions of the Standard Material Transfer Agreement (SMTA) of the International Treaty on Plant Genetic Resources for Food and Agriculture.

Acknowledgements

The research was conducted under the CGIAR Sustainable Animal Productivity (SAPLING) and the genebank initiatives, in Feed and Forage Development (FFD) program at the ILRI forage genebank in Addis Ababa, Ethiopia. CGIAR research is supported by contributions from the CGIAR Trust Fund. The authors would like to thank EMBRAPA, USDA-ARS, KALRO, ICRISAT, and the ILRI forage genebank for making their germplasm and breeding lines available for this study.

Author Contributions

C.S.J. designed the study and supervised the project and the manuscript writing, M.S.M. analysed the data and wrote the manuscript, W.A. and A.M. provided germplasms and suggested some topics for the manuscript, Y.A. and T.B. collected leaf samples and extracted DNA and sent samples for genotyping, A.T.N., A.T.G. and E.H.H. involved in seedling raising and management at Bishoftu site, T.F. supervised the data analysis. All authors contributed to the writing, review and approval of the final manuscript.

Additional Information

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

1. Singh, B. P., Singh, H. P. & Obeng, E. Elephant grass (ed. Singh, B. P.) 271–291 (CAB, 2013).
2. Chemisquy, M. A., Giussani, L. M., Scataglini, M. A., Kellogg, E. A. & Morrone, O. Phylogenetic studies favour the unification of *Pennisetum*, *Cenchrus* and *Odontelytrum* (Poaceae): A combined nuclear, plastid and morphological analysis, and nomenclatural combinations in *Cenchrus*. *Ann. Bot.* **106**, 107–130 (2010).
3. Kativu, S., Mithen, R. *Pennisetum* in Southern Africa. *Plant Genetic Resources Newsletter* 73/74:1-8 (1987).
4. Negawo, A. T., Teshome, A., Kumar, A., Hanson, J. & Jones, C. S. Opportunities for Napier grass (*Pennisetum purpureum*) improvement using molecular genetics. *Agronomy* **7**, 1-21, <https://doi.org/10.3390/agronomy7020028> (2017).
5. Habte, E. *et al.* Productivity and Feed Quality Performance of Napier Grass (*Cenchrus purpureus*) Genotypes Growing under Different Soil Moisture Levels. *Plants* **11**, <https://doi.org/10.3390/plants11192549> (2022).
6. Getiso, A., Mijena, D. & Com, P. Performance evaluation of Napier grass (*Penisetum Purpuruem* (L.) Schumach) accessions under rain fed and irrigation system at Wondo Genet. doi:10.17352/gje.0000141 (2021).
7. Oliveira, M. L. F., Daher, R. F., Menezes, B. R. S. & Vivas, M. Genetic diversity of elephant grass (*Cenchrus purpureus* [Schumach.] Morrone) for energetic production based on quantitative and multi-category traits. *Chil. J. Agric. Res.*, **77(1)**:48-57. (2017).
8. Roslan, A. M., Ibrahim, M. F. & Hassan, M. A. Potential use of *Pennisetum purpureum* for phytoremediation and bioenergy production: a mini review. *Sci. Rep.* **10**:6613. (2020).
9. Romero, J. *et al.* Unraveling candidate genes underlying biomass digestibility in elephant grass (*Cenchrus purpureus*). *MBC Plant Biol*, **19(1)**: 1–12. (2019).
10. Takara, D. & Khanal, S. K. Characterizing compositional changes of Napier grass at different stages of growth for biofuel and biobased products potential. *Bioresour. Technol.* **188**, 103–108 (2015).
11. Yasuda, M. *et al.* Enhancement of ethanol production from Napiergrass (*Pennisetum purpureum* Schumach) by a low-moisture anhydrous ammonia pretreatment. *J. Sustain. Bioenergy Syst*,

3(3):179. (2013).

12. Kebede, G. *et al.* Agronomic performance, dry matter yield stability and herbage quality of Napier grass (*Pennisetum purpureum* (L.) Schumach) accessions in different agro-ecological zones of Ethiopia. *J. Agric. Crop Res.* **5**, 49–65 (2017).
13. Nguyen, L. Van *et al.* Variation in Root Development Response of Napier Grass to Drought Stress. *J. Ecol. Eng.* **22**, 64–74 (2021).
14. Cardoso, J. A. *et al.* Contrasting strategies to cope with drought conditions by two tropical forage C4 grasses. *AoB Plants* **7**, 107 (2015).
15. Asudi, G. O. *et al.* Napier grass stunt disease in East Africa: Farmers' perspectives on disease management. *Crop Prot.* **71**, 116–124 (2015).
16. Jones, P., Devonshire, B. J., Holman, T. J. & Ajanga, S. Napier grass stunt: A new disease associated with a 16SrXI group phytoplasma in Kenya. *Plant Pathol.* **53**, 519 (2004).
17. Farrell, G. & Hillocks, R. J. Pests, diseases and weeds of Napier grass, *Pennisetum purpureum*: A review. *Int. J. Pest Manag.* **48**, 39–48 (2002).
18. Muktar, M. S. *et al.* Genotyping by sequencing provides new insights into the diversity of Napier grass (*Cenchrus purpureus*) and reveals variation in genome-wide LD patterns between collections. *Sci. Rep.* **9**, 1-15 (2019).
19. Yan, Q. *et al.* The elephant grass (*Cenchrus purpureus*) genome provides insights into anthocyanidin accumulation and fast growth. *Mol. Ecol. Resour.* **21**, 526–542 (2021).
20. Martel, E., De Nay, D., Siljak-Yakovlev, S., Brown, S. & Sarr, A. Genome size variation and basic chromosome number in pearl millet and fourteen related *Pennisetum* species. *J. Hered.* **88**, 139–143 (1997).
21. Anderson, W. F., Casler, M. D & Baldwin, B. S. Improvement of perennial forage species as feedstock for bioenergy (ed. Vermerris, W.) 347–376, <https://doi.org/10.1007/978-0-387-70805> (Springer Science + Business Media LLC, 2008).
22. Keerthana, K. *et al.* Identification of superior pearl millet Napier hybrids with enhanced green fodder yield and quality through genetic variability studies. *YMER*, (2022).
23. Robert *et al.* Wild Crop Relatives: Genomic and Breeding Resources, Millets and Grasses (ed. Kole, C.), 217-255 DOI 10.1007/978-3-642-14255-0_13 (Springer-Verlag Berlin Heidelberg, 2011).
24. Keerthana, K., Ganesan, K. N., Iyanar, K., Sivakumar, S. D. & Jeyakumar, P. Gene action and contribution of different traits for enhanced green fodder yield and quality in pearl millet - Napier interspecific hybrids. *Electron. J. Plant Breed.* **13**, 1066–1076 (2022).
25. Kaur, A., Kapoor, R., Vikal, Y., Kalia, A. & Bhardwaj, R. Production of Interspecific Hybrids between Pearl Millet [*Pennisetum glaucum* (L.) R. Br.] × Napier Grass [*Pennisetum purpureum* (K.) Schum] and their Characterization. *Int. J. Curr. Microbiol. Appl. Sci.* **8**, 1308–1313 (2019).
26. Hanna, W., Mathews, B. W., Burns, J. C. & Sollenberger, L. E. Perennial *Pennisetums* Nutrient Cycling in Tropical Pastures: What do we know? View project Optimizing Ecosystem Services of Long-Term

Grass-N and Legume-Based Grassland-Livestock Systems View project.

<https://www.researchgate.net/publication/292768672> (2004).

27. Bizuneh, T. Genetic Diversity Study of Napier Grass (*Cenchrus Purpureus* L.) Collections from Different Part of the World and Progeny Plants. *Thesis* (2021).
28. Harris, K., Anderson, W. & Malik, R. Genetic relationships among napiergrass (*Pennisetum purpureum* Schum.) nursery accessions using AFLP markers. *Plant Genet. Resour. Characterisation Util.* **8**, 63–70 (2009).
29. Burton, G.W. Registration of ‘Merkeron’ Napiergrass. *Crop Science* **29**, 1327 (1989).
30. Hanna, W.W. & Monson, W.G. Registration of dwarf Tift N75 Napiergrass germplasm. *Crop Science* **28**, 870–871 (1988).
31. Bhandari, H. *et al.* Assessment of genetic diversity in crop plants - an overview. *Adv. Plants Agric. Res.* 279–286 DOI: 10.15406/apar.2017.07.00255 (2017).
32. Negawo, A. T. *et al.* Molecular markers as a tool for germplasm acquisition to enhance the genetic diversity of a Napier grass (*Cenchrus purpureus* syn. *Pennisetum purpureum*) collection. *Trop. Grasslands-Forrajes Trop.* **6**, 58–69 (2018).
33. Nei, M. Genetic distance between populations. *Am. Nat.* (1972).
34. Wright, S. The genetical structure of populations. *Annals of Eugenics* **15**, 323-354 (1951).
35. Milligan, B. G. Maximum-likelihood estimation of relatedness. *Genetics* **163**, 1153-1167 (2003).
36. Vanderzande, S. *et al.* High-quality, genome-wide SNP genotypic data for pedigreed germplasm of the diploid outbreeding species apple, peach, and sweet cherry through a common workflow. *PLoS One* **14**, (2018).
37. Cheung, C. Y. K., Thompson, E. A., Wijsman, E. M. Detection of Mendelian consistent genotyping errors in pedigrees. *Genet Epidemiol.* **38**, 291–299, <https://doi.org/10.1002/gepi.21806>, (2014).
38. Wamalwa, N. I. E. *et al.* Screening Napier grass accessions for resistance to Napier grass stunt disease using the loop-mediated isothermal amplification of DNA (LAMP). *Crop Prot.* **98**, 61–69 (2017).
39. Rocha, J. R. do A. S. de C. *et al.* Bioenergetic potential and genetic diversity of elephantgrass via morpho-agronomic and biomass quality traits. *Ind. Crops Prod.* **95**, 485–492 (2017).
40. Azevedo, A. L. S. *et al.* Cross species amplification of *Pennisetum glaucum* microsatellite markers in *Pennisetum purpureum* and genetic diversity of Napier grass accessions. *Crop Sci.* **52**, 1776–1785 (2012).
41. Pattanashetti, S. K. *et al.* Genetic variability in Napier grass (*Pennisetum purpureum*) germplasm conserved at ICRISAT genebank. *Int. Gras. Cong.*, November 20-24, New Delhi, (2015).
42. Kandel, R., Singh, H. P., Singh, B. P., Harris-Shultz, K. R. & Anderson, W. F. Assessment of Genetic Diversity in Napier Grass (*Pennisetum purpureum* Schum.) using Microsatellite, Single-Nucleotide Polymorphism and Insertion-Deletion Markers from Pearl Millet (*Pennisetum glaucum* [L.] R. Br.). *Plant Mol. Biol. Report.* **34**, 265–272 (2016).

43. Kilian, A. *et al.* Diversity arrays technology: a generic genome profiling technology on open platforms. *Methods Mol Biol.* **888**, 67–89, https://doi.org/10.1007/978-1-61779-870-2_5 (2012).
44. Muktar, M. S. *et al.* Insights Into the Genetic Architecture of Complex Traits in Napier Grass (*Cenchrus purpureus*) and QTL Regions Governing Forage Biomass Yield, Water Use Efficiency and Feed Quality Traits. *Front. Plant Sci.* **12**, (2022).
45. Wanjala, B. W. *et al.* Genetic diversity in Napier grass (*Pennisetum purpureum*) cultivars: Implications for breeding and conservation. *AoB Plants* **5**, (2013).
46. Holsinger, K. E. Reproductive systems and evolution in vascular plants. *PNAS*, **97** www.pnas.org (2000).
47. Negawo, A. T. *et al.* Genetic diversity and population structure of a rhodes grass (*Chloris gayana*) collection. *Genes (Basel)*. **12**, (2021).
48. Huff, D.R. RAPD characterization of heterogeneous perennial ryegrass cultivars. *Crop Science* **37**, 557–594 (1997).
49. Xie, X. M., Zhou, F., Zhang, X. Q., Zhang, J. M. Genetic variability and relationships between MT-1 elephant grass and closely related cultivars assessed by SRAP markers. *Journal of Genetics* **88**, 281–290 (2009).
50. Sinche, M. *et al.* Development and characterization of a Napier grass (*Cenchrus purpureus* Schumach) mapping population for flowering-time- and biomass-related traits reveal individuals with exceptional potential and hybrid vigor. *GCB Bioenergy* **13**, 1561–1575 (2021).
51. Menezes, B. R. S. *et al.* Estimates of heterosis parameters in elephant grass (*Pennisetum purpureum* schumach.) for bioenergy production. *Chil. J. Agric. Res.* **75**, 395–401 (2015).
52. Mwendia, S. W., Mwangi, D. M., Wahome, R. G., Wanyoike, M. Assessment of growth rate and yields of three Napier grass varieties in central highlands of Kenya. *E.Afr.agric.For.J.* **74**, 211-217 (2008).
53. Criscioni, P. *et al.* Replacement of alfalfa hay (*Medicago sativa*) with maralfalfa hay (*Pennisetum sp.*) in diets of lactating dairy goats. *Anim. Feed Sci. Technol.* **219**, 1–12 (2016).
54. de Souza, F. R. *et al.* Estimation of outcrossing rate in Napier grass. *Crop Sci.* **59**, 1030–1036 (2019).
55. Cheng, Y. K. Forage breeding in Taiwan. *Asian-Australasian Journal of Animal Sciences* **4**, 203–209 (1991).
56. Gruber, B., Unmack, P. J., Berry, O. F., Georges, A. dartr: An r package to facilitate analysis of SNP data generated from reduced representation genome sequencing. *Mol. Ecol. Resour.* **18**, 691–699. doi:10.1111/1755-0998.12745 (2018).
57. Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., Weir, B.S. A High-performance Computing Toolset for Relatedness and Principal Component Analysis of SNP Data. *Bioinformatics*, doi: 10.1093/bioinformatics/bts610 (2012).
58. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics*, **155**, 945 – 959 (2000).

59. Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of individuals using the software Structure: A simulation study. *Mol. Ecol.* **14**, 2611–2620 (2005).
60. Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281 (2014).
61. Jombart, T. Adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**, 1403–1405. doi: 10.1093/bioinformatics/btn129 (2008).
62. Excoffier, L., Smouse, P. E., & Quattro, J. M. Analysis of molecular variance inferred from metric distances among DNA haplotypes - application to human mitochondrial-DNA restriction data. *Genetics* **131**, 479-491 (1992).
63. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of phylogenetics and evolution in R language. *Bioinformatics* **20**, 289–290 (2004).
64. Sethuraman, A. On inferring and interpreting genetic population structure-applications to conservation, and the estimation of pairwise genetic relatedness. *Graduate Theses and Dissertations* (2013).
65. Buiteveld, J. *et al.* Advanced genebank management of genetic resources of European wild apple, *Malus sylvestris*, using genome-wide SNP array data. *Tree Genet. Genomes* **17**, (2021).

Figures







Figure 1

Genome-wide distribution and polymorphism of the markers. The distribution of the markers across the fourteen chromosomes of the Napier grass genome is shown in A (SNPs) and B (SilicoDARTs). The distribution of polymorphic information content (PIC) (orange) and heterozygosity (He) (blue) values are shown for the SNPs (C) and SilicoDARTs (D).




Figure 2

Polymorphism of SNP markers filtered for genetic diversity analysis. The distribution of polymorphic information content (PIC) and heterozygosity (He) of the markers are shown by orange and blue colors, respectively (A). Their distribution across the fourteen Napier grass chromosomes is shown in B.




Figure 3

Major clusters and sub-clusters of the Napier grass collections and progeny plants used in the study. **(A)** The delta K suggesting three major clusters that are partitioned into sub-clusters. **(B)** Bar plots based on the admixture model in STRUCTURE, for K = 3 and 11. **(C)** Clusters detected by hierarchical clustering. **(D)** Clusters detected by PCA.





Figure 4

UPGMA dendrogram (**A**) depicting the genetic relationships between the five Napier grass collections and progeny plants. In (**B**) the minimum, maximum, and average Nei's genetic distance within each of the collections and progeny plants.




Figure 5

Distinct Napier grass genotypes and potential duplicates detected in each of the five collections and progeny plants.






Figure 6

A gap in Mendelian error counts calculated on 133 progeny plants for which the recorded seed parents were confirmed by IBD analysis. **(A)** for PC1 (parent-child relationship for potential seed parents), **(B)** for PC2 (parent-child relationship for potential pollen parents), **(C)** for PPC (parent-parent-child relationship for possible crosses or as potential complete parents).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTables.xlsx](#)